


## Alternating right bundle branch block or intermittent preexcitation?

### ¿Bloqueo de rama derecha alternante o preexcitación intermitente?

Angel Cueva-Parra<sup>1\*</sup>, Michael Soto-Gamarra<sup>2</sup>, Pedro Iturralde-Torres<sup>1</sup>, and Santiago Nava<sup>1</sup>

<sup>1</sup>Department of Electrophysiology, Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico; <sup>2</sup>Department of Cardiology, Hospital Regional Docente de Cajamarca, Cajamarca, Peru

Right bundle branch block (RBBB) is a very common electrocardiographic feature in patients with Ebstein anomaly (EA) and it is present in more than 94% of these patients<sup>1</sup>. We present the case of a 33-year-old woman who was admitted to the emergency room with palpitations, with BP 95/55 mmHg and HR 155 bpm, her heart rhythm was irregular and she had a systolic murmur in the tricuspid focus. In the electrocardiogram (ECG), there was an irregular wide QRS tachycardia, which was negative in leads V1 and DIII, compatible with preexcited AF by a right postero-septal accessory pathway (AP) (Fig. 1). Electrical cardioversion was performed going into sinus rhythm with intermittent RBBB (Fig. 2). The echocardiogram revealed findings consistent with EA with a tricuspid septal leaflet attachment index of 10 mm/m<sup>2</sup> (Fig. 3). The patient was discharged with propafenone 150 mg QID pending ablation. In 2006, Iturralde et al. reported that the absence of RBBB in patients with EA is strongly associated with the presence of an AP<sup>2</sup>, this is explained because anterograde conduction through a right AP during sinus rhythm masked RBBB. In this case, the preexcitation appeared only in some QRS complexes, causing the RBBB to be masked when it was present; therefore, it is not an alternating RBBB itself, but rather an intermittent preexcitation. In these patients, the pre-excitation pattern is



**Figure 1.** Preexcited AF the negative QRS polarity in leads V1 and III, and the positive QRS polarity in V2 suggests that the anterograde conduction occurs through a right posteroseptal AP.

not always evident, even in some leads the PR interval may be normal (Fig. 2), so the most notable electrocardiographic characteristic will always be the absence of RBBB.

#### \*Correspondence:

Angel Cueva-Parra  
E-mail: angel.cueva.parra@gmail.com  
1405-9940 / © 2022 Instituto Nacional de Cardiología Ignacio Chávez. Published by Permalyer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 29-08-2021


Date of acceptance: 18-04-2022

DOI: 10.24875/ACM.21000279


Available online: 12-01-2023

Arch Cardiol Mex. 2023;93(4):493-495

[www.archivoscardiologia.com](http://www.archivoscardiologia.com)



**Figure 2. A:** sinus rhythm with intermittent preexcitation, note that when there is no preexcitation, the RBBB is observed (stars) and when there is antegrade conduction through the AP (accessory pathway), the RBBB is masked (arrows). **B:** comparison between normal and pre-excited complexes, note that despite the existence of preexcitation the PR interval is normal in some leads.



**Figure 3. A:** apical 4 chamber view which shows tricuspid septal leaflet attachment. **B:** apical 4 chamber view with Doppler color which shows severe tricuspid regurgitation. RV: right ventricle; aRV: atrialized right ventricle; RA: right atrium; LV: left ventricle; LA: left atrium.

## **Funding**

None.

## **Conflicts of interest**

None.

## **Ethical disclosures**

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this study.

**Confidentiality of data.** The authors declare that they have followed the protocols of their work center on the publication of patient data.

**Right to privacy and informed consent.** The authors have obtained the informed consent of the patients and/or subjects referred to in the article.

## **References**

1. Walsh EP. Ebstein's anomaly of the tricuspid valve: a natural laboratory for re-entrant tachycardias. *JACC Clin Electrophysiol.* 2018;4:1271-88.
2. Iturralde P, Nava S, Sálica G, Medeiros A, Márquez MF, Colín L, et al. Electrocardiographic characteristics of patients with Ebstein's anomaly before and after ablation of an accessory atrioventricular pathway. *J Cardiovasc Electrophysiol.* 2006;17:1332-6.