

## Indoor PM<sub>10</sub> and its heavy metal composition at a roadside residential environment, Phitsanulok, Thailand

THUNWADEE SRITHAWIRAT

*Department of Environmental Science, Faculty of Science and Technology, Pibulsongkram Rajabhat University, Phitsanulok 65000, Thailand*

Corresponding author: email: T.srithawirat@psru.ac.th

MOHD TALIB LATIF

*School of Environment and Natural Resource Sciences, Faculty of Science and Technology, and Institute for Environment and Development (LESTARI), University Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

FAZRUL RAZMAN SULAIMAN

*Faculty of Applied Sciences, Universiti Teknologi MARA Cawangan Pahang, 26400 Bandar Tun Abdul Razak Jengka, Pahang, Malaysia*

Received: December 30, 2015; accepted: September 7, 2016

### RESUMEN

Se midieron concentraciones de PM<sub>10</sub> tanto en interiores como exteriores en 10 edificios residenciales de Phitsanulok, Tailandia, durante las temporadas de seca y lluvias de 2014. Además, se analizaron siete metales traza en el PM<sub>10</sub>: Zn, Fe, Pb, Cd, Ni, Cu y Cr. Las concentraciones mensuales intra y extramuros de PM<sub>10</sub> fueron de 41.5 a 105.3 µg m<sup>-3</sup> y de 95.2 a 145.1 µg m<sup>-3</sup>, respectivamente. Las concentraciones de PM<sub>10</sub> fueron significativamente mayores durante la temporada seca, en comparación con la temporada húmeda. Las razones interior/exterior fueron menores a uno, lo cual indica que el material particulado se origina en ambientes exteriores. En general, las concentraciones medias de metales pesados en el PM<sub>10</sub> variaron de 0.2 a 2.7 y de 0.5 a 7.1 µg m<sup>-3</sup> para el ambiente interior y el exterior, respectivamente. En el PM<sub>10</sub> de interiores se encontró una fuerte correlación positiva entre Zn y Cu, Zn y Ni, y Cu y Ni. Asimismo, se encontró una fuerte correlación entre Zn y Ni, Pb y Cu, Cu y Ni, Cd y Ni, y Zn y Cu en exteriores. Los factores de enriquecimiento de Zn, Cu, Pb, Cr y Ni fueron menores a uno, lo cual sugiere que los metales en el PM<sub>10</sub> de interiores se originaron en materiales de la corteza. En cuanto a la valoración de riesgos sanitarios, se determinó mediante una evaluación de riesgos con un sistema integrado de información, que el Cr implica el mayor riesgo de cáncer.

### ABSTRACT

The concentrations of PM<sub>10</sub> were measured both indoors and outdoors at 10 roadside residential buildings in Phitsanulok, Thailand during the dry and wet seasons of 2014. Seven trace metals (Zn, Fe, Pb, Cd, Ni, Cu and Cr) were also analysed in PM<sub>10</sub>. The monthly average concentrations of indoor and outdoor PM<sub>10</sub> were 41.5 to 105.3 µg m<sup>-3</sup> and 95.2 to 145.1 µg m<sup>-3</sup>, respectively. PM<sub>10</sub> concentrations were significantly higher during the dry season compared to the wet season. The indoor/outdoor (I/O) ratios were less than one indicating that the particulate matter originates from the outdoor environment. Overall, the average concentrations of heavy metals in PM<sub>10</sub> ranged from 0.2 to 2.7 µg m<sup>-3</sup> and 0.5 to 7.1 µg m<sup>-3</sup> for the indoor and outdoor environments, respectively. A strong positive correlation in indoor PM<sub>10</sub> was found between Zn and Cu, Zn and Ni, and Cu and Ni. Zn and Ni, Pb and Cu, Cu and Ni, Cd and Ni, and Zn and Cu showed strong positive correlations in the outdoor environment. The enrichment factors of Zn, Cu, Pb, Cr and Ni were less than one suggesting that the metals in indoor PM<sub>10</sub> have originated from crustal materials. For the health risk assessment, Cr was found to have the highest excess cancer risk in an evaluation using an Integrated Risk Information System.

**Keywords:** Indoor air quality, heavy metal, enrichment factor, excess cancer risk, I/O ratio.

## 1. Introduction

The distribution of urbanization in Thailand is imbalanced. The rapid growth of urban areas has created high population densities in large cities. The number of new vehicles registered in Thailand has increased around three times from 2000 to 2011 (DLT, 2014). Traffic emissions are recognized as the major source of air pollution in cities (Jinsart *et al.*, 2002; Zhang and Batterman, 2013). Previous studies have shown that the populations of several cities of Thailand are exposed to high levels of particulate matter (PM) such as PM<sub>10</sub> and PM<sub>2.5</sub> (PM with an aerodynamic diameter less than 10 and 2.5  $\mu\text{m}$ , respectively). These levels often exceed the National Ambient Air Quality Standards (NAAQS) of Thailand (Guo *et al.*, 2014). Several other studies have demonstrated that PM in Thailand is associated with respiratory problems and a wide range of premature mortality causes (Vichit-Vadakan and Vajanapoom, 2011; Qiu *et al.*, 2015). Fine particles are more dangerous than coarse particles. Apart from the size of the particles, other specific physical, chemical, and biological characteristics that can produce harmful health effects include the presence of metals, polycyclic aromatic hydrocarbons (PAHs), and other organic components (Vichit-Vadakan and Vajanapoom, 2011).

Exposure to PM occurs mainly indoors because people spend most of their time inside buildings, especially in their homes (Massey *et al.*, 2013; Du *et al.*, 2011; Yang Razali *et al.*, 2015). Effects of inhaled aerosols depend on the specific chemical species, the concentration, the duration of exposure, and the site of deposition within the respiratory tract (Salma *et al.*, 2002). Particle size is the most important factor determining the location of the deposited particles as well as the mechanism of deposition (Salma *et al.*, 2002; Braniš *et al.*, 2005). In general, airborne fine particles have been considered of greater health significance than other major air pollutants. The heavy metals associated with fine particles have been shown to increase lung cancer and cardiopulmonary injuries in humans (Park *et al.*, 2008). Fine particles carry a higher burden of toxic metals than coarse particles (Shridhar *et al.*, 2010; Massey *et al.*, 2013).

Many studies indicate that indoor air quality is affected by outdoor air intrusion into the indoor

environment (Jones *et al.*, 2000; Kuo and Shen, 2010; Leung, 2015; Tunsaringkarn *et al.*, 2015). Outdoor air pollutants can enter buildings through open doors, open windows, ventilation systems, and cracks in structures. Ventilation has a strong influence on indoor PM concentration, but this may also vary due to the difference in the physical nature of pollutants and their source characteristics (Leung, 2015). Several studies have shown a relationship between indoor and ambient air (Jamriska *et al.*, 1999; Jones *et al.*, 2000; Cyrus *et al.*, 2004). In urban areas, high density residential zones and high traffic volume influence the concentration of air pollutants in indoor environment (Agrawal *et al.*, 2003; Massey *et al.*, 2012).

The measurement of indoor PM and its relation to toxicity are essential in order to determine the exposure risk of inhabitants. Thus, the aim of this study was to measure indoor and outdoor PM<sub>10</sub> concentrations and their chemical composition in roadside residential buildings of Phitsanulok, a rapidly growing province in lower northern Thailand. This is a pioneering investigation on the probable sources of trace metals in this area and the health risks associated with them. The data is expected to provide valuable information for future mitigation and sustainable urban planning.

## 2. Methodology

### 2.1 Sampling site

The Phitsanulok province (16° 50' N, 100° 15' E) (Fig. 1) covers an area of about 10816 km<sup>2</sup> and it encompasses a large city located in the lower north territory of Thailand, with a population of 854 000. Phitsanulok was chosen for this study because it is a rapidly developing area due to the east-west economic corridor linking Thailand with Myanmar, Laos and Vietnam (ADB, 2010). The climate of Phitsanulok is under the influence of monsoon winds of seasonal character. The south-west monsoon (May–October) brings a stream of warm moist air from the Indian Ocean causing abundant rain over the country with temperatures ranging from 23 to 39 °C and relative humidity from 35 to 100%. The northeast monsoon (October–February) brings cold and dry air from anticyclonic conditions on the Chinese mainland with temperatures ranging from 14 to 38 °C and relative humidity from 27 to 98% (TMD, 2014).




Fig. 1. Sampling location.

Five residential buildings (sites 1-5 in Fig. 1) near the Asian Highway 16 (AH16) and another five (site 6-10 in Fig. 1) near the Asian Highway 13 (AH13) in the municipality of Phitsanulok were chosen as sampling sites. The sampling was conducted from January to June 2014. The houses varied in age from five to 30 years to provide a variable sample of typical indoor and outdoor concentrations. The houses are made of bricks and cement.

## 2.2 Sample collection

Portable air samplers (MiniVol™ TAS 5.0, Airmetrics, USA) were used for monitoring the indoor and outdoor mass concentration of PM<sub>10</sub> at a constant flow rate of  $5 \text{ L min}^{-1} \pm 5\%$  with a controller for continuous measurement during the 8-h simultaneous sampling

period (8.00-16.00 LT). The flow rate calibration measurements were performed using a mass flow calibrator during equipment placement and removal activities. The particles were collected on a 47-mm quartz filter paper for chemical analysis. Before and after sampling, the filters were stored in desiccators for 24 h prior to an initial weighing in a clean room under controlled temperature and relative humidity conditions. The indoor sampling instrument was placed at a height of 1.5 m and at locations in the house where the occupants spend most of their time. Inlet heads were positioned as close as possible to head height, in the breathing zone. Outdoor sampling was undertaken 1.5 m away from the boundary of the houses. The device was placed at a height of 1.5 m from the ground. A total of 120 samples were

collected from ten houses per month, 60 from the indoor environment and 60 from the outdoor environment of each house on different dates over a six-month period (January to June, 2014). The average mass concentration of PM<sub>10</sub> was calculated by the gravimetric method. After collecting, the filters were refrigerated at about 4 °C prior to chemical analysis.

### 2.3 Analytical procedure

The samples were digested in a mixture of pure concentrated acids (HNO<sub>3</sub> and HClO<sub>4</sub> in a 4:1 ratio) and diluted with deionized water up to 25 ml, then filtered through a 0.45 µm Millipore filter paper to obtain a clear solution in pre-cleaned volumetric flasks and stored at 4 °C prior to analysis. The chemical compounds of the PM samples were determined via inductively coupled plasma optical emission spectrometry (ICP-OES) (JY 238, Horiba, Japan). All PM samples were analyzed for the following seven elements: Zn, Fe, Pb, Cd, Ni, Cu and Cr.

The limit of detection (LOD) for each of the metals was determined by measuring the signal to noise ratio. The LOD for Zn, Pb, and Fe was 0.10 ng m<sup>-3</sup>; for Cd and Cr, 0.03 ng m<sup>-3</sup>; for Ni, 0.13 ng m<sup>-3</sup>, and for Cu, 0.07 ng m<sup>-3</sup>. Percentage recoveries of the metals using acid digestion ranged between 98 and 117%. The blank filter sampler was monitored using unexposed filter papers which were processed with field samples. Background contamination of heavy metals was determined by subtracting the field blank values from the concentrations. In this study, the background contamination was used to correct measurements.

### 2.4 Risk assessment from carcinogenic metals

Excess cancer risks (ECRs) were calculated using the unit risk and the PM-bound concentration of the metals, which can be used to represent their total concentration (Park *et al.*, 2008). The information of carcinogenic types and unit risks of the metals is available from the US-EPA database for IRIS (Integrated Risk Information System). We assumed that a breathing rate for 8-h occupational exposure = 10 m<sup>3</sup>, and a breathing rate for 24-h continuous exposure = 20 m<sup>3</sup> (occupational exposure = 5 days/week; continuous exposure = 7 days/week) (US-EPA, 1998). The typical method for calculating ECRs is by using the following formula (US-EPA, 1998):

$$\text{Excess cancer risk (inhalation)} = \frac{\text{atmospheric concentration of pollutant } (\mu\text{g m}^{-3}) \times \text{unit risk } (\mu\text{g m}^{-3})^{-1}}{(1)}$$

### 2.5 Statistical analyses

Analysis of the experimental data was performed using the Statistical Package for Social Sciences (SPSS v. 11.5, IBM, USA). Descriptive statistics, including the mean concentrations of PM<sub>10</sub> and standard deviations, were used to investigate the elemental concentrations of the PM<sub>10</sub> samples. Pearson correlations were conducted to evaluate the relationship of the indoor and outdoor PM<sub>10</sub> and its metal concentrations.

## 3. Results and discussion

### 3.1 PM mass concentrations

The average monthly PM<sub>10</sub> concentration and standard deviations obtained in the residential buildings are presented in Table I. The PM<sub>10</sub> mass in indoor and

Table I. Average monthly concentration of indoor and outdoor PM<sub>10</sub> at roadside buildings (in µg m<sup>-3</sup>).

Month	N	AH16			AH13		
		Indoor (I)	Outdoor (I)	I/O ratio	Indoor (I)	Outdoor (O)	I/O ratio
January	10	91.1 ± 9.4	145.1 ± 7.9	0.6	60.6 ± 6.1	103.6 ± 9.1	0.6
February	10	95.4 ± 9.8	125.6 ± 8.5	0.8	41.5 ± 5.3	105.6 ± 5.4	0.4
March	10	80.6 ± 6.1	123.6 ± 9.1	0.7	45.4 ± 6.8	98.6 ± 2.4	0.5
April	10	85.3 ± 5.2	115.6 ± 9.4	0.7	55.2 ± 9.4	95.2 ± 8.5	0.6
May	10	71.9 ± 6.5	95.6 ± 9.4	0.8	61.5 ± 5.3	100.6 ± 10.1	0.6
June	10	105.3 ± 10.2	135.4 ± 8.8	0.8	65.5 ± 8.5	101.6 ± 9.4	0.6
Average		88.3 ± 11.7	123.5 ± 17.1	0.7	54.9 ± 9.6	100.9 ± 3.7	0.6
Minimum		71.9	95.6	0.6	41.5	95.2	0.4
Maximum		105.3	145.1	0.8	65.5	105.6	0.6

N: number of samples.

outdoor samples ranged from 71.9 to 105.3  $\mu\text{g m}^{-3}$  and 95.6 to 145.1  $\mu\text{g m}^{-3}$  for AH16, and from 41.5 to 65.5  $\mu\text{g m}^{-3}$  and 95.2 to 105.6  $\mu\text{g m}^{-3}$  for AH13, respectively. The trends of PM<sub>10</sub> concentrations were higher in outdoor samples compared to indoor samples. Overall, the houses at AH16, a six-lane highway, showed higher average monthly PM<sub>10</sub> concentrations compared to those at AH13 (a four-lane highway), except for the outdoor PM<sub>10</sub> concentration in May. There are significant differences in PM<sub>10</sub> concentrations recorded at the different sampling sites ( $p < 0.05$ ). However, statistical analysis performed with the  $t$ -test shown no significant difference ( $p > 0.05$ ) between the mean concentrations of indoor and outdoor PM<sub>10</sub> in the five houses located on each road site, indicating that similar sources lead to the generation of particulate pollutants in these environments (Massey *et al.*, 2013).

The results indicate that PM<sub>10</sub> in indoor and outdoor environments is mainly affected by particle emissions from the roadside area. Traffic volume can increase in the wider highway (AH16), so it is expected to contribute to higher concentrations of PM<sub>10</sub> inside and outside roadside buildings due to high traffic density and congestion in this area. The higher PM<sub>10</sub> concentrations at the sampling sites may be emitted by vehicles, depending on the locations and prevailing winds. Generally, the contribution from cross-border sources is less significant in urban areas due to the increased distance from the pollution sources (Cheng *et al.*, 2009). However, urban air quality is highly affected by city design. Densely distributed buildings, deep street canyons, and buildings with high ratios of height to road width, can block and weaken the approaching wind, thus reducing air dispersion (Cheng *et al.*, 2009; Li *et al.*, 2009).

As there is no indoor air quality standard for public buildings in Thailand, we compared our results with the indoor air quality standards of other Asian countries. The concentrations of indoor PM<sub>10</sub> presented in this study were below the guideline for Singapore (150  $\mu\text{g m}^{-3}$  for 8-h averaging time) (NEA, 1996). The PM<sub>10</sub> concentrations recorded in this study are classified as good according to the Indoor Air Quality Objectives for Offices and Public Places in Hong Kong (180  $\mu\text{g m}^{-3}$  for 8-h averaging time) but exceeded the excellence class value at 20  $\mu\text{g m}^{-3}$  (EPD, 2003).

### 3.2 Seasonal variation of PM<sub>10</sub>

In this study, the seasonal variation was divided into two seasons: dry (January–March) and wet (April–June) based on the regional meteorological conditions from the Thai Meteorological Department (TMD, 2014). Fig. 2 shows the average seasonal concentrations of PM<sub>10</sub> in indoor and outdoor environments of the AH16 and AH13 sites. The mean variations of PM<sub>10</sub> at the AH16 and AH13 sites show similar seasonal trends. In the dry season, the mean concentrations of PM<sub>10</sub> in indoor and outdoor environments at AH16 were 89.0 and 131.4  $\mu\text{g m}^{-3}$ , respectively. For AH13, the average PM<sub>10</sub> concentrations were 49.2 and 102.6  $\mu\text{g m}^{-3}$ . In the wet season, the mean concentrations of PM<sub>10</sub> in indoor and outdoor environments at AH16 were 87.5  $\mu\text{g m}^{-3}$  and 115.5  $\mu\text{g m}^{-3}$ , respectively. For AH13, the average PM<sub>10</sub> concentrations were 60.7  $\mu\text{g m}^{-3}$  and 99.1  $\mu\text{g m}^{-3}$ , respectively. The overall seasonal trends showed significantly higher PM<sub>10</sub> concentrations in the dry season in comparison




Fig. 2. Seasonal variation of PM<sub>10</sub> in indoor and outdoor environments. The first quartile (Q1 or 25th percentile) is the median of the lower half of the data set. The third quartile (Q3 or 75th percentile) is the median of the upper half of the data set.



to the wet season ( $p < 0.05$ ), except for indoor  $\text{PM}_{10}$  at AH13. Lower  $\text{PM}_{10}$  concentrations were obtained in the wet season due to the washout of particles in the atmosphere (Saha and Krishna Moorthy, 2004). The higher  $\text{PM}_{10}$  concentrations were recorded at the AH16 site, indicating outdoor particulate levels have an impact on the indoor environment via infiltration. Similar results were reported by Chao and Wong (2002). The average indoor  $\text{PM}_{10}$  concentrations in homes near a minor road in Hong Kong were higher than those of the homes far away from the traffic, while they were higher for the homes near a main road than for homes near a minor road. High indoor PM concentrations were observed in homes close to heavy traffic. This indicates that high outdoor PM concentrations originating from vehicle and industrial exhaust, as well as wind-blown dust pose a great impact on indoor PM concentrations (Chao and Wong, 2002).

Comparing this study with the results from other Asian countries, it was found that the average indoor and outdoor  $\text{PM}_{10}$  concentrations of households in rural areas of India were  $242.5$  and  $217.76 \mu\text{g m}^{-3}$  (Massey *et al.*, 2013). Massey *et al.* (2012) also found that annual average concentrations of  $\text{PM}_{10}$  in indoor and outdoor environments in Agra, India were  $247$  and  $255 \mu\text{g m}^{-3}$  in roadside houses, and  $181$  and  $195 \mu\text{g m}^{-3}$  in urban houses. A study in homes of Delhi, India reported that the average indoor and outdoor  $\text{PM}_{10}$  concentrations ranged from  $104$  to  $506 \mu\text{g m}^{-3}$  and from  $100$  to  $598 \mu\text{g m}^{-3}$ , respectively (Kulshreshtha and Khare, 2011). The results from India are about 2-3 times higher than those from this study. Chao and Wong (2002) reported that the mean indoor and outdoor  $\text{PM}_{10}$  concentrations in houses in Hong Kong were found to be  $63.3$  and  $69.5 \mu\text{g m}^{-3}$ , respectively, which are about 1-2 times lower than in this study.

### 3.3 Indoor/outdoor ratios (I/O)

The indoor/outdoor ratio (I/O) has been widely used for evaluating the difference between indoor and outdoor concentrations as an indicator of indoor sources strength (Kulshreshtha and Khare, 2011). Indoor air pollutant concentrations are affected by the infiltration of outdoor pollutants into the homes and by pollutants from indoor sources. Therefore, the I/O ratios of  $\text{PM}_{10}$  were calculated to quantify the impact of outdoor air and indoor sources on indoor air quality. The I/O ratio is defined as the ratio  $C_{\text{in}}/C_{\text{out}}$  where  $C_{\text{in}}$  and  $C_{\text{out}}$  are the indoor and outdoor concentrations

of PM, respectively (Chen and Zhao, 2011). The I/O ratio can vary largely due to factors such as location, building design and different activities of the occupants (Massey *et al.*, 2012).

The I/O ratios of  $\text{PM}_{10}$  were presented in Table I. They ranged from 0.6 to 0.8 and from 0.4 to 0.6 for AH16 and AH13, respectively. The I/O ratios of all the sampling sites were less than one, indicating that the PM originated from vehicle induced emissions from the nearby congested roads, which directly enter the indoor environment of homes increasing their concentration levels (Oosterlee *et al.*, 1996). Pathways of outdoor air pollutants to the indoor environment depend on ventilation and infiltration. The ventilation of a dwelling by fan or air conditioner, or by individual or central air conditioning systems, draw in air from outdoors through fresh air intakes (Leung, 2015). The prevailing wind provides natural ventilation whenever the doors and windows are open. Infiltration can also occur through cracks and leaks in the building, which may be significant for a building with poor sealing (Leung, 2015). Due to these mechanisms, air pollutants from outdoors can penetrate into the indoor environment, and can either be diluted or accumulated according to the ventilation condition (Leung, 2015).

### 3.4 Trace elemental concentrations and enrichment

The chemical elemental concentrations associated with indoor and outdoor  $\text{PM}_{10}$  are given in Table II. The results show the mean concentration of  $\text{PM}_{10}$  and the standard deviation of each metal at individual sites. No trend of monthly changes in metal concentrations was observed for individual metals. The overall trends of indoor and outdoor metal concentrations at AH16 were  $\text{Fe} > \text{Pb} > \text{Zn} > \text{Cu} > \text{Cr} > \text{Ni} > \text{Cd}$  and  $\text{Fe} > \text{Pb} > \text{Cr} > \text{Zn} > \text{Ni} > \text{Cu} > \text{Cd}$ , respectively. For AH13, the concentration trends of indoor and outdoor metal concentrations were  $\text{Zn} > \text{Cr} > \text{Cu} > \text{Fe} > \text{Pb} > \text{Ni} > \text{Cd}$  and  $\text{Pb} > \text{Fe} > \text{Zn} > \text{Cr} = \text{Ni} > \text{Cu} > \text{Cd}$ , respectively. Compared with other metals at AH16, Fe concentrations were the highest in both indoor and outdoor environments. In the present study, Pb concentrations were found to be higher outdoors, compared to the indoor environment. Higher Pb concentrations at the roadside on these sites may be due to the mixing of residual Pb in soil and its resuspension in the air due to motor vehicles and community activities such as outdoor painting, automobile repair shops, battery manufacturers and recycle shops (Srithawirat and

Table II. Statistical analysis of trace metal concentrations in PM<sub>10</sub> at roadside buildings (in  $\mu\text{gm}^{-3}$ ).

Heavy metal	N	AH16				AH13			
		Indoor (I)		Outdoor (I)		Indoor (I)		Outdoor (I)	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD
Fe	12	2.7	0.8	5.8	1.9	0.8	0.4	6.3	4.0
Zn	12	1.3	0.6	0.9	0.9	1.6	1.3	2.1	1.6
Pb	12	1.7	0.7	3.7	1.9	0.6	0.1	7.1	1.7
Cu	12	1.1	0.2	0.6	0.3	1.0	0.5	1.3	0.9
Cr	12	0.9	0.7	1.0	0.9	1.5	0.9	2.0	1.1
Cd	12	0.2	0.1	0.5	0.2	0.2	0.1	0.7	0.1
Ni	12	0.3	0.2	0.7	0.6	0.3	0.2	2.0	0.5

N: number of samples.

Latif, 2015). Soils in older areas of the cities are highly contaminated by lead, largely due to past use of lead additives in gasoline, the use of lead in exterior paints, and industrial lead sources. Laidlaw *et al.* (2012) sustain that automotive traffic turbulence plays a significant role in the resuspension of contaminated roadside soils and dusts, and resuspended soil appears to be a significant underlying source of atmospheric Pb (Zahran *et al.*, 2013).

Mineral and stone materials used in road building may contain Cd, Cu, Fe, Ni, Pb and Zn, which are released to the environment as the road surface wears (Lindgren, 1996). After the prohibition of leaded gasoline in Thailand, lead-based paint could be considered a major exposure source of Pb in houses, as it is very commonly used in existing residences, such as on walls, toys and furniture (Lin *et al.*, 2009). Fergusson and Marsh (1993) suggested that worn construction, furnishing and carpet materials can lead to elevated heavy metal content.

Sources of metals in aerosols are reported differently by different researchers. Lindgren (1996) reported that engine wear releases Cr, Cu, Mn, and Ni. Cr is derived from yellow paints in roads and alloys in automobiles (Madany *et al.*, 1994); it is also emitted by municipal waste burning outdoors (Massey *et al.*, 2013). Zn in outdoor urban environments may be released from galvanized steel road equipment, such as crash barriers, road signs and lamp posts, as a corrosion product (Yuen *et al.*, 2012). Rubber underlays, carpets, and galvanized iron roofing are considered as important sources of Zn in indoor environments (Kim and Fergusson, 1993).

The correlations between chemical elements found in indoor and outdoor environment are presented in Table III. The statistical analysis of data from indoor environments indicates that there are positive and negative correlations at the 99% confidence level. A strong positive correlation was found between Zn and Cu, Zn and Ni, and Cu and Ni in indoor environments ( $r = 0.858, 0.888$  and  $0.916$ ). For outdoor environments, a strong correlation was observed between Zn and Cu, Zn and Ni, Pb and Cu, Cu and Ni, and Cd and Ni ( $r = 0.852, 0.783, 0.788, 0.831$  and  $0.961$ , respectively). The results indicate that these metals may originate from similar sources, such as automobile emissions, street dust and other related industrial activities.

Enrichment factors (EFs) were calculated to determine whether the PM originated from crustal or non-crustal sources. They were estimated for individual elements over the average elemental composition of the upper continental crust (Hernandez *et al.*, 2003). Typically, the EF method normalizes the measured heavy metal content with respect to a sample reference metal such as Fe or Al (Ravichandran *et al.*, 1995). The EFs of an atmospheric element were calculated using the following equation:

$$EF_x = (C_x/C_{ref})_{atmosphere} / (C_x/C_{ref})_{crust} \quad (2)$$

where  $x$  is the concentration of an element measured in the atmosphere and  $C_{ref}$  is the concentration of the reference element. Fe was used in this study as a reference because it is a major constituent of clay minerals and has been successfully used by several researchers. Deely and Fergusson (1994) reported that Fe is an

Table III. Summary on indoor and outdoor of trace metal concentrations in PM<sub>10</sub> correlations among studied variables from ten study locations

Overall (10 stations)	Indoor						
	Fe	Zn	Pb	Cu	Cr	Cd	Ni
Fe	1						
Zn	0.632	1					
Pb	0.856	-0.656	1				
Cu	-0.653	0.858**	-0.920**	1			
Cr	-0.956*	0.138	-0.864	-0.197	1		
Cd	-0.921*	-0.534	0.281	-0.647	0.356	1	
Ni	-0.614	0.888**	-0.630	0.916**	-0.324	-0.546	1
Overall (10 stations)	Outdoor						
	Fe	Zn	Pb	Cu	Cr	Cd	Ni
Fe	1						
Zn	0.657	1					
Pb	0.501	0.659	1				
Cu	0.489	0.852**	0.788**	1			
Cr	0.480	0.497	0.254	0.451	1		
Cd	0.212	0.443	0.998**	0.807	-0.576	1	
Ni	0.514	0.783**	0.967**	0.831**	0.358	0.961**	1

\*correlation is significant at the 0.05 level (2-tailed); \*\*correlation is significant at the 0.01 level (2-tailed).

acceptable normalization element to be used in the calculation of the EF, since its distribution is unrelated to other heavy metals. The average element concentrations in soil were obtained from Taylor and McLennan (1985). EFs close to one point are considered to have a crustal origin, while it is assumed that those greater than 10 have a non-crustal source (Nolting *et al.*, 1999). Further, EFs can also assist in the determination of the degree of metal contamination (Yongming *et al.*, 2006). The contamination categories are recognised on the basis of the EF. EFs of more than five indicate that they are significantly enriched and contaminated (Yongming *et al.*, 2006). For both roadside environments (AH16 and AH13), Zn, Cu, Pb, Cr and Ni in PM<sub>10</sub> showed EFs of less than one which means that the metals analyzed in indoor particles have penetrated from outdoor sources (Kulshrestha *et al.*, 2014). Only the enrichment of Cd was found to be greater than one at all sites, which may be due to indoor dust at these sites, associated with paint and related indoor activities.

### 3.5 Risk assessment from carcinogenic metals

Among the heavy metals in indoor and outdoor PM<sub>10</sub> on the roadside residential areas of this study,

Cr, Cd, Ni and Pb are known carcinogenic agents that can cause serious health risks when people are exposed through the inhalation pathway (US-EPA, 1998). According to the US-EPA classification of carcinogens, Cr (VI) is categorized in group A as a known human carcinogen by exposure through the inhalation route. A previous research has acknowledged that the measured total Cr concentration was assumed to be a mixture of carcinogenic (Cr [VI]) and non-carcinogenic (Cr [III]) agents in a 1:6 concentration ratio in ambient air (US-EPA, 2004). Mancuso (1997), forming the basis for the estimation of cancer potency for Cr compounds, suggests that in an urban area one-seventh of the total chromium concentrations is hexavalent (Cr [VI]). Therefore, in this study the concentration of Cr (VI) was assumed to be one-seventh of the total Cr concentration for the carcinogenic risk assessment (Park *et al.*, 2008; Hieu and Lee, 2010; Massey *et al.*, 2013; Taner *et al.*, 2013). Cd is classified as a group B1 human carcinogen. Ni dust from refineries and nickel subsulfide are also known human carcinogens. The concentrations of PM<sub>10</sub> analyzed in this study may contain nickel subsulfide, nickel oxide and metallic nickel, which



Table IV. Excess cancer risks of PM<sub>10</sub> carcinogenic metals.

Metal	Concentration ( $\mu\text{g m}^{-3}$ )				Inhalation unit risk ( $\mu\text{g m}^{-3}$ ) <sup>-1</sup>	Excess cancer risks (in 1 million)			
	Average		95th percentile			Average		95th percentile	
	Indoor	Outdoor	Indoor	Outdoor		Indoor	Outdoor	Indoor	Outdoor
Cr (VI)	1.2	1.5	1.7	3.9	$1.2 \times 10^{-2}$	14400	18000	20400	46800
Cd	0.2	0.6	0.4	1.3	$1.8 \times 10^{-3}$	360	1080	720	2340
Ni	0.3	1.35	0.7	4.2	$2.4 \times 10^{-4}$	72	324	168	1008

are classified as group A carcinogens. Pb is a probable human carcinogen categorized in group B2, but the ECR for Pb was not calculated due to inadequate evidence on its carcinogenic effect on humans and to its unit risk currently being amended by the US-EPA.

The ECR of PM<sub>10</sub>-bound carcinogenic trace elements in the indoor and outdoor environments for the average values, as well as the 95th percentile values of Cd, Cr and Ni are presented in Table IV. The ECRs calculated for the average values of Cr (VI), Cd and Ni concentrate in indoor and outdoor PM<sub>10</sub> were multiplied by  $10^6$  to compare the risks to a one in a million standard (US-EPA, 1998). Cr showed the highest ECR followed by Cd and Ni in indoor and outdoor environments, due to the much higher unit risk of Cr (VI), ranging from seven to 50 times compared to the unit risk of Cd or Ni. The total ECRs based on the average values of Cd, Cr (VI) and Ni in indoor and outdoor environments were 14 832 and 19 404, respectively. These results indicate that 14 832 and 19 404 people out of one million are at risk of developing cancer from exposure to the carcinogenic metals in the PM<sub>10</sub> fraction of indoor and outdoor particulates in this area. The ECRs trend was found to be higher in outdoor than in indoor environments, indicating that the occupants' exposure to toxic heavy metals in outdoor areas results in an increased cancer risk in comparison to the indoor environment.

#### 4. Conclusions

This study determined the concentrations of PM<sub>10</sub> and their metal composition in indoor and outdoor environments at roadside residential locations in Phitsanulok, Thailand. The results showed that the concentrations of PM<sub>10</sub> ranged from 41.5 to 105.3  $\mu\text{g m}^{-3}$ , and from 95.2 to 145.1  $\mu\text{g m}^{-3}$  for indoor and outdoor environment, respectively. The seasonal variation showed that PM<sub>10</sub> levels were significantly higher

in the dry season in comparison to the wet season ( $p < 0.05$ ). Precipitation washout of PM<sub>10</sub> concentrations in the atmosphere was considered a factor causing lower concentrations during the wet season. The trends of PM<sub>10</sub> concentrations were higher in the outdoor environment in comparison to the indoor environment. The high density traffic and congestion was thought to contribute to the concentrations of PM<sub>10</sub> in roadside buildings. The I/O ratios of all the sampling sites were less than one, indicating that PM originated from vehicle emissions, which enter the indoor environment by ventilation and infiltration. The average concentrations of PM<sub>10</sub> in this study were found to be lower than the values set by the Singapore NEA and the Hong Kong Guidelines for Offices and Public Places. PM-bound heavy metals Zn, Fe, Pb, Cd, Ni, Cu and Cr were detected in PM<sub>10</sub>. Cr was found to have the highest carcinogenic risk among the trace metals in residential areas. This study suggests that there is a need to address the issue of fine particulate monitoring (i.e. particulate matter with diameter size below than 2.5  $\mu\text{m}$  [PM<sub>2.5</sub>]) and its toxic effects should be comprehensively investigated, considering different locations as well as the pattern in resident's activities in order to design control and mitigation strategies.

#### Acknowledgments

The authors would like to acknowledge Pibulsongkram Rajabhat University for financial support with the research grant RDI-1-57-6-13. We also thank to Dr. Rose Norman for proofreading this manuscript.

#### References

- ADB, 2010. Asian Development Bank. Available at: <http://www.adb.org/sites/default/files/publication/27496/gms-action-plan-east-west.pdf> (last accessed in July 2014).
- Agrawal M., B. Singh, M. Rajput, F. Marshall and J. N. B. Bell, 2003. Effect of air pollution on peri-urban

- agriculture: A case study. *Environ. Pollut.* **126**, 323-329, doi:10.1016/S0269-7491(03)00245-8.
- Braniš M., P. Řezáčová and M. Domasová, 2005. The effect of outdoor air and indoor human activity on mass concentrations of PM<sub>10</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> in a classroom. *Environ. Res.* **99**, 143-149, doi:10.1016/j.envres.2004.12.001.
- Chao C. Y. and K. K. Wong, 2002. Residential indoor PM<sub>10</sub> and PM<sub>2.5</sub> in Hong Kong and the elemental composition. *Atmos. Environ.* **36**, 265-277, doi:10.1016/S1352-2310(01)00411-3.
- Chen C. and B. Zhao, 2011. Review of relationship between indoor and outdoor particles: I/O ratio, infiltration factor and penetration factor. *Atmos. Environ.* **45**, 275-288, doi:10.1016/j.atmosenv.2010.09.048.
- Cheng W. C., C.-H. Liu and D. Y. C. Leung, 2009. On the correlation of air and pollutant exchange for street canyons in combined wind-buoyancy-driven flow. *Atmos. Environ.* **43**, 3682-3690, doi:10.1016/j.atmosenv.2009.04.054.
- Cyrs J., M. Pitz, W. Bischof, H. E. Wichmann and J. Heinrich, 2004. Relationship between indoor and outdoor levels of fine particle mass, particle number concentrations and black smoke under different ventilation conditions. *J. Expo. Anal. Environ. Epidemiol.* **14**, 275-283, doi:10.1038/sj.jea.7500317.
- Deely J. M. and J. E. Fergusson, 1994. Heavy metal and organic matter concentrations and distributions in dated sediments of a small estuary adjacent to a small urban area. *Sci. Total Environ.* **153**, 97-111, doi:10.1016/0048-9697(94)90106-6.
- DLT, 2014. Department of Land Transport. Available at: [http://apps.dlt.go.th/statistics\\_web/statistics.html](http://apps.dlt.go.th/statistics_web/statistics.html) (last accessed in July 2014).
- Du L., S. Batterman, E. Parker, C. Godwin, J.-Y. Chin, A. O'Toole, T. Robins, W. Brakefield-Caldwell and T. Lewis, 2011. Particle concentrations and effectiveness of free-standing air filters in bedrooms of children with asthma in Detroit, Michigan. *Build. Environ.* **46**, 2303-2313, doi:10.1016/j.buildenv.2011.05.012.
- EPD, 2003. Environmental Protection Department. Available at: <http://www.iaq.gov.hk/media/8694/cert-guide-eng.pdf> (last accessed in August 2014).
- Fergusson C. and J. Marsh, 1993. Assessing human health risks from ingestion of contaminated soil. *Land. Contam. Reclam.* **1**, 177-185.
- Guo Y., S. Li, B. Tawatsupa, K. Punnasiri, J. J. K. Jaakkola and G. Williams, 2014. The association between air pollution and mortality in Thailand. *Sci. Rep.* **4**, 1-8, doi:10.1038/srep05509.
- Hernandez L., A. Probst, J. L. Probst and E. Ulrich, 2003. Heavy metal distribution in some french forest soils: Evidence for atmospheric contamination. *Sci. Total Environ.* **312**, 195-219, doi:10.1016/S0048-9697(03)00223-7.
- Hieu N. T. and B.-K. Lee, 2010. Characteristics of particulate matter and metals in the ambient air from a residential area in the largest industrial city in Korea. *Atmos. Environ.* **98**, 526-537, doi:10.1016/j.atmosres.2010.08.019.
- Jamriska M., S. Thomas, L. Morawska and B. A. Clark, 1999. Relation between indoor and outdoor exposure to fine particles near a busy arterial road. *Indoor Air* **9**, 75-84, doi:10.1111/j.1600-0668.1999.t01-2-00002.x.
- Jinsart W., K. Tamura, S. Loetkamonwit, S. Thepanondh, K. Karita and E. Yano, 2002. Roadside particulate air pollution in Bangkok. *J. Air Waste Manage. Assoc.* **52**, 1102-1110, doi:10.1080/10473289.2002.10470845.
- Jones N. C., C. A. Thornton, D. Mark and R. M. Harrison, 2000. Indoor/outdoor relationships of particulate matter in domestic homes with roadside, urban and rural locations. *Atmos. Environ.* **34**, 2603-2612, doi:10.1016/S1352-2310(99)00489-6.
- Kim N. and J. Fergusson, 1993. Concentrations and sources of cadmium, copper, lead and zinc in house dust in Christchurch, New Zealand. *Sci. Total Environ.* **138**, 1-21, doi:10.1016/0048-9697(93)90400-Z.
- Kulshreshtha P. and M. Khare, 2011. Indoor exploratory analysis of gaseous pollutants and respirable particulate matter at residential homes of Delhi, India. *Atmos. Pollut. Res.* **2**, 337-350, doi:10.5094/APR.2011.038.
- Kulshreshtha A., D. D. Massey, J. Masih and A. Taneja, 2014. Source characterization of trace elements in indoor environments at urban, rural and roadside sites in a semi arid region of India. *Aerosol Air Qual. Res.* **14**, 1738-1751, doi:10.4209/aaqr.2013.05.0147.
- Kuo H.-W. and H.-Y. Shen, 2010. Indoor and outdoor PM<sub>2.5</sub> and PM<sub>10</sub> concentrations in the air during a dust storm. *Build. Environ.* **45**, 610-614, doi:10.1016/j.buildenv.2009.07.017.
- Laidlaw M. A. S., S. Zahran, H. W. Mielke, M. P. Taylor and G. M. Filippelli, 2012. Re-suspension of lead contaminated urban soil as a dominant source of atmospheric lead in Birmingham, Chicago, Detroit and Pittsburgh, USA. *Atmos. Environ.* **49**, 302-310, doi:10.1016/j.atmosenv.2011.11.030.
- Leung D. Y. C., 2015. Outdoor-indoor air pollution in urban environment: Challenges and opportunity. *Front. Environ. Sci.* **2**, 1-7, doi:10.3389/fenvs.2014.00069.

- Li X.-X., C.-H. Liu and D. Y. C. Leung, 2009. Numerical investigation of pollutant transport characteristics inside deep urban street canyons. *Atmos. Environ.* **43**, 2410-2418, doi:10.1016/j.atmosenv.2009.02.022.
- Lin G. Z., R. F. Peng, Q. Chen, Z. G. Wu and L. Du, 2009. Lead in housing paints: An exposure source still not taken seriously for children lead poisoning in China. *Environ. Res.* **109**, 1-5, doi:10.1016/j.envres.2008.09.003.
- Lindgren Å., 1996. Asphalt wear and pollution transport. *Sci. Total Environ.* **189-190**, 281-286, doi:10.1016/0048-9697(96)05220-5.
- Madany I. M., M. Salim Akhter and O. A. Al Jowder, 1994. The correlations between heavy metals in residential indoor dust and outdoor street dust in Bahrain. *Environ. Int.* **20**, 483-492, doi:10.1016/0160-4120(94)90197-X.
- Mancuso T. F., 1997. Chromium as an industrial carcinogen: Part I. *Am. J. Ind. Med.* **31**, 129-139, doi:10.1002/(SICI)1097-0274(199702)31:2<129::AID-AJIM1>3.0.CO;2-V.
- Massey D., A. Kulshrestha, J. Masih and A. Taneja, 2012. Seasonal trends of PM<sub>10</sub>, PM<sub>5.0</sub>, PM<sub>2.5</sub> and PM<sub>1.0</sub> in indoor and outdoor environments of residential homes located in north-central India. *Build. Environ.* **47**, 223-231, doi:10.1016/j.buildenv.2011.07.018.
- Massey D. D., A. Kulshrestha and A. Taneja, 2013. Particulate matter concentrations and their related metal toxicity in rural residential environment of semi-arid region of India. *Atmos. Environ.* **67**, 278-286, doi:10.1016/j.atmosenv.2012.11.002.
- NEA, 1996. National Environmental Agency, Institute of Environmental Epidemiology, Ministry of the Environment, Singapore. Available at: [https://bca.gov.sg/Greenmark/others/NEA\\_Office\\_IAQ\\_Guidelines.pdf](https://bca.gov.sg/Greenmark/others/NEA_Office_IAQ_Guidelines.pdf) (last accessed in July 2014).
- Nolting R. F., A. Ramkema and J. M. Everaarts, 1999. The geochemistry of Cu, Cd, Zn, Ni and Pb in sediment cores from the continental slope of the banc d'arguin (Mauritania). *Cont. Shelf. Res.* **19**, 665-691, doi:10.1016/S0278-4343(98)00109-5.
- Oosterlee A., M. Drijver, E. Fau-Lebret, E. Lebret, B. Fau-Brunekreef and B. Brunekreef, 1996. Chronic respiratory symptoms in children and adults living along streets with high traffic density. *Occup. Environ. Med.* **53**, 241-247.
- Park E.-J., D.-S. Kim and K. Park, 2008. Monitoring of ambient particles and heavy metals in a residential area of Seoul, Korea. *Environ. Monit. Assess.* **137**, 441-449, doi:10.1007/s10661-007-9779-y.
- Qiu H., L. Tian, K.-F. Ho, V. C. Pun, X. Wang and I. T. S. Yu, 2015. Air pollution and mortality: Effect modification by personal characteristics and specific cause of death in a case-only study. *Environ. Pollut.* **199**, 192-197, doi:10.1016/j.envpol.2015.02.002.
- Ravichandran M., M. Baskaran, P. H. Santschi and T. S. Bianchi, 1995. History of trace metal pollution in sabine-neches estuary, Beaumont, Texas. *Environ. Sci. Technol.* **29**, 1495-1503, doi:10.1021/es00006a010.
- Saha A. and K. Krishna Moorthy, 2004. Impact of precipitation on aerosol spectral optical depth and retrieved size distributions: A case study. *J. Appl. Meteor.* **43**, 902-914, doi:10.1175/1520-0450(2004)043<0902:IOPOAS>2.0.CO;2.
- Salma I., I. Balásházy, R. Winkler-Heil, W. Hofmann and G. Záray, 2002. Effect of particle mass size distribution on the deposition of aerosols in the human respiratory system. *J. Aerosol Sci.* **33**, 119-132, doi:10.1016/S0021-8502(01)00154-9.
- Shridhar V., P. S. Khillare, T. Agarwal and S. Ray, 2010. Metallic species in ambient particulate matter at rural and urban location of Delhi. *J. Hazard. Mater.* **175**, 600-607, doi:10.1016/j.jhazmat.2009.10.047.
- Srithawirat T. and M. T. Latif, 2015. Concentration of selected heavy metals in the surface dust of residential buildings in Phitsanulok, Thailand. *Environ. Earth. Sci.* **74**, 2701-2706, doi:10.1007/s12665-015-4291-0.
- Taner S., B. Pekey and H. Pekey, 2013. Fine particulate matter in the indoor air of barbeque restaurants: Elemental compositions, sources and health risks. *Sci. Total Environ.* **454-455**, 79-87, doi:10.1016/j.scitotenv.2013.03.018.
- Taylor S. and S. McLennan, 1985. *The continental crust: Its composition and evolution*, Blackwell Publisher, Oxford. 312 pp.
- TMD, 2014. Thai Meteorological Department. Available at: <http://www.tmd.go.th> (last accessed in August 2014).
- Tunsaringkarn T., T. Prueksasit, D. Morknong, R. Sawatsing, V. Chinveschakitvanich, A. Rungsiyothin and K. Zapaung, 2015. Indoor air assessment, health risks, and their relationship among elderly residents in urban warrens of Bangkok, Thailand. *Air Qual. Atmos. Health.* **8**, 603-615, doi:10.1007/s11869-014-0302-7.
- US-EPA, 1998. United States Environmental Protection Agency, Integrated Risk Information System, IRIS Substance List. Available at: <http://www.epa.gov/iris/> (last accessed in August 2014).

- US-EPA, 2004. United States Environmental Protection Agency, Region 9, Preliminary Remediation Goals, Air-Water Calculations.
- Vichit-Vadakan N. and N. Vajanapoom, 2011. Health impact from air pollution in Thailand: Current and future challenges. *Environ. Health. Perspect.* **119**, A197-A198, doi:10.1289/ehp.1103728.
- Yang Razali N. Y., M. T. Latif, D. Dominick, N. Mohamad, F. R. Sulaiman and T. Srithawirat, 2015. Concentration of particulate matter, CO and CO<sub>2</sub> in selected schools in Malaysia. *Build. Environ.* **87**, 108-116, doi:10.1016/j.buildenv.2015.01.015.
- Yongming H., D. Peixuan, C. Junji and E. S. Posmentier, 2006. Multivariate analysis of heavy metal contamination in urban dusts of Xi'an, central China. *Sci. Total Environ.* **355**, 176-186, doi:10.1016/j.scitotenv.2005.02.026.
- Yuen J. Q., P. H. Olin, H. S. Lim, S. G. Benner, R. A. Sutherland and A. D. Ziegler, 2012. Accumulation of potentially toxic elements in road deposited sediments in residential and light industrial neighborhoods of Singapore. *J. Environ. Manage.* **101**, 151-163, doi:10.1016/j.jenvman.2011.11.017.
- Zahran S., M. A. S. Laidlaw, S. P. McElmurry, G. M. Filippelli and M. Taylor, 2013. Linking source and effect: Resuspended soil lead, air lead, and children's blood lead levels in Detroit, Michigan. *Environ. Sci. Technol.* **47**, 2839-2845, doi:10.1021/es303854c.
- Zhang K. and S. Batterman, 2013. Air pollution and health risks due to vehicle traffic. *Sci. Total Environ.* **450-451**, 307-316, doi:10.1016/j.scitotenv.2013.01.074.