



Boletín Médico del Hospital Infantil de México

www.elsevier.es/bmhim



CLINICAL CASE

Chronic active EBV infection: the experience of the Samsung Medical Center in South Korea



Tae-Hee Lee, Young-Hyeh Ko *

Department of Pathology, Samsung Medical Center, Sungkyunkwan University, Seoul, Korea

Received 11 November 2015; accepted 3 December 2015

Available online 1 February 2016

KEYWORDS

Epstein-Barr virus;
Children;
Chronic active
infection;
NK lymphocytosis;
Mosquito-bite
hypersensitivity

Abstract

Background: Chronic active EBV infection (CAEBV) of T-cell or NK-cell type is an EBV+ polyclonal, oligoclonal or often monoclonal lymphoproliferative disorder (LPD) recognized as representing the spectrum of EBV-associated T-cell and NK-cell LPD with different clinical presentations; one systemic and two cutaneous disorders including hydroa vacciniforme-like T-cell LPD and mosquito bite hypersensitivity. The systemic form of the disease is characterized by fever, persistent hepatitis, hepatosplenomegaly and lymphadenopathy, which shows varying degrees of clinical severity depending on the immune response of the host and the EBV viral load.

Case reports: We described the clinicopathological findings of two children with CAEBV with a brief review of the literature.

Conclusions: Recognition of the disease is important for adequate management of the patient. EBV analysis should be included in the principal diagnostic tests for febrile children.

© 2015 Hospital Infantil de México Federico Gómez. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Virus de Epstein-Barr;
Niños;
Infección crónica
activa;
Linfocitosis de células
NK;
Hipersensibilidad por
picadura de mosquito

Infección crónica activa por virus de Epstein-Barr: experiencia del Centro Médico Samsung en Corea del Sur

Resumen

Introducción: La infección crónica activa (CA) de células T o células tipo NK por virus de Epstein-Barr (VEB) es un desorden linfoproliferativo (DLP) VEB+ policlonal, oligoclonal o, frecuentemente, monoclonal reconocido como representación del espectro del DLP de células T y células NK asociado con VEB que tiene diversas presentaciones clínicas: un padecimiento sistémico y dos cutáneos que incluyen el DLP de células T que semeja hidroa vacciniforme y la hipersensibilidad por picadura de mosquito. Los síntomas de la enfermedad sistémica incluyen

* Corresponding author.

E-mail addresses: yhko310@skku.edu, younghyeh.ko@samsung.com (Y.-H. Ko).

fiebre, hepatitis persistente, hepatoesplenomegalia y linfadenopatías que muestran diferente grado de severidad clínica, dependiendo de la respuesta inmune del hospedero y de la carga viral del VEB.

Casos clínicos: Se describen los hallazgos clínico-patológicos de dos niños con CAEBV y una breve revisión de la literatura.

Conclusiones: Es importante reconocer esta enfermedad para proporcionar el manejo adecuado al paciente. El análisis de VEB debería incluirse como una de las principales pruebas diagnósticas en niños con fiebre.

© 2015 Hospital Infantil de México Federico Gómez. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Chronic active Epstein-Barr virus infection (CAEBV) is a clinical term initially defined by Straus as a disease related to chronic or persistent infection of EBV. Suggested criteria for diagnosing severe chronic EBV infection includes the following: 1) severe illness of >6 months duration that began as primary EBV infection and is associated with grossly abnormal EBV antibody titers [IgG to viral capsid antigen (VCA)=~1:5120; antibody to early antigen (EA)=~1:640; or antibody to EBV nuclear antigen (EBNA) <2], and 2) histological evidence of major organ involvement such as interstitial pneumonia, hypoplasia of some bone marrow elements, uveitis, lymphadenitis, persistent hepatitis or splenomegaly, and 3) detection of increased quantities of EBV in affected tissues by anticomplementary immunofluorescence for EBNA or nucleic acid hybridization¹. In early reports, the cellular lineage of EBV-infected cells has not drawn attention; however, subsequent reports documented that EBV mainly exists in T-cells^{2,3} but not in B-cells, as revealed by dual staining immunofluorescence analysis³. Just as the accumulation of clinical and laboratory data for CAEBV, it has been recognized that clinical and laboratory findings of CAEBV are of a broader spectrum than initially thought. Because individual cases do not fulfill the diagnostic criteria initially proposed, Kimura and colleagues modified the diagnostic criteria for CAEBV as follows: (1) illness ≥ 3 months duration [EBV-related illness or symptoms including fever, persistent hepatitis, extensive lymphadenopathy, hepatosplenomegaly, pancytopenia, uveitis, interstitial pneumonia, hydroa vacciniforme (HV), and hypersensitivity to mosquito bites (HMB)]; (2) increased amounts of EBV DNA or grossly abnormal levels of EBV antibodies, for instance, detection of EBV DNA in tissues or peripheral blood by Southern blot hybridization; EBV-encoded small RNA-1-positive cells in affected tissues or peripheral blood, $>10^{2.5}$ copies of EBV DNA/ μg of DNA of peripheral blood mononuclear cells; and grossly abnormal levels of EBV antibodies (anti-VCA IgG titers ≥ 5120 or anti-EA IgG titers ≥ 640), and (3) no evidence of previous immunological abnormalities or other recent infection that might explain the observed condition.⁴

Based on the accumulated data, CAEBV of T-cell or NK-cell type has been recently defined as a systemic EBV+ polyclonal, oligoclonal or often monoclonal lymphoproliferative disorder (LPD) characterized by fever,

persistent hepatitis, hepatosplenomegaly and lymphadenopathy, which shows varying degrees of clinical severity depending on the immune response of the host and the EBV viral load.^{5,6}

CAEBV is often accompanied by cutaneous lesions such as severe mosquito bite allergy and hydroa-vacciniforme (HV)-like T-cell lymphoproliferative disease. Mosquito bite hypersensitivity (or mosquito bite allergy, MBH) is a unique cutaneous manifestation of CAEBV infection characterized by abnormally intense local reaction at arthropod bite area associated with systemic symptoms and signs like fever, lymphadenopathy and liver dysfunction. EBV genome is mainly found within NK cells of peripheral blood of MBH patients who often show NK cell lymphocytosis.⁷⁻¹⁰

HV-like LPD is an EBV-associated polyclonal, oligoclonal, or monoclonal cutaneous T-cell lymphoproliferative disease characterized by recurrent vesiculopapular eruptions, mainly on the face and arms. It shows a broad spectrum of clinical aggressiveness and usually a long clinical course with risk to develop systemic lymphoma. As the disease progresses patients develop severe and extensive skin lesions with systemic symptoms including fever, hepatosplenomegaly and lymphadenopathy. Classic HV, severe HV and HV-like T-cell lymphoma constitute a continuum spectrum of EBV-associated HV-like LPD.^{5,11-14}

Recently, the three disorders indicated above have been recognized as representing the spectrum of EBV-associated T-cell and NK-cell LPD with different clinical presentations; one systemic and two cutaneous disorders including HV-like T-cell LPD and MBH.¹⁴

2. Case reports

The patients described in the following two case reports depict typical clinical manifestations of CAEBV with MBH in children. Case 2 was reported previously.¹⁵

2.1. Case 1

A 15-year-old Korean boy was admitted to the hospital with chronic intermittent dermatological and various systemic symptoms and signs since he was 5 years old. Birth weight was normal and adequate for gestational age. At a recent visit to a pediatric department, he showed adequate height of 163.7 cm for his age, but a low body weight of 39.6 kg.




Figure 1 Case 1. (A) Mosquito bite hypersensitivity at 12 years of age. (B) Erythematous scaly skin lesion of the face at 15 years of age. (C) Similar skin lesion in the neck.

At 5 years old, fever, cough, and abdominal distention suddenly developed. White blood cell count was $12.1 \times 10^3/\mu\text{l}$ (lymphocytes 70%, segmented neutrophils 21%, eosinophils 1%, and monocytes 8%). Qualitative fluorescent antinuclear antibody was negative. Following a clinical diagnosis of intestinal perforation, he underwent a subtotal colectomy and ileostomy. One month after surgery, lymphocyte subset counts by flow cytometry showed low numbers of B- and T-cells, with markedly increased NK-cells (CD16+ CD56+ = 68%, normal range 5.6-31%); 1,579/ μl (normal range 100-430/ μl).

He was subsequently hospitalized five times in 3 years because of acute pharyngotonsillitis, upper respiratory infection, bronchiolitis, and pneumonia and experienced recurrent fever two or three times per month. At 10 years old, he was hospitalized because of upper respiratory tract infection. Splenomegaly of 10.76 cm was found incidentally during workup. From the age of 10, intense skin responses manifested at mosquito- or other bug-bitten sites that were characterized by an initial bullous change followed by ulceration, crust formation, and scarring with pigmentation (Fig. 1). Most lesions were <1 cm in diameter and were frequently accompanied by fever. Such cutaneous lesions tended to occur in the summer and on the upper and lower extremities. Laboratory tests performed at the time of severe mosquito bite allergy showed mild elevation of aspartate transaminase/alanine transaminase (AST/ALT) with significant elevation of alkaline phosphatase (200 U/l) and globulin (4.5 to 4.8 g/dl).

A skin biopsy obtained from a lesion was reported as indicating a possibility of Sweet's syndrome. Abdominal sonography indicated splenomegaly. Differential counts of peripheral blood showed low B- and T-cell populations, and NK-cell lymphocytosis [CD16+ CD56+ = 62%; 3520/ μl (normal range = 70-1200/ μl)]. The serum IgE level was increased to 20,370 U/ml although no specific response to various allergen stimuli was identified. Despite no proven family history, a clinical impression of hyper-IgE syndrome was made as the first differential diagnosis. A genetic study that included STAT3 sequencing was performed but failed to find the expected mutation. In considering the clinical course and

NK lymphocytosis, CAEBV or EBV-associated lymphoproliferative disease was considered as other differential diagnoses and relevant tests were established.

Serologic tests showed positivity of EBV VCA-IgG, EBV-early antigen, and negativity of EBV VCA IgM. Using a whole blood sample, real-time quantitative PCR for EBV DNA was performed to determine the viral load in blood (11,450 copies/5 μl of whole blood). Subsequently, he underwent a regular checkup of viral genome copy number, which was found to be consistently high, ranging from 212.5 copies/ μl to 1,562 copies/ μl of whole blood. His skin problem associated with mosquitoes or bugs had occurred intermittently ever since, while intermittent episodes of febrile illness devoid of skin lesion had also persisted. Nonspecific skin problem persisted for 15 months. Physical examination revealed erythematous to brownish scaly patches with yellowish crusts on his face, neck, scalp, upper chest, and upper back. He was treated with topical steroids and oral antihistamines under impression of seborrheic dermatitis for 11 months (Fig. 1). Two months prior to examination, his symptoms became aggravated and he visited a dermatology clinic. Two skin biopsies from his face and neck were taken and EBV in situ hybridization was performed. The biopsies showed increased lymphoplasma cells in the superficial and deep dermis with occasional periadnexal distribution. There was no clear atypia in most lymphoid cells. Immunohistochemistry showed many CD56+ lymphoid cells. There were numerous lymphocytes showing a nuclear positive reaction to EBER in situ hybridization. Retrospectively, EBER in situ hybridization was applied to both the skin biopsy of the lower leg taken for severe mosquito bite allergy and that of the perforated intestine resected 10 years ago (Fig. 2). Many lymphoid cells with a positive reaction were identified, not only in the skin but also in the intestinal wall and sinusoids of the liver and lymph node, which confirmed CAEBV infection with mosquito bite allergy and NK lymphocytosis. The patient was treated conservatively. Two years later he was readmitted because of hemophagocytic lymphohistiocytosis (HLH). He was treated according to the HLH-2004 treatment guideline¹⁶ and responded transiently, but later developed bowel perforation and died of sepsis.




Figure 2 Case 1. (A) Perforated colon at 5 years of age. (B) Microscopic findings show acute and chronic inflammation with granulation tissue. (C) EBER in situ hybridization reveals EBV-positive lymphocytes. (D) EBV-positive lymphocytes are positive for CD3.

2.2. Case 2

A 10-year-old Korean boy presented to the Department of Dermatology complaining of intense skin reactions at mosquito bite sites. He exhibited multiple scattered, punctuated ulcerations on bilateral lower extremities. In addition to the marked cutaneous reaction, he also endorsed systemic symptoms including fever following insect bites throughout his childhood. A skin biopsy was taken and local treatment for the lesion was administered without further systemic evaluation. At 16 years of age, the patient admitted with a palpable mass located on the left neck of 2–3 weeks' duration. On physical examination, a conglomerate of large lymph nodes was present in the left neck, the largest of which measured 7×5 cm. Due to his history of skin lesions, careful inspection of his legs was performed, revealing multiple shallow ulcers with healing scars. Computed tomography and positron emission tomography scan showed multiple poorly enhancing and variable sized homogeneous lymph nodes involving the left cervical region and bilateral inguinal areas. In addition, there were multifocal lesions demonstrating increased 2-deoxy-2-[fluorine-18] fluoro-D-glucose uptake in the skin and subcutaneous layer of bilateral cheeks and buttocks. On hematologic examination, blood cell counts and lactic acid dehydrogenase level were within normal limits. Anti-VCA IgG, anti-EA and anti-EBNA IgG were positive and anti-VCA IgM was negative. The EBV DNA copy number was 529.8 copies/ μ l of whole blood.

Lymph node biopsy was performed revealing widening of the paracortex with infiltration of heterogeneous inflammatory cells including small lymphocytes, histiocytes and many eosinophils. The inflammatory infiltrate was punctuated by scattered large mononuclear or multinuclear Reed-Sternberg (RS)-like atypical cells, which had large eosinophilic nucleoli and thick nuclear membrane. Using immunohistochemical study, the RS-like cells demonstrated a strong membranous stain for CD30 and perinuclear staining for CD15 and were positive for PAX-5 and LMP-1, but negative for CD20, CD3, and EBNA-2. Atypical mononuclear and multinuclear RS-like cells showed a positive signal by EBER in situ hybridization. Aside from the RS-like cells, many small lymphocytes scattered in the background were also positive for EBER in situ hybridization. No abnormal findings were noted in the bone marrow biopsy except for a few EBV-positive small lymphocytes. The skin biopsy from the region of mosquito-bite allergy in 2002 was retrospectively reviewed, revealing severe necrosis of the epidermis and upper dermis. There was infiltration of small lymphocytes, histiocytes and many eosinophils around the blood vessels and hair follicles and the blood vessel lumens were often obliterated by red blood cells (RBCs) and fibrin. Immunohistochemical analysis revealed that the infiltrating cells were heterogeneous in their lineage; some were positive to CD3 and CD4 or CD8 helper or cytotoxic T cells, whereas others were NK cells positive for CD56. Many EBV-positive cells were documented by EBER in situ hybridization. The diagnosis of chronic active EBV infection with mosquito bite hypersensitivity

Table 1 Clinical and laboratory findings of 13 patients with chronic active EBV infection related disease.

Patient	Age at admission	Sex	Diagnosis	Clinical findings at presentation									Onset (yrs of age)
				Fever	Abnormal LFT	HSM	LD	Non-specific skin lesion	MBH	NK lymphocytosis	Others		
1	10	M	CAEBV	+	+	+	+	-	-	UK	-	8	
2	10	M	CAEBV	+	+	+	+	-	-	-	Bowel perforation, cytopenia, HLH	9	
3	14	F	HV-like T-LPD	+	-	-	-	-	-	UK	IgA nephropathy	Infancy	
4	16	F	HV-like T-LPD	+	+	-	+	+	-	UK	-	UK	
5*	10	M	CAEBV with MBH	+	+	-	+	-	+	UK	-	10	
6	16	M	CAEBV with MBH	+	+	+	UK	-	+	UK	Eye angioedema	14	
7*	16	M	CAEBV with MBH	+	+	+	-	+	+	+	Bowel perforation, hypereosinophilia, HLH	5	
8	24	M	CAEBV with MBH	+	+	UK	UK	-	+	UK	HLH, nausea, vomiting	Since childhood	
9	22	M	CAEBV	+	+	+	+	-	-	+	Paroxysmal dyskinesia, severe oral ulcer, pneumonia, thrombocytopenia, HLH	20	
10	27	F	CAEBV	+	+	+	-	-	-	UK	Headache, pancytopenia, lung infiltration	3 months ago	
11	34	M	CAEBV	+	+	+	-	-	-	UK	Cytopenia, HLH, myalgia	33	
12	43	M	CAEBV	+	+	+	+	-	-	UK	-	42	
13	58	F	CAEBV	+	+	+	+	-	-	+	Myalgia, blurred vision	57	

Table 1 (Continued)

Patient	TCR γ Gene rearrangement	EBV load in whole blood/Serology	Treatment	FU (yrs)	Outcome	Causes of death
1	Monoclonal	1143 copies/5 μ l	HLH 2004 protocol	3	Died	Acute hepatic failure, HLH, hepatic encephalopathy
2	polyclonal	7.5 copies/5 μ l	Allo PBSCT	3	Alive	
3	polyclonal	ND	CHOP	13	Died	Sepsis, EBV+ T-cell lymphoma at larynx
4	polyclonal	ND	None	5	Alive	
5*	polyclonal	EA+, VCA IgM+, 35 copies/5 μ l	ABVD	13	Alive	
6	ND	7,000 copies/ μ l/EA+, VCA IgM+	None	2.5	Alive	
7*	polyclonal	1,562 copies/5 μ l, VCA IgM/IgG (-/+), EA+ EBNA+	HLH 2004 protocol	11	Died	HLH, hematochezia
8	ND	ND	IMVP16	10	Died	ANKL
9	polyclonal	150 copies/5 μ l	CHOP, ICE/DEXA	2	Died	HLH
10	polyclonal	6936 copies/5 μ l	UK	1.5/lost to FU	Alive	
11	ND	17,230 copies/5 μ l VCA IgG+, -VCA IgM- EA+, EBNA+	None	1	Died	UK
12	ND	EA+, VCA IgG+, EBNA+	None	0.5	Died	HLH
13	ND	326.40 copies/5 μ l	CVP	lost to FU	Alive	

5*: Case 2, 7*: Case 1.

MBH, mosquito-bite hypersensitivity; LFT, liver function test; LPD, lymphoproliferative disease; FU, follow up; ND, not done; ANKL, aggressive NK cell leukemia; HSM, hepatosplenomegaly; LD, lymphadenopathy; PBSCT, peripheral blood stem cell transplantation; HV, hydroa vacciniforme; HLH, hemophagocytic lymphohistiocytosis; UK, unknown.

and polymorphic lymphoproliferation simulating Hodgkin's lymphoma was made. The patient was treated with ABVD (Adriamycin, Bleomycin, Vinblastin, and Dacarbazine) for six cycles with complete remission. Currently he remains free of disease for 7 years after treatment.

3. Discussion

The clinical course of CAEBV patients depends on the balance between EBV-related factors and host immune function and can be smoldering, progressive, or aggressive. Some patients develop EBV+ T/NK cell lymphoma/leukemia.^{4,17} From 1998 to 2014, 13 patients aged 10–58 years (median age 22 years, nine males and four females) were diagnosed with CAEBV disease at Samsung Medical Center, Korea (Table 1). The common clinical findings included fever (13/13), hepatosplenomegaly (9/12), lymphadenopathy (7/11), NK lymphocytosis (3/4), MBH (4/13), and HV-like LPD (2/13). Some patients presented with bowel perforation, chorea, or brain infarction. In the median follow-up of 36 months, seven patients (54%) died of the disease, two patients (15%) had persistent disease, and two patients (17%) were free of disease. Two patients were lost to follow-up. The causes of death were hemophagocytic syndrome and organ failure in four patients, EBV-positive T cell lymphoma in one patient, and aggressive NK cell leukemia in one patient. B-lineage lymphoproliferation mimicking Hodgkin's lymphoma as seen in case 2 is rare but has been previously reported.¹⁸

CAEBV infection is almost always accompanied by varying degrees of lymphoproliferation. The clonality of EBV and EBV-infected T- or NK-cells varies and may be polyclonal, oligoclonal, or monoclonal. As the disease progresses from polyclonal lymphoproliferation to monoclonal disease, histological atypia increases. Ohshima et al. proposed the categorization of CAEBV into three groups—polymorphous and polyclonal, polymorphous and monoclonal, or monomorphic and monoclonal—based on clonality and histological changes.¹⁹ In the series reported by Ohshima et al., 8/48 patients with CAEBV infection were polyclonal for TCR gene rearrangement and the infiltrated cells displayed polymorphic histomorphology; 15 patients showed polymorphic morphology and biclonal or monoclonal TCR gene rearrangement; and 25 patients showed monomorphic histomorphology and monoclonal TCR gene rearrangement. Patients with the monomorphic and monoclonal type of CAEBV infection had poorer prognoses than those with polymorphic polyclonal or polymorphic monoclonal disease. The survival of the polymorphic/polyclonal and polymorphic/monoclonal groups did not differ significantly. Monomorphic/monoclonal groups of CAEBV by Ohshima et al. may correspond to systemic T-cell LPD according to the 2008 WHO classification.²⁰

In summary, CAEBV is a rare EBV-associated LPD of predominantly T-cell or NK-cell lineage with a broad spectrum of clinical presentation. Recognition of the disease is important for adequate management of the patient. EBV analysis through real-time PCR of a blood sample or EBV in situ hybridization in the affected tissue is important to avoid overlooking these patients and should be included in the principal diagnostic tests for febrile children.

Ethical disclosure

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflict of interest

The authors declare no conflict of interest of any nature.

References

1. Straus SE. The chronic mononucleosis syndrome. *J Infect Dis.* 1988;157:405–12.
2. Jones JF, Shurin S, Abramowsky C, Tubbs RR, Sciotto CG, Wahl R, et al. T-cell lymphomas containing Epstein-Barr viral DNA in patients with chronic Epstein-Barr virus infections. *N Engl J Med.* 1988;318:733–41.
3. Kikuta H, Taguchi Y, Tomizawa K, Kojima K, Kawamura N, Ishizaka A, et al. Epstein-Barr virus genome-positive T lymphocytes in a boy with chronic active EBV infection associated with Kawasaki-like disease. *Nature.* 1988;333:455–7.
4. Kimura H, Morishima T, Kanegane H, Ohga S, Hoshino Y, Maeda A, et al. Prognostic factors for chronic active Epstein-Barr virus infection. *J Infect Dis.* 2003;187:527–33.
5. Ko YH, Kim HY, Oh YH, Park G, Lee SS, Huh J, et al. EBV-associated T and NK cell lymphoproliferative disorders: consensus report of the 4th Asian Hematopathology Workshop. *J Haematopathol.* 2012;5:319–24.
6. Kimura H. Pathogenesis of chronic active Epstein-Barr virus infection: is this an infectious disease, lymphoproliferative disorder, or immunodeficiency. *Rev Med Virol.* 2006;16:251–61.
7. Satoh M, Oyama N, Akiba H, Ohtsuka M, Iwatsuki K, Kaneko F. Hypersensitivity to mosquito bites with natural-killer cell lymphocytosis: the possible implication of Epstein-Barr virus reactivation. *Eur J Dermatol.* 2002;12:381–4.
8. Ishihara S, Ohshima K, Tokura Y, Yabuta R, Imaishi H, Wakiguchi H, et al. Hypersensitivity to mosquito bites conceals clonal lymphoproliferation of Epstein-Barr viral DNA-positive natural killer cells. *Jpn J Cancer Res.* 1997;88:82–7.
9. Ishihara S, Yabuta R, Tokura Y, Ohshima K, Tagawa S. Hypersensitivity to mosquito bites is not an allergic disease, but an Epstein-Barr virus-associated lymphoproliferative disease. *Int J Hematol.* 2000;72:223–8.
10. Kawa K, Okamura T, Yagi K, Takeuchi M, Nakayama M, Inoue M. Mosquito allergy and Epstein-Barr virus-associated T/natural killer-cell lymphoproliferative disease. *Blood.* 2001;98:3173–4.
11. Iwatsuki K, Satoh M, Yamamoto T, Oono T, Morizane S, Ohtsuka M, et al. Pathogenic link between hydroa vacciniforme and Epstein-Barr virus-associated hematologic disorders. *Arch Dermatol.* 2006;142:587–95.
12. Cho KH, Kim CW, Lee DY, Sohn SJ, Kim DW, Chung JH. An Epstein-Barr virus-associated lymphoproliferative lesion of the skin presenting as recurrent necrotic papulovesicles of the face. *Br J Dermatol.* 1996;134:791–6.

13. Cho KH, Lee SH, Kim CW, Jeon YK, Kwon IH, Cho YJ, et al. Epstein-Barr virus-associated lymphoproliferative lesions presenting as a hydroa vacciniforme-like eruption: an analysis of six cases. *Br J Dermatol*. 2004;151:372–80.
14. Attygalle AD, Cabeçadas J, Gaulard P, Jaffe ES, de Jong D, Ko YH, et al. Peripheral T-cell and NK-cell lymphomas and their mimics; taking a step forward—report on the lymphoma workshop of the XVIth meeting of the European Association for Haematopathology and the Society for Hematopathology. *Histopathology*. 2014;64:171–99.
15. Park S, Bahng S, Kim EK, Park SB, Sung YK, Kim WS, et al. Hodgkin's lymphoma arising in a patient with hypersensitivity to mosquito bites: a case report. *J Clin Oncol*. 2010;28:e148–50.
16. Henter JI, Horne A, Aricó M, Egeler RM, Filipovich AH, Imashuku S, et al. HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48:124–31.
17. Kimura H, Hoshino Y, Kanegane H, Tsuge I, Okamura T, Kawa K, et al. Clinical and virologic characteristics of chronic active Epstein-Barr virus infection. *Blood*. 2001;98:280–6.
18. Tamayose K, Egashira M, Sugimoto K, Ando J, Mori S, Oshimi K. Epstein-Barr virus-positive Hodgkin's lymphoma in a patient with chronic active Epstein-Barr virus infection. *Int J Hematol*. 2004;80:199–200.
19. Ohshima K, Kimura H, Yoshino T, Kim CW, Ko YH, Lee SS, et al. Proposed categorization of pathological states of EBV-associated T/natural killer-cell lymphoproliferative disorder (LPD) in children and young adults: overlap with chronic active EBV infection and infantile fulminant EBV T-LPD. *Pathol Int*. 2008;58:209–17.
20. Quintanilla-Martínez L, Kimura H, Jaffe ES. Epstein-Barr virus (EBV) positive T-cell lymphoproliferative disorders of childhood. In: Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et al., editors. *WHO Classification of Tumors of Haematopoietic and Lymphoid Tissues*. Lyon: IARC; 2008. p. 278–80.