



A FEW ISLANDS OF PRELIMINARY RESEARCH AMONG A SEA OF UNKNOWN: MOVING PLANT ECOPHYSIOLOGY FORWARD IN MEGADIVERSE COUNTRIES

*Report on the Symposium "Advances in Plant Ecophysiology," within the VII Mexican Ecology Congress,
Sociedad Científica Mexicana de Ecología, Juriquilla, Querétaro,
Mexico, 3 October 2019*

UNAS CUANTAS ISLAS DE INVESTIGACIÓN PRELIMINAR EN UN MAR DE INCÓGNITAS: AVANZANDO LA ECOFISIOLOGÍA DE PLANTAS EN PAÍSES MEGADIVERSOS

*Reseña del Simposio «Avances en ecofisiología vegetal», acaecido el 3 de octubre de 2019 en el marco del
VII Congreso Mexicano de Ecología, Sociedad Científica Mexicana de Ecología, Juriquilla, Querétaro,
México, octubre 3, 2019*

OSCAR BRIONES¹, CASANDRA REYES-GARCÍA², EDISON A. DÍAZ-ÁLVAREZ³, CLARA TINOCO-OJANGUREN⁴,
ALEJANDRO E. CASTELLANOS⁵, ERICK DE LA BARRERA^{6*}, JOSÉ LUIS ANDRADE²

¹Instituto de Ecología AC, Xalapa, Veracruz, Mexico.

²Unidad de Recursos Naturales, Centro de Investigación Científica de Yucatán, Mérida, Yucatán, Mexico.

³Instituto de Investigaciones Forestales, Universidad Veracruzana, Xalapa, Veracruz, Mexico.

⁴Departamento de Ecología de la Biodiversidad, Instituto de Ecología, Universidad Nacional Autónoma de México, Hermosillo, Sonora, Mexico.

⁵Departamento de Investigaciones Científicas y Tecnológicas, Universidad de Sonora, Hermosillo, Sonora, Mexico.

⁶Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Morelia, Michoacán, Mexico.

*Corresponding author: delabarrera@unam.mx

Keywords: Climate change, Drought, Ecosystem productivity, Environmental pollution, Epiphytes, Food security.

Palabras clave: Cambio climático, contaminación ambiental, epífitas, productividad del ecosistema, seguridad alimentaria, sequía.

Plant physiological ecology is not a new scientific discipline in Mexico. Indeed, ecophysiological research has been conducted in this country since at least 1976, when the late Carlos Vázquez-Yanes, a dear professor of some of us, first published his work on germination of tropical rainforest plants (Vázquez-Yanes 1976, Vázquez-Yanes & Orozco-Segovia 1984). Indeed, Prof. Vázquez-Yanes is considered the founder of Mexican plant ecophysiology. At present, however, the development of this discipline appears lagging relative to other Mexican ecological sciences. Factors contributing to this deceiving perception might include its technological dependence, the elevated costs of some research protocols, and the intrinsic difficulties of combining

extensive fieldwork with a precise knowledge and use of first principles of physics and chemistry, which are essential for understanding physiological processes.

A principal intention of ecological sciences is to explain biodiversity: why are some species and not others found in certain places (Lambers *et al.* 2008). As an experimental science, plant ecophysiology focuses on the interactions among plants, the environment, and other biological species. In this context, the VII Mexican Ecology Congress focused on "addressing the complexity of nature." Some examples of how ecophysiology can address said complexity were discussed during the symposium.



From elucidating fundamental processes to anticipating climate change responses. A basic understanding of physiological functions is still lacking for many species. This is certainly the case for the establishment phase of ferns from the Mexican cloud forest (Oscar Briones, Instituto de Ecología AC, Xalapa, Ver., México). Although cloud forests receive large amounts of rainfall, the soil surface and tree branches are exposed to frequent periods of water scarcity that can cause mortality of gametophytes and young sporophytes of fern species. Experimental evidence with herbaceous, arborescent, and epiphytic ferns shows that the water requirement of spore germination is correlated with the distribution of the adult sporophyte in the forest and that the spores can germinate with very low light availability. In turn, gametophyte growth depends on multiple environmental conditions and resources, with amount of light, relative humidity, thickness of litter layer, and disturbance being most important. The distribution and abundance of sporophytes and spore physiology support the hypothesis that the coexistence of fern species in the cloud forest occurs by differentiation in the regeneration niche. However, ecological interactions during the establishment of gametophytes and young sporophytes are still virtually unknown.

Considering that ample portions of Mexico are arid or semiarid and that climate change scenarios anticipate an increase in air temperature and a decrease in annual precipitation in various regions of the country ([Sáenz-Romero *et al.* 2010](#)), current arid lands can be considered natural laboratories for studying plant responses to impending climate change. Water stress is one of the main selection factors in such regions, giving rise to different coping strategies. Plant responses to drought can actually be placed along an isohydric-anisohydric continuum, a conceptual axis that describes how plants regulate their water potential, based on their water relations strategies ([Tardieu & Simonneau 1998](#)). This framework integrates multiple components of water transport and stomatal regulation of water status, based on which, the anatomical, morphological, and physiological hydraulic characteristics of tree species of the Sonoran Desert with different phenology and age (young and adult plants) have been classified (Clara Tinoco-Ojanguren, Universidad Nacional Autónoma de México, Hermosillo, Son., Mexico), finding a clear expression of the different strategies of desert tree plants living under the same stress conditions. For young and adult plants, isohydric and anisohydric groups were

clearly differentiated by their hydroscape area and phenology; both characteristics showed a high correlation with leaf and stem hydraulic characteristics and other important variables. Studies of several species growing under extreme conditions have provided important information of functional diversity and plant responses to climatic change.

While arid and semiarid lands are commonly associated with low primary productivity, studies across the arid northwestern Mexico over the past two decades have revealed otherwise (Alejandro E. Castellanos, Universidad de Sonora, Hermosillo, Son., Mexico). In particular, high maximum photosynthetic rates, leading to high ecosystem productivity, have been found across different vegetation types, including much simplified socio-ecosystems, such as pastures sown with the exotic buffelgrass. Thus, the current arid environment-low productivity paradigm needs to be revisited, in particular its relation to resource use efficiencies and nutrient stoichiometry interactions ([Castellanos *et al.* 2018](#)).

Ecophysiology in anthropic environments. Human activities, such as industrial manufacturing and the use of motor vehicles, release a substantial amount of pollutants to the environment, with dire consequences on biodiversity and human health ([Rockström *et al.* 2009](#), [Landigan *et al.* 2017](#)). For the particular case of carbon and nitrogen gaseous emissions, pollutants have specific isotopic signatures that depend on the source. For this reason, the spatial distribution of atmospheric pollution is traceable by means of the isotopic composition of the vegetation (Edison A. Díaz-Álvarez, Universidad Veracruzana, Xalapa, Ver., Mexico). For example, plants that take up carbon from anthropic sources show a significant reduction of their $\delta^{13}\text{C}$ values, while the $\delta^{15}\text{N}$ values tend to increase for plants exposed to nitrogen emissions from urban areas ([Díaz-Álvarez & de la Barrera 2018](#)). This is not the case, for instance, for orchids from a cloud forest in the Colombian Andes, which display $\delta^{15}\text{N}$ values indicative of negligible pollution despite their close proximity (less than 50 km) to the megalopolis of Bogotá ([Díaz-Álvarez *et al.* 2019](#)). Biomonitoring of atmospheric emissions can thus be useful for evaluating ecosystem integrity ([Díaz-Álvarez *et al.* 2018](#)).

Another important component of pollution is the release of heavy metals, such as lead, which is normally discharged in tropical coastal areas, where mangroves occur. Mangrove

forests, which are the most productive and diverse natural wetland systems, grow in a complex environmental network governed by salinity and hydro-period. To help restoring affected mangrove areas, we need to make detailed experiments to observe anatomical and physiological changes in controlled environments. From such experiments, mangrove species have been identified that accumulate lead in roots in the absence of salt, with a greater translocation to the leaves under conditions of higher salinity (José Luis Andrade, Centro de Investigación Científica de Yucatán, Mérida, Yuc., Mexico).

Habitat fragmentation is well characterized as having a negative effect on ecosystems and biodiversity ([Aide *et al.* 2013](#), [Haddad *et al.* 2015](#)), but the physiological ecology effects on tropical plants are hardly documented, as discussed for ecosystems within the Yucatan Peninsula (Casandra Reyes-García, Centro de Investigación Científica de Yucatán, Mérida, Yuc., Mexico). Permanent plots set up across the Peninsula show that forests and coastal sand dune scrubs in the north of the Peninsula are experiencing higher population decline in both trees and epiphytic bromeliads, compared to those in the southern region. The northern region is highly fragmented, mainly as a result of urbanization, which is especially grave for coastal dune communities ([Martínez *et al.* 2014](#)). There are differences among susceptible and resilient species. For example, trees with higher water demand, associated with permanent water table access, are having higher mortality. This raises the question whether the water table contamination and excessive aquifer exploitation are reducing tree access and/or affecting water quality. The most affected trees are also those that host the highest numbers of epiphytic bromeliads, in turn affecting their populations.

Parallel to what occurs in urban ecosystems, agricultural landscapes display a reduction of biological species and ecosystem functions stemming from land use change, greenhouse gas emissions, and agrochemical pollution ([McDonald 2010](#)). However, agriculture is also one of the most vulnerable sectors to global environmental change, as agricultural operations greatly depend on certain climatological regularity ([de la Barrera 2016](#)). Considering that food is a topic that everybody has some familiarity with, it can be a useful heuristic for bringing biodiversity to the public discourse, just like "global warming" has become a well-known environmental issue (Erick de la Barrera, Universidad Nacional Autónoma de México, Morelia, Mich., Mexico).

Moving forward. A logical starting point for developing an action plan for further advancement of plant ecophysiology is a two-fold diagnosis: determine the scope and urgency of environmental questions and problems that the discipline can answer. However, despite that the combined work of the research groups participating in the symposium has considered a few hundred different species over the past several decades and various studies describing our current knowledge on plant physiological diversity from different ecosystems have already been published ([Castellanos-Villegas *et al.* 2010](#), [Tinoco-Ojanguren *et al.* 2018](#)), the syntheses presented in the symposium should still be considered preliminary, as they represent a minuscule fraction of plant biodiversity, *i.e.*, less than 3 % of the species from the Sonoran Desert, a region with a high frequency of endemism ([Shreve & Wiggins 1964](#)), and less than 3 % of the species from the Yucatán Peninsula, a region with high rates of deforestation ([Valdez-Hernández *et al.* 2015](#)). It is almost like a "few islands of knowledge among a sea of unknown". We identified a need to train more ecophysicists to work in the different parts of the country. As an initial step, we are releasing a website at <http://ecofisiologia.mx/>, as a point of access for sharing new research, information on graduate programs, and other academic activities.

Acknowledgments

We thank the board of the Sociedad Científica Mexicana de Ecología for facilitating the organization of the Plant Ecophysiology Symposium; our students and coauthors for enriching our ongoing research; and funding by the Consejo Nacional de Ciencia y Tecnología (Grant 284418, to O.B.). We declare that this is an original work, that it has not been previously published, nor is it currently under consideration by another journal; and the authors do not have a conflict of interest. We also declare that said work does not infringe the intellectual property rights of another person, research, or entity. We agree that the manuscript may be reviewed with the plagiarism programs that the journal uses for that purpose. We authorize the Sociedad Botánica de México, A.C. in a non-exclusive manner through its scientific dissemination body Botanical Sciences to correct, edit, publish, preserve, and distribute in their repositories and other digital platforms online and printed our manuscript. In addition, we authorize the work to be made available to users in open access, respecting the author's rights in accordance with the conditions set by the journal's license:

License CC Recognition-Non-Commercial 4.0 International
<https://creativecommons.org/licenses/by-nc/4.0/>

Literature cited

Aide TM, Clark ML, Grau HR, López-Carr D, Levy MA, Redo D, Bonilla-Moheno M, Riner G, Andrade-Núñez MJ, Muñiz M. 2013. Deforestation and reforestation of Latin America and the Caribbean (2001-2010). *Biotropica* **45**: 262-271. DOI: <https://doi.org/10.1111/j.1744-7429.2012.00908.x>

Castellanos-Villegas AE, Bravo LC, Koch GW, Llano J, López D, Méndez R, Rodríguez JC, Romo R, Sisk TD, Yanes G. 2010. Impactos Ecológicos por el Uso del Terreno en el Funcionamiento de Ecosistemas Áridos y Semi-Áridos de Sonora. In: Molina-Freaner FE, Van Devender T, eds. *Diversidad Biológica del Estado de Sonora*. Mexico City: Universidad Nacional Autónoma de México-Comisión Nacional para el uso y Conocimiento de la Biodiversidad. pp. 157-186. ISBN 978-607-02-0427-2

Castellanos AE, Llano-Sotelo JM, Machado-Encinas LI, López-Piña JE, Romo-León JR, Sardans J, Peñuelas J. 2018. Foliar C, N, and P stoichiometry characterize successful plant ecological strategies in the Sonoran Desert. *Plant Ecology* **219**: 775-788. DOI: <https://doi.org/10.1007/s11258-018-0833-3>

de la Barrera E. 2016. COP-eration for global food security. *F1000Research* **5**: 2814. DOI: <https://doi.org/10.12688/f1000research.10323.1>

Díaz-Álvarez EA, de la Barrera E. 2018. Characterization of nitrogen deposition in a megalopolis by means of atmospheric biomonitoring. *Scientific Reports* **8**: 13569. DOI: <https://doi.org/10.1038/s41598-018-32000-5>

Díaz-Álvarez EA, Felix JD, de la Barrera E. 2019. Elemental and isotopic assessment for Colombian orchids from a montane cloud forest: a baseline for global environmental change. *Acta Physiologiae Plantarum* **41**: 99. DOI: <https://doi.org/10.1007/s11738-019-2893-y>

Díaz-Álvarez EA, Lindig-Cisneros R, de la Barrera E. 2018. Biomonitoring of atmospheric nitrogen deposition: potential uses and limitations. *Conservation Physiology* **6**: coy011. DOI: <https://doi.org/10.1093/conphys/coy011>

Haddad NM, Brudvig LA, Cloibert J, Davies KF, González A, Holt RD, Lovejoy TE, Sexton JO, Austin MP, Collins CD, Cook WM, Damshen EI, Ewers RM, Foster BL, Jenkins CN, King AJ, Laurance WF, Levey DJ, Margulies CR, Melbourne BA, Nicholls AO, Orrock JL, Song DX, Townsend JR. 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**: e1500052. DOI: <https://doi.org/10.1126/sciadv.1500052>

Lambers H, Chapin III FS, Pons TL. 2008. *Plant Ecophysiology*. New York: Springer. ISBN: 978-0-387-78341-3

Landrigan PJ, Fuller R, Acosta NJR, Adeyi O, Arnold R, Basu N, Baldé AB, Bertollini R, Bose-O'Reilly S, Boufford JI, Breysse PN, Chiles T, Mahidol C, Coll-Seck AM, Cropper ML, Fobil J, Fuster V, Greenstone M, Haines A, Hanrahan D, Hunter D, Khare M, Krupnick A, Lanphear B, Lohani B, Martin K, Mathiasen KV, McTeer MA, Murray CJL, Ndhimananjara J, Perera F, Potočnik J, Preker A, Ramesh J, Rockström J, Salinas C, Samson LD, Sandilya K, Sly PD, Smith KR, Steiner A, Stewart RB, Suk WA, van Schayck OCP, Yadama GN, Yumkella K, Zhong M. 2017. The Lancet Commission on pollution and health. *The Lancet* **391**: 462-512. DOI: [https://doi.org/10.1016/S0140-6736\(17\)32345-0](https://doi.org/10.1016/S0140-6736(17)32345-0)

McDonald BL. 2010. *Food Security*. Cambridge, UK: Polity Press, ISBN: 978-0745648088

Martínez ML, Mendoza-González G, Silva-Casarín R, Mendoza-Baldwin E. 2014. Land use changes and sea level rise may induce a "coastal squeeze" on the coasts of Veracruz, Mexico. *Global Environmental Change* **29**: 180-188. DOI: <https://doi.org/10.1016/j.gloenvcha.2014.09.009>

Rockström J, Steffen W, Noone K, Persson Å, Chapin III FS, Lambin EF, Lenton TM, Scheffer M, Folke C, Schellnhuber HJ, Nykvist B, de Wit CA, Hughes T, van der Leeuw S, Rodhe H, Sörlin S, Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J, Walker B, Liverman D, Richardson K, Crutzen P, Foley JA. 2009. A safe operating space for humanity. *Nature* **461**: 472-475. DOI: <https://doi.org/10.1038/461472a>

Sáenz-Romero C, Rehfeldt GE, Crookston NL, Duval P, St-Amant R, Beaulieu J, Richardson BA. 2010. Spline models of contemporary, 2030, 2060 and 2090 climates for Mexico and their use in understanding climate-change impacts on the vegetation. *Climatic Change* **102**: 595-623. DOI: <https://doi.org/10.1007/s10584-009-9753-5>

Shreve F, Wiggins IL. 1964. *Vegetation and flora of the Sonoran Desert*. Vols. I & II. Stanford, California: Stanford University Press. ISBN: 978-0804701631

Tardieu F, Simonneau T. 1998. Variability among species of stomatal control under fluctuating soil water status and evaporative demand: modelling isohydric and anisohydric behaviours. *Journal of Experimental Botany* 49: 419-432. DOI: https://doi.org/10.1093/jxb/49.Special_Issue.419

Tinoco-Ojanguren C, Andrade JL, Briones O, Castellanos A. 2018. Functional Diversity in Plants: Implications for Conservation Issues of the Mexican Biodiversity. In: Ortega-Rubio A, ed. *Mexican Natural Resources Management and Biodiversity Conservation*. New York: Springer Nature . pp. 519-554. DOI: <https://doi.org/10.1007/978-3-319-90584-6>

Valdez-Hernández M, González-Salvaterra C, Reyes-García C, Jackson PC, Andrade JL. 2015. Physiological ecology of vascular plants of the Yucatan Peninsula. In: Islebe G, Calmé S, Schmook B, León J, eds. *Biodiversity and Conservation of the Yucatan Peninsula, Mexico*. New York: Springer Ecology Series, pp. 97-129. DOI: https://doi.org/10.1007/978-3-319-06529-8_5

Vázquez-Yanes C. 1976. Estudios sobre la ecofisiología de la germinación en una zona cálido-húmeda de México. In: Gómez-Poma A, Vásquez-Yanes C, Del Amo S, Butanda A, eds. *Investigaciones sobre la Regeneración de Selvas Altas en Veracruz, México*. Ciudad de México: Editorial Continental, pp. 279-387. ISBN: 9684440472

Vázquez-Yanes C, Orozco-Segovia A. 1984. Ecophysiology of seed germination in the tropical humid forests of the world: A review. In: Medina E, Mooney HA, Vásquez-Yanes C, eds. *Physiological Ecology of Plants of the Wet Tropics*. The Hague: Dr. W. Junk Publishers, pp. 37-50. DOI: <https://doi.org/10.1007/978-94-009-7299-5>; ISBN: 978-94-009-7301-5

Associate Editor: Arturo de Nova Vázquez

Author contributions: Oscar Briones organized the symposium and wrote the manuscript, Casandra Reyes-García wrote the manuscript, Edison Díaz-Álvarez wrote the manuscript, Clara Tinoco-Ojanguren organized the symposium and wrote the manuscript, Alejandro E. Castellanos organized the symposium and wrote the manuscript, Erick de la Barrera wrote and coordinated the manuscript, and José Luis Andrade organized the symposium and wrote the manuscript. All authors revised and approved the final version. The author order follows the sequence of presentations during the symposium.