

New record of mollusks from the El Molino mammoth site, Parras, Coahuila, Mexico

Nuevo registro de moluscos del yacimiento mamuts El Molino, Parras, Coahuila, México

Perla Guadalupe **Butrón Xancopinca**^{1,*}, Alexander **Czaja**², Martha **Carolina Aguillón**³,
Rosario **Gómez Núñez**³, Ignacio **Vallejo González**³, José Luis **Estrada Rodríguez**²,

¹ Universidad Michoacana de San Nicolás de Hidalgo, Instituto de Investigaciones en Ciencias de la Tierra, Ciudad Universitaria, 58030, Morelia, Michoacán, Mexico.

² Facultad de Ciencias Biológicas, Universidad Juárez del Estado de Durango, 35010, Gómez Palacio, Durango, Mexico.

³ Laboratorio de Paleontología, Museo del Desierto, Boulevard Carlos Abedrop Dávila #3745, 25022, Saltillo, Coahuila, Mexico.

* Corresponding author: (P.G. Butrón Xancopinca) perlab2010@hotmail.com

How to cite this article:

Butrón Xancopinca, P.G., Czaja, A., Carolina Aguillón, M., Gómez Núñez, R., Vallejo González, I., Estrada Rodríguez, J.L., 2022, New record of mollusks from the El Molino mammoth site, Parras, Coahuila, Mexico: Boletín de la Sociedad Geológica Mexicana, 74 (1), A051021. <http://dx.doi.org/10.18268/BSGM2022v74n1a051021>

Manuscript received: April 4, 2021
Corrected manuscript received: June 26, 2021
Manuscript accepted: October 5, 2021

Peer Reviewing under the responsibility of Universidad Nacional Autónoma de México.

This is an open access article under the CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

ABSTRACT

We present new Late Pleistocene and Holocene records of a land and freshwater malacofauna assemblage from the mammoth bearing site El Molino in Parras, Coahuila, northern Mexico. We identified 19 mollusk taxa, 14 species were found within the Late Pleistocene sediments and 10 species in the different strata belonging to the Holocene. The gastropods *Gastrocopta tappaniana*, *Pupilla hebes* and *Habroconus* sp. are new Late Pleistocene records for Mexico, the first two being previously recorded for United States Pleistocene deposits. New Mexican Holocene fossil records include *Euglesa casertana*, *Galba humilis*, *Gastrocopta cristata*, *Zonitoides arboreus*, *Hawaiiia minuscula*, and *Deroceras laeve*. The habitat requirements of the El Molino site malacofauna assemblage provides additional information on the environmental changes that occurred during the Pleistocene-Holocene transition. Woodland associated, hygrophilic and hydrophilic malacofauna suggest a humid forested and grassland habitat during the Late Pleistocene, which subsequently changed to xeric conditions with the colonization of xerophytic and aridity-tolerant aquatic species during and after the Pleistocene-Holocene transition at the study area.

Keywords: Late Pleistocene, Holocene, freshwater mollusks, terrestrial mollusks, paleoreconstruction.

RESUMEN

Se presentan nuevos registros de un ensamblaje de moluscos terrestres y de agua dulce perteneciente al Pleistoceno tardío y el Holoceno del yacimiento de mamuts El Molino en Parras, Coahuila, ubicado en el norte de México. Se identificaron 19 taxones de moluscos, entre los cuáles 14 especies se distribuyen dentro del Pleistoceno tardío y 10 especies en los distintos estratos del Holoceno. Los nuevos registros de moluscos del Pleistoceno tardío para México incluyen a las especies *Gastrocopta tappaniana*, *Pupilla hebes* y *Habroconus* sp. de las cuáles, las primeras dos habían sido registradas anteriormente en depósitos Pleistocénicos de los Estados Unidos. En el Holoceno de México, se presentan por primera vez los registros fósiles de las especies *Euglesa casertana*, *Galba humilis*, *Gastrocopta cristata*, *Zonitoides arboreus*, *Hawaiiia minuscula* y *Deroceras laeve*. Los requisitos de hábitat del conjunto de malacofauna hallada en el sitio El Molino proponen información adicional sobre los cambios ambientales sucedidos durante la transición Pleistoceno-Holoceno. La presencia de malacofauna higrófila e hidrófila asociada a bosques sugiere un hábitat boscoso húmedo con pastizales durante el Pleistoceno tardío, el cual cambió posteriormente a condiciones xéricas con la colonización de especies xerófitas y acuáticas tolerantes a la aridez durante y después de la transición Pleistoceno-Holoceno en el área.

Palabras clave: Pleistoceno tardío, Holoceno, moluscos de agua dulce, moluscos terrestres, paleoreconstrucción.

1. Introduction

Although Northern Mexico presents extensive areas with sediments of numerous paleolakes that covered the region during the late Quaternary, only a few sites have been studied for their fossil remains. Nevertheless, more studies have been carried out recently, especially from the paleolakes of the Chihuahuan and Sonoran Deserts with species-rich molluscan assemblages (Catto and Bachhuber, 2000; Czaja *et al.*, 2014a, 2014b).

We present a new assemblage of terrestrial and freshwater malacofauna located at the southern portion of the state of Coahuila in northern Mexico. The study area is located among farmlands at Rancho Buena Fe, called the El Molino Mammoth site, approximately 4 km east of the city of Parras de la Fuente (Figure 1). The locality was described in detail by Miller *et al.* (2008) and consists of an abandoned well site approximately eight meters deep by 6 meters wide, excavated in the year 2000 (25° 25'54.8" N, 102° 08'56.52" W; Elevation 1,530 m) (Figures 2a and 2b). The same authors carried out a Carbon14 dating of

the basal sediments of the cross section which yielded an age of 11,740 ± 50 years B.P. (Late Pleistocene). The macrofauna remains found at the site included mammoth (*Mammuthus columbi*), horse (*Equus ssp.*), camels (*Camelops cf. hesternus*) and several small mammal remains (Miller *et al.*, 2008). During the last few years, partial collapses within the well have buried the Late Pleistocene sediments, making them inaccessible for sampling.

This paper focuses on reporting new Late Pleistocene and Holocene records of a continental malacofauna assemblage from the El Molino Mammoth site in Rancho Buena Fe, Parras, Coahuila, as well as to propose additional information on the environmental changes during the Pleistocene-Holocene transition.

2. Materials and methods

Because of the collapsed basal part of the geological strata, Late Pleistocene sediments could not be sampled for this study. However, we used the sediments remaining from the mammoth excava-




Figure 1 Location of the abandoned well at the El Molino Mammoth site in Rancho Buena Fe, Parras, Coahuila, Mexico.




Figure 2 a. The El Molino Mammoth sites abandoned well in present-day Rancho Buena Fe; b. Image of the abandoned well from the year 2000 excavation; c. Geological profile of the Holocene strata sampling points. Black dash lines represent the start of the D strata. Image b is modified from Miller *et al.* (2008).

tion in 2000, which were stored at the laboratory of Paleontology of the Museo del Desierto in Saltillo, Coahuila, Mexico.

At the site, five 1 kg sediment samples were collected from the 3.30 m Holocene strata currently exposed within the abandoned well (Figure 2c). Samples from the C and D strata correspond to the Late Pleistocene-Holocene transition, while the remaining B1, B2 and A samples belong to the Holocene (Miller *et al.*, 2008). The samples were screened through two sieves with 0.5 mm and 0.3 mm mesh sizes. The selection of mollusks was carried out under a stereoscopic microscope, and subsequently washed with 3% hydrogen peroxide in order to

remove mud or any other organic residue. All shells were photographed with a Zeiss AxioCam ERc 5s camera attached to a Zeiss Stemi 2000-C microscope. Mollusk identifications are based on Pilsbry (1946, 1948), Leonard (1950, 1952), Metcalf and Smartt (1997), Nekola (2004), Wethington *et al.* (2009), Nekola and Coles (2010), and Walther *et al.* (2010). The revision was based mainly on biodiversity websites including WoRMS (World Register of Marine Species), and MolluscaBase (MolluscaBase, 2021). The material is part of the University Juárez Malacological Collection (UJMC) and is housed at the Faculty of Biological Science of the Juarez State University of Durango (UJED).

Table 1. Taxonomic list of Late Pleistocene and Holocene malacofauna from the El Molino Mammoth site in Rancho Buena Fe, Parras, Coahuila.

Family	Genus	Species	Late	
			Pleistocene	Holocene
Bivalves				
Sphaeriidae	<i>Euglesa</i>	<i>E. compressa</i>	x	
		<i>E. casertana</i>		x
Aquatic				
Lymnaeidae	<i>Galba</i>	<i>G. humilis</i>		x
		<i>G. obrussa</i>	x	x
Physidae	<i>Physella</i>	<i>P. acuta</i>		x
Planorbidae	<i>Gyraulus</i>	<i>G. parvus</i>	x	
	<i>Ferrissia</i>	<i>F. californica</i>	x	
Terrestrial				
Ellobiidae	<i>Carychium</i>	<i>C. exiguum</i>	x	
Succineidae	<i>Succinea</i>	<i>Succinea</i> sp.	x	x
Cochlicopidae	<i>Cochlicopa</i>	<i>C. lubrica</i>	x	
Gastrocoptidae	<i>Gastrocopta</i>	<i>G. cristata</i>		x
		<i>G. tappaniana</i>	x	
Pupillidae	<i>Pupilla</i>	<i>P. hebes</i>	x	
Valloniidae	<i>Vallonia</i>	<i>V. gracilicosta</i>	x	
Gastrodontidae	<i>Zonitoides</i>	<i>Z. arboreus</i>	x	x
	<i>Glyphyalinia</i>	<i>G. indentata</i>	x	x
Euconulidae	<i>Habroconus</i>	<i>Habroconus</i>	x	
		sp.		
Pristilomatidae	<i>Hawaia</i>	<i>H. minuscula</i>		x
Agriolimacidae	<i>Deroceras</i>	<i>D. laeve</i>	x	x

3. Results

The studied material comprises a total of 19 mollusk taxa belonging to 15 families and 17 genera. Mollusks found within the Late Pleistocene strata include 14 species belonging to 12 families and 14 genera, while the Holocene strata includes 10 species (8 families and 9 genera) (Table 1).

3.1. SYSTEMATIC DESCRIPTIONS

Bivalvia Linnaeus, 1758

Family Sphaeriidae Deshayes, 1855

Genus *Euglesa* Jenyns, 1832

Euglesa casertana (Poli, 1791)

(Figure 3j)

Ecology: Perennial and ephemeral swamp ponds,

streams, rivers and lakes (Herrington, 1962).

Current distribution: United States, Mexico, Cuba, Puerto Rico and from Honduras to Patagonia (Herrington, 1962).

Stratigraphic remarks: The stratigraphic distribution of *E. casertana* at the El Molino Mammoth site is limited to the A strata belonging to the Holocene. This species has also been found in a Late Pleistocene site of San Luis Potosí, Mexico (Arroyo-Cabrales *et al.*, 2008).

Referred material: UJMC 600, 18 specimens.

Measurements: Length: 2.2 mm; diameter: 3 mm.

Euglesa compressa (Prime, 1852)
(Figure 3i)



Figure 3 a. *Deroceras laeve*; b. *Galba humilis*; c. *Ferrissia californica*; d. *Physella acuta*; e. *Gyraulus parvus*; f. *Galba obrussa*; g. *Carychium exiguum*; h. *Hawaiiia minuscula*; i. *Euglesa compressa*; j. *Euglesa casertana*. Scale bars = 2 mm.

Ecology: *E. compressa* is restricted to areas of permanent water, preferring shallow sandy bottoms and rooted vegetation (Czaja *et al.*, 2014b).

Current distribution: Canada and United States. In Mexico it can be found in Chihuahua, Coahuila and Tamaulipas (Burch, 1972; Bequaert and Miller, 1973).

Stratigraphic remarks: The stratigraphic distribution of *E. compressa* at the El Molino Mammoth site is limited to Late Pleistocene strata. The first Pleistocene record for this species in Mexico was reported by Czaja *et al.* (2014b) from paleolake Irritila, Coahuila, northern Mexico.

Referred material: UJMC 601, 44 specimens.

Measurements: Length: 4 mm; diameter: 5 mm.

Gastropoda Cuvier, 1795

Family Lymnaeidae Rafinesque, 1815

Genus *Galba* Schranck, 1803

Galba humilis (Say, 1822)

(Figure 3b)

Ecology: This semi-aquatic species prefers muddy areas along the edges of creeks, lakes, ponds, and swamps (Clarke, 1981; Stewart and Dillon, 2004).

Current distribution: North America. Mexican states of Chihuahua, Coahuila, Hidalgo, Nuevo León, Sonora and Tamaulipas (Mazzotti, 1956; Landeros *et al.*, 1981; Thompson, 1999; Thorp and Rogers, 2016).

Stratigraphic remarks: At the El Molino

Mammoth site, the stratigraphic distribution of *G. humilis* is limited to the Holocene A strata. *Galba humilis* is widely known with its synonymous name, *Lymnaea humilis*.

Referred material: UJMC 602, 14 specimens.

Measurements: Height: 8.5 mm; aperture length: 4 mm.

Galba obrussa (Say, 1825)
(Figure 3f)

Ecology: *Galba obrussa* is common along the edges of small bodies of water such as streams and ponds, preferring areas along river banks or under protected spots (Baker, 1928; Miller, 1966).

Current distribution: From the Atlantic to the Pacific. From Canada to the southern state of Arizona and into northern Mexico, including the states of Durango and Coahuila (Baker, 1928; Mazzotti, 1955; Naranjo-Garcia, 2010).

Stratigraphic remarks: The stratigraphic distribution of *G. obrussa* at the El Molino Mammoth site ranged from the Late Pleistocene up to the Holocene D strata. This species was formerly known and described as *Fossaria obrussa*. This is the first Pleistocene record of this species in Coahuila, previously known only from a Late Pleistocene locality within the Valsequillo basin of central Mexico (Stevens *et al.*, 2012).

Referred material: UJMC 603-604, 11 specimens (Late Pleistocene: 7; Holocene: 4).

Measurements: Height: 4 mm; aperture length: 1.90 mm.

Family Physidae Fitzinger, 1833
Genus *Physella* Haldeman, 1842
Physella acuta (Draparnaud, 1805)
(Figure 3d)

Ecology: *Physella acuta* populations can be found in ponds, reservoirs, and the margins of rivers and streams, especially in disturbed sediments or

disturbed areas rich in organic material environments (Wethington *et al.*, 2009).

Current distribution: Cosmopolitan, can be found across six continents due to their high expansiveness by natural processes and accidental dispersal mediated by humans during the transport of exotic plants to Europe. In Mexico, the species is reported from Coahuila, Durango, Puebla, Aguascalientes, Veracruz and Michoacán (Arroyo-Cabrales *et al.*, 2008; Wethington *et al.*, 2009; Vinarski, 2017; Spyra *et al.*, 2019; Czaja *et al.*, 2020).

Stratigraphic remarks: The stratigraphic distribution of *P. acuta* at the El Molino Mammoth site is limited to the A strata (Holocene). Holocene and Pleistocene records for this species are known from Coahuila and Mexico City (Arroyo-Cabrales *et al.*, 2008; Czaja *et al.*, 2014a, 2014b).

Referred material: UJMC 605, 14 specimens.

Measurements: Height: 5.1 mm; aperture length: 3.85 mm.

Family Planorbidae Rafinesque, 1815
Genus *Gyraulus* Charpentier, 1837
Gyraulus parvus (Say, 1817)
(Figure 3e)

Ecology: *Gyraulus parvus* is usually associated with abundant vegetation in shallow water (Tuthill *et al.*, 1964).

Current distribution: North America. The species distribution in Mexico includes Durango, Morelos, Puebla and Sonora (Arroyo-Cabrales *et al.*, 2008; Thompson, 2011; Czaja *et al.*, 2020).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution of *G. parvus* is limited to Late Pleistocene strata. This is the first record of *G. parvus* for the Pleistocene of Coahuila. This species has also been reported from other Late Pleistocene localities of central Mexico (Arroyo-Cabrales *et al.*, 2008) and southern Mexico (Guerrero-Arenas *et al.*, 2013).

Referred material: UJMC 606, 33 specimens.

Measurements: Diameter: 4.8 mm.

Genus *Ferrissia* Walker, 1903
Ferrissia californica (Rowell, 1863)
 (Figure 3c)

Ecology: *Ferrissia californica* can be found in small bodies of water, often capable of surviving dry conditions (Walther *et al.*, 2010).

Current distribution: Members of the gastropod genus *Ferrissia* have a near-cosmopolitan distribution (Walther *et al.*, 2010).

Stratigraphic remarks: The stratigraphic distribution of *F. californica* at the El Molino Mammoth site is limited to Late Pleistocene strata. *F. californica* has also been found in Pleistocene and Holocene localities within the state of Coahuila (Czaja *et al.*, 2014 a, b).

Referred material: UJMC 607, 5 specimens.

Measurements: Height: 1.2 mm; length: 2.8 mm.

Family Ellobiidae Pfeiffer, 1854
 Genus *Carychium* Müller, 1773
Carychium exiguum (Say, 1822)
 (Figure 3g)

Ecology: *Carychium exiguum* inhabits humid environments such as swampy areas, is strongly hygrophilous and usually found under fallen leaves rocks and logs not far from water (Branson, 1961).

Current distribution: Canada, Colorado, New Mexico and Alabama, and Nuevo León in Mexico (Hubricht, 1985; Contreras-Arquieta, 1995).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution of *C. exiguum* is limited to Late Pleistocene strata. This is the first Pleistocene record of this species from Coahuila. Arroyo-Cabrales *et al.* (2008) recorded *C. exiguum* from a Late Pleistocene locality of San Luis Potosí.

Referred material: UJMC 608, +100 specimens.

Measurements: Height: 2 mm.

Family Succineidae Beck, 1837
 Genus *Succinea* Draparnaud, 1801
Succinea sp.
 (Figure 4c)

Ecology: Members of the Family Succineidae can be found in disturbed areas but prefers freshwater habitats with humid conditions among rocks, logs or leaf litter (Forsyth, 2005).

Current distribution: The Succineidae Family in Mexico is found distributed in Baja California, Central Mexico, Tamaulipas and Veracruz (Naranjo-García and Fahy, 2010).

Stratigraphic remarks: Late Pleistocene and the Holocene A, B1 and D strata. Late Pleistocene records in Mexico include Villa Acuña, Coahuila and Rancho La Amapola, San Luis Potosí (Arroyo-Cabrales *et al.*, 2008).

Referred material: UJMC 609-610, 62 specimens (Late Pleistocene: 18; Holocene: 44).

Measurements: Height: 5.5 mm; aperture length: 3 mm.

Family Cochlicopidae Pilsbry, 1900
 Genus *Cochlicopa* Férussac, 1821
Cochlicopa lubrica (O.F. Müller, 1774)
 (Figure 4a)

Ecology: *Cochlicopa lubrica* is considered a woodland mollusk and is usually found in forested montane habitats among damp under leaves (Judd, 1963; Kolb *et al.*, 1975; Metcalf and Smartt, 1997).

Current distribution: The genus *Cochlicopa* contains a single Mexican species, *C. lubrica*, which inhabits northwest Chihuahua, southern Nuevo León and Durango (Pilsbry, 1953; Bequaert and Miller, 1973; Contreras-Arquieta, 1995; Correa-Sandoval, 2003; Naranjo-García and Fahy, 2010).

Stratigraphic remarks: The stratigraphic distribution of *C. lubrica* at the El Molino Mammoth site is limited to Late Pleistocene strata. Fossil




Figure 4 a. *Cochlicopa lubrica*; b. *Glyphyalinia indentata*; c. *Succinea* sp.; d. *Gastrocopta cristata*; e. *Zonitoides arboreus*; f. *Habroconus* sp.; g. *Pupilla hebes*; h. *Gastrocopta tappaniana*; i. *Vallonia gracilicosta*. Scale bars = 2 mm.

records include sites in Coahuila and the Grava Valsequillo Formation of Puebla (Arroyo-Cabrales *et al.*, 2008).

Referred material: UJMC 611, 12 specimens.

Measurements: Height: 6.5 mm; aperture length: 2.1 mm.

Family Gastrocoptidae Pilsbry, 1918

Genus *Gastrocopta* Wollaston, 1878

Gastrocopta cristata (Pilsbry and Vanatta, 1900) (Figure 4d)

Ecology: *Gastrocopta cristata* is found in wooded slopes near streams and under fallen logs, as well as in grass where moisture conditions are favorable and stable (Leonard, 1950).

Current distribution: Kansas, Oklahoma and Texas to western New Mexico and Arizona. In Mexico it is reported from Sonora (Leonard, 1950; Bequaert and Miller, 1973).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution of *G. cristata* includes the entire Holocene strata. This species has only been recorded in Late Pleistocene

deposits from Rancho La Amapola, San Luis Potosí (Arroyo-Cabrales *et al.*, 2008).

Referred material: UJMC 612, +100 specimens.

Measurements: Height: 3 mm.

Gastrocopta tappaniana (C.B. Adams, 1841)
(Figure 4h)

Ecology: *G. tappaniana* is a hydrophilic species observed in grasslands, lowland forests, wooded slopes and flood plains with poor drainage, usually among leaf litter and below logs and stones (Leonard, 1950; Branson, 1961; Nekola, 2004).

Current distribution: Arizona, New Mexico, Kansas, South Dakota in the United States and Alberta and Manitoba in Canada (Pilsbry, 1948). Not reported from Mexico.

Stratigraphic remarks: The stratigraphic distribution of *G. tappaniana* at the El Molino Mammoth site is limited to Late Pleistocene strata. This is the first fossil record of this species in Mexico.

Referred material: UJMC 613, 23 specimens.

Measurements: Height: 2.8 mm.

Family Pupillidae Turton, 1831

Genus *Pupilla* Fleming, 1828

Pupilla hebes (Ancey, 1881)
(Figure 4g)

Ecology: *Pupilla hebes* can be found in forested mountain habitats consisting of wet meadows (Metcalf and Smartt, 1997).

Current distribution: The species has been reported in the United States from various localities of the western mountains, except in California. In Mexico, it has been found in Chihuahua and Baja California (Metcalf and Smartt, 1997; Miller, 1981).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution of *P*

hebes is limited to Late Pleistocene strata. This is the first fossil record of this species in Mexico.

Referred material: UJMC 614, 9 specimens.

Measurements: Height: 3.1 mm.

Family Valloniidae Morse, 1864

Genus *Vallonia* Risso 1826

Vallonia gracilicosta Reinhardt, 1883
(Figure 4i)

Ecology: Common under bushes protected spots such as under litter, rocks and dead grass. It is not restricted to a wooded habitat and can tolerate dry conditions (Kolb *et al.*, 1975; Metcalf and Smartt, 1997).

Current distribution: Oklahoma, New Mexico and adjacent states. In Mexico, it is recorded only in Nuevo León (Metcalf and Smartt, 1997; Correa-Sandoval, 2003).

Stratigraphic remarks: The stratigraphic distribution of *V. gracilicosta* at the El Molino Mammoth site is limited to Late Pleistocene strata. Arroyo-Cabrales *et al.* (2008), recorded *V. gracilicosta* in two Pleistocene localities in Mexico, one near the municipality of Villa Acuna in Coahuila and the other at Rancho La Amapola, San Luis Potosí.

Referred material: UJMC 615, 62 specimens.

Measurements: Diameter: 2.5 mm.

Family Gastrodontidae Tryon, 1866

Genus *Zonitoides* Lehmann, 1862

Zonitoides arboreus (Say, 1817)
(Figure 4e)

Ecology: *Zonitoides arboreus* occurs in woodlands and is associated with trees as well as living under shaded areas such as bark, leaves and stones (Leonard, 1950; Metcalf and Smartt, 1997).

Current distribution: From Canada through the southern United States, Mexico and Central America. The distribution in Mexico includes Chihuahua,

Nuevo León, San Luis Potosí, Puebla, and Veracruz (Metcalf and Smartt, 1997; Naranjo-García and Fahy, 2010).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution of *Z. arboreus* includes both Late Pleistocene and Holocene strata. The first Pleistocene record for this species was made by Arroyo-Cabrales *et al.* (2008) near the municipality of Villa Acuna in Coahuila.

Referred material: UJMC 616-617, +100 specimens (Late Pleistocene: +100; Holocene: 14).

Measurements: Diameter: 5 mm.

Genus *Glyphyalinia* von Martens, 1892

Glyphyalinia indentata (Say, 1822)
(Figure 4b)

Ecology: This species can be usually found in leaf litter in forests, open meadows, and anthropogenic impacted habitats (Nekola, 2010).

Current distribution: Arizona, New Mexico, Texas, Mexico and Guatemala. In Mexico, it has been reported from Baja California, Durango, Jalisco, Michoacán, Mexico City, Morelos and Puebla (Thompson, 2011).

Stratigraphic remarks: The stratigraphic distribution of *G. indentata* at the El Molino Mammoth site includes Late Pleistocene and Holocene A strata. The first Pleistocene record for this species in Coahuila was made by Arroyo-Cabrales *et al.* (2008) near the municipality of Villa Acuna.

Referred material: UJMC 618-619, 53 specimens (Late Pleistocene: 40; Holocene: 13). **Measurements:** Diameter: 3.5 mm.

Family Euconulidae Baker, 1928

Genus *Habroconus* Cross & P. Fischer, 1872

Habroconus sp.
(Figure 4f)

Ecology: Members of the genus *Habroconus* can be found in wooded and forested habitats, under

rocks, leaf litter, shrubs and trees (Baker, 1930; Veitenheimer-Mendes and Aguiar-Nunes, 2001; Veitenheimer-Mendes and Postal, 2003).

Current distribution: *Habroconus* specimens in Mexico can be found in the states of Jalisco, Quintana Roo, Mexico, Michoacán, Nuevo Leon, Puebla, San Luis Potosí, Sonora, Tamaulipas, Veracruz and Yucatan (Pilsbry, 1919b; Baker, 1930; Thompson, 1967b; Correa-Sandoval, 1997; Araiza y Naranjo-García, 2013; Naranjo-García and Fahy, 2010; Van Devender *et al.*, 2012)

Stratigraphic remarks: The stratigraphic distribution of *Habroconus* sp. at the El Molino Mammoth site is limited to Late Pleistocene strata. This is the first fossil record of this species in Mexico.

Referred material: UJMC 620, 35 specimens.

Measurements: Diameter: 2.8 mm.

Family Pristilomatidae Cockerell,
1891

Genus *Hawaiia* Gude, 1911

Hawaiia minuscula (A. Binney, 1841)
(Figure 3h)

Ecology: *Hawaiia minuscula* is an inhabitant of humid environments, living on leaf mold, beneath the bark of trees and among mosses. It is capable of withstanding long periods of drought and high temperatures (Leonard, 1950).

Current distribution: Canada and United States. In Mexico, the species has been reported from Baja California, Sonora, Tamaulipas, San Luis Potosí, Veracruz, Puebla, Nayarit and Yucatán (Bequaert and Miller, 1973; Metcalf and Smartt, 1997).

Stratigraphic remarks: At the El Molino Mammoth site, the stratigraphic distribution includes the Holocene A, B2, C and D strata. Arroyo-Cabrales *et al.* (2008) reported *H. minuscula* from the Late Pleistocene locality of Rancho La Amapola near the city of San Luis Potosí.

Referred material: UJMC 621, +100 specimens.

Measurements: Diameter: 2 mm.

Table 2. The habitat requirements of the El Molino Mammoth Site Late Pleistocene and Holocene malacofauna (modified from Miller, 1966).

Species	Habitat requirements
<i>C. exiguum</i> <i>G. tappaniana</i> <i>D. laeve</i>	Hygrophilic: moist situations in shaded areas not far from water.
<i>E. compressa</i>	Perennial water: stream or lake with slow to moderate current and shallow spots not affected by seasonal drying.
<i>G. parvus</i>	Shallow quiet water: small body of water with no current or areas of rooted vegetation with little current, both not subjected to significant seasonal drying.
<i>Habroconus</i> sp. <i>C. lubrica</i> <i>Z. arboreus</i> <i>G. indentata</i> <i>G. obrussa</i>	Woodland: moist areas under leaf litter, down timber, among tall marsh grass.
<i>V. gracilicosta</i> <i>P. hebes</i> <i>G. cristata</i> <i>H. minuscula</i>	Sheltered situations: these species are not restricted to a woodland habitat and can tolerate drier conditions.
<i>F. californica</i> <i>E. casertana</i> <i>P. acuta</i> <i>G. humilis</i> <i>Succinea</i> sp.	Shallow quiet water: small body of water that may become dry during part of the year. From relatively dry to humid conditions, preferring the later.

Family Agriolimacidae Wagner, 1935

Genus *Deroceras* Rafinesque, 1820

Deroceras laeve (O.F. Müller, 1774)

(Figure 3 a)

Ecology: *Deroceras laeve* has a circumpolar distribution and is considered a hygrophilous species. It can be found at low elevations among cultivated, urban and marshy areas. In mountains, it occurs along springs, streams and other bodies of water (Bequaert and Miller, 1973; Metcalf and Smartt, 1997; Rowson *et al.*, 2014).

Current distribution: Originally, this species was found native to North America but can be found today on other continents as well. In Mex-

ico, it has been recorded in Mexico City, Puebla and Veracruz (Bequaert and Miller, 1973; Metcalf and Smartt, 1997).

Stratigraphic remarks: The stratigraphic distribution of *D. laeve* at the El Molino Mammoth site includes the Late Pleistocene and Holocene A and C strata. The first Late Pleistocene record for *D. laeve* was reported from Rancho La Amapola near San Luis Potosí by Arroyo-Cabrales *et al.* (2008).

Referred material: UJMC 622-623, 17 specimens (Late Pleistocene: 11; Holocene: 6).

Measurements: Height: 4 mm; width: 2 mm.

4. Discussion

4.1. NEW MOLLUSCAN RECORDS

From the 14 mollusk taxa reported for the Late Pleistocene of the El Molino Mammoth site, all but three species were found distributed around Mexico and the United States Quaternary deposits. For the first time, we report the species *Gastrocopta tappaniana*, *Pupilla hebes* and *Habroconus* sp. for the Pleistocene of Mexico, of which the first two were previously known from Pleistocene deposits within the United States. Today, these species are extinct in the area, with *G. tappaniana* being mostly distributed outside of Mexico, usually from western to north eastern United States. In the case of *P. hebes*, recent distribution includes only

the northern Mexican states of Chihuahua and Baja California, whereas specimens of the genus *Habroconus* can be found from the northern state of Nuevo Leon to the southern state of Yucatan. For the state of Coahuila, new Pleistocene records include *G. tappaniana*, *G. obrussa*, *C. exiguum*, *G. parvus*, *P. hebes*, *D. laeve* and *Habroconus* sp. The remaining species (*E. compressa*, *F. californica*, *Succinea* sp., *C. lubrica*, *V. gracilicosta*, *Z. arboreus* and *G. indentata*) have previously been reported from other Pleistocene sites around the state.

Of the 10 Holocene mollusk taxa reported from the El Molino Mammoth site, three species have been previously recorded from Holocene fossil localities in Mexico. These include *G. obrussa* and *P. acuta* from Coahuila (Czaja *et al.*, 2019) and *Succinea* sp. from Sonora (Copeland, 2011). The




Figure 5 Generalized sediment sequence and malacofauna distribution from the abandoned well at the El Molino Mammoth site.

remaining seven species, *E. casertana*, *G. humilis*, *G. cristata*, *G. indentata*, *Z. arboreus*, *H. minuscula*, and *D. laeve*, are new fossil records for the Holocene of Mexico.

4.2. PALEOECOLOGICAL INTERPRETATIONS

Most of the El Molino Mammoth site mollusk taxa were also present during the early Quaternary in various sites of North America, making possible detailed paleoenvironmental interpretations through the available data associated with their environmental preferences and requirements (Carobene *et al.*, 2018). Generally, species composition and abundance of certain taxa in each assemblage depend on climatic conditions and local ecological factors, especially vegetation cover (Sümegei and Krolopp, 2002; Carobene *et al.*, 2018).

The large amount of terrestrial mollusks from the Late Pleistocene strata of the El Molino Mammoth site is noteworthy (Table 2). Miller (1966) associated the same species (*C. lubrica*, *Z. arboreus* and *G. indentata*) with those from the Great Plains to be woodland associated and commonly found in forested areas among shallow bodies of water.

Additionally, the finding of numerous fragmental remains of fossil wood among the Late Pleistocene sediments confirm the presence of arboreous vegetation at the site. In addition, the area must have also supported grassland patches due to the numerous shells found of *V. gracilicosta*, a species commonly recorded from humid permanent bodies of water with grassland vegetation (McMullen and Zakrzewski, 1972). These habitat requirements well confirm the previous environmental interpretation given by Miller *et al.* (2008) for the site based on the vertebrate fauna. The fossil remains of horses, camels and other mammals found infer woodland and grassland habitats, thus supporting a community of species representing localized woodland interrupted by grassland habitats near a permanent body of water at the site. In the Late Pleistocene and Early Holocene, the climatic conditions of the deserts in northern Mexico were much cooler than at the present (Metcalf *et al.*, 2000). Most of the Late Pleistocene malacofauna of the El Molino Mammoth site did not survive the rapid environmental changes during the Pleistocene-Holocene transition. Notable in the El Molino Mammoth site profile is the




Figure 6 Comparison of the habitat association percentage between Late Pleistocene and Holocene mollusks from the El Molino Mammoth site.

sudden appearance of numerous species typically associated with grasslands such as, *G. cristata* and *H. minuscula*, and the simultaneous disappearance of wetland species (Figure 5). Both presumably reflect the striking change of climate conditions during and after the transition.

The early mid-Holocene was probably warmer and wetter than today, and true desert conditions did not set in until about 4000 yr B.P. (Metcalf *et al.*, 2000). It is likely that the B horizon with its gypsum and calcium carbonate crystals represent these desert condition with dry climates and periods of evaporation and drought, and would likely explain the rapid colonization of the xerophytic elements in the profile that overall represent 60% of the malacofauna during this period (Figure 6). During the final phase of the body of water, there was an increase in the aquatic species *P. acuta* and *G. humilis*, accompanied by the aridity tolerant freshwater bivalve *E. casertana*. Agenbroad and Mead (1994) point to the abundance of *Physella* as indicative of warm waters. On the other hand, *G. humilis* is an excellent colonizer of new habitats and known for invading ponds devoid of vegetation and other species of gastropods (Jokinen, 2005). The ten species of the malacofauna that survived the transition into the Holocene are known for their tolerance of high temperatures (Miller, 1966). During this time, the site consisted of an ephemeral pond with predominant vegetation consisting mainly of short grasses and shrubs.

5. Conclusion

Our findings expand the mollusk data record known for Mexico with the entry of the three new Late Pleistocene gastropods, *Gastrocopta tappaniana*, *Pupilla hebes* and *Habroconus* sp. as well as seven new Holocene mollusk fossils, *E. casertana*, *G. humilis*, *G. cristata*, *G. indentata*, *Z. arboreus*, *H. minuscula*, and *D. laeve*, from the El Molino Mammoth site of Northern Mexico. The Late Pleistocene and Holocene molluscan fauna from the site provide a broader view of the paleoenvironmental changes

that occurred in northern Mexico during this time period. The recorded species confirm the presence of a humid forested and grassland habitat capable of supporting the hygrophilic and hydrophilic malacofauna along with the fossil remains of horses, camels and other mammals. Many mollusk species were not able to survive the strong shift to the more xeric conditions after and during the Pleistocene-Holocene transition. The surviving freshwater mollusks relied on the ephemeral body of water until the long periods of drought resulted in the total evaporation of water from the El Molino Mammoth site.

Contributions of authors

The authors of this research are: Perla Guadalupe Butrón Xancopinca (PGBX), Alexander Czaja(AC), Martha Carolina Aguillón (MCA), Rosario Gómez Núñez (RGN), Ignacio Vallejo González (IVG) and José Luis Estrada Rodríguez (JLER). In addition, its specific contribution is: 1.Conceptualization: PGBX and AC; 2.Data acquisition: PGBX and AC; 3.Methodologic/technical development: PGBX and AC; 4.Writing of the original manuscript: PGBX and AC; 5.Writing of the corrected and edited manuscript: PGBX, AC, RGN, MCA, IVG and JLER; 6.Graphic design: PGBX; 7.Fieldwork: PGBX, AC, RGN, MCA, IVG and JLER; 8.Interpretation: PGBX and AC; 9.Financing: Universidad Juárez del Estado de Durando, Campus Gómez Palacio, Mexico.

Financing

Universidad Juárez del Estado de Durando, Campus Gómez Palacio, Mexico.

Acknowledgements

We thank Jorge Banda, owner of Rancho Buena Fe, for allowing us to access the abandoned well

site. A special thanks to the Paleontologists Rosario Gómez Núñez and Martha Carolina Aguillón as well as Ignacio Vallejo González for their support during the visit to the Laboratory of Paleontology of the Museo del Desierto in Saltillo, Coahuila and for generously providing the Late Pleistocene material utilized in this study. We express our gratitude to Dr. Wade E. Miller and Dr. Jim I. Mead for their comments and suggestions towards the improvement of this manuscript. Transport funding was provided by the authorities of the Universidad Juárez del Estado de Durango.

Conflicts of interest

The authors declare that there is no conflict of interest with other authors, institutions or other third parties about the content of this article.

References

- Agenbroad, L.D., Mead, J.I., 1994, The Hot Springs Mammoth Site: a decade of field and laboratory research in paleontology, geology, and paleoecology: The Mammoth Site of Hot Springs, South Dakota, 457 p. <https://doi.org/10.1017/s0016756897276466>
- Araiza, V., Naranjo-García, E., 2013, Lista sistemática de la malacofauna terrestre del municipio de Atoyac, Veracruz: *Revista Mexicana de Biodiversidad*, 84, 765-773. <https://doi.org/10.7550/rmb.33106>
- Arroyo-Cabrales, J., Carreño, A.L., Lozano-García, S., Montellano-Ballesteros, M., 2008, La diversidad en el pasado, in *Capital natural de México*, 1, Conocimiento actual de la biodiversidad: Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México, 227-262.
- Baker, F.C., 1928, The fresh-water Mollusca of Wisconsin, Part I Gastropoda: *Bulletin of the Wisconsin Geological and Natural History Survey*, University of Wisconsin, 70(2), 507 p.
- Baker, H.B., 1930, Mexican mollusks collected for Dr. Bryant Walker in 1926, Part II Auriculidae, Orthurethra, Heterurethra, and Aulacopoda: *Occasional Papers of the Museum of Zoology, University of Michigan*, 220, 1-45.
- Bequaert, J.C., Miller, W.B., 1973, The mollusks of the arid Southwest with an Arizona Checklist: The University of Arizona Press, Tucson, Arizona, 271 p. <https://doi.org/10.1086/407860>
- Branson, B.A., 1961, The Recent Gastropoda of Oklahoma, III. Terrestrial Species: Pupillidae, Carychiidae, Strobilopsidae and Oligyridae: *Proceedings Oklahoma Academy of Science*, 41, 45-69.
- Burch, J.B., 1972, Biota of Freshwater Ecosystems, Identification Manual Number 3: Freshwater Sphaeriacean clams (Mollusca: Pelecypoda) of North America: U.S. Environmental Protection Agency, 31 p.
- Carobene, D., Harzhauser, M., Mandic, O., Gatto, R., 2018, Taxonomy and palaeoecology of continental Gastropoda (Mollusca) from the Late Pleistocene mammoth-bearing site of bullendorf in NE Austria: *Rivista Italiana di Paleontologia e Stratigrafia*, 3, 509-534. <https://doi.org/10.13130/2039-4942/10616>
- Catto, N., Bachhuber, F.W., 2000, Aeolian geomorphic response to climate change: an example from the Estancia Valley, central New Mexico, in McLaren, S., Kniveton, D. (eds.), *Linking Climate Change to Land Surface Change*, *Advances in Global Change Research*: New York, Springer, 6, 171-192. https://doi.org/10.1007/0-306-48086-7_9
- Clarke, A.H., 1981, The Freshwater Molluscs of Canada: Ottawa, Canada, National Museum of Natural Sciences, National Museums of Canada, 446 p. <https://doi.org/10.5962/bhl.title.132376>
- Contreras-Arquieta, A., 1995, Capítulo 10 Lista malacofaunística preliminar del Estado de Nuevo León, México, in Contreras-Balderas,

- S., González-Saldívar, F., Lazcano-Villarreal, D., Contreras-Arquieta, A. (eds.), *Listado Preliminar de la fauna silvestre del estado de Nuevo León: Monterrey, México*, Consejo Consultivo Estatal para la Preservación y Fomento de la Flora y Fauna Silvestre de Nuevo León, Gobierno del Estado de Nuevo León, 139-149.
- Copeland, A.E., 2011, Stratigraphy and geochronology of La Playa archeological site, Sonora: Tucson, Arizona, The University of Arizona Mexico, Bachelor's thesis, 29 p.
- Correa-Sandoval, A., 1997, Caracoles terrestres (Mollusca: Gastropoda) de Iturbide, Nuevo León, México: *Revista de Biología Tropical*, 44, 137-142.
- Correa-Sandoval, A., 2003, Gasterópodos terrestres del noreste de México: *Revista de Biología Tropical*, 51(3), 507-522.
- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., 2014A, Freshwater mollusks of the Valley of Sobaco, Coahuila, Northeastern Mexico – a sub fossil ecosystem similar to Cuatrociénegas: *Boletín de la Sociedad Geológica Mexicana*, 66(3), 459-469. <https://doi.org/10.18268/bsgm2014v66n3a4>
- Czaja, A., Palacios-Fest, M.R., Estrada-Rodríguez, J.L., Romero-Méndez, U., Alba-Ávila, J.A., 2104b, Inland dunes fauna and flora from Paleolake Irritila in the Comarca Lagunera, Coahuila, northern Mexico: *Boletín de la Sociedad Geológica Mexicana*, 66 (3), 541-551. <https://doi.org/10.18268/bsgm2014v66n3a9>
- Czaja, A., Covich, A.P., Estrada-Rodríguez, J.L., Romero-Méndez, U., Saenz-Mata, J., Meza-Sánchez, I.G., Ávila-Rodríguez, V., Becerra-López, J.L., Aguillón-Gutiérrez, D.R., Castañeda-Gaytán, J.G., 2019, Fossil freshwater gastropods from northern Mexico – A case of a “silent” local extirpation, with the description of a new species: *Boletín de la Sociedad Geológica Mexicana*, 71 (3), 609-629. <http://dx.doi.org/10.18268/BSGM2019v71n3a2>
- Czaja, A., Meza-Sánchez, I.G., Estrada-Rodríguez, J.L., Romero-Méndez, U., Sáenz-Mata, J., Ávila-Rodríguez, V., Becerra-López, J.L., Estrada-Arellano, J.R., Cardoza-Martínez, G.F., Aguillón-Gutiérrez, D.R., Cordero-Torres, D.G., Covich, A.P., 2020, The freshwater snails (Mollusca: Gastropoda) of Mexico: updated checklist, endemism hotspots, threats and conservation status: *Revista Mexicana de Biodiversidad*, 91, 1-22. <https://doi.org/10.22201/ib.20078706e.2020.91.2909>
- Forsyth, R.G., 2005, Terrestrial gastropods of the Peace River Northern Rockies Region of British Columbia: Victoria, Royal British Columbia Museum, 23 p.
- Guerrero-Arenas, R., Jiménez-Hidalgo, E., García-Barrera, P., 2013, New records of temperate mollusks in two Late Pleistocene terrestrial localities from northeastern Oaxaca, Southern Mexico: *Journal of South American Earth Sciences*, 47, 213-219. <https://doi.org/10.1016/j.jsames.2013.07.008>
- Herrington, H.B., 1962, A revision of the Sphaeriidae of North America (Mollusca: Pelecypoda): *Miscellaneous publications, Museum of Zoology, University of Michigan*, 118, 1-74. <https://doi.org/10.2307/2422763>
- Hubricht, L., 1985, The distributions of the native land mollusks of the eastern United States: *Fieldiana, Zoology*, 24, 1-191. <https://doi.org/10.5962/bhl.title.3329>
- Jokinen, E.H., 2005, Pond molluscs of Indiana Dunes National Lakeshore: then and now: *American Malacological Bulletin*, 20, 1-9.
- Judd, W.W., 1963, Molluscs and arthropods from Cardinal Nests at London, Ontario: *The Canadian Entomologist*, 95(3), 327-333. <https://doi.org/10.4039/ent95327-3>
- Kolb, K.K., Nelson, M.E., Zakrzewski, R.J., 1975, The Duck Creek Molluscan Fauna (Illinoian) from Ellis County, Kansas: *Transactions of the Kansas Academy of Science*, 78, 63-74. <https://doi.org/10.2307/3627225>

- Landeros, M. A., Velarde, F.I., Corona, J.L.E., Suazo, F.M., 1981, Determinación de algunos hospederos intermediarios de *Fasciola hepatica*, en la cuenca lechera de Tulancingo, Hgo: Revista Mexicana de Ciencias Pecuarias, 40, 47-51.
- Leonard, A.B., 1950, A Yarmouthian Molluscan Fauna in the Midcontinent Region of the United States: University of Kansas, Paleontological Contributions, Mollusca, 3, 48 p.
- Leonard, A.B., 1952, Illinoian and Wisconsinan molluscan faunas in Kansas: University of Kansas, Paleontological Contributions, Mollusca, 4, 38 p.
- Mazzotti, L., 1955, *Lymnaea obrussa* Say, huésped intermediario de *Fasciola hepática*: Revista del Instituto de Salubridad y Enfermedades Tropicales, 15(3), 163-5.
- Mazzotti, L., 1956, *Lymnaea humilis* (Say), huésped intermediario de *Fasciola hepática*: Revista del Instituto de Salubridad y Enfermedades Tropicales, 16(4), 21-23.
- Metcalf, A.L., Smartt, R.A., 1997, Land Snails of New Mexico: New Mexico Museum of Natural History and Science, Bulletin 10, 145 p.
- Metcalf, A.L., O'Hara, S.L., Caballero, M., Davies, S.J., 2000, Records of Late Pleistocene-Holocene climatic change in Mexico- A review: Quaternary Science Reviews, 19, 699-721. [https://doi.org/10.1016/S0277-3791\(99\)00022-0](https://doi.org/10.1016/S0277-3791(99)00022-0)
- McMullen, T.L., Zakrzewski, R.J., 1972, A new Late Pleistocene fauna from Northeastern New Mexico: New Mexico Geological Society Guidebook, 23, 134-136.
- Miller, B.B., 1966, Five Illinoian molluscan faunas from the southern Great Plains: Malacologia, 4(1), 173-260.
- Miller, W.B., 1981, *Helminthoglypta reederi* spec. nov. (Gastropoda: Pulmonata: Helminthoglyptidae), from Baja California, Mexico: Veliger, 24(1), 46-48.
- Miller, W.E., Delgado de Jesús, C.R., Gómez-Núñez, R., Vallejo-González, J.I., López-Espinosa, J., 2008, Preliminary report of Pleistocene Mammals from the state of Coahuila, Mexico: Natural History Museum of Los Angeles County, 41, 333-357.
- MolluscaBase, 2021, Data base MolluscaBase, accessed at: <http://www.molluscabase.org>
- Naranjo-García, E., Fahy, N.E., 2010, The Lesser Families of Mexican Terrestrial Molluscs: American Malacological Bulletin, 28, 59-80. <https://doi.org/10.4003/006.028.0222>
- Nekola, J.C., 2004, Terrestrial gastropod fauna of northeastern Wisconsin and the southern Upper Peninsula of Michigan: American Malacological Bulletin, 18, 21-44.
- Nekola, J.C., 2010, Acidophilic terrestrial gastropod communities of North America: Journal of Molluscan Studies, 76, 144-156. <https://doi.org/10.1093/mollus/eyp053>
- Nekola, J.C., Coles, B.F., 2010, Pupillid Land Snails of Eastern North America: American Malacological Bulletin, 28(2), 29-57. <https://doi.org/10.4003/006.028.0221>
- Pilsbry, H.A., 1919b, Mollusca from Central America and Mexico: Proceedings of the Academy of Natural Sciences of Philadelphia, 71, 212-223.
- Pilsbry, H.A., 1946, Monograph 3, Land Mollusca of North America (North of Mexico): Proceedings of the Academy of Natural Sciences of Philadelphia, 2(1), 1-520.
- Pilsbry, H.A., 1948, Monograph 3, Land Mollusca of North America (North of Mexico): Proceedings of the Academy of Natural Sciences of Philadelphia, 2(2), 521-1113.
- Pilsbry, H.A., 1953, Inland Mollusca of northern Mexico. II Urocoptidae, Pupillidae, Strobilopsidae, Vallonidae and Cionellidae: Proceedings of the Academy of Natural Sciences of Philadelphia, 105, 133-167.
- Rogers, D.C., 2016, Chapter 11 Phylum Mollusca, in Thorp, J.H., Rogers, D.C. (eds.), Thorp and Covich's Freshwater Invertebrates: Keys to Nearctic Fauna, Fourth edition, 2: San Diego, California, Academic Press, 189-221. <https://doi.org/10.1016/B978-0-12-385028-7.00011-1>

- Rowson, B., Anderson, R., Turner, J.A., Symondson, W.O.C., 2014, The slugs of Britain and Ireland: undetected and undescribed species increase a well-studied, economically important fauna by more than 20%. *PLOS One*, 9(3), 1-21. <https://doi.org/10.1371/journal.pone.0091907>
- Spyra, A., Cieplik, A., Strzelec, M., Babczyńska, A., 2019, Freshwater alien species *Physella acuta* (Draparnaud, 1805) - A possible model for bioaccumulation of heavy metals: *Ecotoxicology and Environmental Safety*, 185, 1-10. <https://doi.org/10.1016/j.ecoenv.2019.109703>
- Stevens, R.E., Metcalfe, S.E., Leng, M.J., Lamb, A.L., Sloane, H.J., Naranjo-García, E., González, S., 2012, Reconstruction of late Pleistocene climate in the Valsequillo Basin (Central Mexico) through isotopic analysis of terrestrial and freshwater snails: *Journal of Palaeogeography, Palaeoclimatology, Palaeoecology*, 319-320, 16-27. <https://doi.org/10.1016/j.palaeo.2011.12.012>
- Stewart, T.W., Dillon Jr., R.T., 2004, Species composition and geographic distribution of Virginia's freshwater gastropod fauna: a review using historical records: *American Malacological Bulletin*, 19, 79-91.
- Sümeği, P., Krolopp E., 2002, Quaternary malacological analysis for modeling of the Upper Weichselian palaeoenvironmental changes in the Carpathian Basin: *Quaternary International*, 9, 53-63. [https://doi.org/10.1016/s1040-6182\(01\)00102-1](https://doi.org/10.1016/s1040-6182(01)00102-1)
- Thompson, F.G., 1967b, The land and freshwater snails of Campeche: *Bulletin of Florida State Museum*, 11, 221-256.
- Thompson, F.G., 1999, An identification manual for the freshwater snails of Florida: *Walkerana*, 10, 1-96.
- Thompson, F.G., 2011, An annotated checklist and bibliography of the land and freshwater snails of Mexico and Central America: *Bulletin of the Florida State Museum*, 50, 1-299.
- Thorp, J.H., Rogers, D.C., 2016, Thorp and Covich's Freshwater Invertebrates, Fourth edition, Volume 2, Keys to Nearctic Fauna: Academic Press, Elsevier, 740 p.
- Tuthill, S.J., Clayton, L., Laird, W.M., 1964, A comparison of a fossil Pleistocene molluscan fauna from North Dakota with a recent molluscan fauna from Minnesota: *American Midland Naturalist*, 71, 344-362. <https://doi.org/10.2307/2423292>
- Van Devender, A.S., Van Devender, R.W., Rivera-Garcia, A., Jimenez-Maldonado, R.E., Van Devender, M.N., 2012, Report on the terrestrial mollusks of the Sierra de la Madre (Oposura), Sonora, Mexico – *The Caracoleros: American Malacological Bulletin*, 30(2), 315-322. <https://doi.org/10.4003/006.030.0211>
- Veitenheimer-Mendes, I.L., Aguiar-Nunes, J., 2001, Moluscos, in Mirapalhete, S.R. (Coord. E Org.), *Flora e Fauna do Parque Natural do Morro do Osso: Secretaria Municipal do Meio Ambiente, Porto Alegre, Brasil*, 48-57.
- Veitenheimer-Mendes, I.L., Postal, M., 2003, Moluscos terrestres e límnicos registrados para a sub-bacia hidrográfica do arroio Itapuã, Rio Grande do Sul, Brasil: *Revista Brasileira de Biociências*, 1(2), 55-69.
- Vinarski, M.V., 2017, The history of an invasion: phases of the explosive spread of the physid snail *Physella acuta* through Europe, Transcaucasia and Central Asia: *Biological Invasions*, 19(4), 1299-1314. <https://doi.org/10.1007/s10530-016-1339-3>
- Walther, A.C., Burch, J.B., Ó Foighil, D., 2010, Molecular Phylogenetic Revision of the Freshwater Limpet Genus *Ferrissia* (Planorbidae: Ancylinae) in North America Yields Two Species: *Ferrissia* (*Ferrissia*) *rivularis* and *Ferrissia* (*Kincaidilla*) *fragilis*: *Malacologia*, 53(1), 25-45. <https://doi.org/10.4002/040.053.0102>
- Wethington, A.R., Wise, J., Dillon Jr., R.T., 2009, Genetic and morphological characterization of the Physidae of South Carolina, with description of a new species: *Nautilus*, 123(4), 282-292.