

Abdominal aortic aneurysm characteristics and outcomes: a single-center retrospective cross-sectional study

Características y resultados de aneurismas aórticos abdominales: estudio transversal y retrospectivo de un solo centro

Gregorio Arellano-Gutiérrez^{1*}, Alan Y. Rodríguez-Andrade², and Iván E. Murillo-Barrios¹

¹Department of Vascular Surgery and Endovascular Therapy, Hospital de Cardiología, Unidad Médica de Alta Especialidad No. 34, Instituto Mexicano del Seguro Social, Monterrey, Nuevo León; ²Department of Surgery, Hospital Regional de Alta Especialidad de Ciudad Victoria "Bicentenario 2010", Ciudad Victoria, Tamaulipas. Mexico

Abstract

Objective: The objective of the study is to evaluate the characteristics and outcomes of patients with abdominal aortic aneurysm (AAA) and its correlation with mortality in the first 30 days after the procedure was performed. **Methods:** Demographic information, clinical and radiological characteristics, as well as outcomes 30 days after the procedure was performed were assessed and compared. Continuous variables were analyzed with Student's t-test and categorical with Chi-square and Fisher's exact test. **Results:** Duration of the procedure ($p = 0.001$), blood loss ($p < 0.001$), age > 75 years ($p = 0.027$), aneurysm size > 65 mm ($p = 0.01$), open surgery ($p = 0.001$), presence of pain ($p = 0.005$), chronic kidney disease ($p = 0.03$), and rupture of the aneurysm ($p < 0.001$) were the factors significantly associated with mortality. **Conclusion:** It is essential that patient characteristics and comorbidities are assessed, as well as factors that may affect the outcomes to predict the prognosis in patients with AAA. At present, no mortality predictive model is universally applicable and highly variable performance across different populations might need a model that adapts to the population of interest.

Keywords: Aortic abdominal aneurysm. Characteristics. Outcomes. Mexico.

Resumen

Objetivo: Evaluar las características y resultados de los pacientes con aneurisma de aorta abdominal y su correlación con la mortalidad en los primeros 30 días después de realizado el procedimiento. **Métodos:** Se evaluó y comparó la información demográfica, las características clínicas y radiológicas, así como los resultados a los 30 días de realizado el procedimiento. Las variables continuas se analizan con la prueba de t de Student y las categóricas con Chi-cuadrado y la prueba exacta de Fisher. **Resultados:** La duración del procedimiento ($p = 0.001$), pérdida de sangre ($p < 0.001$), edad > 75 años ($p = 0.027$), tamaño del aneurisma > 65 mm ($p = 0.01$), cirugía abierta ($p = 0.001$), presencia de dolor ($p = 0.005$), enfermedad renal crónica ($p = 0.03$) y rotura del aneurisma ($p < 0.001$) fueron los factores asociados significativamente a la mortalidad. **Conclusión:** Es fundamental evaluar las características de los pacientes y las comorbilidades, así como los factores que pueden afectar los resultados para predecir el pronóstico en pacientes con aneurisma de aorta abdominal. En la actualidad, ningún modelo predictivo de mortalidad es universalmente aplicable y la alta variabilidad de resultados entre diferentes poblaciones podría necesitar un modelo que se adapte a la población de interés.

Palabras clave: Aneurisma de la aorta abdominal. Características. Resultados. México

*Correspondence:

Gregorio Arellano-Gutiérrez
E-mail: gr.arellano.dr@gmail.com

Date of reception: 18-09-2022

Date of acceptance: 02-03-2023

DOI: 10.24875/CIRU.22000474

Cir Cir. 2023;91(6):730-735

Contents available at PubMed

www.cirugiaycirujanos.com

0009-7411/© 2023 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Abdominal aortic aneurysm (AAA) describes a segmental, full-thickness dilatation of the abdominal aorta exceeding the normal vessel diameter by 50%, although an aneurysm diameter of 3.0 cm is commonly regarded as the threshold¹. The prevalence rates of AAA range from 1.9% to 18.5% in males and 0% to 4.2% in females². Frequent risk factors associated with AAA are age, smoking, gender, chronic obstructive pulmonary disease, chronic kidney disease, hypertension, family history, and coronary artery disease³. They are usually asymptomatic unless complications occur. Patients at the highest risk of aortic rupture are those with larger diameter AAAs. Growth rates increase markedly with aneurysm diameter; for each 0.5 cm increase in AAA diameter, growth rates increased by 0.5 mm/year and rupture rates double^{4,5}. The estimated annual rupture risk according to diameter is 0%, 0-5%, 13-15%, 10-20%, 20-40%, and 30-50% in cases with diameter less than 4 cm, 4-5 cm, 5-6 cm, 6-7 cm, 7-8, and > 8, respectively⁶. This catastrophic event is associated with a mortality of 50–80% and even with emergency surgical repair, mortality is about 40% to 50%^{7,8}; the strongest predictor of in-hospital mortality has been found to be age⁹. There are currently open and endovascular techniques whose objective is to isolate the aneurysm from the circulation, eliminating the risk of rupture. Early complications after the procedure are common. They include endoleaks (persistent blood flow into the aneurysm sac after graft placement), access site complications (arterial rupture and dissection), atheroembolism, lower limbs ischemia, renal failure, and endograft migration.

Recent studies have found a significantly decrease in AAA rupture mortality rates, mainly due to improvements in public health awareness, screening, and management¹⁰⁻¹². In view of the high morbidity and mortality rates, the importance of early diagnosis and treatment to decrease the public health burden, this study is aimed to evaluate the characteristics and outcome of patients with AAAs and its correlation with mortality in the first 30 days after the procedure was performed. To the best of our knowledge, this is the first analysis of the characteristics and outcomes of AAA including Mexican population.

Materials and methods

Study design and setting

This is a retrospective, cross-sectional study conducted at a national vascular surgery referral hospital

in Mexico. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. For this type of study, formal consent is not required. This retrospective study was approved by the institutional ethics review board and a waiver of consent was obtained.

Study population

From January 2016 to December 2021, patients with AAA presenting to the emergency department or in-hospital vascular surgery consultation were studied. For this study, patients with confirmed AAA were enrolled using census sampling method. Patients with an incomplete medical profile and those discharged against medical advice were excluded.

Data collection

Data were collected by reviewing medical records of the medical center. Pre-designed checklists containing medical history, demographic data (age, gender), chief complaint, characteristics of aneurysm (location, size), family history, presence or absence of leakage based on computed tomography scan findings, and treatment approach, as well as outcomes (mortality) (Fig. 1).

Statistical analysis

Data analysis was done using the Statistical Package for the Social Sciences version 25.0 statistical software. Findings were presented as mean \pm standard deviation or frequency (%) for numerical and categorical variables, respectively. The utilized tests were chi-square, independent sample t-test, and Fisher's exact test. $p < 0.05$ was considered statistically significant.

Results

Characteristics of patients

During COVID-19 pandemic, the admission of patients reduced significantly, leaving 56 patients for analysis. Table 1 and Figure 2 show the baseline characteristics of studied patients. The mean age was 71.93 ± 10.21 (22-86). 80.4% (45) were male and 19.6% (11) were female. The most frequent chief




Figure 1. A: CT reconstruction of the abdomen and pelvis shows a 7.1 cm AAA. **B:** sagittal CT with intravascular contrast shows an AAA with mural thrombus on both the anterior (white arrowheads) and posterior (black arrowhead) margin of the aorta.

complaint was pain (51.8%), whereas abdominal pain being the most frequent location (37.5%), followed by lumbar pain (12.5%) and thoracic pain (8.9%). Limb ischemia presented in 7.1% and shock state only in 3.6%. The main duration of pain was 1.65 ± 4.33 months and mean aneurysmal size was $65.42 \text{ mm} \pm 15.26$. Location of the aneurysm was found infrarenal in 89.3%, juxtarenal in 5.4%, and abdomino-thoracic aorta in 5.4%. Synchronous aneurysm presented on the right common iliac artery (14.3%), left common iliac artery (3.6%), and both iliac arteries (10.7%).

Outcomes

Elective procedure was performed in 80.4% (45). Open surgery was performed in 32.1% (18). Of all mortality in open surgery, ruptured aneurysm represented 80% (8). Endovascular procedure was performed in 67.9% (38) with a mortality of 10.5% (4). Ruptured aneurysms represented 19.6% (11) of cases, in which death was seen in 81.8% (9). 25% (14) patients did not survive 30 days after the procedure was performed, in which ruptured aneurysms represented 64.3% (9). Duration of the procedure ($p = 0.001$), blood loss ($p < 0.001$), age > 75 years ($p = 0.027$), aneurysm size $> 65 \text{ mm}$ ($p = 0.01$), open surgery ($p = 0.001$), presence of pain ($p = 0.005$), chronic kidney disease ($p = 0.03$), and rupture of the aneurysm ($p < 0.001$) were among the factors significantly associated with mortality (Table 2).

Table 1. Baseline characteristics of patients

Variables	Frequency, n (%)
Gender	
Male	45 (80.4)
Female	11 (19.6)
Pain location	
Abdominal	21 (37.5)
Lumbar	7 (12.5)
Thoracic	5 (8.9)
Shock state	2 (3.6)
Limb ischemia	4 (7.1)
Medical history	
Smoking	50 (89.3)
Hypertension	43 (76.8)
Ischemic heart disease	13 (23.2)
Diabetes mellitus	12 (21.4)
Chronic kidney disease	9 (16.1)
Dyslipidemia	6 (10.3)
Chronic obstructive pulmonary disease	5 (8.9)
Cerebrovascular accident	3 (5.4)
Iliac artery synchronous aneurysm	
Right common iliac artery	8 (14.3)
Bilateral common iliac artery	6 (10.7)
Left common iliac artery	2 (3.6)
Family history	1 (1.8)
Aneurysm location	
Infrarenal	50 (89.3)
Juxtarenal	3 (5.4)
Abdomino-thoracic	3 (5.4)

Discussion

In this study, we show the characteristics of patients with AAA, the outcomes after the procedure (open or endovascular), and its correlation with mortality in the first 30 days after the procedure was performed (Figs. 3 and 4). According to previous data, smoking is the main risk factor correlated to AAA³. Other risk factors include age, male gender, high blood pressure, coronary artery disease, family history of AAA, high cholesterol, lower extremity peripheral arterial disease, history of a cerebrovascular event, and obesity. Diabetes has been found as a negative association with AAA¹³. This correlates with our study, in which we observed smoking as the most frequent associated factor, followed by hypertension and coronary artery disease. More recent studies have found a positive correlation between chronic obstructive pulmonary disease, chronic kidney disease, and greater incidence of AAA^{14,15}.

The incidence of concomitant common iliac artery aneurysm has been reported from 12% to 40%, and




Figure 2. Comorbidities of studied patients.

CAD: coronary artery disease; DM: type 2 diabetes mellitus; CKD: chronic kidney disease; DLP: dyslipidemia, COPD: chronic obstructive pulmonary disease.

Table 2. Correlation between baseline characteristics, outcomes, and mortality

Variables	Survived, n (%)	Died, n (%)	p value	OR (CI 95%)
Age > 75	13 (59.1)	9 (40.9)	0.02	4.01 (1.1-14.3)
Aneurysm size > 65 mm	11 (55)	9 (45)	0.01	5 (1.3-18.4)
Open surgery	8 (44.4)	10 (55.6)	0.001	10.6 (0.6-42.7)
Female gender	7 (63.6)	4 (36.4)	0.43	0.5 (0.1-2)
Pain	17 (21.8)	12 (41.4)	0.005	8.8 (1.7-44)
Ruptured aneurysm	2 (18.2)	9 (81.8)	< 0.001	36 (5.9-216)
Cerebrovascular accident	1 (33.3)	2 (66.6)	0.15	6.8 (0.5-82)
Hypertension	31 (72.1)	12 (27.9)	0.48	2.1 (0.4-11)
Smoking	38 (76)	12 (24)	0.63	0.6 (0.1-3)
Diabetes mellitus	12 (100)	0 (0)	0.02	0.6 (0.5-0.8)
Chronic kidney disease	4 (44.4)	5 (55.6)	0.03	5.2 (1.1-23)
Ischemic heart disease	11 (84.6)	2 (15.4)	0.48	0.4 (0.09-2.4)
Chronic obstructive pulmonary disease	2 (40)	3 (60)	0.09	5.4 (0.8-36)
Length of stay (days)				
Mean ± DE	13.1 ± 6.9	10.2 ± 10.7	0.24	
Duration of procedure (min)				
Mean ± DE	233.9 ± 17.2	378.5 ± 173.6	0.001	
Blood loss (ml)				
Mean ± DE	467.6 ± 595	2971 ± 2885	< 0.001	




Figure 3. A: diagnostic angiography demonstrating the presence of an infrarenal AAA. B: computed tomography 3D volume rendering after EVAR showing no stent-graft migration, limb occlusion, or endoleak.




Figure 4. Abdominal aortic aneurysm repair with the placement of a dacron graft. Note the aneurysm sac surrounding the graft.

that right sided is more common than left sided¹⁶⁻¹⁸. We report an incidence of 28.6%, also with right sided (14.6%) being more frequent than left (3.6%).

Factors significantly correlated with mortality 30 days after the procedure was performed were the duration of the procedure, blood loss, age > 75 years, aneurysm size > 65 mm, open surgery, presence of pain, chronic kidney disease, and rupture of the aneurysm at presentation. Several predictor models and risk factors for mortality after AAA repair in different clinical situations (elective or urgent; open or endovascular) have been described with contrasting and similar results, but the most consistent data associated with mortality are procedure performed, chronic obstructive pulmonary disease, cerebrovascular disease, renal insufficiency, aneurysm size, increasing age, female sex, serum creatinine level, cardiac disease, previous aortic surgery or stent, abnormal white cell count, and abnormal serum sodium level. Lijftogt et al.¹⁹. conducted a systematic review for mortality risk prediction models for AAA surgery and at present, no predictive model is universally applicable due to the lack of external validation, and they also demonstrated highly variable performance across different populations.

Randomized controlled trials²⁰⁻²² of AAA have shown marked benefits of EVAR with respect to 30-day mortality compared to open repair. Although aneurysm rupture is the most significant complication, aneurysm-related death is a relatively minor determinant of long-term survival in all three randomized trials²³, systematic reviews,²⁴ and retrospective studies^{25,26}; atherosclerotic cardiovascular, end-stage renal disease, and cancer-related death represent a greater threat.

Considering that survival is influenced by these factors, this raises the issue of screening for AAA to ease early elective intervention at a younger age, ideally before the onset of comorbidity and therefore, higher risk of death. Evidence shows that screening reduces aneurysm-related mortality in certain populations, with the strongest recommendation for men aged 65-75 years²⁷. Screening patients below 65 years old are unlikely to be economically viable (due to a low prevalence of disease), although expanded criteria for screening have been recently proposed²⁸. It is therefore imperative that aside from perioperative details, patient characteristics and comorbidities are assessed, as well as factors that may affect the outcome to predict the prognosis in patients with AAA. These data might prove to be of clinical interest and relevance for the assessment of prognostic variables for short and long-term survival.

This study has some limitations. First, it is retrospective and therefore prone to inherent bias with such studies. This was a small retrospective study with only 56 patients, and its statistical power may be limited.

Second, the follow-up period was lost in some patients as to permit an investigation of the long-term clinical outcomes of overall survival. It is also important to note that several types of graft were utilized; however, this reflects real-world clinical practice.

Acknowledgments

The authors would like to acknowledge the vascular surgery team.

Funding

The authors declare that no funding was received.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors obtained approval from the Ethics Committee for analysis and publication of routinely acquired clinical data and informed consent was not required for this retrospective observational study.

References

- Kent KC. Clinical practice. Abdominal aortic aneurysms. *N Engl J Med.* 2014;371:2101-8.
- Ullery BW, Hallett RL, Fleischmann D. Epidemiology and contemporary management of abdominal aortic aneurysms. *Abdom Radiol (NY).* 2018;43:1032-43.
- Altobelli E, Rapacchietta L, Profeta VF, Fagnano R. Risk factors for abdominal aortic aneurysm in population-based studies: a systematic review and meta-analysis. *Int J Environ Res Public Health.* 2018;15:2805.
- Thompson SG, Brown LC, Sweeting MJ, Bown M, Kim L, Glover M, et al. Systematic review and meta-analysis of the growth and rupture rates of small abdominal aortic aneurysms: implications for surveillance intervals and their cost-effectiveness. *Health Technol Assess.* 2013;17:1-118.
- Golledge J. Abdominal aortic aneurysm: update on pathogenesis and medical treatments. *Nat Rev Cardiol.* 2019;16:225-42.
- Brewster DC, Cronenwett JL, Hallett JW Jr., Johnston KW, Krupski WC, Matsumura JS, et al. Guidelines for the treatment of abdominal aortic aneurysms. Report of a subcommittee of the Joint Council of the American Association for Vascular Surgery and Society for Vascular Surgery. *J Vasc Surg.* 2003;37:1106-17.
- Kuivaniemi H, Ryer EJ, Elmoro JR, Tromp G. Understanding the pathogenesis of abdominal aortic aneurysms. *Expert Rev Cardiovasc Ther.* 2015;13:975-87.
- Filardo G, Powell JT, Martinez MA, Ballard DJ. Surgery for small asymptomatic abdominal aortic aneurysms. *Cochrane Database Syst Rev.* 2015;2015:CD001835.
- Visser P, Akkersdijk GJ, Blankensteijn JD. In-hospital operative mortality of ruptured abdominal aortic aneurysm: a population-based analysis of 5593 patients in The Netherlands over a 10-year period. *Eur J Vasc Endovasc Surg.* 2005;30:359-64.
- Bartek MA, Kessler LG, Talbott JM, Nguyen J, Shalhoub S. Washington State abdominal aortic aneurysm-related mortality shows a steady decline between 1996 and 2016. *J Vasc Surg.* 2019;70:1115-22.
- Takagi H, Ando T, Umemoto T, ALICE (All-Literature Investigation of Cardiovascular Evidence) Group. Abdominal aortic aneurysm screening reduces all-cause mortality: make screening great again. *Angiology.* 2018;69:205-11.
- Reimerink JJ, van der Laan MJ, Koelemay MJ, Balm R, Legemate D. Systematic review and meta-analysis of population-based mortality from ruptured abdominal aortic aneurysm. *Br J Surg.* 2013;100:1405-13.
- Kent KC, Zwolak RM, Egorova NN, Riles TS, Manganaro A, Moskowitz AJ, et al. Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals. *J Vasc Surg.* 2010;52:539-48.
- Matsushita K, Kwak L, Ballew SH, Grams ME, Selvin E, Folsom AR, et al. Chronic kidney disease measures and the risk of abdominal aortic aneurysm. *Atherosclerosis.* 2018;279:107-13.
- Takagi H, Umemoto T, ALICE (All-Literature Investigation of Cardiovascular Evidence) Group. A meta-analysis of the association of chronic obstructive pulmonary disease with abdominal aortic aneurysm presence. *Ann Vasc Surg.* 2016;34:84-94.
- Wallinder J, Georgiou A, Wanhainen A, Björck M. Prevalence of synchronous and metachronous aneurysms in women with abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg.* 2018;56:435-40.
- Armon MP, Wenham PW, Whitaker SC, Gregson RH, Hopkinson BR. Common iliac artery aneurysms in patients with abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg.* 1998;15:255-7.
- Bannazadeh M, Jenkins C, Forsyth A, Kramer J, Aggarwal A, Somerset AE. Outcomes for concomitant common iliac artery aneurysms after endovascular abdominal aortic aneurysm repair. *J Vasc Surg.* 2017;66:1390-7.
- Lijftogt N, Luijnenburg TW, Vahl AC, Wilschut ED, Leijdekkers VJ, Fiocco MF, et al. Systematic review of mortality risk prediction models in the era of endovascular abdominal aortic aneurysm surgery. *Br J Surg.* 2017;104:964-76.
- Greenhalgh RM, Brown LC, Kwong GP, Powell JT, Thompson SG, EVAR Trial Participants. Comparison of endovascular aneurysm repair with open repair in patients with abdominal aortic aneurysm (EVAR trial 1), 30-day operative mortality results: randomised controlled trial. *Lancet.* 2004;364:843-8.
- Prinssen M, Verhoeven EL, Buth J, Cuypers PW, van Sambeek MR, Balm R, et al. A randomized trial comparing conventional and endovascular repair of abdominal aortic aneurysms. *N Engl J Med.* 2004;351:1607-18.
- Lederle FA, Freischlag JA, Kyriakides TC, Padberg FT Jr., Matsumura JS, Kohler TR, et al. Outcomes following endovascular vs open repair of abdominal aortic aneurysm: a randomized trial. *JAMA.* 2009;302:1535-42.
- Patel R, Sweeting MJ, Powell JT, Greenhalgh RM, EVAR Trial Investigators. Endovascular versus open repair of abdominal aortic aneurysm in 15-years' follow-up of the UK endovascular aneurysm repair trial 1 (EVAR trial 1): a randomised controlled trial. *Lancet.* 2016;388:2366-74.
- Khushram M, Williman JA, Hider PN, Jones GT, Roake JA. Systematic review and meta-analysis of factors influencing survival following abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg.* 2016;51:203-15.
- Healy GM, Redmond CE, Gray S, Iacob L, Sheehan S, Dowdall JF, et al. Midterm analysis of survival and cause of death following endovascular abdominal aortic aneurysm repair. *Vasc Endovascular Surg.* 2017;51:274-81.
- Bath MF, Gokani VJ, Sidloff DA, Jones LR, Choke E, Sayers RD, et al. Systematic review of cardiovascular disease and cardiovascular death in patients with a small abdominal aortic aneurysm. *Br J Surg.* 2015;102:866-72.
- US Preventive Services Task Force, Owens DK, Davidson KW, Krist AH, Barry MJ, Cabana M, et al. Screening for abdominal aortic aneurysm: US preventive services task force recommendation statement. *JAMA.* 2019;322:2211-8.
- Carnevale ML, Koleilat I, Lipsitz EC, Friedmann P, Indes JE. Extended screening guidelines for the diagnosis of abdominal aortic aneurysm. *J Vasc Surg.* 2020;72:1917-26.