

## Impact of powerful volcanic eruptions and solar activity on the climate above the Arctic Circle

Elena A. Kasatkina\*, Oleg I. Shumilov, Mauri Timonen and Alexander G. Kanatjev

Received: April 26, 2017; accepted: October 09, 2017; published on line: January 01, 2018

### Resumen

En este trabajo, se evalúa la respuesta del crecimiento de los árboles en la región polar (Península de Kola y la Laponia de Finlandia) a las erupciones volcánicas más potentes ( $VEI > 4$ ) durante un periodo de 1445-2005. El análisis se basó en dos cronologías de anillos de árboles de elevada resolución: Loparskaya (1445-2005) y Finnish ( $\sim 7500$  años). Estas cronologías se desarrollaron a partir de muestras de *Pinus sylvestris* L. recogidas cerca de la línea de árboles del norte en la estación de Loparskaya ( $68.63^\circ$  N,  $33.25^\circ$  E) y la Laponia finlandesa ( $68-70^\circ$  N;  $20-30^\circ$  E), respectivamente. El análisis de sobreposición de época indicó una disminución significativa en el crecimiento de los anillos de los árboles polares durante 7 años después de las erupciones. Asimismo, el estudio de los anillos muestra que el nivel de crecimiento de los árboles en la región polar se ve afectado por las más poderosas erupciones volcánicas producidas en latitudes bajas. En cambio, los volcanes islandeses no tienen un impacto significativo en el clima de la Península de Kola y la Laponia finlandesa. La aplicación del análisis wavelet para la cronología del anillo de árboles de Loparskaya permitió identificar la existencia de los ciclos principales de actividad solar (11, 20-25 y  $\sim 100$  años). Además el análisis wavelet reveló una fuerte coherencia en las bandas 8-13 y 20-30 años indicando un posible vínculo entre la actividad solar y el cambio climático a escala regional.

Palabras clave: erupciones volcánicas, datos de los anillos de los árboles, cambio climático, actividad solar.

### Abstract

The tree growth response in polar region (Kola Peninsula and Finnish Lapland) to the most powerful ( $VEI > 4$ ) volcanic eruptions is assessed over a period of 1445-2005. The analysis was based on the Loparskaya (1445-2005) and Finnish supra-long ( $\sim 7500$  years) tree-ring chronologies. These chronologies were developed from *Pinus sylvestris* L. samples collected near the northern tree line at Loparskaya station ( $68.63^\circ$  N,  $33.25^\circ$  E) and Finnish Lapland ( $68-70^\circ$  N;  $20-30^\circ$  E), respectively. A superposed epoch analysis indicated a significant decrease in polar tree-ring growth over 7 years after the eruptions with subsequent recovery to its normal level. The level of tree growth in polar region is affected by most powerful low-latitude volcanic eruptions. By contrast, high-latitude Icelandic volcanoes have no significant impact on the climate of Kola Peninsula and Finnish Lapland. The application of wavelet analysis for the Loparskaya tree ring chronology allowed to identify the existence of the main cycles of solar activity (11, 20-25, and  $\sim 100$  years). Moreover, the wavelet analysis revealed a strong coherence in the 8-13 and 20-30 year bands, indicating a possible link between solar activity and climate change on a regional scale.

Key words: volcanic eruptions, tree-ring data, climate change, solar activity.

---

E.A. Kasatkina\*  
O.I. Shumilov  
A.G. Kanatjev  
Polar Geophysical Institute  
Fersmana str. 14, 184209 Apatity  
Murmansk Oblast, Russia  
\*Corresponding author: oleg@aprec.ru

M. Timonen  
Natural resources Institute Finland (LUKE)  
Rovaniemi, Finland

## Introduction

It is well known that, throughout the Holocene prior the industrial period, global climate changes took place under the influence of only natural factors including solar activity and powerful volcanic eruptions (Shumilov *et al.*, 2000; Shindell *et al.*, 2003; Soon and Baliunas, 2003). These factors strongly affect climate and atmosphere of the Earth (Crowley, 2000; Shumilov *et al.*, 2000; Shindell *et al.*, 2003; Soon and Baliunas, 2003; Velasco and Mendoza, 2008), but which of them (and some others as well) plays a dominant role is currently under discussion. However, it should be noted that considerable contribution to climatic change in recent decades is due to man-made factors (Vecchi and Soden, 2007). Volcanic eruptions emit huge amounts of volcanic dust, sulfur dioxide and water vapor into the atmosphere reducing the atmospheric transparency and preventing the penetration of solar radiation to the earth's surface. All these processes result in breaking the radiation balance of the atmosphere and lead to reduction of the surface temperature in most cases. Sulfate aerosols that enter the stratosphere after powerful volcanic eruptions may exist there for several years (Robock, 2000). Recent powerful volcanic eruptions such as Tambora (1815), Krakatau (1883), Santa Maria (1902), Katmai (1912), Agung (1963), El Chichon (1982), and Pinatubo (1991), caused a cooling in the Northern Hemisphere by 0.2 – 0.8° C for several years (Rampino and Self, 1982; Robock, 2000; Wigley, 2000; Zanchettin *et al.*, 2013). For example, the 1815 Tambora eruption produced the “year without a summer” in 1815 (Rampino and Self, 1982; Robock, 2000). However, volcanic eruptions cause not always and not everywhere cooling events. The climatic response to volcanic eruptions varies regionally and depends on the amount, chemical composition and height of volcanic emissions. Some studies documented that volcanic activity can lead to regional warming and even droughts, as happened, for example, after the 1783 Laki eruption in Iceland, when most of the European territory was covered by an unusually dry fog (Thordarson and Self, 2003; Ogle *et al.*, 2005), and after the 1991 Pinatubo eruption in Indonesia (McCormick *et al.*, 1995; Robock, 2000).

The period of instrumental meteorological observations does not exceed 100 – 120 years and covers only a small number of powerful volcanic eruptions. Therefore, the climatic effects of volcanic activity were studied using several proxy records (the content of sulfuric acidity in ice cores and tree-ring data) (Hammer

*et al.*, 1980; Scuderi, 1990; Briffa *et al.*, 1998; Gervais and MacDonald, 2001; Vaganov and Shiyatov, 2005; Shumilov *et al.*, 2000; 2011; Kasatkina *et al.*, 2013). The possible response of tree-ring growth and even the formation of so-called “frost rings” (Vaganov and Shiyatov, 2005) are associated with a decrease of temperature after major volcanic eruptions. The emission of a large amount of volcanic gases into the atmosphere leads to acid precipitation, which can as well cause deep depression in tree growth (Thordarson and Self, 2003; Ogle *et al.*, 2005). Moreover, the reduction of atmospheric transparency can affect the process of photosynthesis (Scuderi, 1990; Ogle *et al.*, 2005).

It should be noted that the temperature decreases are more considerable when the volcanic eruptions occur during the minima of solar activity. Widely known is the period of global cooling during the Maunder minimum of solar activity (1645 – 1715 AD) which occurred in the Little Ice Age. The main solar factors influencing climate and atmosphere are solar radiation (Lean *et al.*, 1995; Haigh, 1996), cosmic rays and, possibly, cosmic dust influencing the cloud cover of the atmosphere (Shumilov *et al.*, 1996; Svensmark and Friis-Christensen, 1997; Kasatkina and Shumilov, 2005; Kasatkina *et al.*, 2007). Climatic impacts are most clearly expressed in ring width of trees growing in extreme conditions close to the tree line at sites of their biological surviving (Fritts, 1976; Grace *et al.*, 2002). The northern (or Arctic) tree line is the northernmost latitude where trees can grow. The Arctic tree-ring chronologies are of most interest for paleoclimatic studies because the trees growing near the northern tree-line seem to be highly sensitive to the influence of such external factors as temperature and solar radiation.

The aim of this study is to verify a hypothesis concerning the possible regional climatic response to volcanic and solar activity impacts during the last millennium by analyzing polar tree-ring chronologies from Kola Peninsula and Finnish Lapland.

## Geographic setting

The tree-ring data used in this study were collected near the northern tree-line at Loparskaya station, Kola Peninsula (68.63 N, 33.25 E) (Figure 1). The climate of the Kola Peninsula unlike the other polar regions of Russia is influenced by warm air masses of the North Atlantic and cold air of Arctic regions (the Barents Sea, the Kara Sea, and the Taymir Peninsula). The Kola Peninsula involves

three climatic zones: sea coast, central area, and mountain section. The climate of the northern sea coast including the sampling site is influenced by the Barents Sea. The mean temperature is in the range from  $-6$  to  $-12^{\circ}\text{C}$  in the coldest month (February) and from  $12$  to  $13^{\circ}\text{C}$  in the warmest month (July). The annual mean temperature (precipitation) in the region ranges from  $0^{\circ}\text{C}$  ( $600$ - $700$  mm) on the coasts of the Barents and White Seas to  $-2^{\circ}\text{C}$  ( $500$ - $600$  mm) in the central part of the Kola Peninsula. The mountain area is characterized by a cooler summer, a relatively mild winter, and a lot of precipitation with an average value of  $1000$  mm.


The Kola Peninsula is covered by taiga in the south and tundra in the north. The dominant tree species at or near the tree-line include pine (*Pinus sylvestris* L.), spruce (*Picea obovata* Ledeb.), and birch (*Betula pubescens* ssp. *szerepanovii*).

## Methods

A total of 36 timber cores of *Pinus sylvestris* L. were sampled at the northern tree-line, including the oldest living pine with more than 560 years of age. Ring widths were measured with a precision of  $0.01$  mm by using an image analysis system (scanner and relevant software) (Kanatjev *et al.*, 2014). Using these samples, the Loparskaya tree-ring chronology from living trees covering the period from 1445 to 2005 was developed. The data were

processed using modern methods adopted in dendrochronology (cross-dating and standardization) with the help of COFECHA and ARSTAN programs (Holmes, 1983; Cook and Kairiukstis, 1990). In this study we also used the Finnish supra-long ( $\sim 7500$  years) tree-ring chronology collected at Northern Lapland ( $68$ - $70^{\circ}\text{N}$ ;  $20$ - $30^{\circ}\text{E}$ ) (Helama *et al.*, 2008).

Superposed epoch analysis of both chronologies was performed to assess the tree growth response on the most powerful ( $\text{VEI} > 4$ , Volcanic Explosivity Index) volcanic eruptions (<http://www.volcano.si.edu/world/largeeruptions.cfm>). The VEI evaluates volcanic eruptions based on the height of the eruption column and solid material ejected (Newhall and Self, 1982). The most powerful volcanic eruptions can cause lowering of temperature and tree growth for five years and more (Rampino and Self, 1982; Scuderi, 1990; Robock, 2000; Wigley, 2000; Gervais and MacDonald, 2001; D'Arrigo *et al.*, 2013; Zanchettin *et al.*, 2013). In the present study, the analysis was based only on those events that are at least 7 years apart. As a result, a total of 19 events are considered covering the period from 1445 to 2005 (see the Table 1). Analysis of individual events (Huaynaputina in 1600, Laki in 1783 and Tambora in 1815) was also performed. Yearly averages of sunspot numbers were obtained from the National Geophysical Data Center (<ftp://ftp.ngdc.noaa.gov>) for the period of 1700 – 2005.



**Figure 1.** Location of the sampled site (cross) in relation to the northern pine tree line.

To identify the main periodicities and their evolution in time-frequency mode, we applied the continuous wavelet transform (CWT) of a time series  $X$  (Torrence and Compo, 1998):

$$Wx = \frac{1}{\sqrt{s}} \int_{-\infty}^{\infty} X(t)\psi\left(\frac{t-b}{s}\right) dt \quad (1),$$

where  $t$  is the time,  $s$  is the wavelet scale,  $b$  is the shift of the analysis window,  $\psi$  is the base wavelet function. We used here the Morlet function that may be defined as a plane wave modulated by a Gaussian function (Torrence and Compo, 1998):

$$\psi(t) = \pi^{-1/4} e^{i\omega_0 t/s} e^{-t^2/(2s)^2} \quad (2),$$

where  $\omega_0$  is the dimensionless frequency (we use here  $\omega_0=6$ , since it provides a good balance between time and frequency localization). In order to avoid the edge effects, Torrence and Compo (1998) introduced the cone of influence (COI) in which the wavelet power caused by a discontinuity at the edge has dropped by a factor  $e^{-2}$ .

To examine the relationship in time-frequency scale between tree-ring growth and solar activity, the cross-wavelet transform was applied to two time series (Grinsted *et al.*, 2004):

$$W_{LN}(s) = W_L(s)W_N^*(s) \quad (3),$$

where (\*) denotes the complex conjugate,  $L$  and  $N$  indexes denote tree-ring and sunspot number time series respectively. The phase angle of  $W_{LN}$  describes the phase relationship between  $L$  and  $N$  in time frequency space. We standardized (zero mean, unit standard deviation) both time series before analysis. The wavelet coherence (WTC) was used to assess a localized correlation coefficient in time frequency space between tree ring data and sunspot numbers (Grinsted *et al.*, 2004):


$$R^2(s) = \frac{|S(s^{-1}W_{LN}(s))|^2}{S(s^{-1}|W_L(s)|^2) \cdot S(s^{-1}|W_N(s)|^2)} \quad (4),$$

where  $S$  denotes smoothing in both time and scale. The statistical significance levels of CWTs and WTC were estimated using Monte Carlo method against a red noise model (Torrence and Compo, 1998; Grinsted *et al.*, 2004).

## Results and Discussion

In this study, we analyzed climatic impacts of solar and volcanic activity with help of both (Loparskaya and Finnish supra-long) tree-ring chronologies developed on the basis of samples taken near the northern tree line (Figure 1).

Figure 2 shows tree-ring growth responses on the most powerful volcanic eruption of Huaynaputina (VEI=6) in February 1600 (see the Table). For comparison, the upper timberline Cirque Peak (California; 36.5° N, 118.2° E) foxtail pine tree-ring record (Scuderi, 1990) is also presented (see Figure 2c). The 1600 eruption of Huaynaputina in Peru was the largest volcanic eruption in South America spanning the past 2000 years (de Silva and Zielinski, 1998; Fei *et al.*, 2015; Stoffel *et al.*, 2015). Based on ice core and tree-ring records, this eruption injected a large amount of sulfur (up to 32 Mt) into the atmosphere, reducing solar



**Figure 2.** Tree-ring growth response to the volcanic eruption of Huaynaputina (VEI=6) in Peru (1600 AD): **(A)** the Loparskaya (68.63 N, 33.25 E) pine tree-ring chronology, **(B)** the Finnish supra-long (~7500 years) pine tree-ring chronology (68-70° N; 20-30° E), **(C)** the Cirque Peak (California; 36.5° N, 118.2° E) foxtail pine tree-ring chronology (Scuderi, 1990).

**Table 1.** Most powerful volcanic eruptions (VEI>4) since AD 1470.

Volcanoes		Eruption dates	Coordinates	VEI
Sakura-Jima, Japan		November 3, 1471	31.6 N, 130.7 E	5
Agua de Pau, Azores		June 28, 1563	37.8 N, 25.5 W	5
Huaynaputina, Peru	+	February 19, 1600	16.6 S, 70.9 W	6
Parker, Philippines	+	January 4, 1641	6.1 N, 124.9 E	5
Gamkonora, Indonesia		May 20, 1673	1.4 N, 127.5 E	5
Tongkoko, Indonesia	+	1680	1.5 N, 125.2 E	5
Fuji, Japan		December 16, 1707	35.4 N, 138.7 E	5
Katla, Iceland	+	May 11, 1721	63.5 N, 19 W	5
Shikotsu, Japan	+	August 19, 1739	42.7 N, 141.3 E	5
Katla, Iceland		October 17, 1755	63.5 N, 19 W	5
Laki, Iceland		June 8, 1783	64.4 N, 17.3 W	4+
St Helens, USA	+	January 15, 1800	46.2 N, 122.2 W	5
Tambora, Indonesia		April 10, 1815	8.3 S, 118.0 E	7
Galunggung, Indonesia		October 8, 1822	7.3 S, 108.1 E	5
Cosiguina, Nicaragua	+	January 20, 1835	13 N, 87.6 W	5
Askja, Iceland		March 29, 1875	65 N, 17 W	5
Azul, Chile	+	April 10, 1932	35.7 S, 70.8 W	5
Agung, Indonesia		March 17, 1963	8.3 S, 115.5 E	5
Pinatubo, Philippines	+	June 15, 1991	15.1 N, 120.4 E	6

By crosses indicated the major volcanic eruptions that affect the tree growth at the study area.

insolation and resulting in global temperature drop (Hammer *et al.*, 1980; Scuderi, 1990; de Silva and Zielinski, 1998). The eruption of Huaynaputina in 1600 AD had a global impact on human society and dramatically affected the weather and environment in Central Europe, southern Scandinavia, China and the North America (Scuderi, 1990; de Silva and Zielinski, 1998; Verosub and Lippman, 2008; Fei *et al.*, 2015; Stoffel *et al.*, 2015). In Russia, this event caused strong frosts in the summer of 1601 and crop failure which led to severe famine and unprecedented mortality (more than 500,000 people) in 1601 – 1603, social disruption and political unrest called the Time of Troubles (Verosub and Lippman, 2008). Figure 2 shows a significant (more than 25%) reduction in tree growth, which is observed almost simultaneously in all three sites in the year following the eruption. All three curves behave identically and reveal a significant ring width decrease more than ten years (see Figure 2).


The 1815 eruption of Tambora in Indonesia (VEI>7), which was the most powerful over the study period, ejected 80 Mt of sulfuric dioxide into the atmosphere and led to the temperature decrease ( $\sim 0.8^{\circ}$  C) in the

Northern Hemisphere (Sadler and Grattan., 1999; Ogle *et al.*, 2005; Stoffel *et al.*, 2015). In North America and Central Europe, 1816 was called the “year without a summer”. However, the climate in Scandinavia in that year was mild, and no significant temperature drops were observed (Sadler and Grattan., 1999; D’Arrigo *et al.*, 2013). From Figure 3, it is observed that the year 1816 shows no decrease in tree growth relative to the previous year. The decrease began much earlier (in 1808) and lasted until 1822, and again, all three curves are identical (see Figure 3). Some evidence (large acidity peak in ice core) indicate that a powerful climatically effective unknown volcanic eruption occurred in 1908/1909 (Scuderi, 1990; White *et al.*, 1997; D’Arrigo *et al.*, 2012; Guevara-Murua *et al.*, 2014; Wahl *et al.*, 2014). Two other powerful low-latitude volcanic eruptions in 1673 (Gamkonora) and in 1822 (Galunggung) also did not lead to the expected volcanic cooling (see Table 1). According to radiocarbon dating, a powerful volcanic eruption in 1660 ( $\pm 20$  y) in New Guinea may have preceded the eruption of Gamkonora (White *et al.*, 1997; Wahl *et al.*, 2014). In this case, the expected volcanic cooling possibly resulted from increased atmospheric opacity from low-


latitude eruptions in 1673 (Gamkonora), 1815 (Tambora), and 1822 (Galunggung) could be masked on the already existing background provided by the previous eruptions.

It should be noted that all these eruptions occurred during the minima of solar activity, which possibly intensified the climatic response.

Icelandic volcanoes are most closely located to the Kola Peninsula and Finnish Lapland. Therefore, it is more interesting to study the environmental and climatic impacts of their most powerful eruptions on this region. Figure 4 shows tree growth response to the eruption of Laki (VEI= 4+), which began in June 1783. The 1783 eruption of Laki in Iceland continued eight months, ejecting around 122 Mt of sulfur dioxide into the atmosphere and causing an unusually dry fog and drought in most of Europe (Thordarson and Self, 2003; Ogle *et al.*, 2005). According to our data, no deviation in the annual tree growth was observed in 1783 – 1785 (see Figure 4). From Figure 4b, it is seen that the Loparskaya tree-ring record reveals a significant drop (by 50% relative to the previous year) in 1786, three years after the eruption of Laki volcano, therefore it can



**Figure 3.** Tree-ring growth response to the volcanic eruption of Tambora (VEI=7) in Indonesia (1815): (A) the Loparskaya (68.63 N, 33.25 E) pine tree-ring chronology, (B) the Finnish supra-long (~ 7500 years) pine tree-ring chronology (68-70° N; 20-30° E), (C) the Circue Peak (California; 36.5° N, 118.2° E) foxtail pine tree-ring chronology (Scuderi, 1990).




**Figure 4.** Tree-ring growth response to the volcanic eruption of Laki (VEI=4+) in Iceland (1783): (A) the Loparskaya (68.63 N, 33.25 E) pine tree-ring chronology, (B) the Finnish supra-long (~ 7500 years) pine tree-ring chronology (68-70° N; 20-30° E).

hardly be associated with this event. This conclusion is also confirmed by the absence of any depression after 1783 in the annual tree growth according to the Finnish data (Figure 4a). Similar results demonstrating the absence of a decrease in the annual growth immediately after the Laki eruption were obtained in other studies (Sadler and Grattan, 1999; Thordarson and Self, 2003; Ogle *et al.*, 2005). The analysis of climatic response to other powerful eruptions of Icelandic volcanoes (see the Table 1) revealed no changes. As it concerned the Laki eruption, that question had been considered in details in connection with a meteorological situation during the event (Grattan and Pyatt, 1999; Thordarson and Self, 2003; Sonnek *et al.*, 2016). In accordance with these results, one can assume that a stable blocked high-pressure air pattern over Europe concentrated volcanic gases near the Earth's surface and prevented them to spread into the Kola Peninsula.


Figure 5 presents the composite response of tree growth in polar region obtained by the superposed epoch method for 19 volcanic eruptions listed in the Table 1. A significant suppression in tree growth is observed at both polar sites for seven years after the eruption with the subsequent recovery (see Figure 5). This result is consistent, in general, with the findings obtained earlier from dendroclimatic reconstructions of northern Eurasia (Vaganov and Shiyatov, 2005) and North America (Scuderi, 1990). Our analysis showed that the main contribution to the observed decrease in tree growth at both polar sites was made by low-latitude volcanic eruptions (see the Table 1).

As was noted above, global climatic changes occurred under the composite influence of two main natural factors: powerful volcanic eruptions and solar activity. The main solar factors acting on climate and atmosphere are solar radiation (Lean *et al.*, 1995; Haigh, 1996), cosmic rays and possibly, cosmic dust influencing the cloud cover of the atmosphere (Shumilov *et al.*, 1996; Svensmark and Friis-Christensen, 1997; Kasatkina and Shumilov, 2005; Kasatkina *et al.*, 2007). Volcanic factors act during powerful eruptions, while solar factors work continuously depending on cyclic intensity of solar activity. As can be seen from Fig. 6, the long variation in tree-ring indexes seems to demonstrate a certain similarity with the envelope of the 11-year maxima of sunspot numbers ( $r=0.31$ ,  $p<0.05$ ). The Sporer (1416 – 1534), Maunder (1645 – 1715), and Dalton (1801 – 1816) minima of solar activity are accompanied by cooling in the Kola Peninsula and tree growth depression. The significant tree growth depression and cooling from 1780 to 1830 coincide with the Dalton minimum of solar activity and most powerful volcanic eruption of Tambora in 1815 (see Figure 6).



**Figure 5.** Response of polar tree-ring growth to the most powerful (VEI>4) volcanic eruptions since AD 1470 (see the Table) obtained by the superposed epoch method: **(A)** the Loparskaya (68.63 N, 33.25 E) pine tree-ring chronology, **(B)** the Finnish supra-long (~ 7500 years) pine tree-ring chronology (68-70° N; 20-30° E). Standard errors are shown by vertical lines.

For identification of solar cycles in tree-ring records and understanding their evolution in time frequency scale, we applied the Morlet-based continuous wavelet transform (CWT) and the wavelet coherence analysis (WTC). Figure 7 shows the wavelet power spectrum for the (a) Loparskaya tree-ring chronology, the (b) yearly sunspot number, and the (c) WTC between the tree rings and sunspot number. WTC is used as a localized correlation coefficient between two time series in time frequency space (Grinsted *et al.*, 2004). Arrows indicate a phase relationship between the variables (see Figure 7c). The wavelet spectrum for the Loparskaya tree-ring chronology shows significant periodicity around 100 years (Gleissberg solar cycle), which is present all the time (see Figure 7a). The ~ 22 year solar cycle signal (Hale cycle) with less intensity intermittently appears between 1470-1620, 1680-1830, and 1920-1970 (see Figure 7a). The major ~ 11 year solar cycle signal (Schwabe cycle) is highly suppressed along the time interval, weakly appearing between 1500-1570, 1600-1750, and 1870-1900. The CWT for tree-ring data exhibits also a significant (or near the confidence level) power peak in the 2-7 year band, appearing intermittently along all the time interval. This periodicity corresponds to the second harmonic




**Figure 6.** **(A)** Time series of sunspot number (1700-2005) (thin line) and their 11-year maximum envelope (1500-2005) (thick line). The sunspot numbers since the 1500 AD have been reconstructed on the polar aurora data (Shove, 1955). **(B)** The Loparskaya tree-ring width standard chronology of *Pinus sylvestris* L. (1445-2005) (thin line) and its smoothed 23-year moving average. Most powerful volcanic eruptions (VEI>4) since AD 1470 are shown by vertical lines (see the Table).

(5.5 year) of the Schwabe cycle, and, possibly, is also related to the North Atlantic Oscillation impact (Velasco and Mendoza, 2008). Figure 7c shows strong coherences in the 8-13 year band in phase between 1715 and 1730, and in anti phase for the time intervals of 1770-1840 and 1880-1920 (see Fig. 7c). The wavelet coherence also shows a statistically significant high power in the 20-30 year band over a period of 1770-1850 and 1920-2005 with a non-stationary phase relationship, the  $\sim 100$  year is practically outside of the COI (see Fig. 7c).

In general, the wavelet coherence analysis between tree-ring data and sunspot numbers demonstrated a strong coherence in the 8-13 and 20-30 year bands indicating a possible link between solar activity and climate change on a regional scale. Earlier evidences of solar cycle signals were found in a lot of paleoclimatic proxy records from different sites (Baliunas *et al.*, 1997; Kasatkina *et al.*, 2007; Shumilov *et al.*, 2007; Rigozo *et al.*, 2002; Velasco and Mendoza, 2008; Wang and Zhang, 2011; Edvardsson *et al.*, 2016; Sunkara and Tivari, 2016).

## Conclusions

1. Our analysis of tree ring chronologies obtained for the Kola Peninsula and Finnish Lapland revealed the most significant (by 25% relative to the previous year) depression in tree growth after the 1600 eruption of Huaynaputina in Peru (VEI=6), which was the largest volcanic eruption in South America over the past 2000 years.
2. A superposed epoch analysis revealed a significant reduction in polar tree growth, which was observed for seven years after volcanic eruption with a subsequent recovery to its normal level.
3. The level of tree growth in polar region is affected by most powerful low-latitude volcanic eruptions. High-latitude Icelandic volcanoes have no significant impact on the climate of Kola Peninsula and Finnish Lapland.
4. The application of wavelet analysis for the Loparskaya tree ring chronology identified



**Figure 7.** The continuous wavelet power spectrum of the standardized tree-ring index **(A)** and sunspot number **(B)** time series; **(C)** the wavelet coherence between tree-ring indices and sunspot numbers. The relative phase relationship is shown as arrows (with in-phase pointing right, anti-phase pointing left). The 5% significance level against red noise is shown as a thick contour.

the existence of the main cycles of solar activity (11, 20-25, and  $\sim 100$  years). The wavelet coherence analysis revealed a strong coherence in the 8-13 and 20-30 year bands indicating a possible link between solar activity and climate change on a regional scale.

5. Our results demonstrate the possibility of using polar tree-ring widths as indicators of solar and volcanic activity above the Arctic Circle.

## References

- Baliunas S., Frick P., Sokoloff D., Soon W., 1997, Time scales and trends in the Central England temperature data (1659-1990): A wavelet analysis. *Geophys. Res. Lett.*, *24*, 1351-1354.
- Briffa K.R., Jones P.D., Schweingruber F.H., Osborn T.J., 1998, Influence of volcanic eruptions on Northern Hemisphere summer temperature over the past 600 years. *Nature*, *393*, 450-452.
- Cook E.R., Kairiukstis L., 1990, Methods of Dendrochronology. Dordrecht: Kluwer Academic Publishing, 211 pp.
- D'Arrigo R., Wilson R., Anchukaitis K.J., 2013, Volcanic cooling signal in tree ring temperature records for the past millennium. *J. Geophys. Res.*, *118*, 9000-9010.
- Edvardsson J., Adolphi F., Lindholm H.W., Corona C., Muscheler R., Stoffel M., 2016, Periodicities in mid- to late-Holocene peatland hydrology identified from Swedish and Lithuanian tree-ring data. *Quarter. Sci. Rev.*, 200-208.
- Fei J., Zhang D.D., Lee H.F., 2015, 1600 AD Huaynaputina eruption (Peru), abrupt cooling, and epidemics in China and Korea. *Adv. Meteorol.*, 2016, doi: 10.1155/2016/3217038.
- Fritts H.C., 1976, Tree rings and climate. London: Academic Press, 567 pp.
- Gervais B.R., MacDonald G.M., 2001, Tree-ring and summer temperature response to volcanic aerosol forcing at the northern tree-line, Kola Peninsula, Russia. *The Holocene*, *11*, 499-505.
- Grace J., Berninger F., Nagy L., 2002, Impacts of climate change on the tree line. *Annals of Botany*, *90*, 537-544.
- Grattan J.P., Pyatt F.B., 1999, Volcanic eruptions dry fogs and the European palaeoenvironmental record: localized phenomena or hemispheric impacts? *Global Planet. Change*, *21*, 173-179.
- Grinsted A., Moore J.C., Jevrejeva S., 2004, Application of the cross wavelet transform and wavelet coherence to geophysical time series. *Nonlin. Processes Geophys.*, *11*, 561-566.
- Guevara-Murua A., Williams C.A., Hendy E.J., Rust A.C., Cashman K.V., 2014, Observations of a stratospheric aerosol veil from a tropical volcanic eruption in December 1808: is this the Unknown  $\sim 1809$  eruption? *Clim. Past.*, *10*, 1707-1722.
- Haigh J.D., 1996, The impact of solar variability on climate. *Science*, *272*, 981-984.
- Hammer C.U., Clausen H.B., Dansgaard W., 1980, Greenland ice sheet evidence of post-glacial volcanism and its climatic impact. *Nature*, *288*, 230-235.
- Helama S., Mielikainen K., Timonen M., Eronen M., 2008, Finnish supra-long chronology extended to 5634 BC. *Norw. J. Geograph.*, *62*, 271-277.
- Holmes R.L., 1983, Computer-assisted quality control in tree-ring dating and measurement. *Tree-Ring Bulletin.*, *44*, 69-75.
- Kanatjev A.G., Shumilov O.I., Kasatkina E.A., 2014, Software for dendrochronological measurements. *Instruments and Experimental Techniques*, *57*, 214-217.
- Kasatkina E.A., Shumilov O.I., 2005, Cosmic ray-induced stratospheric aerosols: A possible connection to polar ozone depletions. *Ann. Geophys.*, *23*, 675-679.
- Kasatkina E.A., Shumilov O.I., Lukina N.V., Krapiec M., Jacoby G., 2007, Stardust component in tree rings. *Dendrochronologia*, *24*, 131-135.
- Kasatkina E.A., Shumilov O.I., Timonen M., Kanatjev A.G., 2013, Consequences of powerful volcanic eruptions according to dendrochronological data. *Izvestiya, Atmospheric and Oceanic Physics*, *49*, 432-438.
- Lean J., Beer J., Bradley R., 1995, Reconstruction of solar irradiance since 1610, Implications for climate change. *Geophys. Res. Lett.*, *22*, 3195-3198.

- McCormick M.P., Thomason, L.W., Trepte C.R., 1995, Atmospheric effects of the Mt Pinatubo eruption. *Nature*, 373, 399-404.
- Newhall C.G., Self S., 1982, The Volcanic Explosivity Index (VEI): An estimate of explosive magnitude for historical volcanism. *J. Geophys. Res.*, 87, 1231-1238.
- Ogle N., Turney C.S.M., Kalin R.M., O'Donnell L., Butler C.J., 2005, Palaeovolcanic forcing of short-term dendroisotopic depletion: The effect of decreased solar intensity on Irish oak. *Geophys. Res. Lett.*, 32. doi: 10.1029/2004GL021623.
- Rampino M.R., Self S., 1982, Historic eruptions of Tambora (1815), Krakatau (1883), and Agung (1963), their stratospheric aerosols and climatic impact. *Quater. Res.*, 18, 127-143.
- Rigozo N.R., Nordemann D.J.R., Echer E., Zanandera A., Gonzalez W.D., 2002, Solar variability effects studied by tree-ring data wavelet analysis. *Adv. Space Res.*, 29, 1985-1988.
- Robock A., 2000, Volcanic eruptions and climate. *Rev. Geophys.*, 38, 191-219.
- Sadler J.P., Grattan J.P., 1999, Volcanoes as agents of past environmental change. *Global Planet. Change.*, 21, 181-196.
- Schove D.J., 1955, The Suspot cycle, BD 649 to AD 2000. *J. Geophys. Res.*, 60, 127-136.
- Scuderi L.A., 1990, Tree-ring evidence for climatically effective volcanic eruptions. *Quater. Res.*, 34, 67-85.
- Shindell D.T., Schmidt G.A., Miller, R.L., Mann M.E., 2003, Volcanic and solar forcing of climate change during the Preindustrial Era. *J. Climate.*, 16, 4094-4107.
- Shumilov O.I., Kasatkina E.A., Henriksen K., Vashenyuk E.V., 1996, Enhancement of stratospheric aerosols after solar proton event. *Ann. Geophys.*, 14, 1119-1123.
- Shumilov O.I., Kasatkina E.A., Raspopov O.M., Turunen E., Jacoby G., 2000, An estimation of the climate response to the variations in solar and volcanic activity. *Geomagn. Aeron.*, 40, 687-691.
- Shumilov O.I., Kasatkina E.A., Lukina N.V., Kirtsideli I.Yu., Kanatjev A.G., 2007, Paleoclimatic potential of the northernmost juniper trees in Europe. *Dendrochronologia*, 24, 123-130.
- Shumilov O.I., Kasatkina E.A., Mielikainen K., Timonen M., Kanatjev A.G., 2011, Palaeovolcanos, Solar activity and tree-rings from Kola Peninsula (northwestern Russia) over the last 560 years. *Int. J. Environ. Res.*, 5, 855-864.
- de Silva S.L., Zielinski G.A., 1998, Global influence of the AD 1600 eruption of Huaynaputina, Peru. *Nature*, 393, 455-458.
- Sonnek K.M., Martensson T., Vieback E., Tunved P., Grahn H., Schoenberg P., Brannstrom N., Bucht A., 2017, The impacts of a Laki-like eruption on the present Swedish society. *Nat. Hazards*, 88, 1565-1590.
- Soon W., Baliunas S., 2003, Proxy climatic and environmental changes of the past 1000 years. *Clim. Res.*, 23, 89-100.
- Stoffel M., Khodri M., Corona C., Guillet S., Poulain V., Bekki S., Guiot J., Luckman B.H., Oppenheimer C., Lebas N., Beniston M., Masson-Delmotte V., 2015, Estimates of volcanic-induced cooling in the Northern Hemisphere over the past 1,500 years. *Nature Geosci.*, 8, 784-788.
- Sunkara S.L., Tiwari R.M., 2016, Wavelet analysis of the singular spectral reconstructed time series to study the imprints of solar-ENSO-geomagnetic activity on Indian climate. *Nonlin. Processes Geophys.*, 23, 361-374.
- Svensmark H., Friis-Christensen E., 1997, Variation of cosmic ray flux and global cloud coverage – a missing link in Solar-climate relationships. *J. Atm. Terr. Phys.*, 59, 1225-1232.
- Thordarson T., Self S., 2003, Atmospheric and environmental effects of 1783 – 1784 Laki eruption: A review and re-assessment. *J. Geophys. Res.*, 108, 4011-4019.
- Torrence C., Compo G.P., 1998, A practical guide to wavelet analysis. *Bull. Am. Meteorol. Soc.*, 79, 1998.
- Vaganov E.A., Shiyatov S.G., 2005, Dendroclimatic and dendroecological research in Northern Eurasia. *Lesovedenie*, No 4, 18-27.
- Velasco V.M., Mendoza B., 2008, Assessing the relationship between solar activity and

- some large scale climatic phenomena. *Adv. Space Res.*, 42, 866-878.
- Vecchi G.A., Soden B.J., 2007, Global warming and the weakening of the tropical circulation. *J. Clim.*, 20, 4316-4340.
- Verosub K.L., Lippman J., 2008, Global impacts of the 1600 eruption of Peru's Huaynaputina. *EOS Trans.*, 89, 141-142.
- Wahl E.R., Diaz H.F., Smerdon J.E., Ammann C.M., 2014, Late winter temperature response to large tropical volcanic eruptions in temperate western North America: Relationship to ENSO phases. *Global Planet. Change*, 122, 238-250.
- Wang X., Zhang Q.-B., 2011, Evidence of solar signals in tree rings of Smith fir from Sygera Mountain in southeast Tibet. *J. Atm. Sol.-Terr. Phys.*, 73, 1959-1966.
- White D.E., White J.W.C., Steig E.J., Barlow L.K., 1997, Reconstructing annual and seasonal climatic responses from volcanic events since A.D. 1270 as recorded in the deuterium signal from the Greenland Ice Sheet Project 2 ice core. *J. Geophys. Res.*, 102, 19683-19694.
- Wigley T.M.L., 2000, ENSO, volcanoes and record-breaking temperatures. *Geophys. Res. Lett.*, 27, 4101-4104.
- Zanchettin D., Bothe O., Graf H.F., Lorenz S.J., Luterbacher J., Timmreck C., Jungclaus J.H., 2013, Background conditions influence the decadal climate response to strong volcanic eruptions. *J. Geophys. Res.*, 118, 4090-4106.