

# Validation of instrument to comply institutional measures on tuberculosis infection control

Validación del instrumento para el cumplimiento de las medidas institucionales de control de la infección por tuberculosis

Validação do instrumento de conformidade das medidas institucionais de controle de infecções por tuberculose nos prestadores de serviços de saúde

Validation de l'instrument de respect des mesures institutionnelles de lutte contre l'infection tuberculeuse dans les établissements de santé

Helen Johana Ortiz Rojas<sup>1</sup>  Herney Andrés García Perdomo<sup>2</sup> 

DOI: 10.19136/hs.a19n2.3656

Original article

Date received: February 20, 2020

Date approved: April 1, 2020

Corresponding author:

Raquel García Flores. Mailing address: 5 de febrero 818 sur  
Col. Centro Ciudad Obregón, Sonora CP 85000 México.  
e mail: raquel.garcia@itson.edu.mx

## Abstract

**Objective:** To evaluate the validity and reliability of an instrument that inspect compliance with institutional tuberculosis infection control measures of Healthcare Providers in the city of Cali, Colombia.

**Materials and Methods:** Across-sectional study (psychometric type, instrument validation). The development of instrument considered sections from the Guidelines for Preventing the Transmission of Mycobacterium Tuberculosis in Healthcare Settings (CDC), Guidelines for implementation of infection control of tuberculosis in the Americas and Evaluation Control Tools for Hospital-Acquired Infections. The construction of the final instrument was carried out with the support of seven experts according to Delphi method, obtaining an instrument comprising three domains and a final number of 65 questions; each item was evaluated, usefulness, format, and validity of the instrument. Contents were assessed with the Aiken coefficient and reliability with Bangdiwala concordance coefficient. Rversion3.2.0 application was used.

**Results:** The instrument was concocted by 65 items referring to questions regarding administrative, environmental and respiratory measures. It holds acceptable psychometric properties, including good internal structure and suitable reliability.

**Conclusions:** A valid and reliable instrument was obtained to evaluate compliance with institutional measures on tuberculosis infection control in Healthcare Institutions in the city of Cali.

**Key Words:** Tuberculosis; Instrument Validation; Delphi method.

<sup>1</sup> Bacteriologist. Master in Clinical Epidemiology. Hospital Valle Del Lili Foundation. Universidad Libre. Cali, Colombia.

<sup>2</sup> Medical and Surgeon. Specialization in Urology. Master in Clinical Epidemiology. Doctorate in Clinical Medicine and Public Health (PhD). Doctorate in Education. Doctorate in Biomedical Sciences. Department of Surgery/Urology. School of Medicine. Universidad del Valle. Cali, Colombia

## Resumen

**Objetivo:** Evaluar la validez y confiabilidad de un instrumento que inspecciona el cumplimiento de las medidas institucionales de control de infecciones para la tuberculosis en las Instituciones Prestadoras de Salud de la ciudad de Cali, Colombia.

**Materiales y Métodos:** El estudio fue de corte transversal (tipo psicométrico, validación de un instrumento). El desarrollo del instrumento consideró secciones de las Guidelines for the preventing of transmission of Mycobacterium tuberculosis in health care settings (CDC), Guidelines for implementation of infection control of tuberculosis in the Americas y Evaluation Control Tools for Hospital-Acquired Infections. La construcción del instrumento final se realizó con el apoyo de 7 expertos según la metodología Delphi, obteniéndose un instrumento constituido por tres dominios y con un número final de 65 preguntas; donde se evaluó cada uno de los ítems, la utilidad, el formato y la validez del instrumento. Se evaluó el contenido con el coeficiente de Aiken y la confiabilidad con el coeficiente de concordancia de Bangdiwala. Se utilizó el aplicativo R versión 3.2.0.

**Resultados:** El instrumento quedó conformado por 65 ítems que se refieren a preguntas relacionadas con las medidas administrativas, ambientales y respiratorias. Cuenta con las propiedades psicométricas aceptables, entre ellas, una buena estructura interna y una adecuada confiabilidad.

**Conclusiones:** Se obtuvo un instrumento válido y confiable para evaluar el cumplimiento de las medidas institucionales de control de infecciones para la tuberculosis en las Instituciones de Salud de la ciudad de Cali.

**Palabras clave:** Tuberculosis; Validación de instrumento; Método Delphi.

## Resumo

**Objetivo:** Avaliar a validade e a confiabilidade de um instrumento que inspeciona o cumprimento de medidas institucionais de controle de infecção por tuberculose nas instituições de saúde da cidade de Cali, Colombia.

**Materiais e métodos:** O estudo foi transversal (tipo psicométrico, validação de um instrumento). O desenvolvimento do instrumento considerou seções das Guidelines for the preventing of transmission of Mycobacterium tuberculosis in health care settings (CDC), Guidelines for implementation of infection control of tuberculosis in the Americas e Evaluation Control Tools for Hospital-Acquired Infections. A construção do instrumento final foi realizada com o apoio de 7 especialistas, conforme a metodologia Delphi, obtendo um instrumento composto por três domínios e com um número final de 65 perguntas; onde cada um dos itens, a utilidade, o formato e a validade do instrumento foram avaliados. O conteúdo foi avaliado com o coeficiente de Aiken e a confiabilidade com o coeficiente de concordância de Bangdiwala. O aplicativo R versão 3.2.0 foi usado.

**Resultados:** O instrumento foi composto por 65 itens que se referem a questões relacionadas a medidas administrativas, ambientais e respiratórias. Possui propriedades psicométricas aceitáveis, incluindo boa estrutura interna e confiabilidade adequada.

**Conclusões:** Foi obtido um instrumento válido e confiável para avaliar o cumprimento das medidas institucionais de controle de infecção por tuberculose nas instituições de saúde da cidade de Cali.

**Mots-clés:** Tuberculose; Validation des instruments; Méthode Delphi.

## Résumé

**Objectif:** Évaluer la validité et la fiabilité d'un instrument qui inspecte le respect des mesures institutionnelles de contrôle des infections de la tuberculose dans les établissements de santé de la ville de Cali.

**Matériels et méthodes:** L'étude était transversale (type psychométrique, validation d'un instrument). L'élaboration de l'instrument a examiné des sections des Lignes directrices pour la prévention de la transmission de Mycobacterium tuberculosis dans les établissements de santé (CDC), des Lignes directrices pour la mise en œuvre de la lutte contre les infections de la tuberculose dans les Amériques et des outils de contrôle de l'évaluation des infections nosocomiales. La construction de l'instrument final a été réalisée avec le soutien de 7 experts selon la méthodologie Delphi, obtenant un instrument composé de trois domaines et avec un nombre final de 65 questions; où chacun des éléments, l'utilité, le format et la validité de l'instrument ont été évalués. Le contenu a été évalué avec le coefficient Aiken et la fiabilité avec le coefficient de concordance Bangdiwala. L'application R version 3.2.0 a été utilisée.

**Résultats:** L'instrument était composé de 65 éléments qui renvoient à des questions liées aux mesures administratives, environnementales et respiratoires. Il a des propriétés psychométriques acceptables, y compris une bonne structure interne et une fiabilité adéquate.

**Conclusions:** Un instrument valide et fiable a été obtenu pour évaluer le respect des mesures institutionnelles de contrôle des infections de la tuberculose dans les établissements de santé de la ville de Cali.

**Mots-clés:** Tuberculose; Validation des instruments; Méthode Delphi.

## Introduction

Tuberculosis (TB) has been historically linked with adverse socioeconomic conditions. However, in recent decades, other related factors such as Human Immunodeficiency Virus, the emerging of resistant strains, and exacerbated poverty have been reported, particularly in low-income countries. This disease is still a serious public health issue due its economic and clinical burden<sup>1</sup>.

Given its worldwide impact, the World Health Organization (WHO) poses it as a global health priority, calling on countries to develop and implement prevention and control strategies adjusted to each environment's features and needs<sup>2</sup>.

The basic tuberculosis control strategy is interrupting contagion, cutting the transmission chain between host, reservoir, and infectious agent<sup>3</sup>. Therefore, WHO and CDC have developed guidelines on tuberculosis prevention and transmission in healthcare institutions. These guidelines suggest activities to improve practices on tuberculosis prevention and transmission control within such settings. Control measures established by those agencies are divided into administrative, environmental and respiratory protection measures<sup>4,5</sup>.

Tuberculosis events within healthcare facilities have gained importance, by putting patients and health staff at risk, being the most contagion vulnerable settings, as all kinds of patients with several different conditions and illnesses access these facilities. Thus, it is a must for healthcare institutions to adopt and incorporate protective measures for patients, visitors, and health staff<sup>6</sup>.

Countries such as Peru have incorporated measures enacted by these agencies into their own operational plan<sup>7</sup>. In Colombia, there are few studies regarding implementation of tuberculosis control measures in healthcare facilities<sup>8</sup>, which is why it is necessary to carry out processes of research and validation for instruments that provide support on this matter. Therefore, this research article will show the results of the design, construction and validation of this instrument, taking into account the administrative, environmental and respiratory protection components that health establishments must take into account for compliance with infection control measures. Administrative control measures are those measures that define the health establishment with the objective of preventing and controlling the transmission of tuberculosis between health personnel and patients; Environmental control measures are aimed at reducing the concentration of infectious droplet nuclei and finally respiratory protection measures seek to protect health personnel through personal protection elements.

The existence of these measures will help to address the problems related to the risk the tuberculosis infection.

The purpose of this study was to validate an instrument that will assess infection control compliance of Healthcare Providers in Cali (Colombia).

## Materials and methods

A cross-sectional study, previously approved by the Ethics Committee of Libre University.

### Instrument Formulation (domains identification):

The development of the instrument by the author, considered sections of guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health care settings (CDC)<sup>4</sup>, guidelines for implementation of infection control of tuberculosis in the Americas<sup>5</sup>, and evaluation control tools for Hospital-Acquired Infections<sup>9</sup>, designing an easy to apply instrument with answer opportunities to respondents. The initial instrument covered 68 questions within the two domains: Administrative measures, and environmental and respiratory protection measures.

### Construction/ Formation and evaluation of the instrument:

This research methodology development corresponded to a cross-sectional study (psychometric), where an instrument was designed to check perceived validity and content validity, which would allow a measuring phenomenon that should exist but cannot be directly assessed.

### Pilot test:

Once the instrument was built, it was validated by a group of seven experts (two bacteriologists, two nurses, one psychologist, one sociologist, and one statistician) -who knew about tuberculosis and have worked in this field- using the Delphi method. This method is a structured and multi-stage process with an expert panel invited to participate in a series of rounds to identify, clarify and achieve consensus on a particular topic<sup>10-12</sup>. The selected health specialties experts hold a higher education training (Masters). The objective of this test was to evaluate structure and general design of the survey, the utility, content, and validity of each question and its reliability among experts.

### Delphi Method- First round:

The direct delivery of the instrument to selected experts for reading was carried out previously and personally.

The objectives of the study, and work plan for the Delphi Method development were verbally explained. Their consent to participate in the study was provided.

### Delphi Method-Second round (item generation):

In this round, evaluation of the instrument was executed by each expert. The following issues were to be considered:

- Evaluation of items: Each expert qualified each item according to understanding, ambiguity, writing, range and frequency of responses.
- Evaluation on the usefulness of the instrument: Each expert qualified each item according to application time, need for training, and rating convenience.
- Evaluation on the format of the instrument: Each expert assessed that the instrument had clear instructions, response method and appropriate typeface.
- Evaluation on the validity of the instrument (apparent and content validity): For the study of the apparent validity and content, the Aiken Content Validity Coefficient was used where the judges considered each item of the instrument relevant

Each expert evaluated all issues using the following scale: Adequate (3 points), Slightly Adequate (2 points), and Inadequate (1 point).

### Statistical analysis:

*Data quality:* Once information was obtained, a tabulation of experts' answers was made, building a database on Microsoft Excel 2013. We assessed database quality by randomly take 10% of the 3332 data, finding no inconsistencies on audited data (333). Information analysis was performed with R version 3.2.0. statistical package. The analysis was focused on evaluating psychometric properties of the instrument, including evaluation of items, instrument's usefulness and format, validity and reliability.

*Evaluation of items:* On the items' evaluation, they were rated 'excellent' when 70% of experts set it in the 'Adequate' category. Evaluating criteria for each item was:

- Understanding: Items should be easy to understand regardless level of education.
- Ambiguity: The use of words that may have different Interpretations should be avoided.
- Writing: Contents should not be confusing or provide emotional burden. Wording must be clear.
- Range of responses: Identifying items that need more answer opportunities or the possible removal of unnecessary options.

When evaluating the usefulness of the instrument and format, a frequency distribution analysis was carried out according to the answers provided by the experts.

*Evaluation of apparent validity and content:* Apparent validity and contents were assessed using the Aiken coefficient, which is one of the quantifying techniques on item relevance to a content's domain, with the participation of N number of judges or experts, and with ranges going from 0 to 1. 1 being the possible magnitude that indicates perfect agreement among judges regarding higher validity score on evaluated contents<sup>13,14</sup>. A minimum Aiken coefficient standard value was considered as equal to or greater than 0.70, which allowed quantifying the relevance of each item regarding the domain established according to the number of experts<sup>15,16</sup>.

*Reliability regarding the subject* was assessed by Bangdiwala's original and weighted coefficient. The following classification of Bangdiwala's coefficient was scored: 0.81 – 1.0 (Very Good), 0.61-0.80 (Good), 0.41-0.60 (Moderate), 0.21-0.40 (Weak) and 0.0 – 0.20 (Poor). The value of these coefficients oscillates between "0" (Null Agreement) and "1" (Total Agreement)<sup>17</sup>.

*Correlations among judges:* To determine correlations among judges' agreement for topics of understanding, ambiguity, writing, and range of responses on each domain, Spearman's coefficient (rho) was used. This coefficient fluctuates between -1 to 1. A value of -1 to +1 indicates a perfect negative or positive linear relationship. Correlational relationships can be demonstrated through correlograms using circles filled in a clockwise direction for positive values, and counterclockwise for negative values<sup>18</sup>.

**Final instrument construction:** We considered not only statistical analysis results, but also suggestions and recommendations provided by the experts.

## Results

68 items were initially evaluated by the judges, to obtain 65. The first instrument covered two domains (administrative, and environmental/respiratory), which was adjusted to three domains (administrative, environmental, and respiratory).

**Items Analysis.** The results obtained from expert's items evaluation showed that 19 of the 68 items needed adjustments regarding understanding, ambiguity, writing, and range of responses as they were not rated as 'Adequate' by the experts. Questions with any evaluated criteria percentage below 70 were modified according to experts' recommendations and reflected in table 1.

**Table 1.** Items evaluation on Administrative Control Measures Domain

Administrative Control Measures	Items	Understanding	Ambiguity	Writing	Range of Responses
Infection Control Committee issues	Item 3	58%	86%	58%	100%
	Item 4	58%	72%	58%	72%
	Item 8	29%	29%	58%	86%
	Item 9	58%	72%	72%	86%
	Item 11	100%	100%	100%	29%
	Item 13	100%	100%	86%	58%
	Item 14	58%	72%	58%	15%
Health staff issues	Item 17	44%	72%	44%	86%
Diagnosis issues	Item 21	44%	58%	58%	86%
Sample collection, processing and reporting issues	Item 27	58%	44%	58%	100%
	Item 28	72%	72%	86%	44%
Hospitalization issues	Item 34	58%	58%	58%	72%
	Item 35	58%	86%	58%	72%
Patient follow-up issues	Item 38	72%	44%	44%	100%
	Item 44	100%	100%	100%	58%
	Item 51	58%	86%	72%	100%

Source: Personal Reference

### Up next the findings for each domain:

*Administrative Control Measures domain:* On Infection Control Committee, item 6 was deleted (same contents of item 13), so this domain has only 13 questions out of which, items 3, 4, 8, 9, 11, 13 and 14 were modified.

On health staff, experts rated items 15, 16, 18 and 19 with an 'Adequate' percentage. Item 17 only presented low percentage in the experts' evaluation; they recommended separating both questions contained in it.

On diagnosis, the experts rated items 20, 22, and 23 with an 'Adequate' percentage. Only item 21 showed low percentages in the experts' evaluation.

On sample collection, processing and reporting, the experts rated items 24, 25, 26, 29 and 30 with an 'Adequate' percentage, while items 27 and 28 were rated with low percentages, so they were adjusted. Experts agreed on deleting item 30 as it had same contents of item 57.

On hospitalization, the experts rated items 31, 32, 33, 36 and 37 with an 'Adequate' percentage. They recommended to keep the answer option of "Does Not Apply" in the event

of healthcare facilities that don't provide hospitalization services. Items 34 and 35 showed low percentages in the experts' evaluation.

On patients' follow-up part, the experts assessed items 38 to 55; where items 38, 44 and 51 showed low percentages in the experts' evaluation. Following their recommendation item 46 was re-located under Sample Collection Processing and Reporting. Items 47 and 48 merged into a single one. Items 52 and 53 were removed from the instrument as their contents are same of item 42 and 39 respectively.

*On Environmental Control Measures and Respiratory Protection,* experts recommended separating these measures with their respective questions in two different domains. They assessed items 56 to 68, where only items 61, 62 and 68 showed low percentages in their evaluation. Questions with any evaluated criteria percentage below 70 were modified according to experts' recommendations table 2.

### Evaluation of Instrument's format and usefulness:

Regarding the usefulness of the instrument, it was found that 5 of the 7 expert judges rated instrument filling out time as good (71%), given that it has quick measuring scale that

**Table 2.** Items evaluation on Environmental Control Measures and Respiratory Protection

Measures of Environmental Control and Respiratory Protection	Items	Understanding	Ambiguity	Writing	Range of Responses
Environmental Control and Respiratory Protection issues	Item 61	58%	86%	58%	86%
	Item 62	100%	100%	86%	58%
	Item 68	58%	86%	72%	86%

Source: Personal Reference

do not fatigue respondents and saves time. Five experts believe there is no need for training prior the application of the instrument, since items are general questions anyone in charge of a healthcare institution or an Epidemiological Surveillance Committee must know and can answer.

As for the format evaluation of the instrument, six experts stated that the guides in the instrument were clear, and there is no need for an example on them. Overall, the format, has an appropriate, easy to read font type, for respondents.

The following were the experts' recommendations about the instrument:

- Do not use acronyms.
- Change the word information by issue.
- Remove any authorship reference from instrument.
- Use the word Health Institution instead of Health Establishment.
- Place 'Yes' in the answers and then 'No'.
- Classify questions according to logical order.
- Change the phrasing 'level of complexity' by 'level of assistance'.

### Validity Analysis:

It was submitted to the criterion of the judges— seven altogether— the evaluation on understanding, ambiguity, writing and range of responses from each of the items in the instrument. For result systematization, the Aiken coefficient was used, complemented with the use of confidence intervals using the program proposed by Merino and Livia<sup>14</sup>. A minimum standard of an Aiken's V coefficient equal to or greater than 0.70 was used, accepting those items which Aiken's coefficient was above 0.70 as valid.

Items which confidence intervals were above 0.45 were accepted as valid as well.

Table 3 results show that 11 of the 68 items built in the instrument showed Aiken coefficients below the minimum standard value. We reviewed the suggestions made by the judges and recommendations thoroughly for the respective adjustment of those items according to criterion assessed.

### Subject-Related reliability.

Through this analysis, reliability was represented by measuring the experts' scores to aspects of understanding, ambiguity, writing, and range of responses. It was found that values on original agreement coefficients oscillated between 0.66 and 0.81, while weighted agreement values oscillated between 0.83 and 0.91. It was found that original and weighted agreements were higher than the "poor" classification table 4. The aspect showing greater concordance was range of responses with a weighted coefficient of 0.91.

**Correlations between understanding, ambiguity, writing and range of responses in the Administrative measures domain.** Results show that judges tended to agree on aspects of understanding, writing, and ambiguity. Through the Correlogram, it was possible to demonstrate high positive correlations between "understanding" and "writing" ( $\rho = 0.79$ ) and between "understanding" and "ambiguity" ( $\rho = 0.73$ ). Lowest correlation was present between "range of responses" and "ambiguity" ( $\rho = 0.16$ ). It was also evident that judges 'agreeing on "range of responses" was not related to the other aspects figure 1. On the other hand, the environmental and respiratory measures, the results shown pose that judges tended to agree on aspects of understanding, writing, and ambiguity. Through Correlograms, it is possible to demonstrate high positive correlations between "understanding" and "writing" ( $\rho = 0.71$ ), and between "understanding" and "ambiguity" ( $\rho = 0.81$ ). The lowest correlation was between "understanding" and "range of responses" ( $\rho = 0.11$ ) figure 2.

**Final version.** Instrument's final version comprised of 65 items evaluating administrative control measures (items 1-52), environmental control measures (items 53 – 60),

**Table 3.** Aiken V-coefficient for aspects of items' understanding, ambiguity, writing and range of responses (C. Aiken < 0.70)

Items	Understanding	IC 95%	Ambiguity	IC 95%	Writing	IC 95%	Rangeof responses	IC 95%
Item 8	0.57	0.32-0.79	0.43	0.21-0.67	0.79	0.52-0.92	0.93	0.69-0.99
Item 9	0.64	0.39-0.83	0.79	0.52-0.92	0.79	0.52-0.92	0.93	0.69-0.99
Item 11	1.0	0.78-1.0	1.0	0.78-1.0	1.0	0.78-1.0	0.57	0.32-0.79
Item 14	0.64	0.39-0.83	0.79	0.52-0.92	0.71	0.45-0.88	0.14	0.04-0.40
Item 17	0.64	0.39-0.83	0.86	0.60-0.96	0.64	0.39-0.83	0.93	0.69-0.99
Item 21	0.64	0.39-0.83	0.71	0.45-0.88	0.71	0.45-0.88	0.86	0.60-0.96
Item 27	0.71	0.45-0.88	0.64	0.39-0.83	0.79	0.52-0.92	1.0	0.78-1.0
Item 28	0.79	0.52-0.92	0.79	0.52-0.92	0.93	0.69-0.99	0.64	0.39-0.83
Item 34	0.57	0.32-0.79	0.71	0.45-0.88	0.57	0.32-0.79	0.86	0.60-0.96
Item 38	0.71	0.45-0.88	0.57	0.32-0.79	0.57	0.32-0.79	1.0	0.78-1.0
Item 51	0.64	0.39-0.83	0.86	0.60-0.96	0.86	0.60-0.96	1.0	0.78-1.0

Source: Personal Reference

**Table 4.** Average values of original concordance coefficients (BN) and weighted concordance coefficients (BWN) for all four aspects


Aspects	Original concordance coefficients (bw)	Weighted concordance coefficients (bwn)
Understanding	0.71	0.83
Ambiguity	0.75	0.87
Writing	0.66	0.84
Rangeof responses	0.81	0.91

Source: Personal Reference

and respiratory protection control measures (items 61-65) (Supplementary file 1). Analysis of these results allowed to estimate that items' formulation was relevant, coherent, and sufficient, grouped under the right domain, obtaining the needed information from each. Likewise, answering options were improved to make each a more appropriate question.

**Application of the Instrument.** Below are the results obtained in three domains of tuberculosis infection control measures after instrument's application in healthcare facilities. Five Health-Care Institutions were visited (two private, two public, and a mixed entity) according to authorizations from the Ethics Committee of each entity.

**Administrative control Measures:** In connection with the existence of a committee to monitor healthcare-associated infections, four out of the five healthcare institutions stated the existence of such committee. Two of the healthcare institutions surveyed indicated having a tuberculosis control program implemented which helps them monitor their patients according to their level of complexity. On the other hand, four of the five institutions had a professional or individual in charge of the tuberculosis program. Tuberculosis events reported to domestic's Public Health System (SIVIGILA- *by its acronym in Spanish*) are reported by four of the five healthcare institutions. All health care facilities provide education to health staff in the management of the disease. Regarding health staff involved in tuberculosis management,

**Figure 1.** Correlogram. Administrative measures

Source: Personal Reference


all healthcare institutions refer having an interdisciplinary team of nurse's assistants, psychologists, social workers, infection disease specialists, internists, respiratory therapists, and pharmaceutical chemists; all health staff know the tuberculosis care path. And in terms of continuous training there is an education and training record as well. Four of the five healthcare institutions claim having had tuberculosis events among healthcare staff. 80% of these health care facilities do not run a regular PPD testing to their personnel as part of a health staff care follow-up.

As for medical diagnosis, all healthcare institutions surveyed perform this procedure according to patient's signs and symptoms. The identification of respiratory symptomatic or tuberculosis suspected patients is performed by all health care facilities at their emergency service. Only four of the five institutions sort patients diagnosed with confirmed tuberculosis away from other patients.

The Clinical lab in four of the healthcare institutions carry out a smear sample collection and had containers or inputs (laminar flow cabin) sufficient for such collection and processing.

Report delivery is done within a 24-hour period. Three of the healthcare facilities send molecular tests to labs in other healthcare institutions.

Finally, on patient follow-up, four of the five healthcare institutions had an updated record on tuberculosis diagnosed patients, medical consultation is the primary tool for this follow-up to begin. All healthcare facilities fill out an individual patient card and in all of them are keep sufficient drugs of good quality for these patients.

**Figure 2.** Correlogram. Environmental and Respiratory measures

Source: Personal Reference

### Environmental Control Measures:

On this regard, two of the five institutions had a site established for sample collection. All facilities had waiting rooms where TB diagnosed patients had natural or mechanical ventilation, but are not separated from other patients. Only one healthcare institution had central air conditioning. On tuberculosis patient's confinement, all facilities had at least one exclusive bathroom or toilet; four of the five had the right visible confinement signaling. Regarding ventilation system maintenance, only four of the five entities surveyed had it done monthly or every three months.

### Respiratory control Measures:

Health care institutions provide high-efficiency facemask to health staff. Additionally, there are standardized guides or protocols for the use of these Personal Protective Equipment. Health staff is trained on the proper use of these elements.

### Discussion

Measurement is a value appointing process for certain real life events; this process is part of the daily practice to all professionals since scientific knowledge is obtained from it. So much so, that all fields using measurements can take advantage of it to create theories and test hypothesis<sup>19</sup>. The application of surveys or instruments generates measurement and data which once analyzed, can determine important decisions most of the time.

Consequently, it is necessary to have valid and reliable instruments to use in clinical practice and research, able to respond to human needs and conditions, and promote treatments, early diagnoses and therapeutic decisions.

### Summary of main findings:

As for methodological aspects, the expert consensus is highlighted. The level of expertise of the judges on this particular topic, managed to issue assessments on each evaluated aspect, changing instrument items' form and semantics and then, finally building an instrument version 2 to evaluate institutional compliance with tuberculosis infection control measures of healthcare institutions in the city of Cali.

The proposed survey will make it possible to learn whether or not health-care institutions comply with tuberculosis infection control measures and are able to lead research on this matter. The instrument has the validity of appearance and content to be applied in the health sector.

On appearance and contents validity, through Aiken's coefficient obtained relevance of items regarding contents of established domain was quantified, which helps to accurately evaluate what is intended to measure. Only 11 items showed Aiken coefficient below 0.70. Items 6, 30, 52 and 53 were eliminated. The others showed coefficients above 0.70, which according to Merino Soto and Livia Segovia, is a more conservative value. The above provides evidence of content validity of surveyed domains.

Regarding subject reliability, it shows a moderate to very good result, understood as though two or more experts evaluated each item, on understanding, ambiguity, writing and range of responses, with similar results. The aspect showing greater concordance was "range of responses" with a Bangdiwala weighted coefficient of 0.91, followed by "ambiguity" (BWN = 0.87), then "writing" (BWN = 0.84) and finally "understanding" (BWN = 0.83).

Whereas in correlation, findings show that both, administrative measures and environmental and respiratory measures, have concordance among judges as equal in understanding, writing, and ambiguity. There is a positive linear trend between these aspects.

Overall, results of the instrument's validation process confirm compliance with minimum requirements on validity and reliability.

Considering results regarding the application of the instrument, the commitment and control exercised by each healthcare institution surveyed on tuberculosis is evident. It is worth highlighting the work performed by managers of these healthcare facilities, in the face of administrative, environmental and respiratory strategies, being immersed in their institutional operational plan.

### Contrast with literature:

Literature shows several studies linking the Delphi method as a methodological strategy in the formulation and validation of an instrument. In the study of Longobardi-Vasquez and collaborators, the first one developed was submitted to six experts<sup>20</sup>; likewise, in a research carried out by Noriega Bravo and collaborators, they used the Delphi method with the help of seven experts to evaluate the total of items<sup>21</sup>. Hence confirmed, the number of experts used in this research was appropriate.

As exposed by Cruz-Avelar Agles<sup>18</sup>, Mousavi and collaborators<sup>22</sup> and other authors, validity measures the degree to which items seem to measure what is proposed, and should include the judgment or opinions of some experts in the matter. For the appearance and content validation stage, there are many methods used, among them are quoted: Method based on factorial analysis, content validity index, item-objective congruence index, congruence index, content validity coefficient, among others. Aiken's coefficient was chosen to evaluate contents' validity by judges' criterion; since, according to the literature this coefficient is easy to calculate, it guarantees objectivity of the procedure, and allows to obtain values feasibly contrasted with statistics depending on the size of selected judges' sample. The instrument's 68 items were evaluated by the seven experts in all four aspects: understanding, writing, ambiguity, and range of responses, which made it possible to obtain rates which helped modify or eliminate the instrument's item.

On the other hand, different procedures have been used to determine the degree of agreement among judges. The Kappa coefficient became the most widely used agreement index<sup>23</sup>. Despite that, the Bangdiwala weighted coefficient of this research was chosen because it is more stable than Kappa's as sample size decreases. In addition, Bangdiwala's coefficient relies on the number of judges and is more powerful to determine concordance among them<sup>24</sup>.

Different studies evaluate the level of implementation of control measures. Brouwer and collaborators conducted a study in three provinces of Mozambique where they assessed management, administrative, and environmental measures in the 29 healthcare centers, concluding that guidelines on tuberculosis diagnosis and treatment were not present in those facilities<sup>25</sup>.

Another study conducted in Bogotá - Muñoz Sanchez et al., describes the implementation of control measures in 51 healthcare institutions; evidence showed 68% of them had an infection control plan in place, out of which 16% had mechanical or natural ventilation, and 84% had N95 respirators available for their staff<sup>26</sup>.

## Strengths and Weaknesses:

Obtaining this valid and reliable instrument will identify the level of compliance measures for tuberculosis infection control in health-care institutions. Conversely, this instrument becomes a useful and practical tool to guide these entities' directors/managers into incorporating changes or to reshape institutional procedures on tuberculosis prevention and control. Weaknesses for this instrument's validation were the type of questions, and the nominal and dichotomous variables used, which reduced the incorporated statistical analyses associated with these characteristics.

Application of this instrument's final version was only possible in five out of the seven healthcare institutions chosen. For future research on this topic, would be important to choose more facilities with a different inclusion criterion. Most surveys applied were answered by nurses who were members of COVE\* (Epidemiological Surveillance Committee –\*by its acronym in Spanish), the Hospital Committee, or the tuberculosis program, which makes us think that the assisting staff is trained and skilled on tuberculosis and knows all about the approach to this infection.

## Public Health implications:

Finally, it can be stated that it is a ready-to-use instrument, useful to identify compliance of administrative, respiratory, and environmental measures at any level of health care, including facilities in charge of management, diagnosis, and treatment of tuberculosis patients.

On the other hand, the possibility to apply this instrument allows to evaluate the existence of such measures, thus carrying out other studies in that regard which may lead to identify and reduce the risk of tuberculosis infections.

As a conclusions: On the analysis of items' evaluation, it is concluded that the number of items (68) initially considered in the instrument's 1st version, needs to be modified according to the experts, in all four scoring aspects, thus leaving an instrument Version 2 with 65 questions.

In that sense, the instrument's features, such as the number of replies, vocabulary used, the distribution of the questions, are factors regarded within the four aspects evaluated: understanding, ambiguity, writing and range of responses.

Contents' validity and degree of reliability were adequate and acceptable. Only 11 items showed an Aiken coefficient below 0.70. Incidentally, reliability among experts for evaluated aspects showed a moderate to very good concordance between every pair of experts.

The above, allowed obtaining a reliable and valid tool to be used in health institutions, where administrative, environmental, and respiratory control measures are the main tool to prevent contagion of this infection.

## Acknowledgments

The authors wish to thank the seven experts participating in this instrument's evaluation and made it possible to build.

**Declarations.** The authors declare that there is no conflict of interest in the present study.

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## References

1. Estrada Mota I RLJ. Tuberculosis pulmonar, un riesgo latente para los trabajadores de la salud como problema de Salud Pública. *J Negat No Posit Results*. 2019;4(2):197–209. Disponible en: DOI: 10.19230/jonnpr.2833.
2. Organization, World Health. Global tuberculosis report. WHO. 2015. Available in: DOI: 10.1017/CBO9781107415324.004.
3. Vallejo, Miguel Angel V-MAM. Historia Natural de la Enfermedad. In: *Salud Publica*. 2019. p. 31–91.
4. Jensen P a, Lambert L a, Iademarco MF. Guidelines for preventing the transmission of Mycobacterium tuberculosis in health-care settings,. 2005;1–141. Available in: DOI: 10.2307/42000931.
5. Organización Panamericana de la Salud. Lineamientos para la implementación del control de infecciones de tuberculosis en las Américas. 2014;1–84. Disponible en: DOI: 10.1086/203554.
6. Nathavitharana RR, Bond P, Dramowski A, Kotze K, Lederer P, Oxley I, et al. Agents of change: The role of healthcare workers in the prevention of nosocomial and occupational tuberculosis. *Presse Medicale*. 2017. p. 1–17. Available in: DOI: 10.1016/j.lpm.2017.01.014.
7. Alarcón V, Alarcón E, Figueroa C, Mendoza-Ticona A. Tuberculosis en el Perú: Situación epidemiológica, avances y desafíos para su control. *Rev Peru Med Exp Salud Publica*. 2017;34(2):299–310. Disponible en: DOI: 10.17843/rpmesp.2017.342.2384.

8. Muñoz-Sánchez AI, Castro-Cely Y. Medidas de control de tuberculosis en una institución de salud de Bogotá D.C. *Rev Fac Nac Salud Pública*. 2016;34(1):37–47. Disponible en: DOI: 10.17533/udea.rfnsp.v34n1a05.
9. Social M de SP y A. Herramienta de Evaluación del control de infecciones Intrahospitalarias. 2009;25–33.
10. Humphrey-Murto S, de Wit M. The Delphi method—more research please. *J Clin Epidemiol*. 2019;106:136–9. Available in: DOI: 10.1016/j.jclinepi.2018.10.011.
11. Reguant Álvarez M, Torrado Fonseca M. El método Delphi. *REIRE Rev d'innovació i Recer en Educ*. 2016;9(2):87–102. Available in: DOI: 10.1344/reire2016.9.1916.
12. Cruz Ramírez M, Rúa Vásquez JA. Surgimiento y desarrollo del método Delphi: una perspectiva cuantitativa. *Biblios J Librariansh Inf Sci*. 2018;(71):90–107. Disponible en: DOI: 10.5195/biblios.2018.470.
13. Galicia Alarcón L, Balderrama Trapaga J, Edel Navarro R. Validez de contenido por juicio de expertos: propuesta de una herramienta virtual. *Apert Rev Innovación Educ*. 2017;9(2):42–53. Disponible en: DOI: 10.18381/ap.v9n2.993.
14. Merino Soto C, Livia Segovia J. Intervalos de confianza asimétricos para el índice la validez de contenido: Un programa Visual Basic para la V de Aiken. *An Psicol*. 2009;25(1):169–71. Disponible en: DOI: 10.6018/71631.
15. Merino-Soto C. Confidence interval for difference between coefficients of content validity (Aiken's V): A SPSS syntax. *An Psicol*. 2018;34(3):587–90. Available in: DOI: 10.6018/analesps.34.3.283481.
16. Pedrosa I, Suárez-Álvarez J, García-Cueto E. Evidencias sobre la Validez de Contenido: Avances Teóricos y Métodos para su Estimación. *Acción Psicológica*. 2014;10(2):3–18. Disponible en: DOI: 10.5944/ap.10.2.11820.
17. Warrens MJ, de Raadt A. Properties of Bangdiwala's B. *Adv Data Anal Classif*. 2019;13(2):481–93. Available in: DOI: 10.1007/s11634-018-0319-0.
18. Friendly M. Corgrams: explanatory displays for correlation matrices. *Am Stat*. 2002;56(4):316–24. Available in: DOI: 10.1198/000313002533.
19. Cruz-Avelar Agles C-PE. Metodología para la construcción de instrumentos de medición en salud. *Alergia, Asma e Inmunol Pediátricas*. 2017;26(3):100–5.
20. Longobardi-Vasquez G, Diaz-Velez C V-CM. Validación de Instrumento para medir el nivel de conocimiento en el tratamiento de tuberculosis en médicos del primer nivel de atención. *Revista Hispanoamericana de Ciencias de la Salud*. 2016;2(2):140–5.
21. Noriega Bravo V de las M, Pría Barrios M del C. Instrumento para evaluar el clima organizacional en los grupos de control de vectores. *Rev Cuba Salud Pública*. 2011;37(2):116
22. Mousavi SM, Takian A, Tara M. Design and validity of a questionnaire to assess national eHealth architecture (NEHA): A study protocol. *BMJ Open*. 2018;8(12):1–6. Available in: DOI: 10.1136/bmjopen-2018-022885.
23. Martín Andrés A, Álvarez Hernández M. Hubert's multi-rater kappa revisited. *Br J Math Stat Psychol*. 2020;73(1):1–22. Available in: DOI: 10.1111/bmsp.12167.
24. Shankar V, Bangdiwala SI. Behavior of agreement measures in the presence of zero cells and biased marginal distributions. *J Appl Stat*. 2008;35(4):445–64. Available in: DOI: 10.1080/02664760701835052.
25. Brouwer M, Coelho E, das Dores Mosse C, van Leth F. Implementation of tuberculosis infection prevention and control in Mozambican health care facilities. *Int J Tuberc Lung Dis*. 2015;19(1):44–9. Available in: DOI: 10.5588/ijtld.14.0337.
26. Muñoz Sánchez, A. I., Saavedra Cantor, C. J., & Cruz Martínez OA. Control de la infección por tuberculosis en instituciones de salud. *Investig Andin*. 2016;18(33):1686–96. Disponibilidad en: DOI: 10.4046/trd.2016.79.4.234.