

Potential Antimicrobial and Chemical Composition of Essential Oils from *Piper caldense* Tissues

Dayane Silva Rocha,¹ Janete Magali da Silva,² Daniela Maria do Amaral Ferraz Navarro,³ Cláudio Augusto Gomes Camara,¹ Camila Soledade de Lira³ and Clécio Souza Ramos^{1*}

¹ Department of Chemistry, Rural Federal University of Pernambuco, Recife-Pe, Brazil

² Department of Antibiotics, Federal University of Pernambuco, Recife-Pe, Brazil

³ Department of Fundamental Chemistry, Federal University of Pernambuco, Recife-Pe, Brazil

* Tel: +55 81 33206379; e-mail address: clecio.ramos@ufrpe.br

Received September 19th, 2015; Accepted July 8th, 2016.

Abstract. The essential oils from leaves, stems and roots of *Piper caldense* were analyzed by GC-MS. The antibacterial potential of the oils was evaluated against gram-negative bacteria and gram-positive bacteria. The major chemical constituents that were identified from various parts of this plant were α -cardinal, α -muurolol, tujopsan-2- β -ol and δ -cadiene in the leaves, valencene, pentadecane, elina-3,7-11-dieno α -terpineol in the roots and terpine-4-ol, α -terpineol, α -cadinol 2- β -ol in the stems. Tissue oils showed antibacterial activity against the bacteria tested except for *Enterococcus faecalis*. This is the first report of the biological activity and chemical composition essential oil of *P. caldense*.

Key words: *Piper caldense*; Piperaceae, essential oil; antimicrobial activity.

Resumen. Los aceites esenciales de hojas, tallos y raíces de *Piper Caldense* se analizaron mediante CG-EM. El potencial antibacteriano de los aceites se evaluó frente a bacterias gram-negativas y bacterias gram-positivas. Los principales componentes químicos que fueron identificados de varias partes de esta planta fueron α -cardenal, α -muurolol, tujopsan-2- β -ol y δ -cadiene en las hojas, valenceno, pentadecano, elina-3,7-11-dieno α -terpineol en las raíces y terpine-4-ol, α -terpineol, α -cadinol 2- β -ol en los tallos. Aceites del tejido mostraron actividad antibacteriana contra las bacterias analizadas excepto por *Enterococcus faecalis*. Este es el primer informe de la actividad biológica y la composición química del aceite esencial de *P. caldense*.

Palabras clave: *Piper caldense*; Piperaceae, aceite esencial; actividad antimicrobiana.

Introduction

The Piperaceae family is estimated to contain more than 3.600 species in five genera (*Macropiper*, *Zippelia*, *Piper*, *Peperomia* and *Manekia*) widely distributed in tropical and subtropical regions of the world [1]. The genus *Piper* is the largest one, comprising approximately 2000 species [1]. The Piperaceae family is valued due to its biological, chemical, economic and ecological characteristics, attributed to its secondary metabolites such as alkaloids, lignans, neolignans, amides, chromene, polyketides, phenylpropanoids, prenylated benzoic acid, flavonoids and terpenes [2-3]. In particular, the essential oils obtained from the *Piper* species are considered important sources of insecticidal compounds, such as phenylpropanoids safrole, eugenol and dillapiole. The *Piper* species as *P. hispidinervium*, *divaricatum*, *P. callosum* and *P. aduncum* have been considered as natural sources of safrole with biologic potential [4]. The essential oil of *P. divaricatum* tissues and safrole were active against bacteria *Listeria monocytogenes*, *Salmonella Typhimurium* and *Pseudomonas aeruginosa* [5]. The essential oil from the *P. corcovadensis* exhibited a strong oviposition deterrent effect. Even at a concentration of 5 ppm, the oil inhibiting egg laying of pregnant females, and the major constituents

of oil identified were 1-butyl-3,4-methylenedioxybenzene and terpinolene [6].

As part of the systematic study of volatile constituents from the *Piper* species [5,7,8], in the present study, the chemical constituents of essential oils from the *Piper caldense* (Piperaceae) leaf, root and stem were identified. The *P. caldense* is known by the popular name “pimenta d’arda” or “pimenta d’água” and commonly used in Brazilian folk medicine for the treatment of snake, sedation and diseases of the stomach [9-10]. So the medicinal proprieties of plant, using the antimicrobial activity of essential oils from *P. caldense* tissues, were tested. To our knowledge, there are no published reports on the antimicrobial activity and chemical composition of the essential oils from *P. caldense* tissues.

Results and discussion

Chemical characterization

Results obtained in the present study for the qualitative and quantitative analyses of the essential oils are shown in Table 1. The main classes present in the leaf oil were monoterpene

hydrocarbons and sesquiterpene hydrocarbons in a total of 28 compounds, representing 83.4% of the total oil. The leaf oil comprised mainly of α -cardinol (19.0%), α -muurolol (9.0%), tujopsan-2- β -ol (7.4%), δ -cadiene (5.6%), linalool (3.2%), cubenol (3.2%), γ -amorphene (3.1%), α -terpineol (3.0%). Root oil analysis showed 22 volatile compounds, representing 92.0% of the total oil. The major constituents were hydrocarbon pentadecane (35.7%), valencene (10.5%) and elina-3,7-11-diene- α -terpineol (5.4%). An analysis of the essential oil from *P. caldense* stems identified 20 compounds accounting for 83.2% of the total oil. The main compounds were identified as terpine-4-ol (18.5%), α -terpineol (15.3%), α -cadinol 2- β -ol (9.8%). In the *Piper* species, the sesquiterpene α -cardinol identified as the major compound in the leaves and roots of *P. caldense* has been previously reported in *P. cernuum* and *P. dilatatum* [11-12].

Antimicrobial activity

The essential oils of *P. caldense* tissues were tested against three Gram-positive bacteria and three Gram-negative bacteria. The results, presented in Table 2, show that the oil was biologically active against bacteria tested, with inhibition zones ranged from 20 to 33 mm. The most sensitive microorganism was *Bacillus subtilis*, with inhibition zones of 30, 26 and 23 mm for oils from the root, stem, leaf, respectively. The most effective oil was obtained from the root, showing activity against all bacteria except for resistant Gram-negative bacteria *Enterococcus faecalis*.

All compounds having inhibition diameter zones greater than or equal to 12 mm were selected for MIC (Minimum Inhibitory Concentration). According to the literature criteria

Table 1. Relative composition of the volatile oils from the *P. caldense* root, stem and leaf.

Compounds	AI Adams	Relative amount (%)			Compounds	AI Adams	Relative amount (%)		
		Root	Stem	Leaf			Root	Stem	Leaf
Linalool	1095	0.1	2.4	3.2	Spathulenol	1577			1.7
<i>p</i> -ment-2-en-1-ol	1121		1.7		Caryophyllene oxide	1582		6.2	
<i>trans</i> -pinocarveol	1135		3.3		Tujopsan-2- β -ol	1588			7.4
Borneol	1165	0.1	0.3		Diethyl phthalate	1590		4.0	
Terpine-4-ol	1174	0.1	18.5	0.3	Viridiflorol	1592		1.4	
α -terpineol	1186	0.1	15.3	3.0	Cubeban-11-ol	1595			3.4
<i>trans</i> -piperitol	1205		3.1		Rosifoliol	1600			2.7
Geraniol	1249		2.5		β -Himachalene oxide	1615			0.5
<i>trans</i> -piperitone oxide	1252			1.2	10-epi- γ -eudesmol	1773	3.9		
Longifolene	1407	3.2			Caryophylla-4(12),8(13)-dien-5	1632		0.4	
<i>E</i> -Caryophyllene	1417	1.6			Epi- α -cadinol	1638			1.7
β -Duprezianene	1421			1.4	epoxide- <i>allo</i> -Aromadendrene	1639		1.6	
Aromadendrene	1439		1.8		β -eudesmol	1640	2.4		1.5
6,9-Guaiadiene	1442			0.7	Aristol-1(10)-en-12-ol	1643	3.7		
α -humulene	1452	1.5			α -muurolol	1644	0.3	2.1	9.0
9-epi- <i>E</i> -Caryophyllene	1464			0.7	Cubenol	1645			3.2
Dauca-5,8-diene	1471		3.4		α -cadinol	1652	3.7	9.8	19.0
β -chamigrene	1476	1.9			<i>cis</i> -Calamenen-10-ol	1660			1.1
γ -Muurolene	1478			2.0	kuk7-epi- α -eudesmol	1662	3.0		
Amorpha-4,11-diene	1479			1.6	<i>Z</i> - α -Santalol	1674			2.7
<i>cis</i> -eudesma-6,11-diene	1489			0.9	Germacrene-4(15),5,10,(14)-trien-1- α -ol	1685			0.8
δ -selinene	1492		2.2		Eudesma-4(15),7-dien-1 β -ol	1687		3.2	
γ -amorphene	1495			3.1	Farnesol -2Z,6Z	1698	2.8		0.6
Valencene	1496	10.5			Hexacosane	1778	4.0		
Pentadecane	1500	35.7			α -Eudesmol acetate	1794	2.8		
δ -cadinene	1522	2.2		5.6	Total		92.0%	83.2%	83.4%
Selina-3,7(11)-diene	1545	5.4							
Germacrene B	1559	1.5							
<i>E</i> -Nerolidol	1561	1.5		4.4					

Table 2. Antimicrobial activity of essential oils from the *P. caldense* root, stem and leaf using disc diffusion assay. Diameter of inhibition zone (mm) Concentration of 20 mg/mL.

Microorganisms	Root	Stem	Leaf
Gram-negative bacteria			
<i>Escherichia coli</i>	20	0	0
<i>Klebsiella pneumonia</i>	20	0	22
<i>Pseudomonas aeruginosa</i>	27	20	0
Gram-positive bacteria			
<i>Staphylococcus aureus</i>	33	0	26
<i>Bacillus subtilis</i>	30	26	23
<i>Enterococcus faecalis</i>	0	0	0

[13], the extracts and fractions showing a MIC lower than 100, ranging from 100-500, 500-1000 and over 1000 µg/mL are characterized as potent, moderated, weak and not active, respectively. The essential oils obtained from leaves, stems and roots showed moderate to weak activity with MICs of 325 or 750 µg/mL (Table 3). The antimicrobial activity of leaf and root oils can be attributed, at least in part, to the presence of α -cardinol, considering that α -cardinol is known to present bacteriolytic activity against *S. aureus* [14].

Studies previous revealed that the essential oil from *Piper guineense* fruit was active against the gram negative bacteria *E. coli* and *P. aeruginosa*, and identified as having as major constituents the terpenes β -pinene (23%), β -caryophyllene (7%), bicyclogermacrene (6%) [15]. The essential oil from *P. malacophyllum* leaf inhibited growth of *Trichophyton mentagrophytes* and *Cryptococcus neoformans* fungi, with (+)-camphor (38%) as the major constituent identified [16]. The essential oil of *P. ilheusense* leaf was active against the microorganisms *S. aureus*, *E. coli*, *Candida albicans*, *C. krusei*, and *C. parapsilosis*, the chemical compositions of the oil consisting only of sesquiterpenes [17].

In summary, the present study indicates that both essential oils from *P. caldense* tissues showed potential antimicrobial activity; and it was mainly the oil of the roots that was the more active. This is the first report of the biological activities and chemical composition of the essential oil of *P. caldense*.

Table 3. MIC of essential oils from the *P. caldense* root, stem and leaf.

Microorganisms	Root	Stem	Leaf	Gentamicin
Gram-negative bacteria				
µg/mL				
<i>E. coli</i>	325	-	-	5.0
<i>K. pneumonia</i>	325	-	750	5.0
<i>P. aeruginosa</i>	325	325	325	5.0
Gram-positive bacteria				
µg/mL				
<i>S. aureus</i>	325	-	750	5.0
<i>B. subtilis</i>	750	325	325	5.0

Experimental

Plant material

P. caldense leaf, stem and root were collected in Recife, Pernambuco state, northeastern Brazil, in January 2014. Botanical identification was made by Dr. Ângela M. M. Freitas (Department of Forest Sciences – Pernambuco Federal Rural University) and a voucher specimen was deposited in the Sérgio Tavares Herbarium of that university (HTS 18180).

Obtaining and analysis of the essential oils

The essential oils from fresh leaves, roots and stems were obtained by hydrodistillation using a Clevenger-type apparatus. 100 g of fresh plant tissue and 250 mL of H₂O were used and the distillation was carried out for 2 h after the mixture reached boiling. Traces of water present in the essential oils were removed by treatment with Na₂SO₄. The samples were kept at -20 °C in a freezer until further analysis. Yields were calculated from weight of fresh material. The essential oils were analyzed by GC-MS (60-240 °C at 3 °C min. rate) in a Shimadzu GCMS-QP5050A instrument using fused-silica capillary column (30 m × 0.25 mm i.d. × 0.25 µm.) coated with DB-5. MS spectra were obtained using electron impact at 70 eV with a scan interval of 0.5 s and fragments from 40 to 550 Da (all the samples were analyzed in triplicate).

Quantitative analysis of the chemical constituents was performed by flaming ionization gas chromatography (FID), using a Perkin-Elmer instrument, under the same conditions and with the same column as reported for the GC-MS. Identification of the components was based on GC retention indices with reference to a homologous series of C₇-C₃₀ n-alkanes calculated using the Van den Dool & Kratz [18] equation and by computer matching against the mass spectral library of the GC-MS data system (NIST 98 and Wiley) as well as other published mass spectra [19].

In vitro assay for antimicrobial activity

The antimicrobial activities of essential oils from the *P. caldense* leaf, stem and root were tested against the following microorganisms: Gram-positive bacteria: *S. aureus* (ATCC 6538), *B. subtilis* (ATCC 6633) and *E. faecalis* (ATCC 6057); Gram-negative bacteria: *E. coli* (ATCC 25922), *K. pneumoniae* (ATCC29665) and *P. aeruginosa* (ATCC by the Antibiotics Department of Pernambuco Federal University (UFPEDA) and maintained in nutrient agar (NA), stored at 4 °C. The antibacterial activity was reported preliminarily utilizing the disc diffusion method. In this method, disks containing known amounts of an antimicrobial agent were placed on the surface of an agar plate that had been inoculated with a standardized suspension of each microorganism tested. Paper discs with only ethanol were used as negative controls. All experiments were carried out three times and repeated if the results differed. A stock solution (10 mg/mL) of each test sample was prepared in ethanol

as the solvent. Furthermore, the serial dilution of the test compounds was carried out and the concentrations used ranged from 2500 to 19.5 µg/mL. Test samples at various concentrations were added to culture media in test tubes. All strains were provided different strains were inoculated at 108 bacteria/mL concentration. Tryptic soy agar and nutrient agar were utilized as culture media. The tubes were incubated at 37 °C for 24-48 h and then examined for the presence or absence of growth of the organisms. The MIC values were obtained from the lowest concentration of the test samples where the tubes remained clear, indicating that the bacterial growth was completely inhibited at this concentration. Gentamicin was used as antibacterial substance.

Acknowledgments

This work was funded by grants from FACEPE and CNPq. DSR thanks CAPES for providing a scholarship. The authors acknowledge Dr. Ângela M. M. Freitas (Department of Forest Sciences) for the identification of plant material.

References

1. Suwanphakdee, C.; Simpson, D. A. *Kew Bulletin*, **2012**, 67, 707-711.
2. Scott, I. M.; Jensen, H. R.; Philogène, B. J. R.; Arnason, J. T. *Phytochem. Rev.* **2008**, 7, 65-75.
3. do Nascimento, J. C.; De Paula, V. F.; David, J. M.; David, J. P. *Química Nova* **2012**, 35, 2288-2311.
4. Lima, L. M.; Rev. *Virtual Quim.* **2015**, 7, 495-538.
5. Barbosa, Q. P. S.; Guimarães, E. F.; Nascimento, D. C. O.; Lima-Filho, J. V.; Câmara, C. A. G.; Ramos CS. *Química Nova*, **2012**, 35, 1806-1808.
6. da Silva, M. F. R.; Bezerra-Silva, P. C.; de Lira, C. S.; de Lima Albuquerque, B. N.; Neto, A. C. A.; Pontual, E. V.; Maciel, J. R.; Paiva, P. M. G.; Navarro, D. M. D. A. F. *Exp. Parasitol.* **2016**, 165, 64-70.
7. Ramos, C.S.; Silva, A. M.; Lopes, M. S.; Batista-Pereira, L. G.; Kato, M. J. *Nat. Prod. Commun.* **2012**, 7, 1103-1106.
8. Moraes, M. M., Da Silva, T. M. G.; Da Silva, R. R.; Ramos, C. S.; Câmara, C. A. G. B. *Latinoam. Caribe PL.* **2014**, 13, 270-277.
9. Cardozo, J. E. L.; Chaves, M. C. O. *Pharmaceut. Biol.* **2003**, 41, 216-218.
10. Guimarães, E. F.; Monteiro, D. *Rodriguésia* **2006**, 57, 567-587.
11. Andrade, E. H. A.; Alves, C. N.; Guimarães, E. F.; Carreira, L. M. M.; Maia, J. G. S.; *Biochem. Syst. Ecol.* **2011**, 39, 669-675.
12. Costantin, M.B.; Sartorelli, P. T.; Limberger, R. J. P.; Henriques, A. T.; Steppe, M. P.; Ferreira, M. J.; Ohara, M. J.; Emerenciano, V. J.; Kato, M. J.; *Planta med.* **2001**, 67, 771-773.
13. Holecz, F. B.; Pessini, G. L.; Sanches, N. R.; Cortez, D. A. G.; Nakamura, C. V.; Dias-Filho, B. P. *Mem. do Inst. Oswaldo Cruz*, **97**, 1027-1031.
14. Guerrini, A.; Sacchetti, G.; Grandini, A.; Spagnoletti, A.; Asanza, M.; Scalvenzi, Laura; *Evid Based Complement Alternat Med.*, **2016**, 1638342
15. Oyedeji, O. A.; Adeniyi, B. A.; Ajayi, O.; König, W. A. *Phytother. Res.* **2005**, 9, 362-364.
16. Santos, T. G.; Rebelo, R.; Dalmarco, E. M.; Guedes, A.; Gasper, A. L.; Cruz, A. B.; Schmit, Ana Paula, C.; Rosana C. B.; Steindel, M.; Nunes, R. K.). *Química Nova*, **2012**, 35, 477-481.
17. Oliveira, R.; Assis, A.; Silva, L.; Andrioli, J.; Oliveira, F. *Chemistry of Natural Compounds*, **2016**, 52, 331-333
18. Kovats, E. S. *Adv. Chromatogr.* **1965**, 1, 229-247.
19. Adams, PR. **1995**. Allured Publishing Corporation, Carol Stream, Illinois.