



Associated psychological factors to chronic obstructive pulmonary disease: a narrative review

Factores psicológicos asociados a la enfermedad pulmonar obstructiva crónica: una revisión narrativa

Andrea Hernández-Pérez,*[‡] Inés Vargas-Núñez,[‡] Rogelio Pérez-Padilla,*[‡] Alejandra Ramírez-Venegas*

*Instituto Nacional de Enfermedades Respiratorias Ismael Cosío Villegas, Mexico City, Mexico;

[‡]Universidad Nacional Autónoma de México, Mexico City, Mexico.

ABSTRACT. Chronic obstructive pulmonary disease (COPD) is the third leading cause of global death, and the disease burden is further increased by multiple comorbidities, including some psychological disorders, like anxiety and depression, among others. This interplay between mental health and COPD has become a subject of intense study in recent years. Our objective was to describe the psychological factors associated with COPD, the interplay of factors leading to psychological impacts, and to analyze the effective treatment and intervention alternatives available. A narrative literature search was performed in PubMed and Cochrane Library, by using a snow-ball search technique to identify papers published on the subject. A search was conducted for relevant original articles with emphasis on years 2015-2021. Relevant was the emphasis of a multidisciplinary team to support patients with COPD, experts on psychological factors associated with the disease, especially on anxiety and depression. Treatments available, including psychological interventions demonstrated an improvement in quality of life, adherence to treatments, reduced hospitalizations, symptomatology and exacerbations.

Keywords: COPD, mental health, anxiety, depression, psychiatric disorder, risk factors.

INTRODUCTION

Currently, chronic obstructive pulmonary disease (COPD) is the fourth leading cause of death across the globe, while the World Health Organization (WHO) predicts that COPD will become the third leading cause of death by 2030.¹ Similarly as other chronic conditions, COPD negatively affects

RESUMEN. La enfermedad pulmonar obstructiva crónica (EPOC) es la tercera causa principal de muerte a nivel mundial, y la carga de la enfermedad aumenta por múltiples comorbilidades, incluidos algunos trastornos psicológicos, como ansiedad y depresión, entre otros. Esta interacción entre la salud mental y la EPOC se ha convertido en un tema de intenso estudio en los últimos años. Nuestro objetivo fue describir los factores psicológicos asociados con la EPOC, la interacción de factores que conducen a un impacto en la enfermedad y analizar las alternativas de tratamiento e intervención efectivas disponibles. Se realizó una búsqueda de literatura narrativa en PubMed y Cochrane Library, utilizando una técnica de búsqueda de bola de nieve para identificar artículos publicados sobre el tema. Se realizó una búsqueda de artículos originales relevantes con énfasis en los años 2015-2021. Fue relevante el énfasis de un equipo multidisciplinario de apoyo a los pacientes con EPOC, resulta relevante la evaluación de factores psicológicos asociados a la enfermedad, especialmente ansiedad y depresión. Los tratamientos disponibles, incluidas las intervenciones psicológicas, demostraron una mejora en la calidad de vida, adherencia a los tratamientos, reducción de hospitalizaciones, sintomatología y exacerbaciones.

Palabras clave: EPOC, salud mental, ansiedad, depresión, trastorno psiquiátrico, factores de riesgo.

quality of life and the disease burden is further increased by multiple comorbidities, including cardiovascular disease and cerebrovascular disease. Additionally, patients with COPD are two to three times more likely to have problems of mental health than the general population.² Patients with comorbid physical and mental ill health are less likely to be identified, diagnosed, and treated.³

Correspondence:

Dra. Andrea Hernández-Pérez

Instituto Nacional de Enfermedades Respiratorias Ismael Cosío Villegas, Mexico City, Mexico.

E-mail: andrea.hde@gmail.com

Received: 10-VI-2021; accepted: 04-X-2021.

How to cite: Hernández-Pérez A, Vargas-Núñez I, Pérez-Padilla R, Ramírez-Venegas A. Associated psychological factors to chronic obstructive pulmonary disease: a narrative review. *Neumol Cir Torax.* 2022; 81 (1): 35-40. <https://dx.doi.org/10.35366/105530>

Among patients with COPD, various psychological aspects of functioning have been investigated but so far with little consequence for clinical practice.⁴ In the most recent Clinical Practice Guidelines, anxiety and depression are included as frequent comorbidities in COPD and are associated with poor health status and prognosis. Importantly, comorbidities with symptoms also associated with COPD may be overlooked e.g., heart failure and lung cancer (breathlessness) or depression (fatigue and reduced physical activity).⁵

The objective of the present review was to integrate the psychological factors associated with COPD and analyze the effective treatment alternatives available. A search was conducted in January and February 2021 with no language restrictions under the following criteria: [COPD OR chronic obstructive pulmonary disease] [COPD-mental health], [psychopathology-COPD], [depression-depressive disorder COPD], and [anxiety- anxiety disorders COPD] to identify manuscripts related to the objective. Inclusion criteria were the diagnosis of COPD and at least one psychological factor with a clearly indicated assessment. We searched for other potentially relevant studies by screening the reference lists and citations of included studies. Finally, all authors reviewed, analyzed using the GRADE system, and discussed the articles to provide recommendations. We present the following article in accordance with the narrative review reporting checklist.⁶

Anxiety

The morbidity rate of anxiety in non-hospitalized settings of COPD patients has varied from 13 to 46%, and in hospitalized settings from 10 to 55%.⁷

The coexistence of anxiety disorders and COPD has been shown in a worsening of symptoms, especially the shortness of breath or cough, but also leads to a higher rate of hospital care and increased doses of drugs such as bronchodilators, inhaled corticosteroids, and antibiotics. As an additional result, it causes a higher incidence and severity of adverse effects, accompanied by a worse overall prognosis.^{8,9}

According to the retrospective analysis of a randomized controlled trial found that more than half of trial participants (of a total 1200 responses) reported mood disturbance during the study, participants reported symptoms of anxiety and low mood at least once during the previous 12 months.¹⁰

The available evidence suggests that less than one-third of COPD patients with anxiety are receiving appropriate treatment. Untreated comorbid anxiety and depression in patients with COPD have devastating consequences, as they tend to overwhelm the coping strategies of COPD patients and their caregivers, and may increase healthcare utilisation.¹¹

Table 1: Summary of publications regarding the association between psychological factors and pulmonary obstructive chronic disease.

Outcome measures/Reference	Odds ratio - risk ratio - adjusted odds ratio - hazard ratio	Other factors found
Depression and emergency hospital admissions ¹²	OR = 2.63; 95% CI (1.48-4.66)	Even mild symptoms and moderate to severe symptoms of depression increase the risk of use of urgent care, these findings are independent of severity of disease and comorbidity of other chronic physical conditions
Depression and attendances at emergency departments	OR = 2.78; 95% CI (1.55-4.99)	
COPD and the likelihood of suicide ¹³	OR = 1.90; 95% CI (1.27-2.48)	Similarly, risk factors associated with suicide, such as mental disorders, are underdiagnosed and undertreated
Comorbid depression and risk of mortality ¹⁴	RR = 1.83; 95% CI (1.00-3.36).	COPD increases the risk of developing depression (RR = 1.69; 95% CI 1.45-1.96)
Bipolar disorder and COPD ¹⁵ Schizophrenia and COPD ¹⁶	OR 1.55; 95% CI (1.45-1.65) OR 1.57; 95% CI (1.43-1.72)	The most important confounding factor the contribution of which must be evaluated to clarify the nature of the association between COPD and major mental illness is the smoking
Stress with acute care use in COPD	AOR = 2.51; 95% CI (1.06-5.98)	The high stress group had a 2.5-fold increased adjusted odds of acute care use compared to the low stress group
Physical activity-depression in COPD ¹⁷ PA-anxiety in COPD	HR = 0.85; 95% CI (0.75-0.95) HR = 0.89; 95% CI (0.79-1.00)	The study found more physically active patients [with higher PA levels equivalent to 2.5 metabolic equivalent tasks (METs), e.g., those who report walking outside vs. those who do not] had 15% and 11% lower risks of developing depression and anxiety, respectively, compared to physically inactive patients

OR = Odds ratio, RR = risk ratio, AOR = adjusted odds ratio, HR = hazard ratio, COPD = pulmonary obstructive chronic disease, PA = physical activity, CI = confidence intervals.




Figure 1:

Illustrative model of biopsychosocial factors that interact in pulmonary obstructive chronic disease and its adverse health outcomes. Modified from Latent Health Risk Classes Associated with pulmonary obstructive chronic disease.²² COPD = pulmonary obstructive chronic disease.

Depression

In a prospective longitudinal study of 355 patients with COPD it was found that depression was a predictor of emergency care in COPD, independent of severity of disease or physical comorbidity. Even mild symptoms of depression increase the risk of care by more than twofold, and moderate to severe symptoms of depression increase the risk by nearly five times (*Table 1*).¹²

A systematic review has shown that COPD patients are 1.9 times more likely to commit suicide than people without COPD (*Table 1*).¹³ However, it is well known that the association between clinically relevant depression and COPD is bidirectional: a meta-analysis demonstrated that not only COPD increases the risk of developing depression (relative risk RR, 1.69; 95% CI, 1.45-1.96) but also, depression increases the risk of COPD adverse outcomes and mortality (RR, 1.43; 95% CI, 1.20-1.71) (*Table 1*).¹⁴ In a similar manner, an association between COPD, schizophrenia and bipolar disorder has been described. A systematic review found that patients suffering from schizophrenia were significantly more likely to have comorbid COPD as were patients suffering from bipolar disorder.¹⁵ It has also been described that acute care use and stress are associated in COPD. These associations are more pronounced in the low-income more high stress population who disproportionately contribute to health care utilization.¹⁶ It has been reported as a protective factor to physical activity, in COPD patients, those who perform a great physical activity are less likely to develop symptoms of depression or anxiety in the long term (*Table 1*).¹⁷

Moreover, as a risk factor, the evidence shows that treatment adherence in patients with COPD has a

significant negative correlation with depression. In a study about medication adherence among patients with COPD treated in a primary general hospital during the COVID-19 pandemic showed that COPD with possible depression represented 31% of all cases (191 patients), and possible depression was an independent risk factor for poor treatment adherence over the past two months.¹⁸

These research data highlight the importance of the proper assessment of the screening tools when measuring depression-anxiety and its appropriate intervention in COPD patients.

Other psychiatric disorders

In addition to mood disorders, patients with COPD are more susceptible to other psychiatric disorders.

One third of patients with COPD reported post-traumatic stress disorder (PTSD) symptoms and met criteria for PTSD. In a study with multivariable analysis model, the presence of two or more exacerbations led to a near twofold increase in the prevalence of post-traumatic stress symptoms related to PTSD (PR, 1.71; $p = 0.015$) specially in those requiring hospitalization (PR, 1.13; $p = 0.030$). Overall, these findings suggest that psychological domains should be addressed along with exacerbations in COPD patients.¹⁹

Moreover, a study demonstrated a reduced survival in COPD if psychiatric comorbidity was present including substance addiction, schizophrenia, bipolar affective disorder, depressive episode, depressive disorder, anxiety episode, behavioral syndromes associated with physiological disturbances and disorders of personality.²⁰

An high score on the type A personality scale, and an increase in risk propensity were associated with dyspnea,

and a decrease in empathy score was predictor of number of infections in men with COPD.²¹

The biopsychosocial factors associated with COPD and their associated adverse health outcomes are illustrated in *Figure 1*, modified from latent health risk classes associated with COPD.²² *Figure 2* shows some of the most common symptoms in COPD with their associated psychological factors, based on comorbidities of chronic obstructive pulmonary disease.

Psychological interventions for patients with COPD

The first-line treatment recommended in the COPD clinical practice guidelines is cognitive behavioral therapy (CBT). The CBT has recollated scientific evidence related to its effectiveness and has proven to be a cost-effective alternative. A meta-analysis and systematic review which analyzed randomized controlled trials to evaluate the effect of CBT on anxiety and depression in patients with COPD, showed that CBT can effectively improve anxiety and depression, CBT can change patients' wrong cognitive ideas through communication and emotional management, eliminate patients' negative emotions such as negative pessimism and low self-esteem, help patients form a healthy lifestyle and improve their quality of life.²³

According to a meta-analysis CBT can serve as a complementary therapy to improve anxiety, depression, visits to emergency departments and quality of life in COPD patients and deserves more widespread application in clinical practice.²⁴

A Cochrane review analyzing the benefits of psychological therapies for the treatment of anxiety disorders in patients with COPD pointed out that studies were heterogeneous, treatment efficacy was inconclusive, and overall quality of evidence was low.²⁵ Therefore, well-controlled randomized trials are needed. In another more recent Cochrane review indicated that psychological therapies (using a CBT-based approach) may be effective for treating COPD-related depression; the patients improved more in the intervention (CBT) groups compared to: 1) no intervention (attention placebo or standard care), 2) educational interventions, and 3) a co-intervention (pulmonary rehabilitation). However, the effect sizes were small and quality of the evidence very low due to clinical heterogeneity and risk of bias.²⁶

Group psychological treatments, aimed at improving functional social support, have been found to promote expansion in peer networks, since social comparison and interaction with other people with COPD offer learning opportunities, a sense of validation of the lived experience and an opportunity to make new friends, promoting the sense of belonging and identity to a group.²⁷

In a randomized controlled trial conducted by Farver-Vestergaard et al.,²⁸ on a group intervention that integrates

mindfulness meditation with elements of cognitive-behavioral therapy, called mindfulness-based cognitive therapy for patients with COPD; was shown to be capable of achieving a statistically significant and lasting effect on psychological distress, indicating that it may be an effective complement to standard programs for patients with COPD.

The National Institute for Clinical Excellence (NICE) advises clinicians to 'ask people with COPD if they experience breathlessness, as they find it frightening. If they do, consider including a cognitive behavioral component in their self-management plan to help them manage anxiety.'²⁹

On the other hand, a meta-analysis examined the efficacy of digital interventions on the outcomes of psychological comorbidities (depression and anxiety) related to a specific group of chronic diseases in adult populations (cardiovascular disease, stroke, diabetes, and COPD) revealing an overall moderate and significant effect on the depression outcome. However the effect on anxiety was small and non-significant.³⁰

A survey eliciting clinician and patient perspectives on what is appropriate for ongoing health crisis, involved clinicians (n = 55) and patients with COPD (n = 19) and respondents agreed that there are activities appropriate for remote provision such as: planned activities (gathering patient information on COPD and health status, providing counseling on smoking cessation, and providing education




Figure 2: A proposed mapping of pulmonary obstructive chronic disease symptomatology and psychological factors associated (based on comorbidities of chronic obstructive pulmonary disease). COPD = pulmonary obstructive chronic disease.

on COPD or developing a self-management plan) and urgent care activities (triaging patients for face-to-face care and initiating use of rescue packs).³¹

CONCLUSIONS

Psychological comorbidities are very common in COPD, and adversely affect well-being, and quality of life. Multidisciplinary interventions including nurses, psychologists, rehabilitators, occupational therapists, home care services especially working as a team are effective in improving the condition of patients and the prognosis of evolution.

Still to be defined is to measure the impact of comprehensive care of patients with COPD, in quality of life, in prevention of exacerbations, reduction of dyspnea, adherence to treatment and survival. Thus, awareness of the importance of timely screening for these conditions should be promoted among clinicians and heightened attention should be paid to modifiable risk factors.

Psychosocial factors related to COPD should be considered for comprehensive care. Clinicians should strengthen patient supervision and monitoring; encourage integrated medical teams including pharmacists, nurses, and psychologists; and improve disease management, as well consider humane and empathetic care with patients, further studies are needed and should be focused on enhancement of adjustment to illness, preserving mental health and quality of life related to health.

Acknowledgments

We thank the Council of Science and Technology Mexico (CONACYT) for a doctoral grant that AH-P received [granter number: 286181].

REFERENCES

1. WHO | Burden of COPD. [Accessed January 24, 2021] Available in: <https://www.who.int/respiratory/copd/burden/en/>
2. Divo M, Celli BR. Multimorbidity in patients with chronic obstructive pulmonary disease. *Clin Chest Med.* 2020;41(3):405-419. doi: 10.1016/j.ccm.2020.06.002.
3. Management of Physical Health Conditions in Adults with Severe Mental Disorders WHO GUIDELINES.
4. Rzadkiewicz M, Nasiłowski J. Psychosocial interventions for patients with severe COPD-an up-to-date literature review. *Med.* 2019;55(9):597. doi: 10.3390/medicina55090597.
5. 2021 GOLD Reports - Global Initiative for Chronic Obstructive Lung Disease - GOLD. [Accessed January 16, 2021] Available in: <https://goldcopd.org/2021-gold-reports/>
6. Green BN, Johnson CD, Adams A. Writing narrative literature reviews for peer-reviewed journals: secrets of the trade. *J Chiropr Med.* 2006;5(3):101-117. doi: 10.1016/S0899-3467(07)60142-6.
7. Yohannes AM, Kaplan A, Hanania NA. Anxiety and depression in chronic obstructive pulmonary disease: recognition and management. *Cleve Clin J Med.* 2018;85(2):S11-S18. doi: 10.3949/ccjm.85.s1.03.
8. Pooler A, Beech R. Examining the relationship between anxiety and depression and exacerbations of COPD which result in hospital admission: a systematic review. *Int J COPD.* 2014;9:315-330. doi: 10.2147/COPD.S53255.
9. Uchmanowicz I, Jankowska-Polańska B, Motowidło U, Uchmanowicz B, Chabowski M. Assessment of illness acceptance by patients with COPD and the prevalence of depression and anxiety in COPD. *Int J Chron Obstruct Pulmon Dis.* 2016;11(1):963. doi: 10.2147/COPD.S102754.
10. Whelan ME, Velardo C, Rutter H, Tarassenko L, Farmer AJ. Mood monitoring over one year for people with chronic obstructive pulmonary disease using a mobile health system: Retrospective analysis of a randomized controlled trial. *JMIR mHealth uHealth.* 2019;7(11):e14946. doi: 10.2196/14946.
11. Yohannes AM, Alexopoulos GS. Depression and anxiety in patients with COPD. *Eur Respir Rev.* 2014;23(133):345-349. doi: 10.1183/09059180.00007813.
12. Blakemore A, Dickens C, Chew-Graham CA, et al. Depression predicts emergency care use in people with chronic obstructive pulmonary disease: a large cohort study in primary care. *Int J COPD.* 2019;14:1343-1353. doi: 10.2147/COPD.S179109.
13. Sampaio MS, Vieira W de A, Bernardino Í de M, Herval ÁM, Flores-Mir C, Paranhos LR. Chronic obstructive pulmonary disease as a risk factor for suicide: a systematic review and meta-analysis. *Respir Med.* 2019;151:11-18. doi: 10.1016/j.rmed.2019.03.018.
14. Atlantis E, Fahey P, Cochrane B, Smith S. Bidirectional associations between clinically relevant depression or anxiety and COPD: a systematic review and meta-analysis. *Chest.* 2013;144(3):766-777. doi: 10.1378/chest.12-1911.
15. Zareifopoulos N, Bellou A, Spiropoulou A, Spiropoulos K. Prevalence of comorbid chronic obstructive pulmonary disease in individuals suffering from schizophrenia and bipolar disorder: a systematic review. *COPD J Chronic Obstr Pulm Dis.* 2018;15(6):612-620. doi: 10.1080/15412555.2019.1572730.
16. Parekh TM, Cherrington AL, Bhatia S, et al. The association of low income and high stress with acute care use in COPD patients. *Chronic Obstr Pulm Dis.* 2020;7(2):107-117. doi: 10.15326/jcopdf.7.2.2019.0165.
17. Yu T, Ter Riet G, Puhan MA, Frei A. Physical activity and risk of comorbidities in patients with chronic obstructive pulmonary disease: a cohort study. *npj Prim Care Respir Med.* 2017;27(1):36. doi: 10.1038/s41533-017-0034-x.
18. Zhang H-Q, Lin J-Y, Guo Y, Pang S, Jiang R, Cheng QJ. Medication adherence among patients with chronic obstructive pulmonary disease treated in a primary general hospital during the COVID-19 pandemic. *Ann Transl Med.* 2020;8(18):1179-1179. doi: 10.21037/atm-20-6016.
19. Teixeira PJZ, Porto L, Kristensen CH, Santos AH, Menna-Barreto SS, Do Prado-Lima PAS. Post-traumatic stress symptoms and exacerbations in COPD patients. *COPD J Chronic Obstr Pulm Dis.* 2015;12(1):90-95. doi: 10.3109/15412555.2014.922063.
20. Sikjær MG, Løkke A, Hilberg O. The influence of psychiatric disorders on the course of lung cancer, chronic obstructive pulmonary disease and tuberculosis. *Respir Med.* 2018;135:35-41. doi: 10.1016/j.rmed.2017.12.012.
21. Witusik A, Mokros L, Kuna P, Nowakowska-Domagala K, Antczak A, Pietras T. Type a behavior pattern, impulsiveness, risk propensity, and empathy as predictors of dyspnea and number of infections in men

- with chronic obstructive pulmonary disease: a cross-sectional study. *Med Sci Monit.* 2018;24:3832-3839. doi: 10.12659/MSM.907742.
22. Stollefson M, Wang MQ, Balanay JAG, Wu R, Paige SR. Latent health risk classes associated with poor physical and mental outcomes in workers with copd from central Appalachian U.S. States. *Int J Environ Res Public Health.* 2020;17(18):1-18. doi: 10.3390/ijerph17186798.
 23. Zhang X, Yin C, Tian W, Lu D, Yang X. Effects of cognitive behavioral therapy on anxiety and depression in patients with chronic obstructive pulmonary disease: A meta-analysis and systematic review. *Clin Respir J.* 2020;14(10):891-900. doi: 10.1111/crj.13226.
 24. Ma RC, Yin YY, Wang YQ, Liu X, Xie J. Effectiveness of cognitive behavioral therapy for chronic obstructive pulmonary disease patients: a systematic review and meta-analysis. *Complement Ther Clin Pract.* 2020;38. doi: 10.1016/j.ctcp.2019.101071.
 25. Usmani ZA, Carson K V., Heslop K, Esterman AJ, De Soyza A, Smith BJ. Psychological therapies for the treatment of anxiety disorders in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev.* 2017;2017(3):CD010673. doi: 10.1002/14651858.CD010673.pub2.
 26. Pollok J, Van Agteren JEM, Esterman AJ, Carson-Chahhoud K V. Psychological therapies for the treatment of depression in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev.* 2019;2019(3): CD012347. doi: 10.1002/14651858.CD012347.pub2.
 27. Russell S, Ogunbayo OJ, Newham JJ, et al. Qualitative systematic review of barriers and facilitators to self-management of chronic obstructive pulmonary disease: Views of patients and healthcare professionals. *NPJ Prim Care Respir Med.* 2018;28(1):1-13. doi: 10.1038/s41533-017-0069-z.
 28. Farver-Vestergaard I, O'Toole MS, O'Connor M, et al. Mindfulness-based cognitive therapy in COPD: A cluster randomized controlled trial. *Eur Respir J.* 2018;51(2):1702082. doi: 10.1183/13993003.02082-2017.
 29. Hopkinson NS, Molyneux A, Pink J, Harrisingh MC. Chronic obstructive pulmonary disease: Diagnosis and management: Summary of updated NICE guidance. *BMJ.* 2019;366. doi: 10.1136/bmj.l4486.
 30. Maisto M, Diana B, Di Tella S, et al. Digital interventions for psychological comorbidities in chronic diseases-a systematic review. *J Pers Med.* 2021;11(1):1-14. doi: 10.3390/jpm11010030.
 31. Wu F, Burt J, Chowdhury T, et al. Specialty COPD care during COVID-19: Patient and clinician perspectives on remote delivery. *BMJ Open Respir Res.* 2021;8(1): e000817. doi: 10.1136/bmjresp-2020-000817.

Conflict of interest: The authors declare to have no conflict of interest directly or indirectly related to the manuscript contents.

Funding: This research has not received any specific grant from agencies in the public, commercial, or non-profit sectors.