

**SILICON INDUCED POSITIVE
RESPONSE AT BIOCHEMICAL AND
GENE EXPRESSION LEVEL IN
TOMATO PLANTS INOCULATED
WITH *Fusarium oxysporum***

**EL SILICIO INDUCE RESPUESTAS
POSITIVAS A NIVEL BIOQUÍMICO Y
DE EXPRESIÓN DE GENES EN
PLANTAS DE TOMATE INOCULADAS
CON *Fusarium oxysporum***

López-Pérez, M.C.; F. Pérez-Labrada; Y. González-García y A. Juárez-Maldonado

SILICON INDUCED POSITIVE RESPONSE AT BIOCHEMICAL AND GENE
EXPRESSION LEVEL IN TOMATO PLANTS INOCULATED WITH *Fusarium oxysporum*

EL SILICIO INDUCE RESPUESTAS POSITIVAS A NIVEL BIOQUÍMICO Y DE
EXPRESIÓN DE GENES EN PLANTAS DE TOMATE INOCULADAS CON *Fusarium
oxysporum*



Silicon induced positive response at biochemical and gene expression level in tomato plants inoculated with *Fusarium oxysporum*

El silicio induce respuestas positivas a nivel bioquímico y de expresión de genes en plantas de tomate inoculadas con *Fusarium oxysporum*

López-Pérez, M.C.;
F. Pérez-Labrada;
Y. González-García
y A. Juárez-Maldonado

SILICON INDUCED
POSITIVE RESPONSE AT
BIOCHEMICAL AND GENE
EXPRESSION LEVEL IN
TOMATO PLANTS
INOCULATED WITH
Fusarium oxysporum

EL SILICIO INDUCE
RESPUESTAS POSITIVAS A
NIVEL BIOQUÍMICO Y DE
EXPRESIÓN DE GENES EN
PLANTAS DE TOMATE
INOCULADAS CON
Fusarium oxysporum

POLIBOTÁNICA

Instituto Politécnico Nacional

Núm. 57: 213-236. Enero 2024

DOI:
10.18387/polibotanica.57.13

Mari Carmen López-Pérez / mcloper@outlook.com

<https://orcid.org/0000-0002-7694-5102>

Unidad Académica Navojoa, Universidad Estatal de Sonora, 810
Manlio Fabio Beltrones Blvd., Navojoa 85875, Sonora, México

Fabián Pérez-Labrada / fabperlab@outlook.com

<https://orcid.org/0000-0003-3540-303X>

Departamento de Botánica, Universidad Autónoma Agraria Antonio Narro,
Saltillo 25315, México

Yolanda González-García / yolanda_glezg@hotmail.com

Centro de Agricultura Protegida, Facultad de Agronomía,
Universidad Autónoma de Nuevo León, General Escobedo 66050, México

Antonio Juárez-Maldonado / antonio.juarez@uaaan.edu.mx

<https://orcid.org/0000-0003-3061-2297>

Departamento de Botánica, Universidad Autónoma Agraria Antonio Narro,
Saltillo 25315, México

ABSTRACT: Tomato plants infested by *Fusarium oxysporum* (Fol) show limitations in yield and fruit quality. The application of silicic acid could be used as an alternative to mitigate damage caused by this pathogen. In this work, the effect of silicic acid on "Rio Grande" variety tomato plants inoculated with Fol was studied. Silicic acid was applied via foliar weekly on tomato plant (ten applications). Treatments consisted of silicic acid + Fol, silicic acid, Fol and an absolute control. The incidence and severity of Fol wilt, content of antioxidant compounds (lycopene, β carotene, chlorophylls, vitamin C, reduced glutathione, phenols, flavonoids, total antioxidants and proteins), enzymatic activity (APX, CAT, GPX and SOD), PAL, salicylic and jasmonic acid, and the expression of genes involved in the synthesis of *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* and *ABA*, were determined. ASi+Fol-treated plants, compared to Fol-inoculated plants, disease incidence Fol wilt was reduced by 32% and severity by 22%, also presented an increase in chlorophyll *a*, *b* and total (49.1, 59.4 and 46.7%, respectively), as well as flavonoids (24.6%), the plants overexpressed the genes that participate in the synthesis of *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* and *ABA* at 31 dat (days after transplant), induced a reduction in reduced glutathione, antioxidant capacity by DPPH, CAT, jasmonic acid and the expression of all the genes evaluated in this work at 73 dat. The plants treated with ASi, in comparison with Fol, presented an increase in proteins, lycopene, vitamin C (30.3, 65.0, 52.7%, respectively), chlorophylls and the expression of *PAL5-3* (31 dat), *GAME1* and *ET* (73 dat). In the case Fol, a higher concentration of vitamin C, phenols, DPPH, PAL, CAT, GPX, salicylic and jasmonic acid was observed, and increase in the expression of *JA*, *GAME1*, *PAL5-3*, and *ABA* (73 dat), but it also induced a reduction in the concentration of flavonoids, chlorophylls and the expression of the six genes determined in this work at 31 dat. On the other hand, T0 plants, compared to Fol, showed an increase in proteins, lycopene, flavonoids (33.1, 62.0, 27.8%, respectively), as well as, chlorophyll, and the expression of all the genes analyzed in 31 dat, except for *ET*. The foliar application of silicic acid decreased the negative effects caused by Fol through

the increase of antioxidant compounds and changes in the expression of genes related to the defense system in tomato plants.

Key words: antioxidant compounds, biotic stress, defense genes expression, enzymatic activity, phytohormones.

RESUMEN: Las plantas de tomate infestadas por *Fusarium oxysporum* (Fol) muestran limitaciones en rendimiento y calidad de frutos. La aplicación de ácido silícico podría ser una alternativa para mitigar los daños causados por este patógeno. En este trabajo se estudió el efecto del ácido silícico en plantas de tomate variedad “Rio Grande” inoculadas con Fol. Se aplicó ácido silícico vía foliar semanalmente en plantas de tomate (diez aplicaciones). Los tratamientos consistieron en ácido silícico + Fol, ácido silícico, Fol y un control absoluto. Se determinó la incidencia y severidad de la marchitez de Fol, el contenido de compuestos antioxidantes (licopeno, β caroteno, clorofilas, vitamina C, glutatión reducido, fenoles, flavonoides, antioxidantes totales y proteínas), actividad enzimática (APX, CAT, GPX y SOD), PAL, ácido salicílico y jasmónico, y la expresión de genes que participan en la síntesis de *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* y *ABA*. En las plantas tratadas con ASi+Fol, en comparación con las plantas inoculadas con Fol, se redujo la incidencia de la enfermedad en un 32% y la severidad en un 22%. También presentaron un incremento de clorofila *a*, *b* y total (49.1, 59.4 y 46.7%, respectivamente), así como de flavonoides (24.6%), las plantas sobreexpresaron los genes que participan en la síntesis de *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* y *ABA*, a los 31 ddt, indujeron una reducción de glutatión reducido, capacidad antioxidante por DPPH, CAT, ácido jasmónico y la expresión de todos los genes evaluados en este trabajo a los 73 ddt. Las plantas tratadas con ASi, en comparación con Fol, presentaron un incremento de proteínas, licopeno, vitamina C (30.3, 65.0, 52.7%, respectivamente), clorofilas y la expresión de *PAL5-3* (31 ddt), *GAME1* y *ET* (73 ddt). En el caso de Fol, se apreció mayor concentración de vitamina C, fenoles, DPPH, PAL, CAT, GPX, ácido salicílico y jasmónico, y ligeramente un incremento de la expresión de *JA*, *GAME1*, *PAL5-3* y *ABA* (73 ddt), pero también indujo una reducción en la concentración de flavonoides, clorofilas y la expresión de los seis genes determinados en este trabajo a 31 ddt. Por su parte las plantas T0, comparado con Fol, se observó aumento de proteínas, licopeno, flavonoides (33.1, 62.0, 27.8%, respectivamente), así como clorofilas, y la expresión de todos los genes analizados a los 31 ddt, a excepción de *ET*. La aplicación foliar de ácido silícico disminuyó los efectos negativos causados por Fol a través del aumento de compuestos antioxidantes y cambios en la expresión de los genes relacionados con el sistema de defensa en plantas de tomate. **Palabras clave:** actividad enzimática, compuestos antioxidantes, estrés biótico, expresión de genes de defensa, fitohormonas.

INTRODUCTION

Fusarium oxysporum (Fol) causes plant wilting disease characterized by severe vascular damage and root rot, attacking a wide variety of crops of high human consumption, in tomato (*Solanum lycopersicum* L.) can induce serious yield losses (de Lamo & Takken, 2020). This pathogen increases reactive oxygen species (ROS) causing the antioxidant system to be unable to reduce such accumulation, which causes mutation of nucleic acids, protein oxidation and cellular damage (Sousa *et al.*, 2015). These events cause a reduction in the availability, absorption, distribution and use of nutrients in plants (Romero *et al.*, 2011), as well as photosynthesis decrease (Mo *et al.*, 2017), thus reducing the growth and productivity of tomato plants as well as the quality of the fruits (Leyva-Mir *et al.*, 2013). To mitigate this damage chemical control is the main management measure of this phytopathogenic disease (Solarte *et al.*, 2012). Unfortunately, the indiscriminate application of chemicals has originated the development of resistant fungal populations (Ramírez *et al.*, 2020). Thus, the implementation of alternatives to decrease the incidence and severity of Fol wilt of the disease induced by *F. oxysporum* to avoid crop losses is of great importance.

One of the most abundant elements in the lithosphere is silicon (Si). In soil solution, Si is present as silicic acid (H_4SiO_4 , $\text{pK}_a = 9.25$), which pH lower than 9, is the only form absorbed by the root (López-Pérez *et al.*, 2018; Paye *et al.*, 2018; van den Berg *et al.*, 2021). Uptake this element into crop is accomplished by rejective, passive and active processes, is transported via the xylem through the transpiration current with the help of transporters, however, it is species-specific and complex (Frick *et al.*, 2020; H. Sun *et al.*, 2020). Supplement of this mineral is important because optimizes plants growth, productivity and quality parameters, in addition to biotic stress resistance (Dawa *et al.*, 2020; Malhotra *et al.*, 2016). The function of silicic acid against diseases induced by fungi, bacteria, and virus is given by induction of multiple mechanisms (Hoffmann *et al.*, 2020). Physical barrier is occurs through silicic acid deposition and polymerization in cell structures (Islam *et al.*, 2020), biochemical defense is constitutes by phenylalanine ammonia lyase (PAL); who has an important role in the synthesis of lignin and phenolic compounds, and antioxidant enzymes production as peroxidase (POD), ascorbate peroxidase (APX), catalase (CAT), superoxide dismutase (SOD) (Gulzar *et al.*, 2021), in addition to non-enzymatic antioxidants, as phenolic compounds, flavonoids, among others (Jafari *et al.*, 2015), as well as, phytohormones, and expression of pathogenesis-induced plant defense-related genes, as salicylic, jasmonic, abscisic acid, among others (Bhatt & Sharma, 2018). In different research studies, the effect of Si supplementation in agricultural crops has been reported. The calcium silicate application controlled the disease caused by *Fusarium solani* in *Piper nigrum* (D'Addazio *et al.*, 2020). The same pathogen also was controlled with the supply of silicon dioxide nanoparticles to *Daucus carota* (Siddiqui *et al.*, 2020). Ouzounidou *et al.* (2016) found that potassium silicate supplementation caused increased antioxidant content in *Cucumis sativus*. Similarly, Madany *et al.* (2020) found positive effect of silicon nanoparticles on *Lycopersicon esculentum* by enhanced enzymatic and non-enzymatic antioxidant activity.

Considering this information, the study aimed to evaluate the application of silicic acid, on antioxidant capacity, salicylic and jasmonic acid and defense genes expression in tomato crop inoculated with *F. oxysporum*.

METHODS

Crop development

The development of the experiment was carried out in a greenhouse multi tunnel type, with polyethylene cover. As a biological material, tomato seedlings (*Solanum lycopersicum* L.) cv. "Rio Grande" was used, which were transplanted in 10 L black polyethylene bags, which contained substrate of peat moss and perlite in 1:1 proportion. For the nutritional management of the tomato crop a Steiner fertilizer solution was applied (Steiner, 1961). The tomato crop was developed during 119 days after transplanting (dat).

Silicic acid application

The application of silicic acid (ASi, 99.9%, Sigma-Aldrich, St. Louis, MO, USA) consisted of foliar spraying of this compound on the abaxial and adaxial sides of the whole plant; it was applied to following treatments Asi and Asi+Fol (18 plants each). A total of 10 applications of ASi were carried out, two applications were made prior to the inoculation of the pathogens (7 and 14 dat), and eight subsequent applications with intervals of one week. The concentration of silicic acid used was 110 g ha^{-1} per application.

Inoculation of *Fusarium oxysporum*

Fusarium oxysporum strain (Fol) was used as inoculum; strain from the Laboratory of Molecular Parasitology, Universidad Autónoma Agraria Antonio Narro (Limon-Corona *et al.*, 2022). To carry out the inoculation of *F. oxysporum* (Fol), it was first reactivated, with the objective of increasing the viability of the latent inoculum (Coninck *et al.*, 2020), in potato dextrose agar (PDA), and incubated for 15 days at 27°C (Benhamou & Bélanger, 1998). For the increase of the pathogen conidia a semiliquid medium was used mixing potato, dextrose, PDA, streptomycin

and tomato seedling; in this medium two mycelial discs ($\Theta = 6$ mm) were sowing and it was incubated at 28 °C in an orbital shaker at $168 \times g$ 125 rpm by 15 days. At 21 days after transplantation, the tomato plants of the Fol and ASi+Fol treatments were inoculated at a concentration of 1×10^6 spores mL^{-1} resuspended in distilled water, applying 60 mL of the conidia solution on perforations in the substrate (10 cm deep and 3 cm away from the stem) (Benhamou & Bélanger, 1998). Fol inoculum was not applied to the plants of treatments ASi and T0.

Description of treatments

Four treatments were evaluated ASi+Fol, ASi, Fol and T0. ASi+Fol treatment consisted of the inoculation of Fol plus the foliar application of silicic acid. ASi treatment consisted of the foliar application of silicic acid. Fol treatment consisted of the inoculation of Fol without Asi application. Finally, T0 (absolute control) treatment was not inoculated with Fol and Asi was not sprayed. A total of 18 plants were used per treatment.

Incidence and severity of *Fusarium oxysporum* wilt disease

Incidence and severity of Fol wilt parameters were recorded at 43, 54, 58, 61, 66, 70, 72, 74, 76, 78, 82 y 85 dat. Disease incidence on tomato plants was determined visually by the presence or absence of symptomatic plants. Therefore, zero represented asymptomatic plants and 1 represented symptomatic plants. All data were expressed in percentage. The severity of Fol wilt disease was determined following the visual scale of Diener & Ausubel, (2005) and González-García *et al.* (2023); which is described as follows: 0: plants indistinguishable from mock; 1 = wilting of basal leaves (from the fifth true leaf); 2 chlorosis of basal leaves (from the fifth true leaf) and wilting of young leaves; 3 = wilting of most of the leaves, diffuse desiccation and yellowing; 4 = dead plant. Disease severity is calculated as a percentage using the following equation:

$$\text{Disease severity } \% = \frac{100 \times VS}{4} \quad (1)$$

Biochemical variables

The biomolecules: proteins, catalase, ascorbate peroxidase, superoxide dismutase, glutathione peroxidase, phenylalanine ammonium lyase, reduced glutathione, ABTS, and DPPH antioxidant capacity were extracted of leaf tissue. For this purpose, after 73 days of transplantation, random plants were selected, and the third fully expanded young leaf was taken for biochemical analysis. The tissue was lyophilized, macerated in a mortar and stored (-80 °C) for subsequent analyses. For the enzymatic and non-enzymatic determination, 200 mg of lyophilized and macerated leaves of each treatment and 20 mg of polyvinylpyrrolidone were weighed. After this, 1.5 mL of phosphate buffer with a pH of 7-7.2 (0.1 M) were added, and the mixture was then subjected to micro-centrifugation at $16,128 \times g$ 12,000 rpm for 10 min at 4 °C. The supernatant was filtered with a 0.45 μm pore size nylon membrane (Ramos *et al.*, 2010).

Total protein quantification was determined by the method of Bradford, (1976), the results were expressed as mg g^{-1} dry weight (DW). Catalase (EC 1.11.1.6) activity was quantified by measuring 2 reaction times according to Dhindsa *et al.* (1981), the results were expressed as U g^{-1} TP (U per gram of total proteins) of DW, where U is equal to the mmol equivalent of H_2O_2 consumed per milliliter per minute. Ascorbate peroxidase (EC 1.11.1.1) was determined according to Nakano & Asada, (1981), the results were expressed as U g^{-1} TP of DW, where U is equal to the μmol equivalent of oxidized ascorbate per milliliter per minute. Superoxide dismutase (EC 1.15.1.1) was determined was carried out using the Cayman kit (SOD Assay Kit 706002®, Cayman Chemical, Ann Arbor, Michigan, USA), the results were expressed as U mL^{-1} of DW, where U is defined as the amount of enzyme needed to exhibit 50% dismutation of the superoxide radical. Glutathione peroxidase (EC 1.11.1.9) was conducted according to Flohé & Günzler, (1984) technique, using H_2O_2 as a substrate, the results were expressed as U g^{-1} TP of DW, where U is equal to the mmol equivalent of reduced glutathione per milliliter per minute. Phenylalanine ammonia lyase (EC 4.3.1.5) was determined according to Sykłowska-Baranek *et al.* (2012), the results were expressed as $\text{U } 100 \text{ g}^{-1}$ TP of DW, where U is equal to production of 1 μmol cinnamic acid per milliliter per minute. Reduced glutathione (GSH) was performed

colorimetrically by reaction with DTNB (Xue *et al.*, 2001), the results were expressed as $\mu\text{mol g}^{-1}$ of DW.

ABTS (2,2'-azino-bis(3-Ethylbenzotiazoline-6-Sulphonic Acid)) was performed according to the methodology of Re *et al.* (1999), the results were expressed as equivalents vitamin C $\mu\text{g g}^{-1}$ of DW. DPPH (2,2-diphenyl-1-picrylhydrazyl) was performed using the method of Brand-Williams *et al.* (1995), the results were expressed as equivalents vitamin C $\mu\text{g g}^{-1}$ of DW.

Phenols were determined according to the methodology of Yu & Dahlgren, (2000), the results were expressed as equivalents gallic acid mg g^{-1} of DW. Flavonoids was performed with the method of Arvouet-Grand *et al.* (1994), the results were expressed as equivalents of quercetin mg g^{-1} of DW. Vitamin C was determined by the titration method with 2,6 dichlorophenolindophenol (Padayatt *et al.*, 2001), the results were expressed as $\text{mg } 100 \text{ g}^{-1}$ fresh weight.

The content of lycopene, β -carotene, and *a y b* chlorophylls were determined according to Nagata & Yamashita (1992). Total chlorophyll was the sum of Chl *a* and Chl *b*. All data are expressed as $\text{mg } 100 \text{ g}^{-1}$ of DW.

Endogenous phytohormones

To determine salicylic acid (SA), 1 mL of the extraction solution (89% water, 10% methanol and 1% acetic acid) was added to 50 mg of the foliar tissue. Subsequently, were centrifuged at $16,000 \times g$ (10 minutes). The supernatant was extracted and filtered (0.45 μm filter) and degassed by sonication. A liquid chromatograph (Agilent compact 1120) equipped with variable wavelength UV detector was used. The chromatographic conditions were for stationary phase: column C-18 polaris of 250 mm in length. The mobile phase has a gradient of 50% phase A (water 94.9%, acetonitrile 5%, and formic acid 0.1%) and 50% phase B (acetonitrile 94.9%, water 5%, and formic acid 0.1%), at a flow of 0.8 mL min^{-1} for a run time of 13 min. It was read at a wavelength of 250 nm (Forcat *et al.*, 2008). The results were expressed as mg kg^{-1} of DW.

Jasmonic acid (JA) was determinate mixing 100 mg of leaf tissue and 900 μL of the extraction solution (95% methanol and 5% ethyl acetate), vortexed, sonicated (10 min), and centrifuged at $16,000 \times g$ (10 min). The supernatant was filtered, the solvent was subsequently evaporated in dry oven at $50 \text{ }^\circ\text{C}$, and the residues were re-suspended in 500 μL of mobile phase (60 % methanol, 39% water, and 1% acetic acid). Same equip was used to determine jasmonic acid (JA). The flow was 0.60 mL min^{-1} using a Hypersil ODS column of $25 \text{ cm} \times 4.6 \text{ mm} \times 5 \mu\text{m}$, at a wavelength of 230 nm with an analysis time of 20 min (Kramell *et al.*, 1995; Michelena *et al.*, 2001). The results were expressed as mg kg^{-1} of DW.

Determination of gene expression by qPCR

The RNA was extracted using 100 mg fresh leaf tissue and TRIzol reagent (TRI Reagent[®], MRC, TR 118), purified with chloroform, and precipitated with isopropanol, as described in Cui *et al.* (2004). RNA quantification was carried out in a UV-Vis spectrophotometer (Thermo Scientific Model G10S) at an absorbance of 260 and 280 nm, and RNA quality was analyzed by denaturing electrophoresis in agarose gel. 2 μg of total-RNA were used by synthesis of the cDNA was performed using a commercial kit (Promega, Madison, Wisconsin, USA). The primers correspond to an endogenous gene *ACT* (actin) and six study genes: *PRI* (salicylic acid), *JA* (jasmonic acid), *GAME1* (phytoalexins), *PAL5-3* (phenylalanine ammonia lyase), which were designed in the software AMPLIFIX, OLIGOANALIZER and Primer-BLAST. For the *ET* (ethylene) and *ABA* (abscisic acid), the sequences cited by Anstead *et al.* (2010) and Nitsch *et al.* (2009), were considered, respectively (Table 1).

Table 1. Primers sequence of analyzed genes.
Tabla 1. Secuencia de cebadores de los genes analizados.

Gene	(Forward primer 5'-3')	(Reverse primer 5'-3')	T _m (°C) Annealing
<i>ACT</i>	CCCAGGCACACAGGTGTTAT	CAGGAGCAACTCGAAGCTCA	60 °C
<i>PR1</i>	AAGTAGTCTGGCGCAACTCA	GTCCGATCCAGTTGCTACA	60 °C
<i>JA</i>	TGGTTCGTCGACTTCGTCAT	CTCGGCCTTGAGAGAGTTCA	60 °C
<i>GAME1</i>	TCGTTGTTCCCTGGGTTACCT	TCCGAAACTCGAACTTTCTCG	58 °C
<i>PAL5-3</i>	GGAGGAGAATTTGAAGAATGCTGTG	TCCCTTTCCACCACTTGTAGC	60 °C
<i>ET</i>	TGTCCCAAGCCAGACTTGAT	TGCCATCTTGTTGAGCAATC	60 °C
<i>ABA</i>	CTTATTTGGCTATCGCTGAACC	CCTCCAACCTCAAACCTCATTGC	60 °C

The primers were prepared at a concentration of 15 pmol mL⁻¹. The quantification method used was a standard relative curve, therefore, for each quantification analysis a standard curve per gene was included using a 1:5 dilution.

qPCR reactions were analyzed in an Applied Biosystems StepOne™ Equipment version 2.3 by the standard relative curve method, measuring the fluorescence intensity of SYBR Green. The qPCR reaction for all genes was performed in a total volume of 20 µL. Regarding *ACT* gene, 10 µL of SYBR® Select Master Mix (Applied Biosystems, Foster City, California, USA) were added along with 0.10 µL of first forward (72 nM), 0.08 µL of first reverse (60 nM), 1 µL of cDNA and 8.82 µL of nuclease-free water. Regarding the *PR1* gene, 10 µL of Master Mix were added along with 0.03 µL of first forward (20 nM), 0.05 µL of first reverse (40 nM), 1 µL of cDNA diluted at a ratio of 1:5, and 8.92 µL of nuclease-free water. Regarding the *JA* gene, 10 µL of Master Mix, 0.05 µL of first forward (40 nM), 0.08 µL of first reverse (60 nM), 1 µL of cDNA and 8.87 µL of nuclease-free water were added. *ABA* gene, 10 µL of Master Mix, 4 µL of first forward (3000 nM), 4 µL of first reverse (3000 nM), 1 µL of cDNA and 1 µL of nuclease-free water were added. As for *ET*, *PAL5-3* and *GAME1* genes, 10 µL of Master Mix, 0.13 µL of first forward (100 nM), 0.13 µL of first reverse (100 nM), 1 µL of cDNA (*ET* and *PAL5-3*) or 1 µL of cDNA diluted 1:5 (*GAME1*) and 8.74 µL of nuclease-free water were added. The qPCR was run with the following program in the thermal cyclers: Hot Start, 10 min at 95 °C and PCR (40 cycles), 15 s at 95 °C and 1 min at temperature annealing.

Statistical analysis

The experiment was established in completely random design. To estimate the incidence and severity of Fol wilt the disease 18 replicates per treatment were considered; and for biochemical, secondary metabolites and gene expression five replicates were considered per treatment. An analysis of variance and Fisher's Least Significant Difference test ($p \leq 0.05$) were performed using the InfoStat software (v2019).

RESULTS

Incidence and severity of Fol wilt disease

Statistical differences were found between treatments in incidence of Fol wilt in tomato plants, throughout the experiment, except for the first sampling (Figure 1A). There was no incidence of Fol wilt the disease in silicic acid (ASi) treatment and absolute control (T0). As for Fol treatments (Fol and ASi+Fol) a certain degree of incidence of Fol wilt was observed throughout the evaluation dates but increased as culture developed during time. Particularly, it was observed that ASi application decreased incidence of Fol wilt as ASi+Fol treatment had a lower incidence of Fol wilt than the Fol treatment from inoculation day to day 66 after transplanting. Best effect of ASi was obtained at 61 days after transplanting (dat), since incidence of Fol wilt was reduced by approximately 24%. After 70 dat, both infested treatments reached 100% incidence.

The first symptoms of the severity of phytopathogen were observed at 43 dat and increased progressively during culture development (Figure 1B). Throughout the experiment ASi+Fol treatment consistently decreased Fol severity. At 78 dat, the greatest decrease in severity was observed, with approximately 21% less incidence of wilt than Fol treatment, and 12% at 85 dat.




Figure 1. Incidence (A) and severity (B) of Fol wilt disease in tomato plants with or without ASi application. ns: not significant, *, ***: statistical differences at $p < 0.05$, < 0.0001 respectively. $n = 18$. ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control.

Figura 1. Incidencia (A) y gravedad (B) de la enfermedad de la marchitez por *Fusarium* en plantas de tomate con o sin aplicación de ASi (ácido silícico). ns: no significativo, *, ***: diferencias estadísticas a $p < 0.05$, < 0.0001 respectivamente. $n = 18$. ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto.

Biochemical Responses

Results showed statistical differences in total protein content, carotenoids, and chlorophylls in tomato leaves (Table 2). There was no difference in protein content between ASi and T0 treatments, whereas Fol and ASi+Fol treatments significantly decreased proteins by 33% and 31% respectively.

Lycopene content in tomato leaves inoculated with Fol was $3.1 \text{ mg } 100 \text{ g}^{-1} \text{ DW}$; which corresponds to 51, 62 and 65% less than the content registered in the plants of the ASi+Fol, T0 and ASi treatments, respectively (Table 2). There was no statistically significant difference between the ASi+Fol, ASi and T0 treatments. In the case, β -carotene, there were no statistically significant differences between treatments, only the ASi+Fol treatment presented the highest content, with approximately 59% more than T0.

Regarding chlorophyll content (*a*, *b* and total), results were consistent. The total chlorophyll a content in the plants inoculated with Fol was 92.1 , 40.3 and $132.4 \text{ mg } 100 \text{ g}^{-1} \text{ DW}$, respectively; this quantity was 49.1 , 40.6 and 46.7% , respectively, lower than the content registered in the plants of the ASi+Fol treatments. There was no significant statistical difference between the ASi+Fol, ASi and T0 treatments (Table 2).

Table 2. Total protein, carotenoids, and chlorophyll content in leaves of tomato plants infested and not-infested with *F. oxysporum* and with or without ASi applications**Tabla 2.** Contenido total de proteínas, carotenoides y clorofila en hojas de plantas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico).

Treatments	Proteins (mg g ⁻¹ DW)	Lycopene (mg 100 g ⁻¹ DW)	β-carotene (mg 100 g ⁻¹ DW)	Chlorophyll a (mg 100 g ⁻¹ DW)	Chlorophyll b (mg 100 g ⁻¹ DW)	Total Chlorophyll (mg 100 g ⁻¹ DW)
ASi+Fol	19.1 ^b	6.3 ^{ab}	41.9 ^a	180.8 ^a	67.8 ^a	248.6 ^a
ASi	26.7 ^a	9.0 ^a	30.5 ^{ab}	206.4 ^a	71.3 ^a	277.7 ^a
Fol	18.6 ^b	3.1 ^b	28.8 ^{ab}	92.1 ^b	40.3 ^b	132.4 ^b
T0	27.8 ^a	8.3 ^a	26.3 ^b	172.0 ^a	50.8 ^{ab}	222.7 ^a
CV (%)	23.0	41.0	25.8	22.1	23.2	22.8

n= 5. Averages with same letter per column are statistically equal (LSD Fisher, $p \leq 0.05$). ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control. DW: dry weight. CV: coefficient of variation (%).

n= 5. Medias con la misma letra por columna son estadísticamente iguales (LSD Fisher, $p \leq 0.05$). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto. DW: peso seco. CV: coeficiente de variación (%).

Significant differences between treatments were observed in vitamin C, flavonoids, phenols and antioxidant compounds, as well as in DPPH antioxidant capacity (Table 3). Vitamin C content no differences were found between the plants inoculated with Fol and the plants treated with ASi+Fol, however, the plants inoculated with Fol were 52.7 and 33.1% higher than the content presented in the plants treated with ASi and T0. The reduced glutathione was higher in plants inoculated with Fol, 69.5 and 6.6% higher than ASi+Fol and ASi plants. Phenol content did not present differences between the plants treated with Fol, ASi+Fol and ASi, only an increase of 32% stood out in the plants inoculated with Fol, with respect to T0. Regarding Flavonoid content, showed a reduction of 24.6, 27.8 and 30.4% in plants inoculated with Fol, compared to plants treated with ASi+Fol, T0 and ASi respectively. ABTS did not present differences between treatments, while antioxidant capacity, evaluated by DPPH increased 3.1% in Fol compared to ASi+Fol (Table 3).

Table 3. Non-enzymatic antioxidant compounds in leaves of tomato plants infested and not-infested with *F. oxysporum* and with or without ASi applications at 73 dat.**Tabla 3.** Compuestos antioxidantes no enzimáticos en hojas de plantas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico) a los 73 días después del trasplante (dat).

Treatments	Vitamin C (mg 100 g ⁻¹ FW)	Reduced glutathione (μmol g ⁻¹ DW)	Phenols (mg g ⁻¹ DW)	Flavonoids (mg g ⁻¹ DW)	ABTS (μg g ⁻¹ DW)	DPPH (μg g ⁻¹ DW)
ASi+Fol	123.2 ^{ab}	5.9 ^b	18.4 ^{ab}	28.9 ^a	7.7 ^a	52.4 ^b
ASi	96.8 ^b	8.5 ^{ab}	17.2 ^{ab}	31.3 ^a	7.8 ^a	54.2 ^a
Fol	147.8 ^a	10.0 ^a	20.4 ^a	21.8 ^b	8.9 ^a	54.0 ^a
T0	110.9 ^b	8.9 ^{ab}	15.2 ^b	30.2 ^a	9.5 ^a	53.7 ^a
CV (%)	20.3	27.8	11.9	9.6	21.9	1.1

n= 5. Averages with the same letter per column are statistically equal (LSD Fisher, $p \leq 0.05$). ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control. ABTS: 2,2'-azino-bis(3-ethylbenzthiazolin-6-sulfonic acid). DPPH: 2,2-diphenyl-1-picrylhydrazyl. CV: coefficient of variation (%). FW: fresh weight. DW: dry weight.

n= 5. Las medias con la misma letra por columna son estadísticamente iguales (LSD Fisher, $p \leq 0.05$). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto. ABTS: ácido 2,2'-azino-bis(3-etilbenzthiazolin-6-sulfónico). DPPH: 2,2-difenil-1-picrilhidrazil. CV: coeficiente de variación (%). FW: peso fresco. DW: peso seco.

Except for APX y SOD enzyme, there were statistical differences in PAL, CAT and GPX between treatments (Table 4). PAL no differences were found between the plants inoculated with Fol and the plants treated with ASi+Fol, however, the plants inoculated with Fol presented 3.2% higher activity than the plants treated with ASi and T0. The activity of the CAT enzyme was higher in the plants inoculated with Fol in 85.1 and 108.4 %, with respect to the plants of the ASi+Fol and T0 treatments. The enzymatic activity of GPX was similar in the plants of the Fol, ASi+Fol and ASi treatments, only an increase of 193% was observed in the plants inoculated with Fol, with respect to T0.

Table 4. Enzymatic activity in leaves of tomato plants infested and not-infested with *F. oxysporum* and with or without ASi application at 73 dat.

Tabla 4. Actividad enzimática en hojas de plantas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicación de ASi (ácido silícico) a los 73 días después del trasplante (dat).

Treatments	PAL (U 100 g ⁻¹ TP)	APX (U g ⁻¹ TP)	CAT (U g ⁻¹ TP)	GPX (U g ⁻¹ TP)	SOD (U ml ⁻¹)
ASi+Fol	4.0 ^{ab}	15.9 ^a	513.2 ^b	48.7 ^a	9.6 ^b
ASi	3.2 ^b	9.6 ^a	623.9 ^{ab}	41.3 ^a	24.6 ^a
Fol	5.5 ^a	14.3 ^a	949.7 ^a	58.5 ^a	16.1 ^{ab}
T0	3.2 ^b	16.5 ^a	455.7 ^b	19.6 ^b	25.3 ^a
CV (%)	22.3	56.7	37.1	26.4	46.2

n=5. Averages with same letter per column are statistically equal (LSD Fisher, $p \leq 0.05$). ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control. PAL: phenylalanine ammonia lyase. APX: ascorbate peroxidase. CAT: catalase. GPX: glutathione peroxidase. SOD: superoxide dismutase. TP: total protein. CV: coefficient of variation (%).

n=5. Las medias con la misma letra por columna son estadísticamente iguales (LSD Fisher, $p \leq 0.05$). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto. PAL: fenilalanina amonio liasa. APX: ascorbato peroxidasa. CAT: catalasa. GPX: glutatión peroxidasa. SOD: superóxido dismutasa. TP: proteína total. CV: coeficiente de variación (%).

Endogenous phytohormones

Salicylic acid content in tomato leaves was modified by treatments in two out of three samplings, but no statistical differences were obtained between treatments at 31 dat (Table 5). At 73 dd, when the plants appeared an advanced severity (Figure 1B), the plants of the Fol, ASi+Fol and ASi treatments, did not emerge statistically significant directions, only the plants inoculated with Fol presented 247.5% higher concentration than the plants T0.

The content of JA could only be quantified at 31 and 73 dat, with statistical differences between treatments in both cases (Table 5). Systematically, an increase of JA was observed as culture developed and consequently the severity of the pathogen increased (Figure 1B). At 31 dat, when the disease had not yet proliferated, only JA was quantified in the ASi+Fol and Fol treatments, the latter being significantly higher (245.6%). At 73 dat, once the incidence and severity of Fol wilt increased (Figure 1), an increase JA was observed in all treatments, except for T0. Once again, Fol generated the highest concentration of JA, surpassing the ASi+Fol treatment by 81.3 %, and the ASi treatment by 234.3%.

Table 5. Content of secondary metabolites related to defense systems in leaves of tomato plants infested and not-infested with *F. oxysporum* and with or without ASi applications

Tabla 5. Contenido de metabolitos secundarios relacionados con los sistemas de defensa en hojas de plantas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico).

Treatments	Salicylic acid (mg kg ⁻¹ DW)			Jasmonic acid (mg kg ⁻¹ DW)		
	15 dat	31 dat*	73 dat	15 dat	31 dat*	73 dat
ASi+Fol	Nq	35.1 ^a	171.6 ^a	Nq	5.7 ^b	72.1 ^b
ASi	52.6 ^a	23.9 ^a	72.3 ^{ab}	Nd	Nd	39.1 ^c
Fol	Nq	32.2 ^a	163.0 ^a	Nq	19.7 ^a	130.7 ^a
T0	21.9 ^b	43.9 ^a	46.9 ^b	Nd	Nd	Nd
CV (%)	42.8	49.6	66.2	-	58.3	33.1

n=5. Averages with same letter per column are statistically equal (LSD Fisher, $p \leq 0.05$). ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control. CV: coefficient of variation (%). DW: dry weight. *: 10 days after inoculation of Fol. Nq: not quantified. Nd: not detected (is present but conjugated, not separated).

n=5. Las medias con la misma letra por columna son estadísticamente iguales (LSD Fisher, $p \leq 0.05$). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto. CV: coeficiente de variación (%). DW: peso seco. *: 10 días después de la inoculación con Fol. Nq: no cuantificado. Nd: no detectado (presente pero conjugado, no separado).

Gene expression

PR1 and *JA* genes, which encode salicylic acid and jasmonic acid respectively, showed changes in their expression due to treatments (Figure 2). At 31 dat, only Fol treatment modified *PR1* and *JA* gene expression, repressing them 0.7- and 0.34-fold change respectively in comparison to ASi+Fol. However, at 73 dat all treatments increased *PR1* gene expression, however the combination of ASi+Fol generated the largest increase with 18.7-fold change, while ASi and T0 decreased it 2.2- and 2.5-fold change, respectively, in comparison to Fol (Figure 2A). This trend is consistent with what was observed in SA quantification, where T0 had the lowest amount of this metabolite and Fol treatments showed the highest amount (Table 5).

In the case of *JA* gene at 73 dat, in comparison at 31 dat, a similar trend was observed since all treatments increased its expression. However, ASi+Fol, ASi and Fol decreased 2.9-, 5.5- and 5.7-fold change, respectively, in comparison to Fol (Figure 2B). Results consistent with JA determination, since Fol generated the highest amount of this metabolite, followed by the combinations ASi+Fol and ASi (Table 5).




Figure 2. Relative expression of salicylic acid (*PR1*) (A) and jasmonic acid (*JA*) (B) genes in tomato leaves infested and not-infested with *F. oxysporum* and with or without ASi applications. ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control.

Figura 2. Expresión relativa de los genes de ácido salicílico (*PR1*) (A) y ácido jasmónico (*JA*) (B) en hojas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto.

At 31 dat, application of the treatments practically showed no changes in *GAME1* gene expression (Figure 3A). However, at 73 dat, the expression of this gene was repressed 0.1-fold change in ASi+Fol, and overexpressed ASi and T0 (1.3- and 0.7-fold change, respectively) compared to Fol.

In the *PAL5-3* gene, at 31 dat, a overexpression of this gene was observed by the ASi+Fol and T0 treatments, equivalent to 0.4- and 0.6-fold change, respectively, compared to Fol (Figure 3B). However, at 73 dat all treatments repression this gene, ASi+Fol was 0.2-fold change less, T0 was 1.8-fold change, and ASi generated the highest expression equivalent to 0.1-fold change more than Fol.




Figure 3. Relative expression of phytoalexins (*GAME1*) (A) and phenylalanine ammonia lyase (*PAL5-3*) (B) genes in tomato leaves infested and not-infested with *F. oxysporum* and with or without ASi applications. ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control.

Figura 3. Expresión relativa de los genes de fitoalexinas (*GAME1*) (A) y fenilalanina amonio liasa (*PAL5-3*) (B) en hojas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico). ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto.

The expression of *ET* gene at 31 dat increased slightly by ASi+Fol (0.5-fold change), while T0 treatment decreased 0.1-fold change in comparison to Fol. However, at 73 dat all treatments overexpressed this gene: ASi+Fol, 5.5-fold change, ASi generated the highest expression with 7.1-fold change, except T0 (-1.7-fold change), greater than Fol (Figure 4A). In the case of *ABA* gene, at 31 dat ASi+Fol treatment generated a overexpression of 0.6-fold change in comparison to Fol, whereas ASi+Fol treatment increased the expression in 0.30-fold change, whereas T0 treatment increased the expression in 0.3-fold change (Figure 4B). At 73 dat, all treatments decreased *ABA* gene expression just like what was observed with the *ET* gene (T0) (Figure 4A). T0 treatment generated the lowest expression of the *ABA* gene with 4.5-fold change less, compared to Fol, followed by ASi treatment with 3.3-fold change. ASi+Fol treatment also decreased *ABA* gene expression but not in the same magnitude as T0 and ASi treatments, since it only increased 0.06-fold change in comparison to Fol (Figure 4B).




Figure 4. Relative expression of ethylene (*ET*) (A) and abscisic acid (*ABA*) (B) genes in tomato leaves infested and not-infested with *F. oxysporum* and with or without ASi applications. T0: absolute control. ASi+Fol: silicic acid + *F. oxysporum*. ASi: silicic acid. Fol: *F. oxysporum*. T0: absolute control.

Figura 4. Expresión relativa de los genes de etileno (*ET*) (A) y ácido abscísico (*ABA*) (B) en hojas de tomate infestadas y no infestadas con *F. oxysporum*, con o sin aplicaciones de ASi (ácido silícico). T0: control absoluto. ASi+Fol: ácido silícico + *F. oxysporum*. ASi: ácido silícico. Fol: *F. oxysporum*. T0: control absoluto.

DISCUSSION

Incidence and severity of Fol wilt disease

Cao *et al.* (2017) mentioned that when stress is prolonged, self-regulation in tomato is exceeded, and the positive effects of Si are attenuated, as observed here after 70 dat, where both infested treatments reached 100% incidence. Furthermore, without the use of fungicides, it is practically impossible to completely suppress the disease (Bakhat *et al.*, 2018) which explains the results observed here (Figure 1A). Once the tissue is infected, the xylem vessels is obstructed by fungal structures or polysaccharides (Singh *et al.*, 2017), there is also toxins secretion by this pathogen, this cause wilting and necrosis of the leaves, and dark brown vascular discoloration (Pirayesh *et al.*, 2018). At the same time there is an excessive ROS production, causing oxidative stress, that generate DNA and metabolism damage, eventually culminating in the apoptosis (Aybeke, 2017). These symptoms produced together the high severity levels of Fol recorded in tomato plants under study (Figure 1B).

On the other hand, several authors have reported the different action mechanisms of silicic acid to mitigate the incidence and severity of Fol wilt disease, achieving a better plants development (Figure 1). In this work it was observed that the ASi and T0 treatments did not present symptoms of the disease, since were not inoculated with Fol, however the plants inoculated with Fol showed wilting of the leaves, likewise, in the plants treated with ASi+Fol but with a lower degree of incidence and severity (up to 24 and 21%, respectively) (Figure 1).

It was found that Si can activate the biochemical mechanisms of plants, such as increased antioxidant activity and phytohormones (Bakhat *et al.*, 2018), in addition to pathogenicity related genes expression (Bhatt & Sharma, 2018). Once in the plant, Si can polymerize into amorphous silica and accumulate in the trichomes or cuticles of the leaves, forming a physical defense, which prevents the passage of pathogens (Dann & Le, 2017; Hoffmann *et al.*, 2020). For example, silicic acid induces biochemical barriers, and once polymerized in amorphous silica, generates physical barriers (Bathoova *et al.*, 2021; Dann & Le, 2017). In addition, its accumulation in cell walls, intracellular spaces and leaf trichomes generates firmer and more rigid structures (Carré-Missio *et al.*, 2014; Ferreira *et al.*, 2015). In the cytoplasm, it induces biochemical barriers such as metabolic pathways activation, JA (Bakhat *et al.*, 2018; Ning *et al.*, 2014), SOD and CAT activity (Cao *et al.*, 2017) increase, flavonoid and *PAL5-3* expression (Rahman *et al.*, 2015). Silicic acid also participates in the overexpression or repression of pathogenicity related genes (Brunings *et al.*, 2009).

Biochemical Responses

F. oxysporum produces toxins such as fusaric acid that causes oxidative stress induced by ROS, causing damage to the membrane, and reducing photosynthetic pigment content in tomato leaf tissue (Singh *et al.*, 2017). In this work, proteins and lycopene clearly were affected by Fol, regardless of whether silicic acid is applied (Table 2). Opposite case, the chlorophylls were not reduced with the ASi+Fol treatment, it is possible that the application of ASi in Fol inoculated tomato plants protected against pigment degradation and enhance the photosynthetic capacity (S. Sun *et al.*, 2022), because Chlorophyll content is related to higher photosynthetic capacity in plants under stress (González-García *et al.*, 2023). It has been observed that Si applied on leaves to wheat under hydric stress increased chlorophyll concentration, improving light use efficiency (Maghsoudi *et al.*, 2015). In *Glycine max*, Na_2SiO_3 application promoted Fe mobilization and maintenance in leaves, generating greater protection to chlorophyll degradation (Gonzalo *et al.*, 2013).

Previously, it has been reported that pathogen infection in plants induces oxidative stress, which results in a higher production of antioxidant defense compounds (Pei *et al.*, 2020), as observed here: reduced glutathione, phenols and DPPH (Table 3). Whan *et al.* (2016) found that *F. oxysporum* infection in *Gossypium hirsutum* increased the phenolic compounds concentration, as seen here. Regarding silicon, Hajiboland *et al.* (2017a) have also observed that when applied to *Nicotiana rustica* plants, the concentration of phenolic compounds increases, only when they are subjected to mechanical stress. In the plant, silicic acid binds with hemicelluloses, pectin and polyphenols forming cell walls, in addition to stimulating a higher concentration of these components and lignin (Babalar *et al.*, 2016). This strengthens the plants against attack by pathogens.

Similarly, it has been suggested that Si role against the attack of *F. oxysporum* in banana crop was possibly due to the potentiation of the phenylpropanoid pathway (Fortunato *et al.*, 2014), which is related to the production of phenolic metabolites such as flavonoids, lignin, etc. which are synthesized from L-phenylalanine that produces trans-cinnamic acid derived from PAL content (Hajiboland *et al.*, 2017b; Hussain *et al.*, 2021). This would explain the increase in flavonoid content in plants treated with ASi+Fol. Madany *et al.* (2020), for example, suggested that Si NP favored PAL activity, and thus to the increase in content of lignin. However, the results of this study do not show a difference with the application of ASi+Fol with respect to Fol in

phenols and PAL (Table 3). On the other hand, treatment with Fol increased the PAL (Table 4) and also the content of total phenols respect T0 (Table 3).

It is known that *F. oxysporum* colonization can trigger defense mechanisms, for example, increased callose and lignin deposits in the cell structures or increased ROS and, consequently, greater antioxidant activity (Pei *et al.*, 2020). Dorneles *et al.* (2017) got a higher CAT activity, as observed here (Table 4). However, CAT was not affected in the plants treated with ASi+Fol, possibly due to a low activity of this enzyme in H₂O₂ elimination (Dallagnol *et al.*, 2015), as well as Habibi, (2015) also did not observe changes in CAT in canola under water stress. Which suggests less ROS concentration in this treatment, and therefore less stress (Hu *et al.*, 2020), since in addition to low CAT activity, low concentrations of reduced glutathione and DPPH were also observed in ASi+Fol treatment.

As already mentioned, the infection of pathogens to plants induces an increase in oxidative stress, triggering antioxidant defense (Kang *et al.*, 2014) producing antioxidant enzymes such as GPX and other antioxidant compounds. Silicon can also function as elicitor in plants and generates changes the defense-related enzymes activity and together with other mechanisms lead to induced resistance (Sakr, 2021; M. Wang *et al.*, 2017). This explains the increase of GPX observed in this work in treatments with ASi or Fol (Table 4).

Likewise, SOD activity has a major importance in the defense against pathogens, since it regulates the concentration of ROS in cells (Hao *et al.*, 2011; Sattar *et al.*, 2020). It was found that increase SOD activity was caused by a higher generation of H₂O₂, conferred by *Podospaera xanthii*, in melon plants added with Si (Dallagnol *et al.*, 2015), however, these results do not agree with those reported in this study (Table 4). In this regard, it has been reported that Si-treated plants exhibited reduction in SOD and CAT activities in *Cercospora sojina* infected soybean (Telles Nascimento *et al.*, 2016). Which suggests a lower concentration of ROS in ASi+Fol treatment (Hu *et al.*, 2020), in addition, other antioxidants (*e. g.* GPX) could also be involved in the elimination of ROS. In this case, the positive effect observed on the incidence and severity of Fol wilt may also be the result of the efficacy of other mechanisms, such as phytohormones synthesis and modification of gene expression.

Secondary metabolites

Since SA accumulation is key in the defense of tomato plants after the attack of Fol (Di *et al.*, 2017) the observed increase is directly related to the defense system against Fol (Table 5). Salicylic acid exacerbated the systemic acquired resistance (SAR) conferred by Fol infection in *Musa acuminata* (Z. Wang *et al.*, 2015). In a previous study, it has been observed that the application of Na₂SiO₃ showed lower concentrations of JA in *Oryza sativa* than mechanical stress treatment alone (Kim *et al.*, 2014); similar findings to those obtained in this work (Table 5).

The results obtained here (Table 5) demonstrate that Fol infection triggers the defense mechanism of tomato plants, increasing the SA and JA generation (Jang *et al.*, 2018; Vivancos *et al.*, 2015). This can be caused by ROS accumulation such as H₂O₂ that induce the activity of the lipid hydroperoxidase and benzoic acid hydrolase enzymes, that stimulate the synthesis of jasmonic acid and that transform benzoic acid into salicylic acid, respectively (Dorneles *et al.*, 2017; Fagerstedt *et al.*, 2010). Silicic acid application can also induce the SA and JA biosynthesis (Jang *et al.*, 2018), however, the intensity of the response is less than that caused by *F. oxysporum*, and it may take longer to appear as observed here (Table 5).

Gene expression

The *PR1* and *JA* genes expression responses are derived from the increase in salicylic and jasmonic acids observed in treatments with Fol and ASi+Fol (Table 5). This is because SA and JA are hormones that regulate the expression of these defense responses related genes (Han & Kahmann, 2019). It highlights the production of PAL and various antioxidant compounds (Tables 3 and 4) involved in the ROS reduction.

The results show that Fol triggers the plant defense system, such as the expression of related genes. Although silicic acid application also induces SA and JA production (Table 5), these markers did not have a very strong impact on *PR1* and *JA* genes expression (Figure 2), unlike other authors' reports (Cai *et al.*, 2009). Gulzar *et al.* (2021) cited that silicon amendment in *Lycopersicon esculentum* inoculated with *Alternaria solani*, had an up-regulation the *PR3*, *LOXD* and *JERF3* (JA marker genes). Manivannan & Ahn, (2017) documented that Si treatment-induced resistance was manifested by activating the SAR, through the regulation of the jasmonic acid pathway and genes related in the hypersensitivity response. Therefore, the positive effect observed on the incidence and severity of Fol wilt may occur due to the regulation of other pathogenesis-related genes, or due to the increase in antioxidant compounds (Tables 2-4), or even structural modifications (Dann & Le, 2017; Vivancos *et al.*, 2015).

The gene *GAME1* is related to phytoalexins, which are mainly phenolic compounds that accumulate when infections by phytopathogenic fungi such as *F. oxysporum* appear, participating in the defense against them (Jeandet *et al.*, 2002). However, the results showed an opposite effect with the inoculation of Fol, since this gene expression was repressed (Figure 3A). Opposite case, the clear effect on the expression of *GAME1* gene by the silicic acid application is consistent with other authors' reports, who found that production of flavonol phytoalexins in *Cucumis sativus* and momilactones in *Oryza sativa*, respectively, is stimulates by silicon (Fawe *et al.*, 1998; Rodrigues *et al.*, 2004). Therefore, the production of phytoalexins induced by the silicic acid application could have affected the progress of the incidence and severity of Fol wilt observed in tomato (Figure 1).

Overexpression of *PAL5-3* is important to regulate *Fusarium* infection, since it leads to a greater accumulation of lignin and synthesis of salicylic acid to trigger SAR (Iqbal *et al.*, 2005). Therefore, tolerance to this pathogen can be increased when these circumstances occur, as observed in this study (Figures 1 and 3B). Zvirin *et al.* (2010) demonstrated that *PAL5-3* expression in *Cucumis sativus* increased as colonization of *F. oxysporum* progressed. In addition, Cruz *et al.* (2015) observed in wheat plants inoculated with *Pyricularia oryzae* an increase in *PAL* expression independently of the application of silicic acid, however the highest level of expression was in the silicic acid treatment. These results agree with the observed, since although all the treatments increased the expression of the *PAL5-3* gene, the treatment of only silicic acid presented the highest overexpression (Figure 3B). Kuai *et al.* (2017) and Shetty *et al.* (2011) obtained higher transcripts of this gene when applying silicon; this led to an increase in lignin and phenolic compounds, reducing the severity of pathogens, similar to observed here (Table 3, Figure 1). These results indicate that indeed the application of silicon, in this case as silicic acid, increases the expression of the *PAL5-3* gene. This translates into greater tolerance to pathogens such as Fol, through different mechanisms that can be biochemical (phenolic compounds) or mechanical as physical barriers (lignin).

These results seem to indicate that silicic acid application has a more pronounced effect on the expression of *ET* gene, in comparison with the expression induced by Fol (Figure 4A). Commonly the attack of pathogens, like Fol, in plants, triggers the production of antimicrobial compounds as well as stress hormones such as SA, JA and ethylene (Table 5, Figure 4A) (Islam *et al.*, 2019). However, silicon also confers the activation of genes involved with plant stress, specifically in the form of silicic acid, it induces defense reactions that improves the production of stress hormones (Fauteux *et al.*, 2005). This was observed mainly with the ethylene related *ET* gene (Figure 4A). This finding agree with that of Van Bockhaven *et al.* (2015) who reported that silicon intervenes specifically in the ET signaling pathway. Similarly, Ghareeb *et al.* (2011) demonstrated that Si-induced resistance is mediated by the ET and JA signaling pathways in tomato plants infected with *Ralstonia solanacearum*. This suggests that in the case of tomato, the application of silicic acid increases tolerance to Fol (Figure 1) by regulating the genes related to phytoalexins (gene *GAME1*, Figure 3A) and the production of ethylene (Figure 4A).

In the case of *ABA* gene, apparently Fol induces a greater expression of this gene in comparison with the application of silicic acid (Figure 4B). Although the reports mention that the attack of pathogens in plants involves the production of ethylene (Cai *et al.*, 2009; Fauteux *et al.*, 2005), the results obtained in tomato indicated that Fol induces mainly the expression of the gene related to abscisic acid (Figure 4B). This can be derived from the antagonism between ABA and ethylene, in addition to the negative effect of ABA to pathogens tolerance (Mauch-Mani & Mauch, 2005). This is consistent with our results, since *ABA* gene expression increased with Fol inoculation in tomato especially at 73 dat, when the severity of pathogen is higher (more than 60%) (Figure 1B). While at 31 dat when no incidence of Fol wilt was found (Figure 1A), there were no significant effects in *ABA* gene expression (Figure 4B). Overall, it is clear that there is a complex interaction between *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* and *ABA* genes that result in different defense responses and tolerance to Fol in tomato (Anderson *et al.*, 2004; de Lamo & Takken, 2020).

CONCLUSIONS

Stress by *F. oxysporum* in tomato plants increased antioxidant enzymatic activity and enzymatic activity antioxidant compounds in leaves, mainly vitamin C, total phenols, PAL, CAT and GPX. It was also observed that the inoculation of Fol in tomato increased the content of salicylic and jasmonic acid (247 and 81.3%, respectively). Additionally, Fol induces increased *JA*, *GAME1*, *PAL5-3* and *ABA* genes expression, at 73 dat; however, it caused a lower concentration chlorophyll and flavonoids, and expression of *PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* and *ABA* gene, at 31 dat.

Silicic acid application to tomato plants stressed by *F. oxysporum* (ASi+Fol), reduced the incidence and severity of Fol wilt (until 70 and 85 dat, respectively); since it increased the content of antioxidant compounds in leaves, mainly flavonoids (24.6%) and chlorophyll (248.6%). In addition to this, led an overexpression of genes associated with to salicylic acid, jasmonic acid, phytoalexins, phenylalanine ammonia lyase, ethylene and abscisic acid synthesis mainly (*PRI*, *JA*, *GAME1*, *PAL5-3*, *ET* and *ABA*), which, as a whole, increased the tolerance to Fol in tomato; this effect of silicic acid was observed in the first days of the transplant, since these genes were subsequently decreased, as well as reduced glutathione, DPPH, CAT and jasmonic acid was reduced.

LITERATURE CITED

- Anderson, J. P., Badruzsaufari, E., Schenk, P. M., Manners, J. M., Desmond, O. J., Ehlert, C., Maclean, D. J., Ebert, P. R., & Kazan, K. (2004). Antagonistic Interaction between Abscisic Acid and Jasmonate-Ethylene Signaling Pathways Modulates Defense Gene Expression and Disease Resistance in Arabidopsis. *The Plant Cell*, 16(12), 3460–3479. <https://doi.org/10.1105/tpc.104.025833>
- Anstead, J., Samuel, P., Song, N., Wu, C., Thompson, G. A., & Goggin, F. (2010). Activation of ethylene-related genes in response to aphid feeding on resistant and susceptible melon and tomato plants. *Entomologia Experimentalis et Applicata*, 134(2), 170–181. <https://doi.org/10.1111/j.1570-7458.2009.00945.x>
- Arvouet-Grand, A., Vennat, B., Pourrat, A., & Legret, P. (1994). Standardization of propolis extract and identification of principal constituents. *Journal de Pharmacie de Belgique*, 49(6), 462–468.
- Aybeke, M. (2017). *Fusarium* infection causes genotoxic disorders and antioxidant-based damages in *Orobanche* spp. *Microbiological Research*, 201, 46–51. <https://doi.org/10.1016/j.micres.2017.05.001>
- Babalar, M., Edrisi, B., & Naderi, R. (2016). Evaluation of the Mechanical Strength of Gerbera Flower Stem in Response to Silicon and Salicylic Acid Application. *Journal of Ornamental Plants*, 6(3), 163–171.
- Bakhat, H. F., Bibi, N., Zia, Z., Abbas, S., Hammad, H. M., Fahad, S., Ashraf, M. R., Shah, G.

- M., Rabbani, F., & Saeed, S. (2018). Silicon mitigates biotic stresses in crop plants: A review. *Crop Protection*, *104*, 21–34. <https://doi.org/10.1016/j.cropro.2017.10.008>
- Bathoova, M., Švubová, R., Bokor, B., Neděla, V., Tihlaříková, E., & Martinka, M. (2021). Silicon triggers sorghum root enzyme activities and inhibits the root cell colonization by *Alternaria alternata*. *Planta* *253*:2, 1–14. <https://doi.org/10.1007/S00425-020-03560-6>
- Benhamou, N., & Bélanger, R. R. (1998). Benzothiadiazole-mediated induced resistance to *Fusarium oxysporum* f. sp. *radicis-lycopersici* in tomato. *Plant Physiology*, *118*(4), 1203–1212. <https://doi.org/10.1104/pp.118.4.1203>
- Bhatt, D., & Sharma, G. (2018). Role of silicon in counteracting abiotic and biotic plant stresses. *International Journal of Chemical Studies*, *6*(2), 1434–1442.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, *72*(1–2), 248–254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, *28*(1), 25–30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
- Brunings, A. M., Datnoff, L. E., Ma, J. F., Mitani, N., Nagamura, Y., Rathinasabapathi, B., & Kirst, M. (2009). Differential gene expression of rice in response to silicon and rice blast fungus *Magnaporthe oryzae*. *Annals of Applied Biology*, *155*(2), 161–170. <https://doi.org/10.1111/j.1744-7348.2009.00347.x>
- Cai, K., Gao, D., Chen, J., & Luo, S. (2009). Probing the mechanisms of silicon-mediated pathogen resistance. *Plant Signaling and Behavior*, *4*(1), 1–3. <https://doi.org/10.4161/psb.4.1.7280>
- Cao, B. li, Wang, L., Gao, S., Xia, J., & Xu, K. (2017). Silicon-mediated changes in radial hydraulic conductivity and cell wall stability are involved in silicon-induced drought resistance in tomato. *Protoplasma*, *254*(6), 2295–2304. <https://doi.org/10.1007/s00709-017-1115-y>
- Carré-Missio, V., Rodrigues, F. A., Schurt, D. A., Resende, R. S., Souza, N. F. A., Rezende, D. C., Moreira, W. R., & Zambolim, L. (2014). Effect of foliar-applied potassium silicate on coffee leaf infection by *Hemileia vastatrix*. *Annals of Applied Biology*, *164*(3), 396–403. <https://doi.org/10.1111/aab.12109>
- Coninck, E., Scauflaire, J., Gollier, M., Liénard, C., Foucart, G., Manssens, G., Munaut, F., & Legrève, A. (2020). *Trichoderma atroviride* as a promising biocontrol agent in seed coating for reducing *Fusarium* damping-off on maize. *Journal of Applied Microbiology*, *129*(3), 637–651. <https://doi.org/10.1111/jam.14641>
- Cruz, M. F. A., Debona, D., Rios, J. A., Barros, E. G., & Rodrigues, F. A. (2015). Potentiation of defense-related gene expression by silicon increases wheat resistance to leaf blast. *Tropical Plant Pathology*, *40*(6), 394–400. <https://doi.org/10.1007/s40858-015-0051-7>
- Cui, X., Tao, X., Xie, Y., Fauquet, C. M., & Zhou, X. (2004). A DNA Associated with Tomato Yellow Leaf Curl China Virus Is Required for Symptom Induction. *Journal of Virology*, *78*(24), 13966–13974. <https://doi.org/10.1128/JVI.78.24.13966>
- Dallagnol, L. J., Rodrigues, F. A., Pascholati, S. F., Fortunato, A. A., & Camargo, L. E. A. (2015). Comparison of root and foliar applications of potassium silicate in potentiating post-infection defences of melon against powdery mildew. *Plant Pathology*, *64*(5), 1085–1093. <https://doi.org/10.1111/ppa.12346>
- Dann, E. K., & Le, D. P. (2017). Effects of Silicon Amendment on Soilborne and Fruit Diseases of Avocado. *Plants*, *6*(4), 51. <https://doi.org/10.3390/plants6040051>
- Dawa, K., Zaghoul, M., I. Ahmed, H., & Hamad, K. (2020). Impact of Foliar Application with Iron, Zinc, Silicon Nano Particles and Yeast on Growth, Yield and Water Use Efficiency of Tomato Plants under Water Stress Conditions. *Journal of Plant Production*, *11*(6), 523–530. <https://doi.org/10.21608/jpp.2020.106331>
- de Lamo, F. J., & Takken, F. L. W. (2020). Biocontrol by *Fusarium oxysporum* Using Endophyte-Mediated Resistance. *Frontiers in Plant Science*, *11*, 37. <https://doi.org/10.3389/fpls.2020.00037>
- Dhindsa, R. S., Plumb-Dhindsa, P., & Thorpe, T. a. (1981). Leaf senescence correlated with

- increase levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *Journal of Experimental Botany*, 32(1), 93–101. <https://doi.org/10.1093/jxb/32.1.93>
- Di, X., Gomila, J., & Takken, F. L. W. (2017). Involvement of salicylic acid, ethylene and jasmonic acid signalling pathways in the susceptibility of tomato to *Fusarium oxysporum*. *Molecular Plant Pathology*, 18(7), 1024–1035. <https://doi.org/10.1111/mpp.12559>
- Diener, A. C., & Ausubel, F. M. (2005). Resistance to *Fusarium oxysporum* 1, a dominant *Arabidopsis* disease-resistance gene, is not race specific. *Genetics*, 171(1), 305–321. <https://doi.org/10.1534/genetics.105.042218>
- Dorneles, K. R., Dallagnol, L. J., Pazdiora, P. C., Rodrigues, F. A., & Deuner, S. (2017). Silicon potentiates biochemical defense responses of wheat against tan spot. *Physiological and Molecular Plant Pathology*, 97, 69–78. <https://doi.org/10.1016/j.pmpp.2017.01.001>
- D'Addazio, V., Silva, J. V. G., Jardim, A. S., Longue, L. L., Santos, R. A. A., Fernandes, A. A., Silva, M. B., Silva, D. M., Santos, T. A., Schmildt, E. R., Pfenning, L. H., & Falqueto, A. R. (2020). Silicon improves the photosynthetic performance of black pepper plants inoculated with *Fusarium solani* f. sp. *piperis*. *Photosynthetica*, 58(3), 692–701. <https://doi.org/10.32615/ps.2019.182>
- Fagerstedt, K. V., Kukkola, E. M., Koistinen, V. V. T., Takahashi, J., & Marjamaa, K. (2010). Cell Wall Lignin is Polymerised by Class III Secretable Plant Peroxidases in Norway Spruce. *Journal of Integrative Plant Biology*, 52(2), 186–194. <https://doi.org/10.1111/j.1744-7909.2010.00928.x>
- Fauteux, F., Rémus-Borel, W., Menzies, J. G., & Bélanger, R. R. (2005). Silicon and plant disease resistance against pathogenic fungi. *FEMS Microbiology Letters*, 249(1), 1–6. <https://doi.org/10.1016/j.femsle.2005.06.034>
- Fawe, A., Abou-Zaid, M., Menzies, J. G., & Bélanger, R. R. (1998). Silicon-Mediated Accumulation of Flavonoid Phytoalexins in Cucumber. *Phytopathology*, 88(5), 396–401. <https://doi.org/10.1094/PHYTO.1998.88.5.396>
- Ferreira, H. A., Nascimento, C. W. A. do, Datnoff, L. E., Nunes, G. H. de S., Preston, W., Souza, E. B. de, & Mariano, R. de L. R. (2015). Effects of silicon on resistance to bacterial fruit blotch and growth of melon. *Crop Protection*, 78, 277–283. <https://doi.org/10.1016/j.cropro.2015.09.025>
- Flohé, L., & Günzler, W. A. (1984). Assays of glutathione peroxidase. *Methods in Enzymology*, 105, 114–120. [https://doi.org/10.1016/S0076-6879\(84\)05015-1](https://doi.org/10.1016/S0076-6879(84)05015-1)
- Forcat, S., Bennett, M. H., Mansfield, J. W., & Grant, M. R. (2008). A rapid and robust method for simultaneously measuring changes in the phytohormones ABA, JA and SA in plants following biotic and abiotic stress. *Plant Methods*, 4(1), 16. <https://doi.org/10.1186/1746-4811-4-16>
- Fortunato, A. A., da Silva, W. L., & Rodrigues, F. Á. (2014). Phenylpropanoid Pathway Is Potentiated by Silicon in the Roots of Banana Plants During the Infection Process of *Fusarium oxysporum* f. sp. *cubense*. *Phytopathology*, 104(6), 597–603. <https://doi.org/10.1094/PHYTO-07-13-0203-R>
- Frick, D., Remus, R., Sommer, M., Augustin, J., & von Blanckenburg, F. (2020). Silicon isotope fractionation and uptake dynamics of three crop plants: laboratory studies with transient silicon concentrations. *Biogeosciences*, 17, 6475–6490. <https://doi.org/10.5194/bg-2020-66>
- Ghareeb, H., Bozsó, Z., Ott, P. G., Repenning, C., Stahl, F., & Wydra, K. (2011). Transcriptome of silicon-induced resistance against *Ralstonia solanacearum* in the silicon non-accumulator tomato implicates priming effect. *Physiological and Molecular Plant Pathology*, 75(3), 83–89. <https://doi.org/10.1016/j.pmpp.2010.11.004>
- González-García, Y., Escobar-Hernández, D. I., Benavides-Mendoza, A., Morales-Díaz, A. B., Olivares-Sáenz, E., & Juárez-Maldonado, A. (2023). UV-A Radiation Stimulates Tolerance against *Fusarium oxysporum* f. sp. *lycopersici* in Tomato Plants. *Horticulturae*, 9(4), 499. <https://doi.org/10.3390/horticulturae9040499>
- Gonzalo, M. J., Lucena, J. J., & Hernández-Apaolaza, L. (2013). Effect of silicon addition on soybean (*Glycine max*) and cucumber (*Cucumis sativus*) plants grown under iron

- deficiency. *Plant Physiology and Biochemistry*, 70, 455–461. <https://doi.org/10.1016/j.plaphy.2013.06.007>
- Gulzar, N., Ali, S., Shah, M. A., & Kamili, A. N. (2021). Silicon supplementation improves early blight resistance in *Lycopersicon esculentum* Mill. by modulating the expression of defense-related genes and antioxidant enzymes. *3 Biotech*, 11(5), 1–13. <https://doi.org/10.1007/S13205-021-02789-6>
- Habibi, G. (2015). Contrastive response of *Brassica napus* L. to exogenous salicylic acid, selenium and silicon supplementation under water stress. *Archives of Biological Sciences*, 67(2), 397–404. <https://doi.org/10.2298/ABS140411006H>
- Hajiboland, R., Bahrami-Rad, S., & Poschenrieder, C. (2017). Silicon modifies both a local response and a systemic response to mechanical stress in tobacco leaves. *Biologia Plantarum*, 61(1), 187–191. <https://doi.org/10.1007/s10535-016-0633-3>
- Hajiboland, Roghieh, Moradtalab, N., Eshaghi, Z., & Feizy, J. (2017). Effect of silicon supplementation on growth and metabolism of strawberry plants at three developmental stages. *New Zealand Journal of Crop and Horticultural Science*, 46(2), 144–161. <https://doi.org/10.1080/01140671.2017.1373680>
- Han, X., & Kahmann, R. (2019). Manipulation of phytohormone pathways by effectors of filamentous plant pathogens. *Frontiers in Plant Science*, 10, 822. <https://doi.org/10.3389/fpls.2019.00822>
- Hao, Z., Wang, L., He, Y., Liang, J., & Tao, R. (2011). Expression of defense genes and activities of antioxidant enzymes in rice resistance to rice stripe virus and small brown planthopper. *Plant Physiology and Biochemistry*, 49(7), 744–751. <https://doi.org/10.1016/j.plaphy.2011.01.014>
- Hoffmann, J., Berni, R., Hausman, J. F., & Guerriero, G. (2020). A review on the beneficial role of silicon against salinity in non-accumulator crops: Tomato as a model. *Biomolecules*, 10(9), 1284. <https://doi.org/10.3390/biom10091284>
- Hu, J., Li, Y., & Jeong, B. R. (2020). Silicon Alleviates Temperature Stresses in Poinsettia by Regulating Stomata, Photosynthesis, and Oxidative Damages. *Agronomy*, 10(9), 1419. <https://doi.org/10.3390/agronomy10091419>
- Hussain, S., Shuxian, L., Mumtaz, M., Shafiq, I., Iqbal, N., Brestic, M., Shoaib, M., Sisi, Q., Li, W., Mei, X., Bing, C., Zivcak, M., Rastogi, A., Skalicky, M., Hejnak, V., Weiguo, L., & Wenyu, Y. (2021). Foliar application of silicon improves stem strength under low light stress by regulating lignin biosynthesis genes in soybean (*Glycine max* (L.) Merr.). *Journal of Hazardous Materials*, 401, 123256. <https://doi.org/10.1016/j.jhazmat.2020.123256>
- Iqbal, M. J., Yaegashi, S., Ahsan, R., Shopinski, K. L., & Lightfoot, D. A. (2005). Root response to *Fusarium solani* f. sp. *glycines*: Temporal accumulation of transcripts in partially resistant and susceptible soybean. *Theoretical and Applied Genetics*, 110(8), 1429–1438. <https://doi.org/10.1007/s00122-005-1969-9>
- Islam, W., Naveed, H., Zaynab, M., Huang, Z., & Chen, H. Y. H. (2019). Plant defense against virus diseases; growth hormones in highlights. *Plant Signaling & Behavior*, 14(6), 1596719. <https://doi.org/10.1080/15592324.2019.1596719>
- Islam, W., Tayyab, M., Khalil, F., Hua, Z., Huang, Z., & Chen, H. Y. H. (2020). Silicon-mediated plant defense against pathogens and insect pests. *Pesticide Biochemistry and Physiology*, 168(April), 104641. <https://doi.org/10.1016/j.pestbp.2020.104641>
- Jafari, S. R., Arvin, S. M. J., & Kalantari, K. M. (2015). Response of cucumber (*Cucumis sativus* L.) seedlings to exogenous silicon and salicylic acid under osmotic stress. *Acta Biologica Szegediensis*, 59(1), 25–33.
- Jang, S.-W., Kim, Y., Khan, A. L., Na, C.-I., & Lee, I.-J. (2018). Exogenous short-term silicon application regulates macro-nutrients, endogenous phytohormones, and protein expression in *Oryza sativa* L. *BMC Plant Biology*, 18(1), 4. <https://doi.org/10.1186/s12870-017-1216-y>
- Jeandet, P., Douillet-Breuil, A. C., Bessis, R., Debord, S., Sbaghi, M., & Adrian, M. (2002). Phytoalexins from the vitaceae: Biosynthesis, phytoalexin gene expression in transgenic plants, antifungal activity, and metabolism. *Journal of Agricultural and Food*

- Chemistry*, 50(10), 2731–2741. <https://doi.org/10.1021/jf011429s>
- Kang, G., Li, G., & Guo, T. (2014). Molecular mechanism of salicylic acid-induced abiotic stress tolerance in higher plants. *Acta Physiologiae Plantarum*, 36(9), 2287–2297. <https://doi.org/10.1007/s11738-014-1603-z>
- Kim, Y.-H., Khan, A. L., Waqas, M., Jeong, H.-J., Kim, D.-H., Shin, J. S., Kim, J.-G., Yeon, M.-H., & Lee, I.-J. (2014). Regulation of jasmonic acid biosynthesis by silicon application during physical injury to *Oryza sativa* L. *Journal of Plant Research*, 127(4), 525–532. <https://doi.org/10.1007/s10265-014-0641-3>
- Kramell, R., Atzorn, R., Schneider, G., Miersch, O., Brückner, C., Schmidt, J., Sembdner, G., & Parthier, B. (1995). Occurrence and identification of jasmonic acid and its amino acid conjugates induced by osmotic stress in barley leaf tissue. *Journal of Plant Growth Regulation*, 14(1), 29–36. <https://doi.org/10.1007/BF00212643>
- Kuai, J., Sun, Y., Guo, C., Zhao, L., Zuo, Q., Wu, J., & Zhou, G. (2017). Root-applied silicon in the early bud stage increases the rapeseed yield and optimizes the mechanical harvesting characteristics. *Field Crops Research*, 200, 88–97. <https://doi.org/10.1016/j.fcr.2016.10.007>
- Leyva-Mir, S. G., González-Solano, C. M., Rodríguez-Pérez, J. E., & Montalvo-Hernández, D. (2013). Behavior of advanced lines of tomato (*Solanum lycopersicum* L.) to phytopathogens at Chapingo, Mexico | Comportamiento de líneas avanzadas de tomate (*Solanum lycopersicum* L.) a fitopatógenos en Chapingo, México. *Revista Chapingo, Serie Horticultura*, 19(3), 301–313. <https://doi.org/10.5154/r.rchsh.2012.12.070>
- Limon-Corona, M. L., Flores-Olivas, A., Hernández-Castillo, F. D., González-Merino, A. M., Gonzalez-Morales, S., Virgen-Calleros, G., Cerda-García, P. A., & Hernández-Martínez, R. (2022). Efecto de consorcios microbianos en el patosistema *Solanum lycopersicum* - *Fusarium oxysporum* f. sp. *lycopersici*. *Ecosistemas y Recursos Agropecuarios*, 9(3), 3221. <https://doi.org/10.19136/era.a9n3.3221>
- López-Pérez, M. C., Pérez-Labrada, F., Ramírez-Pérez, L. J., Juárez-Maldonado, A., Morales-Díaz, A. B., González-Morales, S., García-Dávila, L. R., García-Mata, J., & Benavides-Mendoza, A. (2018). Dynamic Modeling of Silicon Bioavailability, Uptake, Transport, and Accumulation: Applicability in Improving the Nutritional Quality of Tomato. *Frontiers in Plant Science*, 9, 1–12. <https://doi.org/10.3389/fpls.2018.00647>
- Madany, M. M. Y., Saleh, A. M., Habeeb, T. H., Hozzein, W. N., & AbdElgawad, H. (2020). Silicon dioxide nanoparticles alleviate the threats of broomrape infection in tomato by inducing cell wall fortification and modulating ROS homeostasis. *Environmental Science: Nano*, 7(5), 1415–1430. <https://doi.org/10.1039/C9EN01255A>
- Maghsoudi, K., Emam, Y., & Ashraf, M. (2015). Influence of foliar application of silicon on chlorophyll fluorescence, photosynthetic pigments, and growth in water-stressed wheat cultivars differing in drought tolerance. *Turkish Journal of Botany*, 39(4), 625–634. <https://doi.org/10.3906/bot-1407-11>
- Malhotra, C. C., Kapoor, R., & Ganjewala, D. (2016). Alleviation of abiotic and biotic stresses in plants by silicon supplementation. *Scientia Agriculturae*, 13(2), 59–73. <https://doi.org/10.15192/PSCP.SA.2016.13.2.5973>
- Manivannan, A., & Ahn, Y.-K. (2017). Silicon Regulates Potential Genes Involved in Major Physiological Processes in Plants to Combat Stress. *Frontiers in Plant Science*, 8, 1–13. <https://doi.org/10.3389/fpls.2017.01346>
- Mauch-Mani, B., & Mauch, F. (2005). The role of abscisic acid in plant – pathogen interactions. *Current Opinion in Plant Biology*, 8(4), 409–414. <https://doi.org/10.1016/j.pbi.2005.05.015>
- Michelena, G., Carrera, E., Bell, A., Altuna, B., & Almeida, G. (2001). Procesos de separación y decoloración del ácido jasmónico a partir de *Botryodiplodia theobromae* cepa 715. *Tecnología, Ciencia, Educación (IMIQ)*, 16(1), 12–19.
- Mo, Z., Lei, S., Ashraf, U., Khan, I., Li, Y., Pan, S., Duan, M., Tian, H., & Tang, X. (2017). Silicon fertilization modulates 2-acetyl-1-pyrroline content, yield formation and grain quality of aromatic rice. *Journal of Cereal Science*, 75, 17–24. <https://doi.org/10.1016/j.jcs.2017.03.014>

- Nagata, M., & Yamashita, I. (1992). Simple Method for Simultaneous Determination of Chlorophyll and Carotenoids in Tomato Fruit. *NIPPON SHOKUHIN KOGYO GAKKAISHI*, 39(10), 925–928. <https://doi.org/10.3136/nskkk1962.39.925>
- Nakano, Y., & Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology*, 22(5), 867–880.
- Ning, D., Song, A., Fan, F., Li, Z., & Liang, Y. (2014). Effects of Slag-Based Silicon Fertilizer on Rice Growth and Brown-Spot Resistance. *PLoS ONE*, 9(7), e102681. <https://doi.org/10.1371/journal.pone.0102681>
- Nitsch, L. M. C., Oplaat, C., Feron, R., Ma, Q., Wolters-Arts, M., Hedden, P., Mariani, C., & Vriezen, W. H. (2009). Abscisic acid levels in tomato ovaries are regulated by LeNCED1 and SICYP707A1. *Planta*, 229(6), 1335–1346. <https://doi.org/10.1007/s00425-009-0913-7>
- Ouzounidou, G., Giannakoula, A., Ilias, I., & Zamanidis, P. (2016). Alleviation of drought and salinity stresses on growth, physiology, biochemistry and quality of two *Cucumis sativus* L. cultivars by Si application. *Revista Brasileira de Botanica*, 39(2), 531–539. <https://doi.org/10.1007/s40415-016-0274-y>
- Padayatt, S. J., Daruwala, R., Wang, Y., Eck, P. K., Song, J., Koh, W. S., & Levine, M. (2001). Vitamin C: from molecular actions to optimum intake. *Handbook of Antioxidants*. CADENAS, E.; PACKER, L. (Eds) 2nd Edition. CRC Press. Washington DC, EE. UU, 117–145. <https://doi.org/doi:10.1201/9780203904046.pt3>
- Paye, W., Tubana, B., Harrell, D., Babu, T., Kanke, Y., & Datnoff, L. (2018). Determination of Critical Soil Silicon Levels for Rice Production in Louisiana Using Different Extraction Procedures. *Communications in Soil Science and Plant Analysis*, 49(17), 2091–2102. <https://doi.org/10.1080/00103624.2018.1495731>
- Pei, Y., Zhu, Y., Jia, Y., Ge, X., Li, X., Li, F., & Hou, Y. (2020). Molecular evidence for the involvement of cotton GhGLP2, in enhanced resistance to *Verticillium* and *Fusarium* Wilts and oxidative stress. *Scientific Reports*, 10(1), 1–15. <https://doi.org/10.1038/s41598-020-68943-x>
- Pirayesh, S., Zamanizadeh, H., & Morid, B. (2018). Molecular identification of physiological races of *Fusarium oxysporum* f. sp. *lycopersici* and radialis *lycopersici* causal agent of fusarium wilt of tomato in Iran. *Journal of Agricultural Science and Technology*, 20(1), 193–202.
- Rahman, A., Wallis, C. M., & Uddin, W. (2015). Silicon-Induced Systemic Defense Responses in Perennial Ryegrass Against Infection by *Magnaporthe oryzae*. *Phytopathology*, 105(6), 748–757. <https://doi.org/10.1094/PHYTO-12-14-0378-R>
- Ramírez, P. G., Ramírez, D. G., Mejía, E. Z., Ocampo, S. A., Díaz, C. N., & Rojas Martínez, R. I. (2020). Extracts of *Stevia rebaudiana* against *Fusarium oxysporum* associated with tomato cultivation. *Scientia Horticulturae*, 259, 108683. <https://doi.org/10.1016/j.scienta.2019.108683>
- Ramos, S. J., Faquin, V., Guilherme, L. R. G., Castro, E. M., Ávila, F. W., Carvalho, G. S., Bastos, C. E. A., & Oliveira, C. (2010). Selenium biofortification and antioxidant activity in lettuce plants fed with selenate and selenite. *Plant, Soil and Environment*, 56(12), 584–588.
- Re, R., Pellegrini, N., Proteggente, A., & Pannala, A. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology And medicine*, 26(9-10), 1231-1237. <http://www.sciencedirect.com/science/article/pii/S0891584998003153>
- Rodrigues, F. Á., McNally, D. J., Datnoff, L. E., Jones, J. B., Labbé, C., Benhamou, N., Menzies, J. G., & Bélanger, R. R. (2004). Silicon enhances the accumulation of diterpenoid phytoalexins in rice: A potential mechanism for blast resistance. *Phytopathology*, 94(2), 177–183. <https://doi.org/10.1094/PHYTO.2004.94.2.177>
- Romero, A., Munévar, F., & Cayón, G. (2011). Silicon and plant diseases. A review. *Agronomía Colombiana*, 29(3), 473–480.
- Sakr, N. (2021). Soluble Silicon Controls *Fusarium* Head Blight in Bread and Durum Wheat Plants. *Gesunde Pflanzen* 73(4), 479-493. <https://doi.org/10.1007/S10343-021-00568-0>

- Sattar, A., Cheema, M. A., Sher, A., Ijaz, M., Wasaya, A., Yasir, T. A., Abbas, T., & Hussain, M. (2020). Foliar Applied Silicon Improves Water Relations, Stay Green and Enzymatic Antioxidants Activity in Late Sown Wheat. *Silicon*, 12(1), 223–230. <https://doi.org/10.1007/s12633-019-00115-7>
- Shetty, R., Frette, X., Jensen, B., Shetty, N. P., Jensen, J. D., Jorgensen, H. J. L., Newman, M.-A., & Christensen, L. P. (2011). Silicon-Induced Changes in Antifungal Phenolic Acids, Flavonoids, and Key Phenylpropanoid Pathway Genes during the Interaction between Miniature Roses and the Biotrophic Pathogen *Podosphaera pannosa*. *Plant Physiology*, 157(4), 2194–2205. <https://doi.org/10.1104/pp.111.185215>
- Siddiqui, Z. A., Hashmi, A., Khan, M. R., & Parveen, A. (2020). Management of bacteria *Pectobacterium carotovorum*, *Xanthomonas campestris* pv. *carotae*, and fungi *Rhizoctonia solani*, *Fusarium solani* and *Alternaria dauci* with silicon dioxide nanoparticles on carrot. *International Journal of Vegetable Science*, 26(6), 547–557. <https://doi.org/10.1080/19315260.2019.1675843>
- Singh, V. K., Singh, H. B., & Upadhyay, R. S. (2017). Role of fusaric acid in the development of ‘Fusarium wilt’ symptoms in tomato: Physiological, biochemical and proteomic perspectives. *Plant Physiology and Biochemistry*, 118, 320–332. <https://doi.org/10.1016/j.plaphy.2017.06.028>
- Solarte, R. D., Osorio, O., Hurtado, A. M., & Mejia, D. F. (2012). Evaluación del Bioinsumo de Fique Pulverizado (*Furcraea* spp) para el Control *in vitro* de *Phytophthora infestans* en papa (*Solanum tuberosum* L). *Informacion Tecnologica*, 23(3), 77–86. <https://doi.org/10.4067/S0718-07642012000300010>
- Sousa, E. O., Miranda, C. M. B. A., Nobre, C. B., Boligon, A. A., Athayde, M. L., & Costa, J. G. M. (2015). Phytochemical analysis and antioxidant activities of *Lantana camara* and *Lantana montevidensis* extracts. *Industrial Crops and Products*, 70, 7–15. <https://doi.org/10.1016/j.indcrop.2015.03.010>
- Steiner, A. A. (1961). A universal method for preparing nutrient solutions of a certain desired composition. *Plant and Soil*, 15(2), 134–154. <https://doi.org/10.1007/BF01347224>
- Sun, H., Duan, Y., Mitani-Ueno, N., Che, J., Jia, J., Liu, J., Guo, J., Ma, J. F., & Gong, H. (2020). Tomato roots have a functional silicon influx transporter but not a functional silicon efflux transporter. *Plant, Cell & Environment*, 43(3), 732–744.
- Sun, S., Yang, Z., Song, Z., Wang, N., Guo, N., Niu, J., Liu, A., Bai, B., Ahammed, G. J., & Chen, S. (2022). Silicon enhances plant resistance to Fusarium wilt by promoting antioxidant potential and photosynthetic capacity in cucumber (*Cucumis sativus* L.). *Frontiers in Plant Science*, 13: 1011859. <https://doi.org/10.3389/fpls.2022.1011859>
- Syklowska-Baranek, K., Pietrosiuk, A., Naliwajski, M. R., Kawiak, A., Jeziorek, M., Wyderska, S., Łojkowska, E., & Chinou, I. (2012). Effect of l-phenylalanine on PAL activity and production of naphthoquinone pigments in suspension cultures of *Arnebia euchroma* (Royle) Johnst. *In Vitro Cellular and Developmental Biology - Plant*, 48(5), 555–564. <https://doi.org/10.1007/s11627-012-9443-2>
- Telles Nascimento, K. J., Debona, D., Silveira, P. R., Silva, L. C., DaMatta, F. M., & Rodrigues, F. Á. (2016). Silicon-Induced Changes in the Antioxidant System Reduce Soybean Resistance to Frogeye Leaf Spot. *Journal of Phytopathology*, 164(10), 768–778. <https://doi.org/10.1111/jph.12497>
- Van Bockhaven, J., Steppe, K., Bauweraerts, I., Kikuchi, S., Asano, T., Höfte, M., & De Vleeschauwer, D. (2015). Primary metabolism plays a central role in moulding silicon-inducible brown spot resistance in rice. *Molecular Plant Pathology*, 16(8), 811–824. <https://doi.org/10.1111/mpp.12236>
- van den Berg, B., Pedebos, C., Bolla, J. R., Robinson, C. V., Baslé, A., & Khalid, S. (2021). Structural Basis for Silicic Acid Uptake by Higher Plants. *Journal of Molecular Biology*, 433(21), 167226. <https://doi.org/10.1016/j.jmb.2021.167226>
- Vivancos, J., Labbé, C., Menzies, J. G., & Bélanger, R. R. (2015). Silicon-mediated resistance of Arabidopsis against powdery mildew involves mechanisms other than the salicylic acid (SA)-dependent defence pathway. *Molecular Plant Pathology*, 16(6), 572–582. <https://doi.org/10.1111/mpp.12213>

Recibido:
21/abril/2023

Aceptado:
9/enero/2024

- Wang, M., Gao, L., Dong, S., Sun, Y., Shen, Q., & Guo, S. (2017). Role of Silicon on Plant–Pathogen Interactions. *Frontiers in Plant Science*, 8, 701. <https://doi.org/10.3389/fpls.2017.00701>
- Wang, Z., Jia, C., Li, J., Huang, S., Xu, B., & Jin, Z. (2015). Activation of salicylic acid metabolism and signal transduction can enhance resistance to *Fusarium* wilt in banana (*Musa acuminata* L. AAA group, cv. Cavendish). *Functional and Integrative Genomics*, 15(1), 47–62. <https://doi.org/10.1007/s10142-014-0402-3>
- Whan, J. A., Dann, E. K., & Aitken, E. A. B. (2016). Effects of silicon treatment and inoculation with *Fusarium oxysporum* f. sp. *vasinfectum* on cellular defences in root tissues of two cotton cultivars. *Annals of Botany*, 118(2), 219–226. <https://doi.org/10.1093/aob/mcw095>
- Xue, T., Hartikainen, H., & Piironen, V. (2001). Antioxidative and growth-promoting effect of selenium on senescing lettuce. *Plant and Soil*, 237(1), 55–61. <https://doi.org/10.1023/A:1013369804867>
- Yu, Z., & Dahlgren, R. A. (2000). Evaluation of Methods for Measuring Polyphenols in Conifer Foliage. *Journal of Chemical Ecology* 26(9), 2119–2140. <https://doi.org/10.1023/A:1005568416040>
- Zvirin, T., Herman, R., Brotman, Y., Denisov, Y., Belausov, E., Freeman, S., & Perl-Treves, R. (2010). Differential colonization and defence responses of resistant and susceptible melon lines infected by *Fusarium oxysporum* race 1·2. *Plant Pathology*, 59(3), 576–585. <https://doi.org/10.1111/j.1365-3059.2009.02225.x>