

PHYSALAEMUS CARRIZORUM (ANURA, LEPTODACTYLIDAE): FIRST RECORD FROM PARAGUAY

PHYSALAEMUS CARRIZORUM (ANURA, LEPTODACTYLIDAE): PRIMER REGISTRO PARA PARAGUAY

FLAVIA NETTO^{1,2*} & FRANCISCO BRUSQUETTI²

¹Itaipu Binacional, División de Áreas Protegidas, Dirección de Coordinación Ejecutiva, Av. Monseñor Rodríguez 150, Ciudad del Este, Alto Paraná, Paraguay.

²Instituto de Investigación Biológica del Paraguay, Del Escudo 1607, CP 1425, Asunción, Paraguay

*Correspondence: flanettosisa@gmail.com

Received: 2022-12-01. Accepted: 2022-12-26. Published: 2023-02-15.

Editor: María Laura Ponssa, Argentina.

Paraguay: Itapúa Department: Alto Verá District: reserve for National Park San Rafael, Urutaú trail (26.631117° S, 55.664883° W, Datum WGS 84, elevation 268 m a.s.l.). Two males collected on 5 September 2008 and 9 December 2008. Both specimens were collected in pitfall traps located on a secondary forest next to rocky bed stream. Collection was authorized by the Ministry of Environment and Sustainable Development of Paraguay (MADES, #57038/2008). Specimens were anesthetized with lidocaine, fixed in 10% formalin followed by long-term storage in 70% ethyl alcohol. Voucher specimens are deposited in the Herpetological collection of the Instituto de Investigación

Biológica del Paraguay (IIBP-H), Asunción, Paraguay under the numbers IIBP-H1347 and IIBP-H1373, both collected by Flavia Netto. This is the first record of *Physalaemus carrizorum* from Paraguay. We extend the distribution of the species in 91 km from nearest records in Argentina, Province of Misiones, Department of Cainguás, 2 de Mayo locality (Cardozo & Pereyra, 2018; Fig. 1).

Physalaemus comprises 50 recognized species (Frost, 2021) widely distributed in South America east of the Andes. *Physalaemus gracilis* species group was proposed by Lynch (1970)




Figura 1. Mapa que muestra el nuevo registro de *Physalaemus carrizorum* de Paraguay y localidades previamente conocidas de Argentina.

Figure 1. Map showing the new record of *Physalaemus carrizorum* from Paraguay and previously known localities from Argentina.




Figura 2. Detalle de los principales caracteres diagnósticos de *Physalaemus carrizorum*, espécimen IIBP-H1347. (A) vista ventral, la barra blanca corresponde a 10 mm; (B) vista dorsal, las flechas rojas resaltan las glándulas inguinales, la barra blanca corresponde a 10 mm; (C) vista lateral, pliegue supratimpánico resaltado en cuadrado rojo; (D) detalles de la vista lateral, las flechas rojas resaltan el pliegue supratimpánico.

Figure 2. Details of main diagnostic characters of *Physalaemus carrizorum*, specimen IIBP-H1347. (A) ventral view, white bar corresponds to 10 mm; (B) dorsal view, red arrows highlight inguinal glands, white bar correspond to 10 mm; (C) lateral view, red square highlight supratympanic fold; (D) details of lateral view, red arrows highlight supratympanic fold.

and supported in Nascimento et al. (2005) and Lourenço et al. (2015). Currently, six species are included in this group: *P. carrizorum* Cardozo and Pereyra, 2018, *P. barrioi* Bokermann, 1967, *P. evangelistai* Bokermann, 1967, *P. gracilis* (Boulenger, 1883), *P. jordanensis* Bokermann, 1967 and *P. lisei* Braun and Braun, 1977. *Physalaemus carrizorum*, historically misidentified as *P. gracilis*,

was described based on external morphology, advertisement call, and 16S genetic distance. Cardozo and Pereyra (2018) mentioned that the species only occurs in Argentina, Misiones Province, located in the Atlantic Forest domain, and that it is likely to occur in neighboring Brazil.

Several authors (Boettger, 1885; Boulenger, 1894; Peracca, 1895; Bertoni, 1914; 1939) have mentioned the presence of *P. gracilis* in Paraguay, which could be referring to *P. carrizorum*, however until now there were no voucher specimens to support its occurrence in the country. The presence of *P. carrizorum* (as *P. gracilis*) in Paraguay was considered probable by Brusquetti and Lavilla (2006) due to geographic proximity localities of Argentinians records (Misiones) and southeastern Brazil (Rio Grande do Sul, Paraná and Santa Catarina), and because of habitat similarity.

Identification of the specimens from Paraguay was confirmed based on Cardozo and Pereyra (2018). Both specimens present the character states that diagnose *P. carrizorum* from all the species of *Physalaemus* not belonging to the *P. gracilis* group (except *P. riograndensis*, from *P. biligonigerus* group): a median stripe defined by the absence of melanocytes on throat, chest, and/or abdomen (Fig. 2A); from *P. riograndensis*: presence of inguinal glands (absent in *P. riograndensis*) (Fig. 2B), larger body size (snout-vent length of IIBP-H1347 is 28.57 mm, of IIBP-H1373 is 28.52 mm; smaller than 20.5 mm in *P. riograndensis*, Cardoso & Pereyra, 2018) and dorsal skin texture near smooth (tuberculate in *P. riograndensis*) (Fig. 2B); from the remaining species of the *P. gracilis* group: supratympanic fold developed (poorly developed in *P. barrioi*, Provete et al., 2012; not evident in the rest of the species) (Figs. 2C and D).

Although the mentioned specimens were collected in an area proposed as reserve for a national park, currently is divided between more than 60 private owners with a great and increasingly serious risk of deforestation. Taking into account that this is the only known population of *P. carrizorum* on the right bank of the Paraná River, and that is a small and isolated population, studies on genetic diversity and its protection should be a priority.

Acknowledgements.— We thank the owners of PROCOSARA Christine Kohli and Hans Hostettler for allowing access to their properties and for protecting the area. We thank the Ministry of Environment and Sustainable Development of Paraguay (MADES) for the collection permits and the Consejo Nacional de Ciencia y Tecnología (CONACYT, Paraguay, Programa Nacional de Incentivo a Investigadores – PRONII), for financial support.

CITED LITERATURE

- Bertoni, A. de W. 1914. Fauna Paraguaya. Catálogos sistemáticos de los vertebrados del Paraguay. Peces, batracios, reptiles, aves y mamíferos conocidos hasta 1913. Descripción Física y Económica del Paraguay 59:1-83.
- Bertoni, A. de W. 1939. Catálogos sistemáticos de los vertebrados del Paraguay. Revista de la Sociedad Científica del Paraguay 4:3-59.
- Boettger, O. 1885. Liste von Reptilien und Amphibien aus Paraguay. Zeitschrift für Naturforschung 58:213-248.
- Boulenger, G.A. 1894. Second report on additions to the lizard collections in the Natural History Museum. Proceedings of the Zoological Society London 1894:722-736.
- Brusquetti, F. & E.O. Lavilla. 2006. Lista comentada de los anfibios de Paraguay. Cuadernos de Herpetología 20:3-79.
- Cardozo, D.E. & M.O. Pereyra. 2018. A new species of *Physalaemus* (Anura, Leptodactylidae) from the Atlantic Forest of Misiones, northeastern Argentina. Zootaxa 4387:580-590.
- Frost, D.R. 2021. Amphibian Species of the World: an Online Reference. Version 6.1 (November 2022). Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA.
- Lynch, J.D. 1970. Systematic status of the American leptodactylid frog genera *Engystomops*, *Eupemphix*, and *Physalaemus*. Copeia 1970:488-496.
- Lourenço, L.B., C.P. Targueta, D. Baldo, J. Nascimento, P.C. Garcia, G.V. Andrade, C.F.B. Haddad & S.M. Recco-Pimentel. 2015. Phylogeny of frogs from the genus *Physalaemus* (Anura, Leptodactylidae) inferred from mitochondrial and nuclear gene sequences. Molecular Phylogenetics and Evolution 92:204-216.
- Nascimento, L.B., U. Caramaschi & C.A.G. Cruz. 2005. Taxonomic review of the species groups of the genus *Physalaemus* Fitzinger, 1826 with revalidation of the genera *Engystomops* Jiménez de la Espada, 1872 and *Eupemphix* Steindachner, 1863 (Amphibia, Anura, Leptodactylidae). Arquivos do Museu Nacional 63:297-320.
- Peracca, M.G. 1895. Viaggio del dott. Alfredo Borelli nella Republica Argentina en el Paraguay. Bolletino dei Musei di Zoologia ed Anatomia Comparata 10:1-32.
- Provete, D.B., M.V. Garey, L.F. Toledo, J. Nascimento, L.B. Lourenço, D.C. Rossa-Feres & C.F.B. Haddad. 2012. Redescription of *Physalaemus barrioi* (Anura: Leiuperidae). Copeia 2012:507-518.

