



Digenean parasites of Ciconiiform birds from Argentina

Digéneos parásitos de aves Ciconiiformes de Argentina

Fabiana B. Drago* and Lía I. Lunaschi

Laboratorio de Helmintología, División Zoología Invertebrados, Museo de La Plata, Facultad de Ciencias Naturales y Museo, Paseo del Bosque S/Nº, 1900 La Plata, Buenos Aires, Argentina.

*Correspondent: fdrago@fcnym.unlp.edu.ar

Abstract. The helminthological survey of 13 specimens of 5 ciconiiform species revealed the presence of 10 species of intestinal digeneans: *Sphincterodiplostomum musculosum* Dubois, 1936 in *Tigrisoma lineatum* and *Ardea alba*; *Tylodelphys elongata* (Lutz, 1928) in *T. lineatum*; *Apharyngostrigea ardearum* (Lutz, 1928) and *Dolichorchis lacombeensis* Lunaschi and Drago, 2006 in *Ardea cocoi* and *Ardea alba*; *Posthodiplostomum nanum* Dubois, 1937 in *A. alba* and *T. lineatum*; *Ascocotyle diminuta* (Stunkard and Haviland, 1924) in *A. alba*; *Petasiger* sp.1 in *A. cocoi* and *Paryphostomum segregatum* Dietz, 1909, *Strigea vaginata* (Brandes, 1888) and *Petasiger* sp. 2 in *Coragyps atratus*. *Syrigma sibilatrix* was free of digeneans. The reports of *A. ardearum*, *T. elongata*, *S. musculosum* and *D. lacombeensis* represent new host records; the other reports are new geographical records. Moreover, the specimens described as *Apharyngostrigea brasiliensis* Szidat, 1929 by Labriola and Suriano (1998) were analyzed and transferred to *A. ardearum*.

Key words: Digenea, Ardeidae, Cathartidae, Strigidae, Diplostomidae, Echinostomatidae, Heterophyidae.

Resumen. El estudio parasitológico de 13 ejemplares pertenecientes a 5 especies de aves Ciconiiformes reveló la presencia de 10 especies de digéneos intestinales: *Sphincterodiplostomum musculosum* Dubois, 1936 en *Tigrisoma lineatum* y *Ardea alba*; *Tylodelphys elongata* (Lutz, 1928) en *T. lineatum*; *Apharyngostrigea ardearum* (Lutz, 1928) y *Dolichorchis lacombeensis* Lunaschi y Drago, 2006 en *Ardea cocoi* y *A. alba*; *Posthodiplostomum nanum* Dubois, 1937 en *A. alba* y *T. lineatum*; *Ascocotyle diminuta* (Stunkard y Haviland, 1924) en *A. alba*; *Petasiger* sp.1 en *A. cocoi* y *Paryphostomum segregatum* Dietz, 1909, *Strigea vaginata* (Brandes, 1888) y *Petasiger* sp. 2 en *Coragyps atratus*. *Syrigma sibilatrix* no se encontró parasitada por digéneos. El presente estudio aporta nuevos registros de hospedadores para *A. ardearum*, *T. elongata*, *S. musculosum* y *D. lacombeensis* y nuevos registros geográficos para las restantes especies. Además, los organismos descritos como *Apharyngostrigea brasiliensis* Szidat, 1929 por Labriola y Suriano (1998) fueron analizados y transferidos a *A. ardearum*.

Palabras clave: Digenea, Ardeidae, Cathartidae, Strigidae, Diplostomidae, Echinostomatidae, Heterophyidae.

Introduction

Although 30 species of ciconiiform birds (14 Ardeidae, 8 Threskiornithidae, 3 Ciconiidae, 5 Cathartidae) are distributed Argentina, the information about its digenean fauna is scarce. At present, 8 ardeid species [*Tigrisoma lineatum* (Boddaert), *Bubulcus ibis* (L.), *Butorides striata* (L.), *Ardea alba* L., *Ardea cocoi* L., *Egretta thula* (Molina), *Ixobrychus involucris* (Vieillot), *Nycticorax nycticorax* (L.)], 4 threskiornithid species [*Ajaia ajaja* (L.), *Phimosus infuscatus* (Lichtenstein), *Plegadis chihi* (Vieillot), *Theristicus caudatus* (Boddaert)] and 2 ciconiid species [*Ciconia maguari* (Gm) and *Mycteria americana*

L.] have been reported as hosts of digeneans (Lunaschi et al., 2007; Drago and Lunaschi, 2008; Lunaschi and Drago, 2009). Most of these records were made on the basis of hosts proceeding from the Humid Pampas ecoregion or under captivity in Zoological Gardens.

The aim of this paper is to increase the knowledge of the diversity of digenean parasites from ciconiiform birds collected in the Humid Chaco and Humid Pampas ecoregions.

Material and methods

Thirteen birds were hunted with guns between 1999 and 2006 from 3 localities of Formosa Province (Bellaco stream

[26°14'S; 58°07'W] and La Marcela farm [26°17'35"S; 59°06'67"W], Pirané department, and Clorinda [25°17'S, 57°43'W], Pilcomayo department), and 1 from Buenos Aires Province (Lacombe lagoon [35° 49'S; 57° 49'W], Lezama). The bird species analyzed were *A. alba*, *A. cocoi*, *T. lineatum*, *Syrigma sibilatrix* (Temminck) and *Coragyps atratus* (Bechstein). The viscera were fixed in 10% formalin and examined for parasites in laboratory. The digenean specimens were stained with hydrochloric carmine, dehydrated and mounted in Canada balsam. The taxonomy of digeneans is given in accordance to Gibson et al. (2002), Jones et al. (2005) and Bray et al. (2008); identification of birds follows Barnett and Pearman (2001), and the parasitic indexes used were based on Bush et al. (1997). All digeneans were deposited in the Helminthological Collections of the Museo de La Plata (MLP), La Plata, Argentina, and the hosts in the Ornithological Collections of the Museo de La Plata (MLP).

Results

Family Strigeidae

Strigea vaginata (Brandes, 1888) Szidat, 1928

Taxonomic summary

Host: *Coragyps atratus*, black vulture (Cathartidae).
Locality: La Marcela farm, Pirané, Formosa Province.
Site of infection: intestine.
Prevalence and mean intensity: 67% (2 of 3); 3 (1-5).
Voucher specimen deposited: MLP 5964.

Remarks

This species was briefly described as *Holostomum vaginatum* Brandes, 1888 from *Cathartes* sp. (Cathartidae) in Brazil, and transferred to *S. vaginata* by Szidat (1928). Later, it was reported parasitizing other Brazilian birds: *C. atratus*, *Sarcoramphus papa* (L.), *Cathartes burrovianus urubitinga* Pelzeln, (Cathartidae), *Cariama cristata* (L.) (Cariamidae), and *Spizaetus ornatus* (Audin) (Accipitridae) (Szidat, 1929; Travassos et al., 1969). *Strigea vaginata* was also found parasitizing *C. atratus* and an indeterminate accipitrid (cited as “gavilán” and “gavião”) in Venezuela (Lutz, 1928; Dubois, 1970a); *Cathartes aura* (L.) in Cuba (Dubois and Macko, 1972); *Theristicus*

caudatus, *Cercibis oxycerca* (Spix) (Threskiornithidae), and *Caracara plancus* (Miller) (Falconidae) in Colombia (Dubois, 1978, 1981). According to Szidat (1929) and Dubois (1968), the specimens collected in the black vulture from Formosa Province were identified as *S. vaginata* mainly by the presence of an enormous copulatory bursa and genital cone, occupying almost half of the hindbody. This is the first report of *S. vaginata* in Argentina.

***Apharyngostrigea ardearum* (Lutz, 1928) Dubois, 1968**
 (Syn. *Apharyngostrigea brasiliiana* Szidat, 1929 of Labriola and Suriano (1998): 3)

Taxonomic summary

Hosts and localities: *Ardea alba* from Lacombe lagoon, Lezama, Buenos Aires Province and Clorinda, Formosa Province; *Ardea cocoi* from Lacombe lagoon.

Site of infection: intestine.

Prevalence and mean intensity: *A. alba*, Lacombe lagoon: 50% (2 of 4); 10 (8-12); *A. alba*, Clorinda: 1 of 1; 33; *A. cocoi*: 50% (1 of 2); 3.

Voucher specimen deposited: *A. alba* MLP 5965 from Buenos Aires Province, MLP 5966 from Formosa Province; *A. cocoi* MLP 5967.

Other material: specimens from *A. alba* (MLP 5968) and *B. ibis* (MLP 5969) from Monte Lagoon, Buenos Aires Province [of Labriola and Suriano (1998)].

Remarks

The morphological features of the specimens studied herein correspond to the description of *A. ardearum* given by Dubois (1981). This species was previously reported parasitizing herons from Venezuela by Lutz (1928) (as *Strigea ardearum* Lutz, 1928), *Ardea herodias herodias* L. and *Nycticorax n. hoactli* (Gmelin) from Cuba by Pérez Viguera (1944) [as *Apharyngostrigea cornu* (Zeder, 1800)]. Moreover, *Ardea purpurea* L. was reported as experimental host in Guadalupe (Dubois, 1981). At present, in Argentina only 1 species of *Apharyngostrigea* Ciurea, 1927, *Apharyngostrigea simplex* (Johnston, 1904), has been reported by Ostrowski de Núñez (1989) parasitizing *E. thula* from Buenos Aires Province.

In addition, specimens described as *A. brasiliiana* were reported parasitizing *A. alba*, *A. a. egretta* and *B. ibis* from Buenos Aires Province (Boero et al., 1972; Labriola and Suriano, 1998). Nevertheless, this species was previously transferred by Dubois (1964) to *Parastrigea* Szidat, 1928.

The specimens poorly described by Boero et al. (1972) do not show the main diagnostic characters of *Parastrigea* (e.i. vitelline follicles in forebody accumulated in 2 symmetrical masses localized in lateral expansions of dorsal lobe of holdfast organ). Unfortunately, their correct identification cannot be reviewed because no specimens were deposited at the appropriate time. The re-examination of the specimens studied by Labriola and Suriano (1998) revealed that they did not conform to the generic diagnosis of *Parastrigea* and their correct identification corresponds to *Apharyngostrigea*. Since the morphology and the metric characters of these specimens correspond to the description by Dubois (1981), we transfer these specimens to *A. ardearum*. The finding of this species in *A. cocoi*, *A. alba* and *B. ibis* represents new hosts and Argentina a new geographic distribution for this parasite species.

Family Diplostomidae

Sphincterodiplostomum musculosum Dubois, 1936

Taxonomic summary

Host: *Tigrisoma lineatum*, rufescent tiger-heron (Ardeidae).
Locality: Bellaco stream, Pirané, Formosa Province.
Site of infection: intestine.
Prevalence and intensity of infection: 1 of 1; 8.
Voucher specimen deposited: MLP 5970.

Remarks

The monospecific genus *Sphincterodiplostomum* Dubois, 1936 is characterized by the presence of a dorsal invagination at the level of the posterior testis equipped with a sphincter. *Sphincterodiplostomum musculosum* was described based on immature specimens collected from the intestine of *Agamia agami* (Gmelin) (Ardeidae) from Brazil (Dubois, 1936, 1938). Later, unencysted metacercariae in the orbit of the eye of *Hoplías malabaricus* (Bloch) (Erythrinidae) and mature specimens parasitizing *A. alba* from northeastern Argentina were described by Szidat (1969) and Lunaschi and Drago (2006a), respectively. The finding of adult specimens of *S. musculosum* in *T. lineatum* represents a new host record.

Tylodelphys elongata (Lutz, 1928) Dubois, 1937

Taxonomic summary

Host: *Tigrisoma lineatum*, rufescent tiger-heron (Ardeidae).
Locality: Bellaco stream, Pirané, Formosa Province, Argentina.
Site of infection: intestine.
Prevalence and intensity of infection: 1 of 1; 1.
Voucher specimen deposited: MLP 5971

Remarks

Tylodelphys elongata originally described as *Alaria elongata* Lutz, 1928 in Venezuela (Lutz, 1928), was reported parasitizing podicipedids, *Tachybaptus dominicus* (L.) from Cuba, Brazil, Venezuela and USA and *Podilymbus podiceps* (L.) from Cuba and USA (Travassos et al., 1969; Dubois, 1970b; Dubois and Macko, 1972; Storer, 2000). It was also recovered in ciconiids, *Jabiru mycteria* (Lichtenstein) from Brazil and Venezuela, and *M. americana* from Venezuela (Travassos et al., 1969; Gomes and Oliveira Rodrigues, 1981). The finding of *T. elongata* in *T. lineatum* represents a new host record, the first record in ardeid birds and the first report of this species in Argentina.

Dolichorchis lacombeensis Lunaschi and Drago, 2006

Taxonomic summary

Hosts: *Ardea cocoi*, *A. alba*.
Locality: Lacombe lagoon, Lezama, Buenos Aires Province.
Site of infection: intestine.
Prevalence and mean intensity: *A. cocoi*: 100% (2 of 2); 45 (14-76); *A. alba*: 50% (2 of 4); 2.
Voucher specimen deposited: *A. cocoi* MLP 5972; *A. alba* MLP 5973

Remarks

This species was described recently by Lunaschi and Drago (2006b) parasitizing *A. cocoi* from Buenos Aires Province. The finding of this diplostomid in *A. alba*

represents a new host record.

***Posthodiplostomum nanum* Dubois, 1937**

Taxonomic summary

Hosts and localities: *Ardea alba* from Lacombe lagoon, Lezama, Buenos Aires Province and Clorinda, Formosa Province; *Tigrisoma lineatum* from Bellaco stream, Formosa Province.

Site of infection: intestine.

Prevalence and mean intensity: *A. alba*, Lacombe lagoon: 50% (2 of 4); 9.5 (2-17); *A. alba*, Clorinda: 1 of 1; 66; *T. lineatum*: 1 of 1; 10.

Voucher specimen deposited: *A. alba* from Buenos Aires Province, MLP 5974; from Formosa Province, MLP 5977; *T. lineatum* MLP 5975.

Remarks

A wide range of birds has been reported as host of *P. nanum* in the Neotropical Region, *Butorides virescens* (L.) and *Butorides striata striata* (L.) (Ardeidae) from Brazil; *Butorides virescens maculata* (Boddaert) from Cuba and Colombia, *Pitangus sulphuratus* (L.) (Tyrannidae) and *Tigrisoma lineatum marmoratum* (Vieillot) (Ardeidae) from Venezuela, *Busarellus nigricollis* (Latham) (Accipitridae) and *B. striata* from Paraguay (Dubois, 1970a, 1970b, 1985, 1988; Gomes and Oliveira Rodrigues, 1981; Lamothe-Argumedo and Jaimes Cruz, 1982). In Argentina, it was reported in the Humid Pampas ecoregion parasitizing ardeid, *A. a. egretta*, *B. striata*, *I. involucris*, and threskiornithid, *P. chihi*, (Boero et al., 1972; Ostrowski de Núñez, 1973; Digiani, 2000). The finding of *P. nanum* parasitizing *T. lineatum* and *A. alba* in the Humid Chaco ecoregion represents a new geographical record.

***Posthodiplostomum* sp.**

Taxonomic summary

Hosts: *Ardea cocoi*.

Locality: Lacombe lagoon, Lezama, Buenos Aires Province.

Site of infection: intestine.

Prevalence and intensity of infection: 50% (1 of 2); 6.

Voucher specimen deposited: MLP 6013.

Remarks

The 6 specimens collected are immature and probably belong to *P. nanum*.

Family Echinostomatidae

***Paryphostomum segregatum* Dietz, 1909**

Taxonomic summary

Host: *Coragyps atratus*, black vulture (Cathartidae).

Locality: La Marcela farm, Pirané, Formosa Province.

Site of infection: intestine.

Prevalence and intensity of infection: 33% (1 of 3); 18.

Voucher specimen deposited: MLP 6010.

Remarks

Paryphostomum segregatum possesses a distribution restricted to the Neotropical Region, and was found in Brazil, Venezuela, Guyana and Paraguay parasitizing cathartid, *C. atratus*, *Coragyps atratus foetens* (Lichtenstein), *Cathartes aura* (L.), *Cathartes aura ruficollis* Spix, *C. b. urubitinga* and *S. papa* (Vevers, 1923; Lutz, 1928; Caballero and Diaz-Ungria, 1958; Travassos et al., 1969; Kostadinova et al., 2002). Particularly, in Argentina, it was reported in the Humid Pampas ecoregion parasitizing *Phalacrocorax brasilianus brasilianus* (Gmelin) by Ostrowski de Núñez (1968). The finding of *P. segregatum* parasitizing *C. atratus* in the Humid Chaco ecoregion represents a new geographical record.

***Petasiger* sp. 1.**

Taxonomic summary

Host: *Ardea cocoi*

Locality: Lacombe lagoon, Lezama.

Site of infection: intestine.

Prevalence and intensity of infection: 50% (1 of 2); 1.

Voucher specimen deposited: MLP 6012.

Remarks

A single adult was found in the small intestine of *A. cocoi*. This specimen was not in sufficiently good condition for species determination, but can be distinguished by having a well developed head collar with 19 spines.

Petasiger sp. 2.

Taxonomic summary

Host: *Coragyps atratus*, black vulture (Cathartidae).

Locality: La Marcela farm, Pirané, Formosa Province.

Site of infection: intestine.

Prevalence and intensity of infection: 33% (1 of 3); 2.

Voucher specimen deposited: MLP 6011.

Remarks

Only 2 specimens of *Petasiger* sp. were found in the small intestine of *C. atratus*. These echinostomatids can be distinguished from specimens obtained from *A. cocoi* by having a well developed head collar with 27 spines. Further studies are needed to identify it to the species level.

Family Heterophyidae

Ascocotyle diminuta (Stunkard and Haviland, 1924)

Taxonomic summary

Host: *Ardea alba*

Locality: Lacombe lagoon, Lezama, Buenos Aires Province.

Site of infection: intestine.

Prevalence and intensity of infection: 25% (1 of 4); 1.

Voucher specimen deposited: MLP 5976

Remarks

This species was previously found in ardeid birds, *A. alba* and *A. herodias* from Mexico and USA (Sepúlveda et al., 1999; Scholz et al., 2001). In Argentina, it was reported parasitizing to *E. thula* from a Zoological Garden in Buenos Aires Province by Ostrowski de Núñez (1993). This is the first record of *A. diminuta* parasitizing wild birds from Argentina.

Final comments

Host-parasite relationships. Among the 10 digenean species collected, the diplostomids were the group best represented (4 species), followed by the echinostomatids (3 species), strigeids (2 species), and heterophyids (1 species). In the Humid Pampas ecoregion, infected birds harbored a mean of 3 (2-4) digenean species; the most prevalent species were *A. ardearum* and *P. nanum* (67%); whereas, in the Humid Chaco ecoregion, the birds harbored a mean of 2.5 (2-3) digenean species, the most prevalent species were *P. nanum* and *S. musculosum* (33%).

The fish eating ardeids, *A. alba*, *A. cocoi* and *T. lineatum* harbored 7 digenean species whose life cycles include fishes, whereas *Syrigma sibilatrix* was found free of digeneans; this may be related to their terrestrial habitats and diet, which include predominantly arthropods, and occasionally amphibians and reptiles (Franz et al., 2007). Moreover, the great egrets from both ecoregions sampled were found parasitized by *A. ardearum* and *P. nanum*, whereas *A. diminuta* and *D. lacombeensis* were only recovered in the Humid Pampas ecoregion and *S. musculosum* was only found in the Humid Chaco. This suggests that this host species shows a little regional difference in feeding. Finally, the cathartid *C. atratus*, was found parasitized by species that were not found in the ardeids; these differences can be explained by their scavenging habits.

Acknowledgements

The authors express their gratitude to Dr Juan T. Timi, Laboratorio de Parasitología, Departamento de Biología, FCEyN, UNdMP for the loan of material described by Labriola and Suriano; Ing. Agr. Marcelo Martínez Leanes and Mr. Roberto Aranda for their help and hospitality at Centro de Pesca San Jorge, and Dr. Carlos Montoya at

Formosa Province. The present study was funded by CIC (Res. N° 578/08).

Literature cited

- Barnett, J. M. and M. Pearman. 2001. Lista comentada de las aves argentinas. Lynx Edicions, Barcelona. 164 p.
- Boero, J. J., J. E. Led and E. Brandetti. 1972. El parasitismo de la fauna autóctona. Revista de la Facultad de Agronomía y Veterinaria, Buenos Aires 1:17-29.
- Bray, R., D. Gibson and A. Jones. 2008. Keys to the Trematoda. Vol. 3. CAB International, London. 824 p.
- Bush, A. O., K. D. Lafferty, J. M. Lotz and A. W. Shostak. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. Journal of Parasitology 83:575-583.
- Caballero, E. and C. Díaz-Ungria. 1958. Intento de un catálogo de los tremátodos digéneos registrados en territorio venezolano. Memoria de la Sociedad de Ciencias Naturales La Salle 18:19-36.
- Gomes, D. C. and H. de Oliveira Rodrigues. 1981. Trematoda. In Aquatic Biota of Mexico, Central America and the West Indies, S. H. Hurlbert and A. Villalobos Figueroa (eds.). San Diego State University. 116-128 p.
- Digiani, M. C. 2000. Digeneans and cestodes parasitic in the white-faced ibis *Plegadis chihii* (Aves: Threskiornithidae) from Argentina. Folia Parasitologica 47:195-204.
- Drago, F. B. and L. I. Lunaschi. 2008. Description of a new species of *Tylodelphys* (Digenea, Diplostomidae) in the wood stork, *Mycteria americana* (Aves, Ciconiidae) from Argentina. Acta Parasitologica 53:263-267.
- Dubois, G. 1936. Nouveaux principes de classification des Trématodes du groupe des Strigeida (Notes préliminaires). Revue Suisse de Zoologie 43:507-515.
- Dubois, G. 1938. Monographie des Strigeida (Trematoda). Mémoires de la Société Neuchâteloise des Sciences Naturelles 6:1-535.
- Dubois, G. 1964. Du statut de quelques Strigeata La Rue, 1926 (Trematoda). I. Bulletin de la Société Neuchâteloise des Sciences Naturelles 87:27-71.
- Dubois, G. 1968. Synopsis des Strigeidae et des Diplostomidae (Trematoda). Mémoires de la Société Neuchâteloise des Sciences Naturelles 10:1-258.
- Dubois, G. 1970a. Les Strigeata (Trematoda) de la collection A. Lutz. Memórias do Instituto Oswaldo Cruz 68:169-196.
- Dubois, G. 1970b. Synopsis des Strigeidae et des Diplostomatidae (Trematoda). Mémoires de la Société Neuchâteloise des Sciences Naturelles 10:259-727.
- Dubois, G. 1978. Notes Helminthologiques. IV. Strigeidae Railliet, Diplostomidae Poirier, Proterodiplostomidae Dubois et Cyathocotylidae Poche (Trematoda). Revue Suisse de Zoologie 85:607-615.
- Dubois, G. 1981. Notes Helminthologiques V: Strigeidae Railliet, Diplostomidae Poirier et Proterodiplostomidae Dubois. Revue Suisse de Zoologie 88:227-232.
- Dubois, G. 1985. Quelques Strigeoidea (Trematoda) récoltés chez des oiseaux du Paraguay par la Mission Claude Weber, automne 1983, du Muséum d'Histoire Naturelle de Genève. Revue Suisse de Zoologie 92:641-648.
- Dubois, G. 1988. Quelques Strigeoidea (Trematoda) récoltés au Paraguay par les expéditions du Muséum d'Histoire Naturelle de Genève, au cours des années 1979, 1982 et 1985. Revue Suisse de Zoologie 95:521-532.
- Dubois, G. and J. Macko. 1972. Contribution à l'étude des Strigeata La Rue, 1926 (Trematoda: Strigeida) de Cuba. Annales de Parasitologie Humaine et Comparée 47:51-75.
- Franz I, I. Rohling Ghizoni, J. L. Berguer Albuquerque, A. Barcellos, C. Britz Hassdenteufel, F. Lohman Arend and C. Martins-Ferreira. 2007. Predação da cobra d'água *Helicopsis infrataeniatus* (Serpentes, Colubridae) pela maria-faceira *Syrigma sibilatrix* (Aves, Ardeidae) no sul do Brasil. Biotemas 20:135-137.
- Gibson D., A. Jones and R. Bray. 2002. Keys to the Trematoda. Vol. 1. CAB International, London. 521 p.
- Jones, A, R. Bray, and D. Gibson. 2005. Keys to the Trematoda. Vol. II. CAB International, London. 745 p.
- Kostadinova A, C. Vaucher and D. I. Gibson. 2002. Redescriptions of two echinostomes from birds in Paraguay, with comments on *Drepanocephalus* Dietz, 1909 and *Paryphostomum* Dietz, 1909 (Digenea: Echinostomatidae). Systematic Parasitology 53:147-158.
- Labriola, J. B. and D. M. Suriano. 1998. Digeneans of bird (Ardeidae) from the Monte lake, Buenos Aires, Argentina. Physis 56:1-7.
- Lamothe-Argumedo, R. and B. Jaimes Cruz. 1982. Trematoda. Parasitic stages. In Aquatic Biota of Mexico, Central America and the West Indies, S. H. Hurlbert and A. Villalobos Figueroa (eds.). San Diego State University. 73-84 p.
- Lunaschi, L. I. and F. B. Drago. 2006a. First report of adult specimens of *Sphincterodiplostomum musculosum* (Digenea, Diplostomidae). Parasitology International 55:7-10.
- Lunaschi, L. I. and F. B. Drago. 2006b. Description of a new species of *Dolichorchis* (Digenea: Diplostomidae) in the cocoi heron, *Ardea cocoi* (Aves, Ardeidae), from Argentina. Acta Parasitologica 51:47-50.
- Lunaschi, L. I. and F. B. Drago. 2009. Digenean parasites of six species of birds from Formosa province, Argentina. Revista Mexicana de Biodiversidad 80:39-46.
- Lunaschi, L. I., F. Cremona and F. B. Drago. 2007. Checklist of digenean parasites of birds from Argentina. Zootaxa 1403:1-36.
- Lutz, A. 1928. Estudios de zoología y parasitología Venezolanas. Universidad Central de Venezuela, Caracas. 133 p.
- Ostrowski de Núñez, M. 1968. Estudios sobre la fauna

- parasitaria del biguá, *Phalacrocorax o. olivaceus*. I. Trematodes pertenecientes a las familias Cathaemasidae y Echinostomatidae. *Revista del Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Parasitología*, 1:131-152.
- Ostrowski de Núñez, M. 1973. Sobre el ciclo biológico de *Posthodiplostomum nanum* Dubois, 1937 (Trematoda, Diplostomatidae). *Physis* 32:121-132.
- Ostrowski de Núñez, M. 1989. The life history of a trematode, *Apharyngostrigea simplex* (Johnston, 1904), from the ardeid bird *Egretta thula* in Argentina. *Zoologischer Anzeiger* 222:322-336.
- Ostrowski de Núñez, M. 1993. Life history studies of heterohyid trematodes in the Neotropical Region: *Ascocotyle* (*Phagicola*) *diminuta* (Stunkard & Haviland, 1924) and *A. (P.) angrense* Travassos, 1916. *Systematic Parasitology* 24:191-199.
- Pérez Viguera, I. 1944. Trematodes de la Super-Familia Strigeoidea; descripción de un género y siete especies nuevas. *Revista de la Universidad de la Habana* 52-54:294-314.
- Scholz, T., M. L. Aguirre-Macedo and G. Salgado-Maldonado. 2001. Trematodes of the family Heterophyidae (Digenea) in Mexico: A review of species and new host and geographical records. *Journal of Natural History* 35:1733-1772.
- Sepúlveda, M. S., M. G. Spalding, J. M. Kinsella and D. J. Forrester. 1999. Parasites of the great egret (*Ardea albus*) in Florida and a review of the helminths reported for the species. *Journal of the Helminthological Society of Washington* 66:7-13.
- Storer, R. W. 2000. The metazoan Parasite fauna of grebes (Aves: Podicipediformes) and its relationship to the birds biology. *Miscellaneous Publications, Museum of Zoology, University of Michigan* 188:1-90.
- Szidat, L. 1928. Zur Revision der Trematodengattung *Strigea* Abildgaard. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene* 105:204-215.
- Szidat, L. 1929. Beiträge zur Kenntnis der Gattung *Strigea* (Abildg.). I Allgemeiner Teil: Untersuchungen über die Morphologie, Physiologie und Entwicklungsgeschichte der Holostomiden Nebst Bemerkungen über Die Metamorphose Der Trematoden Und Die Phylogenie Derselben. *Zeitschrift für Parasitenkunde* 1:612-764.
- Szidat, L. 1969. Structure, development, and behaviour of new strigeatoid. *Metacercariae from subtropical fishes of South America*. *Journal of the Fisheries Research Board of Canada* 26:753-86.
- Travassos, L., J. F. Teixeira de Freitas and A. Kohn. 1969. Trematódeos do Brasil. *Memórias do Instituto Oswaldo Cruz* 67:1-886.
- Vevers, G. M. 1923. Some new and little known helminths from British Guiana. *Journal of Helminthology* 1:35-45.