

Potential economic impact assessment for cattle parasites in Mexico. Review

Evaluación del impacto económico potencial de los parásitos del ganado bovino en México. Revisión

Roger Iván Rodríguez-Vivas^a, Laerte Grisi^{bt}, Adalberto Angel Pérez de León^c, Humberto Silva Villela^d, Juan Felipe de Jesús Torres-Acosta^a, Hugo Fragoso Sánchez^e, Dora Romero Salas^f, Rodrigo Rosario Cruz^g, Fabián Saldierna^h, Dionisio García Carrasco^h

ABSTRACT

Here, economic losses caused by cattle parasites in Mexico were estimated on an annual basis. The main factors taken into consideration for this assessment included the total number of animals at risk, potential detrimental effects of parasitism on milk production or weight gain, and records of condemnation on livestock byproducts. Estimates in US dollars (US\$) were based on reported yield losses in untreated animals. These estimates reflect the major effects on cattle productivity of six parasites, or parasite group. The potential economic impact (US\$ millions) was: gastrointestinal nematodes US\$ 445.10; coccidia (*Elmeria* spp.) US\$ 23.78; liver fluke (*Fasciola hepatica*) US\$ 130.91; cattle tick (*Rhipicephalus microplus*) US\$ 573.61; horn fly (*Haematobia irritans*) US\$ 231.67; and stable fly (*Stomoxys calcitrans*) US\$ 6.79. Overall, the yearly economic loss due to the six major parasites of cattle in Mexico was estimated to be US\$ 1.41 billion. Considering that the national cattle herd registered in 2013 included 32.40 million head, the estimated yearly loss per head was US\$ 43.57. The limitations of some of the baseline studies used to develop these estimates, particularly when extrapolated from local situations to a national scale, are acknowledged. However, the general picture obtained from the present effort demonstrates the magnitude and importance of cattle parasitism in Mexico and the challenges to maximize profitability by the livestock industry without adapting sustainable and integrated parasite control strategies.

KEY WORDS: Potential economic impact, Bovines, Endoparasites, Ectoparasites, México.

RESUMEN

Las pérdidas económicas causadas por parásitos del bovino en México se calcularon anualmente. Los principales factores considerados para esta evaluación incluyeron el número total de animales a riesgo, los posibles efectos dañinos del parasitismo sobre la producción de leche o ganancia de peso, y los decomisos de subproductos pecuarios. Las pérdidas económicas fueron estimadas en dólares americanos (US\$) y se basaron en las pérdidas del rendimiento productivo de animales no tratados. Estas estimaciones reflejan los principales efectos de seis parásitos o grupos de parásitos sobre la producción bovina. El impacto económico potencial (millones de dólares) fue: nematodos gastrointestinales US\$ 445.10; coccidias (*Elmeria* spp.) US\$ 23.78; duelas del hígado (*Fasciola hepatica*) US\$ 130.91; garrapatas (*Rhipicephalus microplus*) US\$ 573.61; mosca de los cuernos (*Haematobia irritans*) US\$ 231.67; y mosca de los establos (*Stomoxys calcitrans*) US\$ 6.79. En general, las pérdidas anuales causadas por los seis principales parásitos del bovino en México se estimaron en US\$ 1.41 mil millones. Considerando que la población bovina nacional en 2013 fue de 32.40 millones de bovinos, la pérdida anual estimada fue de US\$ 43.57 por animal. Se hace mención de las limitaciones que tienen el uso de algunas referencias para las estimaciones, particularmente cuando se extrapolan situaciones locales a una escala nacional. Sin embargo, el resultado general obtenido en este estudio demuestra la magnitud e importancia del parasitismo en el ganado de México y los desafíos para maximizar la rentabilidad de la industria ganadera sin recurrir al uso de estrategias de control integrado sustentable de parásitos.

PALABRAS CLAVE: Impacto económico potencial, Bovinos, Endoparásitos, Ectoparásitos, México.

Recibido el 29 de mayo de 2015. Aceptado el 28 de septiembre de 2015.

^a Facultad de Medicina Veterinaria y Zootecnia. Campus de Ciencias Biológicas y Agropecuarias. Universidad Autónoma de Yucatán. Km. 15.5 carretera Mérida-Xmatkuil, 97100 Mérida, Yucatán, México.

^b Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brazil.

^c USDA-ARS Knippling-Bushland U.S. Livestock Insects Research Laboratory, and Veterinary Pest Genomics Center, Kerrville, TX, USA.

^d Champion Farmoquímico Ltda, Anápolis, GO, Brazil.

^e SENASICA-SAGARPA, México.

^f Facultad de Medicina Veterinaria y Zootecnia. Universidad Veracruzana, Veracruz, México.

^g Centro Nacional de Investigaciones Disciplinarias en Parasitología Veterinaria. INIFAP, México.

^h Laboratorios Virbac México S.A. de C.V.

* Autor de correspondencia: rvivas@uady.mx.

INTRODUCTION

Animal agriculture is an important sector of the economy in Mexico, which ranks 7th globally in terms of the number of cattle produced for commercial purposes⁽¹⁾. In 2010, bovines comprised the second highest livestock species raised in the country after poultry⁽²⁾. Despite increases in production forecasts due to improved genetics and feeding among other things^(3,4), parasitic diseases remain an important factor impacting the productivity of cattle in Mexico⁽⁵⁾. Furthermore, parasitic diseases can hinder the international trade of cattle raised in Mexico. For example, cattle intended for export are inspected at US ports of entry and are returned to Mexico if found infested with cattle fever ticks⁽⁶⁾.

The national herd is estimated to comprise 32,402,461 head of cattle (dairy= 2,420,289; beef= 29,992,172)⁽⁷⁾ distributed in almost 2 million square kilometers of land. Veracruz is the state with the largest herd that includes around 4 million head. Cattle graze in almost 2 million square kilometers of the territory, distributed in four pastoral regions according to agro-climatic conditions: a) Arid and semiarid, covering portions of nine states (Baja California, Baja California Sur, Sonora, Chihuahua, Coahuila, Nuevo León, Durango, Zacatecas and San Luis Potosí); b) Temperate, including nine states and the Federal District (Aguascalientes, Guanajuato, Hidalgo, Jalisco, México, Morelos, Puebla, Querétaro and Tlaxcala) and areas surrounding Mexico City; c) Dry tropical, comprising six states (Colima, Guerrero, Michoacán, Nayarit, Sinaloa, and Tamaulipas), and d) Humid tropical, extending across six states (Campeche, Chiapas, Quintana Roo, Veracruz, Tabasco, and Yucatán). Production systems and costs are largely dependent on the climatic conditions of the respective regions⁽⁸⁾.

Occurrence of internal and external parasites in cattle throughout Mexico is favored by the predominance of hot humid and sub-humid climates in tropical and subtropical latitudes. Cattle production provides an important economic activity for the large expanse of land in Mexico, and the United States (US) is the primary export market with an average of 1 million cattle imported by the US from Mexico between 1989 and 2009⁽⁹⁾.

Economic loss due to parasites in some countries where the cattle industry is also an important economic activity has been estimated. Drummond *et al.*⁽¹⁰⁾ calculated that annual losses due to cattle parasites in United States of America approximated US\$ 2,260 million when the national herd comprised 124.7 million head⁽¹¹⁾. This figure represented 10 % of the production value and sales of the cattle industry at that time. Estimates of annual economic loss for specific pests ranged from US\$ 29.7 million for scabies and mange mites up to US\$ 730.3 million for horn flies. Recently, Grisi *et al.*⁽¹²⁾ reassessed the potential economic impact of internal and external cattle parasites in Brazil and estimated it to be at least US\$ 13.96 billion annually.

A preliminary assessment revealed that the economic impact of parasitic diseases on cattle production in Mexico is uncertain^(5,13,14,15) and which recognized the challenges involving the estimation of economic loss on the productivity of cattle due to parasitic diseases related to multifactorial variables influenced by geographic area and climatic conditions, management practices, and herd size and density. This type of effort has been further complicated by the threat of parasiticide resistance in several populations of endo- and ectoparasites of cattle. Thus, the results from the study presented here represent a systematic approach to ascertain economic losses due to cattle parasitism in Mexico.

The present study aimed to calculate economic losses caused by cattle parasites in Mexico, including the main ecto- and endo-parasite species known to have a significant effect on the productivity of bovine populations in Mexico. Consideration was given to specific cases where parasiticide resistance increases the cost of efforts to control cattle parasites.

Methodology used

Endoparasites considered for the economic evaluation included: gastrointestinal nematodes (GIN), coccidia (*Eimeria* spp.), and liver fluke (*Fasciola hepatica*). The cattle tick (*Rhipicephalus microplus*), horn fly (*Haematobia irritans*), and stable fly (*Stomoxys calcitrans*) are the external parasites included in this analysis. Other parasites affecting cattle in Mexico were excluded because

they: a) Were not reported to be of national economic importance, b) Are limited to small regions of the country, (c) Are associated with reports of enzootic instability, d) Are not associated with robust economic and epidemiological data in Mexico, e) Are missing economic and epidemiological information. Parasites falling on those categories include *Dictyocaulus viviparus*, *Neospora caninum*, *Toxoplasma gondii*, *Cryptosporidium parvum*, *Tritrichomonas foetus*, *Dermatobia hominis*, *Tabanidae*, and chewing lice.

The method to estimate the potential economic impact caused by each parasite species or group considered the number of animals at risk, data available on milk production loss, weight gain loss in beef cattle, or official reports on condemnation of livestock byproducts (i.e., livers)⁽¹²⁾. In most cases the economic analysis used data originated from local studies, generated primarily between 2000 and 2015. It must be clarified that a limitation of this approach is that estimates addressing data gaps were extrapolated from selected studies conducted in other countries on parasitic economic damage to cattle; whenever possible by making comparisons between studies conducted in Mexico under similar environmental conditions. Thus, the potential economic impact assessment for cattle parasites estimated here is dependent on a wide variety of climatic factors. Grisi *et al*⁽¹²⁾ validated this methodology to reassess the potential economic impact of cattle parasites in Brazil. Because information on production effects was obtained from control animals, i.e. parasitized animals kept untreated in the respective studies, or the estimated populations at risk of being affected by a particular parasitic disease, the resulting estimates of economic impact represent losses in untreated cattle or the estimated bovine population at risk. Thus, the estimates presented here do not necessarily represent the actual impact of cattle parasitism in Mexico but the potential losses expected in the absence of parasite control measures.

POTENTIAL IMPACT OF GASTROINTESTINAL NEMATODES

Helminthic infections are an important cause of disease and productivity loss in livestock production

worldwide. Moreover, tropical and temperate climatic conditions favor the presence and incidence of GIN, potentiating the effect on animal health and reducing livestock production. This problem is not only caused by clinical parasitism; non-visible effects of the subclinical phase also provoke deficiencies in nutrient use, which can cause a major production loss^(16,17).

Impact of GIN on cattle production has been extensively studied worldwide. The effect of GIN on milk production was studied by Lima and Grisi⁽¹⁸⁾ in the state of Rio de Janeiro, Brazil where milk production of grazing cows medicated with albendazole at parturition was compared with untreated controls; an increase of 51.90 kg of milk per cow was reported in the medicated group during the 90-d study period, which corresponded to 0.58 kg of milk/cow/d. A similar study conducted in Holand⁽¹⁹⁾ reported an increase of 0.44 kg of milk/cow/d. Milk yield after anthelmintic treatment on pastured dairy cattle in the Netherlands was estimated to increase by 1.0 kg/cow/d⁽²⁰⁾. When Kabaka *et al*⁽²¹⁾ evaluated the impact of GIN infection on cattle from two districts of Kenya, they found a daily loss of 1.4 kg/cow/d. Vercruyse *et al*⁽²²⁾ evaluated 11 commercial dairy farms in Belgium to study the effect of eprinomectin treatment at calving on milk production; in this study eprinomectin treatment resulted in an average increase of 0.97 kg in daily milk yield. Considering that the average loss due to GIN in grazing cows is 0.87 kg of milk cow/d compared to animals with medication, the potential losses due to gastrointestinal worms for grazing dairy cattle population at risk in Mexico would be US\$ 189,212,709 (Table 1). In spite of the evident economic benefit that could represent the control of GIN in grazing dairy cows, it is important to remember that most anthelmintic drugs cannot be used in those animals while they are producing milk for human consumption. Thus, other control methods, based on alternative control measures ought to be used to reduce the economic impact.

Under the hot sub-humid conditions of Yucatan, Mexico, there was a superior average weight gain of 32.5 and 15.8 kg when the animals (*Bos indicus*) were treated with netobimin following two strategic programs (on August and October, or on August and

Table 1. Economic losses due to gastrointestinal nematodes among dairy and beef cattle in Mexico in 2013

	Population at risk (head)*	Milk loss (kg/cow/d)	Yearly milk loss** (kg/cow)	Total herd yearly loss*** (liters)	Price to producer (US\$/liter)	Potential loss for dairy cattle (US\$)
Dairy cattle	1'735,408	0.87	265.35	443'961,328.8	0.42	189'212,709
	SIAP-SAGARPA ⁽⁷⁾	0.58			SIAP- SAGARPA ⁽⁷⁾	
		Lima and Grisi ⁽¹⁸⁾				
		0.44				
		Ploeger <i>et al</i> ⁽¹⁹⁾				
		1.0				
		Charlier <i>et al</i> ⁽²⁰⁾				
		1.4				
		Kabaka <i>et al</i> ⁽²¹⁾				
		0.97				
		Verschave <i>et al</i> ⁽²²⁾				
*72% of the total dairy cattle in Mexico. **Considering a 305-d lactation period.***1 kg of milk = 0.971 liters of milk.						
	Population at risk+ (head)	Yearly weight loss (kg/ head)	Affected herd yearly loss (kg)	Price to producer (US\$/kg)	Potential loss (US\$)	Potential loss for calves (US\$)
Beef and dairy calves (<1 yr old)	7'290,553	43.6/2=21.8 (only six month a year were considered)	158'934,070	1.61		255'883,853
	SIAP-SAGARPA ⁽⁷⁾	49.6 and 24.0/yr		SIAP- SAGARPA ⁽⁷⁾		
		Domínguez-Alpizar <i>et al</i> ⁽²⁴⁾				
		130.3/yr				
		Quiroz-Romero <i>et al</i> ⁽²⁵⁾				
		104.0/yr				
		Quiroz-Romero <i>et al</i> ⁽²⁶⁾				
+22.5% of the total cattle production in Mexico ⁽²⁾						
Total loss by gastrointestinal nematodes						445'096,562.6

January). In a period of 240 d, dewormed calves obtained an additional weight gain of 0.136 kg/d (49.6 kg/yr) or 0.07 kg/d (24.0 kg/yr) respectively, compared to non-treated calves⁽²³⁾. The most common genera of GIN identified in this study area were *Haemonchus*, *Cooperia* and *Trichostrongylus*⁽²⁴⁾.

In Chiapas, Mexico (hot and humid tropical region), an increase of 30.7 kg was obtained during a 90-d experiment (0.350 kg/d, 127.7 kg/yr) after deworming calves with ivermectin+abamectin⁽²⁵⁾. Another study performed in Veracruz, Mexico also under hot and humid tropical conditions, the use of ivermectin+abamectin to control GIN in calves resulted in an additional weight gain of 20.0 kg (0.285 kg/d) and 25.03 kg (0.357 kg/yr) during a 70-d experiment (130.3 kg/yr and 104.0 kg/yr, respectively) compared with non-treated calves⁽²⁶⁾. The most common genera of GIN identified in this study

were *Haemonchus*, *Cooperia*, *Trichostrongylus*, *Oesophagostomum* and *Ostertagia*.

Studies have shown that the productivity of calves in tropical parts of Mexico is most affected by GIN during the rainy season^(24,27). Thus, for estimation purposes a rainy period of 6 mo was applied to derive potential losses caused by GIN in calves from Mexico. Considering an average loss of 87.1 kg of weight gain calves/yr without medication, the potential losses in the calf population caused by GIN in 6 mo would be US\$ 255'883,853 (Table 1). Overall, annual losses in cattle due to GIN can potentially reach US\$ 445,096,562.

POTENTIAL IMPACT OF *EIMERIA* SPP

Major weight loss associated with eimeriosis in cattle is likely to occur within the first year of life when animals are most susceptible to *Eimeria* infection⁽²⁸⁾. Cattle eimeriosis reduces food

consumption, body weight, feed efficiency and may cause 24 % mortality. An estimated 77 million young cattle in the United States are infected by coccidia during the first year of their life, 4 million will be treated for eimeriosis, and 80,000 cattle naturally infected with *Eimeria*, could die from the disease⁽²⁹⁾. Annual economic losses due to eimeriosis in the US have been estimated at US\$ 62 million⁽³⁰⁾. Animals housed in large numbers with bad hygienic practices are more likely to contract the disease. Therefore, feedlot and dairy cattle are the most susceptible. Eimeriosis in feedlot cattle is associated with stress caused by shipping, changes in ration as well as weather conditions, and overcrowding⁽³¹⁾. Fitzgerald⁽³²⁾ suggested that as a rule of thumb expenses associated with eimeriosis should be considered to be US\$ 2.60 per calf/yr. Dairy farmers in Estonia were estimated to incur annual losses of generated income ranging from 8 to 9 % and assigned a loss of US\$ 6.23 per calf/yr due to eimeriosis⁽³³⁾.

Under hot subhumid tropical conditions, Rodríguez-Vivas *et al*⁽³⁴⁾ studied the factors associated with bovine eimeriosis in calves and found *Eimeria* oocysts in 87.8 % of the samples. The most common species identified were *E. bovis*, *E. zuernii*, *E. ellipsoidalis* and *E. auburnensis*. These authors⁽¹⁴⁾ mentioned that these species are the most frequently found in cattle, and are reported among the most pathogenic⁽³⁵⁾. A further study by Rodríguez-Vivas *et al*⁽¹³⁾ investigated the prevalence of gastrointestinal parasites in cattle of different ages in the south of Mexico. Prevalence of *Eimeria* spp represented 71.57 % of the fecal samples tested. Quezada *et al*⁽³⁶⁾ estimated the prevalence of *Eimeria* spp in post-weaner calves in a dry region of Mexico and found 62 % of the animals to be positive. INEGI⁽²⁾ reported that in Mexico 22.5 % of the cattle population are bovines <12 mo of age. According to the total population of cattle in Mexico (32'402,461) reported by SIAP-SAGARPA⁽⁷⁾, and considering that 22.5 % of the national bovine herd is <12 mo of age, it was estimated that 7'290,553 calves are at risk of being affected by eimeriosis. Based on an average of 73.8 % prevalence of *Eimeria* spp^(13,34,36) among young cattle from different regions of Mexico, it was calculated that 5,380,428 young animals have been

affected, each costing US\$ 4.42/animal⁽³³⁾ to treat eimeriosis⁽³³⁾. Therefore, the estimated annual losses due to eimeriosis in Mexican cattle total US\$ 23'781,491.

POTENTIAL IMPACT OF *FASCIOLA HEPATICA*

Liver fluke transmission depends on the presence of the snail intermediate host. In Mexico *Lymnaea humilis*, *L. cubensis* and *L. bulimoides* have been identified as intermediate host of *F. hepatica*⁽³⁷⁾. Therefore, the distribution of the parasite is limited to those geographic areas where the appropriate snail species are present⁽³⁸⁾. Liver flukes are enzootic in 29 of the 32 states of Mexico, especially where high annual rainfall and large areas of poorly drained pasture provide suitable habitats for the lymnaeid snails. Therefore, the economic impact of bovine fasciolosis caused by *F. hepatica* is relevant in areas where the parasite is enzootic or in non-enzootic areas where infected cattle are introduced⁽³⁹⁾.

According to the four pastoral regions where cattle are raised in Mexico, the average (range) rates of liver condemnation due to fasciolosis from all cattle slaughtered are: Arid and semiarid region 2.89 %, (0.2 to 5.3 %) ^(39,40); Temperate region 11.63 % (1.8 to 30.0 %) ^(39,40); Tropical dry 6.9 % (2.0 to 11.63 %) ⁽³⁹⁾; Tropical humid (Yucatan and Quintana Roo States are considered free of *F. hepatica*) 12.5 % (0.2 to 35.8 %) ^(39,41). The overall estimation of annual losses suffered by the cattle industry of Mexico due to liver condemnation as a result of fasciolosis caused by *F. hepatica* is US\$ 4,239,667 (Table 2).

Simpson and Courtney⁽⁴²⁾ conducted a survey in Florida, US, that showed the specific benefits of liver flukes control. Animals treated with the appropriate antiparasitic drug were 8.2 to 9.9 kg heavier when culled cows were considered, and 13.6 to 20.4 kg heavier when calves were treated at weaning. Another study conducted⁽⁴³⁾ at a commercial cow-calf operation in Louisiana, showed that US calves from cows receiving treatments for both flukes and nematodes had an average weight gain advantage of 8.9 kg in 205-d adjusted weaning weights compared to that of calves from cows receiving treatment for nematodes only. Because all treatment groups were maintained on the same conditions, the authors

inferred that administration of flukicides to cows had a positive effect on productivity by increasing milk production, which resulted in heavier calves.

Recently, Charlier *et al*⁽⁴⁴⁾ conducted a trial on 11 dairy herds exposed to *F. hepatica* to assess the effect of closantel treatment at dry-off (or 80-42 d before calving in first-calving heifers) on milk production parameters. The authors found that closantel treatment resulted in a noticeable increase of 303 kg in a 305-d milk production (0.99 kg of milk/d/animal). Based on this result it was estimated that the annual loss in milk production due to infection with *F. hepatica* is US\$ 83'420,131 (Table 2). The estimated loss associated with the effects of *F. hepatica* infection in beef and dairy cattle that livestock producers in Mexico suffer totals US\$ 130'906,653.

POTENTIAL IMPACT OF CATTLE TICKS

Ticks and the pathogens they transmit to domestic animal species, humans, and wildlife are widely distributed throughout the world, particularly in tropical and subtropical regions. It has been estimated that 80 % of the world's cattle population is exposed to tick infestation⁽⁴⁵⁾. Ticks are responsible for a variety of losses, caused by the direct effect of attachment, injection of toxins, or through the morbidity and mortality associated with tick-borne pathogens⁽⁴⁶⁾, and secondary problems as the enhancement of transmission of *Dermatophilosis*, myiasis⁽⁴⁷⁾, or udder damage by tick species belonging to the genus *Amblyomma*⁽⁴⁸⁾. The estimated annual global costs associated with ticks and tick-transmitted pathogens range between US\$ 13.9 billion and US\$ 18.7 billion⁽⁴⁹⁾. Recently, Grisi *et al*⁽¹²⁾ estimated the annual costs associated

Table 2. Economic losses due to liver confiscation produced by *Fasciola hepatica* to cattle production in Mexico in 2013

Regions	Population at risk*	Slaughtered animals	% of liver confiscation	Total liver confiscated	Price to producer (US\$/liver)	Potential loss (US\$)	
Arid and semiarid	7'638,751 (23.57%)	2'073,309	2.89 Ibarra <i>et al</i> ⁽³⁹⁾	59,919	5.64	337,943	
Temperate	6'600,403 (20.37%)	1'791,825	11.63 Ibarra <i>et al</i> ⁽³⁹⁾	208,389	5.64	1'175,314	
Tropical dry	8'750,368 (27.00%)	2'375,026	6.9 Ibarra <i>et al</i> ⁽³⁹⁾	163,877	5.64	924,266	
Tropical humid	9'412,939 (29.06%)	2'556,231	12.5 Ibarra <i>et al</i> ⁽³⁹⁾	319,529	5.64	1'802,144	
Liver confiscation by fasciolosis						4,239,667	
Animal group	Animal population at risk* (head)	Animal population with fasciolosis** (head)	Yearly weight loss (kg/head)		Total yearly weight loss (kg)	Price to producer (US\$/kg)	Potential loss (US\$)
Cow	15'960,309	1'353,434	9.05 Simpson and Courtney ⁽⁴²⁾		12'248,577	1.61 SIAP-SAGARPA ⁽⁷⁾	19'720,208
Calves	8'085,510	685,651	12.95 Simpson and Courtney ⁽⁴²⁾		8'879,180	1.61 SIAP-SAGARPA ⁽⁷⁾	14'295,479
Feedlot	7'862,068	666,703	17.0 Simpson and Courtney ⁽⁴²⁾ 8.9 Malone <i>et al</i> ⁽⁴³⁾ 8.6 Hicks <i>et al</i> ⁽⁹⁴⁾		5'733,645	SIAP-SAGARPA ⁽⁷⁾	9'231,168
Weight loss						43,246,855	
Dairy cattle	Population at risk* (head)	Animal population with fasciolosis**	Milk loss (kg/cow/d)	Yearly milk loss** (kg/cow)	Total herd yearly loss (liters)	Price to producer (US\$/liter)	Potential loss for dairy cattle (US\$)
	SIAP-SAGARPA ⁽⁷⁾	8.48% Ibarra <i>et al</i> ⁽³⁹⁾	0.99 Charlier <i>et al</i> ⁽²⁰⁾			0.42 SIAP-SAGARPA ⁽⁷⁾	
	2'370,289	663,680	657,043 Milk loss	302	198'619,360	83'420,131	83'420,131 83'420,131 130'906,653
Total fasciolosis loss							130'906,653

*Animals from three states considered free of *F. hepatica* were removed from the animal population at risk. **Only 8.48% of animal at risk were considered suffering fasciolosis⁽³⁹⁾.

with the cattle tick *Rhipicephalus (Boophilus) microplus* (*R. microplus*) in the Brazilian cattle industry, to be up to US\$ 3.24 billion. The national cattle fever tick control campaign in Mexico⁽⁵⁰⁾ reported that 65 % of the national territory is infested with *R. microplus*⁽¹⁵⁾. Based on this information, the estimated number of cattle at risk of *R. microplus* infestation is 24'973,983 cattle.

Losses on milk production caused by the cattle ticks *R. microplus*, were recently estimated by Rodrigues and Leite⁽⁵¹⁾ for the state of Minas Gerais, Brazil. These authors estimated that ticks were responsible for a reduction of 90.24 L of milk per cow per lactation period. Based on this information, the estimated loss to the dairy herd of Mexico is US\$ 68'878,694 (Table 3).

Estimates of the losses inflicted by cattle ticks on beef cattle were based on Jonsson⁽⁵²⁾, who reported daily losses of 1.18 and 1.37 g per tick per animal for *Bos indicus* x *Bos taurus* cattle and *B. taurus*, respectively. An average of 70.3 adult ticks per cattle were reported during studies on population dynamics of *R. microplus* infesting *B. taurus* x *B. indicus* cattle in Yucatan⁽⁵⁾. Alonso-Diaz et al⁽⁵³⁾ found an average of 14.0 and 35.4 *R. microplus* adult ticks per animal in two cattle crossbreed genotypes in Veracruz, Mexico. An average of 14.5 adult ticks per criollo cattle⁽⁵⁴⁾ were found. Meanwhile, others⁽⁵⁵⁾ found that *B. taurus*

cattle had an average of 94 ticks/animal, while Gomes et al⁽⁵⁶⁾ reported 3.3 ticks/per animal for *B. indicus* cattle. Considering that the beef cattle herd is composed of 16.80 % of *B. indicus*, 27.98 % of *B. taurus*, and 52.22 % of *B. indicus* x *B. taurus* crosses⁽²⁾, and the potential infestation rate for this bovine population, the estimated economic loss caused by this tick on beef cattle is US\$ 504'729,382. Therefore, the total economic loss attributable to *R. microplus* in the Mexican cattle herd could approach US\$ 573'608,076 (Table 3).

R. microplus is the vector of the hemoprotezoans *Babesia bovis* and *B. bigemina*, which cause bovine babesiosis, and the intraerythrocytic rickettsia *Anaplasma marginale* that causes anaplasmosis^(57,58). Babesiosis and anaplasmosis are responsible for important economic losses in the cattle industry of tropical and subtropical areas of the world. Annual losses (million) of US\$ 500, 16.9, 5.1, 5.4, 6.8, 21.6, 19.4, 57.2, 3.1, and 0.6 caused by babesiosis and anaplasmosis have been reported for Brazil, Australia, Kenya, Zimbabwe, Tanzania, South Africa, China, India, Indonesia, and Philippines, respectively^(59,60). The economic impact is a consequence of direct losses like mortality, abortions, reduction in meat and milk yield, and indirect effects associated with the application of disease and tick control measures. In Argentina, 62 % and 38 % of the financial loss in dairy cattle affected by

Table 3. Economic losses due to the cattle tick *Rhipicephalus microplus*, related to milk and beef production in Mexico in 2013

	Population at risk*	Lactation loss (liters)	Yearly milk loss (liters)	Price to producer (US\$/liter)	Potential loss (US\$)
Dairy cattle	1'734,735	90.24 Rodrigues and Leite ⁽⁵¹⁾	156'542,486	0.44 SIAP-SAGARPA ⁽⁷⁾	68'878,694
Beef cattle**	Population at risk 23'239,248	Daily weight loss (g/tick/head)	Yearly weight loss (kg)	Price to producer (US\$/kg)	Potential loss (US\$)
<i>B. taurus</i>	3'904,184	1.37 Jonsson ⁽⁵²⁾	183'514,997	1.61 SIAP-SAGARPA ⁽⁷⁾	295'459,145
<i>B. indicus</i>	6'502,345	1.18 Jonsson ⁽⁵²⁾	111'742,243	1.61 SIAP-SAGARPA ⁽⁷⁾	179'905,011
<i>B. indicus</i> x <i>B. taurus</i>	12'832,719	1.18 Jonsson ⁽⁵²⁾	18'239,271	1.61 SIAP-SAGARPA ⁽⁷⁾	29'365,226
					504'729,382
				Total potential loss	573'608,076

*Cattle at risk of tick infestation^(7,50). **16.80 % of *Bos indicus*, 27.98 % of *Bos taurus*, and 52.22 % of *Bos indicus* x *Bos taurus* cattle in Mexico. ***Considering the following mean daily tick burdens: *B. taurus*-94 ticks⁽⁵⁵⁾, *B. indicus* x *B. taurus*-39.9 ticks^(5,53), and *B. indicus*-3.3 ticks⁽⁵⁶⁾.

babesiosis were due to direct and indirect effects, respectively⁽⁶¹⁾. Babesiosis and anaplasmosis are prevalent wherever *R. microplus* and *R. annulatus* occur in Mexico⁽⁶²⁻⁶⁶⁾, and 75 % of the national cattle herd is considered to be at risk of becoming infected with *B. bovis* and *B. bigemina*⁽⁶⁷⁾. However; we were unable to estimate the economic impact of babesiosis and anaplasmosis on the cattle industry in Mexico because key epidemiological data, such as description of specific enzootic instability regions, mortality and incidence of babesiosis and anaplasmosis outbreaks, and number of abortions due to clinical cases and seasonal occurrence of outbreaks were not found during the search of the scientific literature.

POTENTIAL IMPACT OF *HAEMATOBIA IRRITANS*

The horn fly, *Haematobia irritans*, is one of the most economically important pests of cattle worldwide^(12,65). It is an obligate hematophagous ectoparasite that feeds almost exclusively on cattle. Because of *H. irritans* feeding behavior and infestation levels tend to be high, cattle expend a great degree of energy in defensive behavior, and reduce the grazing time, feeding efficiency and milk production in the case of dairy cattle⁽⁶⁷⁾.

Losses due to horn fly infestation, based on the studies in Brazil⁽⁶⁸⁾ and Argentina⁽⁶⁹⁾, are estimated as an average yearly weight loss of 3.25 kg per cow and 28.0 g per day (305 d of lactation, 8.54 kg per cow), respectively. Estimated losses in calves also were based on Bianchin and Alves⁽⁶⁸⁾ in Brazil and Steelman *et al*⁽⁷⁰⁾ in the US, which estimated an average yearly weight loss of 2.0 and 8.1 kg per calf, respectively. Yearly weight losses of steer and heifers were estimated as 12.19 kg per animal⁽⁷¹⁾. Losses in milk production caused by *H. irritans* in dairy farms were estimated as 27.0 kg decrease in milk production per cow per year in the US⁽⁷²⁾.

In general, *Haematobia irritans* in Mexico affects cattle during 7 mo of the year⁽⁷³⁾. Cattle in the state of Veracruz (hot humid tropic) experienced high *H. irritans* infestation (70 to 121 flies/animal) between August and November⁽⁷³⁾. Galindo-Velasco *et al*⁽⁷⁴⁾ reported that cattle in the state of Colima, including sub-humid tropic and sub-tropic environments, had high horn fly infestations (120 to 236 flies per animal) during 6 mo of the year. However, in the state of Tamaulipas high horn fly infestation levels were observed in cattle throughout the year with population peaks higher than 200 flies/animal detected in September, April, May, and

Table 4. Economic losses due to the horn fly *Haematobia irritans*, related to beef cattle production in Mexico in 2013

	Live animals at risk* (head)	Yearly weight loss per head (kg)	Total yearly Weight loss ** (kg)	Price to producer (US\$/kg)	Potential loss (US\$)
Steers/heifers	7'776,590	12.19 Bianchin <i>et al</i> ⁽⁷¹⁾	57'667,951	1.61 SIAP-SAGARPA ⁽⁷⁾	92'845,401
Calves	7'290,553	5.05 2.0 Bianchin and Alves ⁽⁶⁸⁾ 8.1 Steelman <i>et al</i> ⁽⁷⁰⁾	22'397,186	1.61 SIAP-SAGARPA ⁽⁷⁾	36'059,469
Cows	17'335,316	5.89 3.25 Bianchin and Alves ⁽⁶⁸⁾ 8.54 Guglielmo <i>et al</i> ⁽⁶⁹⁾	62'113,881	1.61 SIAP-SAGARPA ⁽⁷⁾	100'003,348
Dairy cows	1'734,735	Milk loss (liter/cow/yr) 27.0 Kunz <i>et al</i> ⁽⁷²⁾	Total yearly loss (liters)** 4'102,995	Price to producer (US\$/liter) 0.42 SIAP-SAGARPA ⁽⁷⁾	Potential loss (US\$) 2'757,212
	32'402,461			Total potential loss	231'665,430

* From the total cattle production in Mexico⁽⁷⁾, 22.5 % was calves, 24.0 % steers/heifers, and 53.5 % adults (>3 yr)⁽²⁾. **7.3 mo per year were considered for *H. irritans* affect on cattle^(73,74,75).

June⁽⁷⁵⁾. Considering the beef cattle population at risk, losses due to horn fly parasitism in Mexico are estimated to be US\$ 231'665,430 (Table 4). No estimates of horn-fly losses on dairy cattle or beef cattle pregnancy rates are provided here. These issues need further investigation.

POTENTIAL IMPACT OF *STOMOXYS CALCITRANS*

The stable fly, *Stomoxys calcitrans*, is another economically important pest of livestock worldwide. The stable fly problem can be exacerbated in areas where high organic content, like hay waste residues, are present at or nearby pastures where cattle graze⁽⁷⁶⁻⁷⁸⁾. Heavy stable fly infestations can result in a reduction of weight gain and feed efficiency of confined and grazing beef cattle⁽⁷⁹⁾. Taylor *et al*⁽⁷⁶⁾ and Grisi *et al*⁽¹²⁾ reported that the stable fly has an economic impact on US and Brazilian cattle production of US\$ 2,211.0 and 335.5 million per year, respectively. Kunz *et al*⁽⁷²⁾ estimated a potential loss due to stable flies in dairy and beef cattle of 0.1 kg per steer per day in feedlots and a 27.0 kg decrease in milk production per cow per year. Taylor *et al*⁽⁷⁶⁾ estimated an annual per animal production loss of 139.0 kg milk for dairy cows, and 6.0, 26.0, and 9.0 kg body weight for preweaning calves, pastured stockers, and feeder cattle, respectively.

Cruz-Vázquez *et al*⁽⁸⁰⁾ studied the annual variation of *S. calcitrans* adult stage infestation in Aguascalientes, Mexico and they found seasonal behavior of infestation with higher infestations between mid-summer and mid-fall period (2.5 mo) with abundance ranging from 4 to 40 flies per animal. Based on this information, economic losses

due to the stable fly in Mexico were estimated to be US\$ 6'786,792 (Table 5).

POTENTIAL ECONOMIC IMPACT OF ALL STUDIED PARASITES

Annual potential economic loss due to the six major endo- and ectoparasites of cattle in Mexico considered here was estimated to be US\$ 1,411'845,004. Considering that the national cattle herd registered in 2013 included 32'402,461 head⁽⁷⁾, the estimated yearly loss per head caused by the selected cattle parasites was US\$ 43.57. By comparison, the annual loss per head caused by a similar group of cattle parasites in Brazil, except the cattle grub (*Dermatobia hominis*) and New World screwworm fly (*Cochliomyia hominivorax*), was estimated to be US\$ 65.49⁽¹²⁾.

The limitations of some of the baseline studies used to develop these estimates, particularly when extrapolated from local situations to a national scale⁽¹²⁾, are acknowledged. However, the general picture obtained from the present effort demonstrates the magnitude and importance of cattle parasitism in Mexico and the unfeasibility of a profitable livestock industry without adapting sustainable and integrated parasite control strategies. This approach is required given the favorable environmental conditions for livestock parasites in grazing areas where cattle are raised in Mexico.

PARASITICIDE RESISTANCE

The main strategy for cattle parasite control in Mexico is based on the use of chemicals such as

Table 5. Economic losses due to the stable fly *Stomoxys calcitrans* among dairy and feedlot cattle in Mexico in 2013

	Population at risk (head)	Milk loss (L/cow/yr)	Total yearly loss (L)	Price to producer (US\$/L)	Potential loss (US\$)
Dairy cattle SIAP-SAGARPA ⁽⁷⁾	1'734,735	27.0 Kunz <i>et al</i> ⁽⁷²⁾	1'561,261	0.42 SIAP-SAGARPA ⁽⁷⁾	655,729
	Population at risk (head)	Loss at feedlots (g/head/d)	Total loss at feedlots (kg)	Price to producer (US\$/kg)	Potential loss (US\$)
Feedlot cattle SIAP-SAGARPA ⁽⁷⁾	7'776,590	100.0 Kunz <i>et al</i> ⁽⁷²⁾	3'784,607	1.62 SIAP-SAGARPA ⁽⁷⁾	6'131,063
				Total potential loss	6'786,792

* 2.5 mo per year were considered to affect *S. calcitrans* to cattle⁽⁸⁰⁾.

anthelmintics, coccidiostats, flukicides, acaricides, insecticides and endectocides⁽⁸¹⁾. The indiscriminate use of these chemicals has resulted in the development of resistance in populations of internal and external parasites of cattle, which impacts the livestock industry in Mexico⁽⁸²⁻⁸⁷⁾.

Parasiticide resistance is a multifaceted problem that not only impacts the ability of producers to mitigate the impact of parasites on farm animal production, but it also poses risks to the environment and public health^(88,89). Resistance to conventional acaricides and macrocyclic lactones, including organophosphates (OPs), pyrethroids (SPs), amitraz, fipronil, and ivermectin among cattle fever tick populations has been identified as one of the most common problems in cattle from Mexico^(83,87). Resistance to OPs first developed in the 1980s in Mexico, and resistance to SPs emerged in the 1990s. Amitraz was introduced along with SPs to control OPs-resistant ticks in 1986. Amitraz was not widely used initially due to its higher cost, but its application became more prevalent and intensive after SPs resistance was discovered in 1993⁽⁸⁵⁾. The first case of amitraz resistance in *R. microplus* from Mexico was confirmed in 2001 in the state of Tabasco⁽⁸⁶⁾. The first reports of *R. microplus* populations resistant to ivermectin and fipronil in Mexico were made in 2010 and 2013, respectively^(87,88). Rodríguez-Vivas *et al.*⁽⁸⁹⁾ studied 217 field populations of *R. microplus* and determined the prevalence of farms with resistance to SPs, OPs and amitraz in the southern part of the country. They found that resistance to SPs like cypermethrin, flumethrin, and deltamethrin was one of the most serious problems faced by livestock producers in the Mexican tropics where 66 to 96 % of the farms had cattle infested with *R. microplus* resistant to SPs. The presence of *R. microplus* populations resistant to ivermectin has been documented in the states of Yucatan and Veracruz where resistance prevalence ranges from 41 to 100 %, with most of the resistance level categorized as low. Resistance to multiple classes of antiparasitic drugs, including OPs, SPs, amitraz and ivermectin, appears to be an emerging problem among *R. microplus* populations in Mexico⁽⁸³⁾.

Resistance to ivermectin in GIN has been identified in cattle from Mexico. Initial reports of

ivermectin-resistant GIN in Mexico were obtained for infected cattle in the states of Campeche⁽⁹⁰⁾, Yucatán⁽⁸⁴⁾, and Veracruz⁽⁹¹⁾. Studies performed in Mexico suggest a high frequency of cattle herds with IVM-resistant GIN (>71 % of the surveyed herds), and they involved species in the genera *Haemonchus* and *Cooperia*.

Horn fly populations resistant to insecticides have been documented in Mexico. Insecticide resistance was reported for horn fly populations in the states of Tamaulipas, Veracruz, San Luis Potosi, Sinaloa, Aguascalientes, and Nuevo León^(92,93). The development of resistance to permethrin in horn flies from Veracruz and Nuevo León apparently was associated with strong selection pressure and the lack of integrated parasite management practices because in the farms SPs were used commonly to control *R. microplus* too^(92,93).

Antiparasitic resistance in endo- and ectoparasite populations affecting the health and productivity of cattle in Mexico increases the cost of parasite control and decreases the profitability of livestock producers. The use of commercial antiparasitic drugs to control endo- and ectoparasites in cattle is expected to continue increasing in Mexico due primarily to: availability and ease of use for endectocidal products, increasingly accessible prices especially for generic versions of macrocyclic lactones, and a dearth of products containing active ingredients with new modes of action⁽⁸³⁾.

Studies assessing the actual economic impact of parasiticide resistance on the profitability of cattle producers in Mexico are needed. The establishment of integrated parasite control programs adapted to each of the four main agroecological regions could reduce the impact of endo- and ectoparasites on the health and productivity of the national cattle herd⁽⁸³⁾. An adaptive and sustainable approach will help mitigate the burden of diseases caused by the six major endo- and ectoparasites considered here, which was estimated to cost the cattle industry of Mexico US\$ 1.41 billion annually and a yearly loss per head of US\$ 43.57.

ACKNOWLEDGMENTS

Thanks to Champion Farmoquímico Ltda, Anápolis, GO, Brazil, and Laboratorio Virbac, Mexico, for facilitating organizational meeting by our group to discuss the economic impact of cattle parasites, which resulted in initiative to produce this publication. We would like to dedicate this paper to the memory of Prof. Laerte Grisi, a brilliant scientist and a kind and generous colleague.

LITERATURA CITADA

- Organización de las Naciones Unidas para la Agricultura y la Alimentación (FAO). Ganadería bovina en América Latina: Escenarios 2008-2009 y tendencias del sector. Santiago de Chile. 2010.
- INEGI (Instituto Nacional de Estadística y Geografía) Censo Agropecuario, www.inegi.org.mx. 2007.
- Larios-Sarabia N, Ramírez-Valverde R, Núñez-Domínguez R, García-Muñiz JG, Ruíz-Flores A. Technical, social and economic characterization of companies in the registered Jersey cattle herd in México. *Agric Soc Desarrollo* 2011;8:229-247.
- Callejas-Juárez N, Aranda-Gutiérrez H, Rebollar-Rebollar S, de la Fuente-Martínez ML. Situación económica de la producción de bovinos de carne en el estado de Chihuahua, México. *Agronomía Mesoamericana* 2014;25:133-139.
- Rodríguez-Vivas RI, Quiñones AF, Ramírez CG. Epidemiología, prevención y control de la babesiosis bovina. En: *Enfermedades de importancia económica en producción animal*. Rodríguez-Vivas RI editor. México DF: McGraw-Hill-UADY; 2005:205-225.
- Pérez de León AA, Rodríguez-Vivas RI, Guerrero FD, García-Vázquez Z, Temeyer KB, Domínguez-García DI, *et al*. Acaricide resistance in *Rhipicephalus (Boophilus) microplus*: impact on agro-biosecurity and cattle trade between Mexico and the United States of America. *Proc Int Symp Pest Resistance Arthropods* 2013;3:18-35.
- SIAP-SAGARPA. Servicio de Información Agroalimentaria y Pesquera. Inventario ganadero. 2013.
- García, E. Modificaciones al sistema de clasificación climática de Köppen. Instituto de Geografía de la Universidad Nacional Autónoma de México. Quinta ed. México, DF. 2004.
- Peel DS, Johnson RJ, Mathews KH. Cow-calf beef production in Mexico. A report from the economic research service. USDA. www.ers.usda.gov. 2010.
- Drummond RO, Lambert G, Smally HE Jr, Terrill CE. Estimated losses of livestock to pests. In: Pimentel D. *Handbook of pest management in agriculture*. Boca Raton, USA: CRC Press 1981;1:111-127.
- National Agriculture Statistics Service - NASS, United States Department of Agriculture – USDA, Cattle inventory: executive briefing 2011. http://www.nass.usda.gov/Newsroom/Executive_Briefings/2011/07_22_2011.pdf. Accessed Nov 03, 2015.
- Grisi L, Cerqueira Leite R, de Souza Martins JR, Medeiros de Barros AT, Andreotti R, Duarte Cançado PH, *et al*. Reassessment of the potential economic impact of cattle parasites in Brazil. *Braz J Vet Parasitol* 2014;3(2):150-156.
- Rodríguez-Vivas RI, Cob-Galera LA, Domínguez-Alpizar JA. Frecuencia de parásitos gastrointestinales en animales domésticos diagnosticados en Yucatán, México. *Rev Biomed* 2001;12:19-25.
- Rodríguez-Vivas RI, Ojeda-Chi MM, Pérez-Cogollo LC, Rosado-Aguilar JA, Ramírez-Cruz GT, Guemes-Ceballos AA. Epidemiología, diagnóstico y control de la coccidiosis bovina. Capítulo 4. *Epidemiología de enfermedades parasitarias en animales domésticos*. Quiroz RH. *et al*, editores. México, DF: AMPAVE; 2011:52-66.
- Bautista-Garfias CR, Martínez-Ibañez F. Experiences on the control of cattle tick *Rhipicephalus (Boophilus) microplus* in Mexico. In: *Ticks: Disease, management and control*. Moges Woldemeskel editor. Nova Science Publishers, Inc. 2012:205-216.
- Vercruysse J, Claerebout E. Treatment vs non treatment of helminth infections in cattle: defining the threshold. *Vet Parasitol* 2001;98:195-214.
- Molento MB, Fortes FS, Pondelek DA, Borges F de A, Chagas AC, Torres-Acosta JF, Geldhof P. Challenges of nematode control in ruminants: focus on Latin America. *Vet Parasitol* 2011;180(1-2):126-32.
- Lima MM, Grisi L. Verminose subclínica em vacas em lactação no Estado do Rio de Janeiro. *Horiz Vet* 1984;19(4):37-40.
- Ploeger HW, Schoenmaker GJW, Kloosterman A, Borgsteede FHM. Effect of anthelmintic treatment of dairy cattle on milk production related to some parameters estimating nematode infection. *Vet Parasitol* 1989;34(3):239-253.
- Charlier J, Höglund J, Von Samson-Himmelstjerna G, Dorny P, Vercruysse J. Gastrointestinal nematode infections in adult dairy cattle: impact on production, diagnosis and control. *Vet Parasitol* 2009;164(1):70-79.
- Kabaka WM, Gitau GK, Kitala PM, Maingi N, VanLeeuwen JA. The prevalence of gastrointestinal nematode infection and their impact on cattle in Nakuru and Mukurweini districts of Kenya. *Ethiop Vet J* 2013;17(1):95-104.
- Verschave SH, Vercruysse J, Forbes A, Opsomer G, Hostens M, Duchateau L, Charlier J. Non-invasive indicators associated with the milk yield response after anthelmintic treatment at calving in dairy cows. *BMC Vet Res* 2014;10:264.
- Domínguez-Alpizar JL, Rodríguez-Vivas RI, Cob GLA. Evaluación de dos programas estratégicos de control de nematodos gastrointestinales en becerros de la zona centro del estado de Yucatán, México. *Rev Biomed* 1996;7:27-34.
- Domínguez-Alpizar J, Rodríguez-Vivas VRI, Honhold N. Epizootiología de los parásitos gastrointestinales en bovinos del Estado de Yucatán. *Vet Méx* 1993;24(3):189-193.
- Quiroz-Romero H, Chavarría-Martínez B, Hernández-Suárez A, Ochoa-Galván P, Cruz- Pérez J, Cruz-Mendoza I. Efecto de una nueva formulación de ivermectina+abamectina de larga duración contra nematodos gastrointestinales y la diferencia en ganancia de peso en bovinos. *Vet Méx* 2009;40(2):157-165.
- Quiroz-Romero H, Chavarría-Martínez B, Carrillo J, Pérez M, Ochoa-Galván P. Efecto prolongado de ivermectina+abamectina contra nematodos y la diferencia de peso en bovinos en Veracruz, México. *Rev Ibero-Latinoam Parasitol* 2009;68(2):180-184.

27. Encalada-Mena LA, Corbala-Bermejo JA, Vargas-Magaña JJ, García-Ramírez MJ, Uicab-Brito L, del Río-Rodríguez J. Prevalencia de nematodos gastroentéricos de becerros en sistemas de doble propósito del municipio de Escárcega, Campeche, México. *Agrociencia* 2009;43(6):569-576.
28. Bangoura B, Dauschies A. Parasitological and clinical parameters of experimental *Eimeria zuernii* infection in calves and influence on weight gain and haemogram. *Parasitol Res* 2007;100:1331-1340.
29. Fitzgerald PR. The significance of bovine coccidiosis as a disease in the United States. *Bovine Pract* 1975;10:28.
30. Williams L. Dealing with the frustration of the war on coccidiosis. *DVM* 1984;15:38.
31. Ernst JV, Benz GW. Intestinal coccidiosis in cattle. The veterinary clinics of North America/parasites: epidemiology and control. Philadelphia, PA: W.B. Saunders Company; 1986.
32. Fitzgerald P. The economic impact of coccidiosis in domestic animals. *Adv Vet Sci Comp Med* 1980;24:121-143.
33. Lassena B, Østergaard S. Estimation of the economic effects of *Eimeria* infections in Estonian dairy herds using a stochastic model. *Prev Vet Med* 2012;106:258-265.
34. Rodríguez-Vivas RI, Domínguez-Alpizar JL, Torres-Acosta JF. Epidemiological factors associated to bovine coccidiosis in calves (*Bos indicus*) in a sub-humid tropical climate. *Rev Biomed* 1996;7:211-218.
35. Hasbullah H, Akiba Y, Takano H, Ogimoto K. Seasonal distribution of bovine coccidia in beef herd in the University farm. *Jpn J Vet Sci* 1990;52:1175-1179.
36. Quezada CA, Ramos GJI, Figueroa VC, Rivas CRR, Martínez de la Rosa R, Trillo MV. Prevalencia de coccidia en becerras Holstein en la etapa de desarrollo. *CULCYT* 2013;10(49):54-60.
37. Landeros y Valdez MA, Ibarra VF, Escudero CJL, Milian SF. Determinación de algunos hospederos intermediarios de *Fasciola hepatica*, en la cuenca lechera de Tulancingo Hidalgo. *Tec Pecu Méx* 1981;40:47-51.
38. Kaplan RM. *Fasciola hepatica*: A review of the economic impact in cattle and considerations for control. *Vet Therap* 2001;2(1):40-50.
39. Ibarra VF, Vera MY, Munguía XJ. Epidemiología de la fasciolosis animal y humana. En: *Epidemiología de enfermedades parasitarias en animales domésticos*. Capítulo 9. Quiroz RH, *et al* editores. México, DF: AMPAVE; 2011:137-172.
40. Castellanos HAA, Escutia SI, Quiroz RH. Frecuencia de fasciolosis hepática en bovinos sacrificados en las plantas tipo inspección federal en México de los años 1979 a 1989. *Vet Mex* 1992;23:339-342.
41. Rangel LJ, Martínez DE. Pérdidas económicas por decomiso de hígados y distribución geográfica de la fasciolosis bovina en el estado de Tabasco, México. *Vet Méx* 1994;25:327-331.
42. Simpson JR, Courtney CH. Liver flukes in Florida: Prevalence, economics and management practices on ranches surveyed. Gainesville, University of Florida. Florida Coop Ext Serv. Institute of Food and Agricultural Sciences. 1990.
43. Malone J, Loyacano A, Armstrong D, Archbald L. Bovine fascioliasis: Economic impact and control in gulf coast cattle based on seasonal transmission. *Bovine Pract* 1982;17:126-133.
44. Charlier J, Hostens M, Jacobs J, Van Ranst B, Duchateau L, Vercruysse J. Integrating Fasciolosis control in the dry cow management: The effect of closantel treatment on milk production. *PLoS ONE* 2012;7(8):e43216.
45. FAO. Ticks and tick borne disease control. A practical field manual. Volume I. Tick control. Rome. 1984.
46. Esteve-Gasent MD, Perez de Leon AA, Romero-Salas D, Feria-Arroyo TP, Pation R, Castro-Arellano I, *et al*. Pathogenic landscape of transboundary zoonotic diseases in the Mexico-US border along the Rio Grande. *Front Public Hlth* 2014;17(177):1-23.
47. Reck J, Marks FS, Rodrigues RO, Souza UA, Webster A, Leite RC, Gonzales JC, *et al*. Does *Rhipicephalus microplus* tick infestation increase the risk for myiasis caused by *Cochliomyia hominivorax* in cattle? *Prev Vet Med* 2014;113:59-62.
48. Estrada-Peña A, Salman M. Current limitations in the control and spread of ticks that affect livestock: A review. *Agriculture* 2013;3:221-235.
49. De Castro JJ, James AD, Minjauw B, DiGiulio G, Permin A, Pegram RG, *et al*. Long-term studies on the economic impact of ticks on Sanga cattle in Zambia. *Exp Appl Acarol* 1997;21:3-19.
50. SAGARPA-SENASICA. Situación zoonosaria la campaña para el control de la garrapata *Boophilus* spp en México. 2014.
51. Rodrigues DS, Leite RC. Economic impact of *Rhipicephalus (Boophilus) microplus*: estimate of decreased milk production on a dairy farm. *Arq Bras Med Vet Zootec* 2013;65(5):1570-1572.
52. Jonsson N. The productivity effects of cattle tick (*Boophilus microplus*) infestation on cattle, with particular reference to *Bos indicus* cattle and their crosses. *Vet Parasitol* 2006;137:1-10.
53. Alonso-Díaz MA, López SBJ, Leme de Magalhães-Labarthe AC, Rodríguez-Vivas RI. Infestación natural de *Boophilus microplus* Canestrini, 1887 (Acari: Ixodidae) en dos genotipos de bovinos en el trópico húmedo Mexicano. *Vet Méx* 2007;38(4):503-509.
54. González-Cerón F, Becerril-Pérez CM, Torres-Hernández G, Díaz-Rivera P. Garrapatas que infestan regiones corporales del bovino criollo lechero tropical en Veracruz, México. *Agrociencia* 2009;43:11-19.
55. Smith RD, Evans DE, Martins JR, Ceresér VH, Correa BL, Petracchia C, *et al*. Babesiosis (*Babesia bovis*) stability in unstable environments. *Ann N Y Acad Sci* 2000;916:510-520.
56. Gomes A, Honer MR, Schenk MA, Curvo JB. Populations of the cattle tick (*Boophilus microplus*) on purebred Nellore, Ibage and Nellore X European crossbreds in the Brazilian savanna. *Trop Anim Hlth Prod* 1989;21(1):20-24.
57. Pérez de León AA, Vannier E, Almazán C, Krause P. Tick-borne protozoa. In: Sonenhine DE, Roe RM editors. *Biology of ticks*. 2nd ed, New York, USA: Oxford University Press; 2014(2):147-179.
58. Aubry P, Geale DW. A review of bovine anaplasmosis. *Transbound Emerg Dis* 2011;5(8):1-30.
59. Bock R, Jackson L, de Vos A, Jorgensen W. Babesiosis of cattle. *Parasitology* 2004;129 Suppl:S247-269.
60. Grisi L, Massard CL, Moya Borja GE, Pereira JB. Impacto económico das principais ectoparasitoses em bovinos no Brasil. *Hora Vet* 2002;21(125):8-10.
61. Guglielmo, AA, Aguirre DH, Späth EJA, Gaido AB, Mangold AJ, de Rios LG. Long-term study of incidence and financial loss due to cattle babesiosis in an Argentinian dairy farm. *Vet Parasitol* 1992;12(3-4):307-312.

62. Solorio RJ, Rodríguez VRI, Pérez E, Wagner G. Management factors associated with *Babesia bovis* seroprevalence in cattle from eastern Yucatan, Mexico. *Prev Vet Med* 1999;40:261-269.
63. Rodríguez-Vivas RI, Mata MY, Pérez GE, Wagner G. The effect of management factors on the seroprevalence of *Anaplasma marginale* in *Bos indicus* cattle in the Mexican tropics. *Trop An Hlth Prod* 2004;36:135-143.
64. Pérez de León AA, Teel PD, Auclair AN, Messenger MT, Guerrero F, Schuster G, Miller R. Integrated strategy for sustainable cattle fever tick eradication in U.S.A. is required to mitigate the impact of global change. *Front Physiol* 2012;3:195.
65. Xu Q, Guerrero FD, Palavesam A, Pérez de León AA. Use of electroporation as an option to transform the horn fly, *Haematobia irritans*: a species recalcitrant to microinjection. *Insect Sci* 2015;00:1-9. doi: 10.1111/1744-7917.12207.
66. Foil LD, Hogsette JA. Biology and control of tabanids, stable flies and horn flies. *Revue Scientifique et Technique* 1994;13:1125-1158.
67. Bautista-Garfias CR, Castañeda-Arriola R, Álvarez-Martínez JA, Rojas-Martínez C, Figueroa-Millán JV, Rodríguez-Lozano A. The simultaneous vaccination of bovines with *Lactobacillus casei* and the bivalent vaccine against bovine babesiosis induces a better protection against *Babesia bovis* and *B. bigemina* transmitted by ticks in extreme field conditions. *Vet Méx* 2012;43:189-200.
68. Bianchin I, Alves RGO. The hornfly, *Haematobia irritans*: behaviour and damage caused in Nellore cows and pre-weaning calves. *Pesq Vet Bras* 2002;22(3):109-113.
69. Guglielmo AA, Volpogni MM, Quaino OR, Anziani OS, Mangold AJ. Long-term study of *Haematobia irritans* (Diptera: Muscidae) seasonal distribution in central Argentina with focus on winter fly abundance. *Parasitol* 2001;8(4):369-373.
70. Steelman CD, Brown AHJr, Gbur EE, Tolley G. Interactive response of the horn fly (Diptera: Muscidae) and selected breeds of beef cattle. *J Econ Entomol* 1991;84(4):1275-1282.
71. Bianchin I, Koller WW, Alves RGO, Detmann E. Effects of the horn fly, *Haematobia irritans* (L.) (Diptera: Muscidae) in the weight gain on Nellore cattle. *Cienc Rural* 2004;34(3):885-890.
72. Kunz SE, Murrell KD, Lambert G, James LF, Terrill CE. Estimated losses of livestock to pests. In: Pimentel D. *Handbook of pest management in agriculture*. 2nd ed. Boca Raton, USA: CRC Press; 1991:69-98.
73. Alonso-Díaz MA, Acosta-Rodríguez R, Maldonado-Simán E. Dinámica poblacional de *Haematobia irritans* en bovinos del trópico mexicano. *Rev Cient FCV-LUZ; XVII* 2007;(4):330-334.
74. Galindo-Velasco E, Cruz-Vázquez C, Lezama-Gutiérrez R, Reyes-Velázquez W, Aguilar-Espinoza S, Pescador-Rubio A. Fluctuación poblacional de *Haematobia irritans* (Diptera: Muscidae) en un hato bovino en Tecmán, Colima, México. *Vet Méx* 2008;39(2):181-186.
75. Almazán-García C, Castillo-Salas S, Loredó-Osti J, García-Vázquez Z, Ramírez-Valverde R, Bermúdez-Villanueva L. Dinámica poblacional de *Haematobia irritans* en un hato de bovinos de Soto la Marina, Tamaulipas, México. *Vet Méx* 2001;32(2):149-152.
76. Taylor DB, Moon RD, Mark DR. Economic impact of stable flies (Diptera: Muscidae) on dairy and beef cattle production. *J Med Entomol* 2012;49(1):198-209.
77. Liu SS, Li AY, Lohmeyer KH, Pérez de León AA. Effects of pyriproxyfen and buprofezin on immature development and reproduction in the stable fly. *Med Vet Entomol* 2012;26:379-385.
78. Solórzano JA, Gilles J, Bravo O, Vargas C, Gomez-Bonilla Y, Bingham GV, Taylor DB. Biology and trapping of stable flies (Diptera: Muscidae) developing in pineapple residues (*Ananas comosus*) in Costa Rica. *J Insect Sci* 2015;15(1):145.
79. Campbell JB, Skoda SR, Berkebile DR, Boxler DJ, Thomas GD, Adams DC, Davis R. Effects of stable flies (Diptera: Muscidae) on weight gains of grazing yearling cattle. *J Econ Entomol* 2001;94(3):780-783.
80. Cruz-Vázquez C, Martínez RS, Vitela MI, Ramos PM, Quintero MMT, García VZ. Variación anual de la infestación por *Stomoxys calcitrans* (L.) (Diptera: Muscidae) en tres establos lecheros de Aguascalientes, México. *Téc Pecu Méx* 2000;38(2):135-142.
81. Martínez MI, Cruz M. The use of agricultural and livestock chemical products in the cattle- ranching area of Xico, central Veracruz, Mexico, and their possible environmental impact. *Acta Zool Mex* 2009;25:673-681.
82. Canul-Ku HL, Rodríguez-Vivas RI, Torres-Acosta JFJ, Aguilar-Caballero AJ, Pérez-Cogollo LC, Ojeda-Chi MM. Frequency of cattle herds with ivermectin resistant nematodes in the hot sub-humid tropics of Mexico. *Vet Parasitol* 2012;183:292-298.
83. Rodríguez-Vivas R, Pérez-Cogollo LC, Rosado-Aguilar JA, Ojeda-Chi MM, Trinidad-Martínez I, Miller RJ, et al. *Rhipicephalus microplus* resistant to acaricides and ivermectin in cattle farms of Mexico. *Braz J Vet Parasitol* 2014;23(2):113-122.
84. Alegría-López MA, Rodríguez-Vivas RI, Torres-Acosta JFJ, Ojeda-Chi MM, Rosado-Aguilar JA. Use of ivermectin as endoparasiticide in tropical cattle herds generates resistance in gastrointestinal nematodes and the tick *Rhipicephalus microplus* (Acari: Ixodidae). *J Med Entomol* 2015; 1-8. doi: 10.1093/jme/tju02.
85. Fragoso-Sánchez H, García-Vázquez Z, Tapia-Pérez G, Ortiz-Najera A, Rosario-Cruz R, Rodríguez-Vivas RI. Response of Mexican *Rhipicephalus (Boophilus) microplus* ticks to selection by amitraz and genetic analysis of attained resistance. *J Entomol* 2011;8(3):1812-5670.
86. Soberanes NC, Santamaría MV, Fragoso HS, García VZ. First case reported of amitraz resistance in the cattle tick *Boophilus microplus* in Mexico. *Téc Pecu Méx* 2002;40(1):81-92.
87. Pérez-Cogollo LC, Rodríguez-Vivas RI, Ramírez-Cruz GT, Miller RJ. First report of the cattle tick *Rhipicephalus microplus* resistant to ivermectin in Mexico. *Vet Parasitol* 2010;168(1-2):165-169.
88. Miller RJ, Almazán C, Ortiz-Estrada M, Davey RB, George JE, Pérez de León A. First report of fipronil resistance in *Rhipicephalus (Boophilus) microplus* of Mexico. *Vet Parasitol* 2013;191(1-2):97-101.
89. Rodríguez-Vivas RI, Rivas AL, Chowell G, Fragoso SH, Rosario CR, García Z, Smith SD, Williams JJ, Schwageret SJ. Spatial distribution of acaricide profiles (*Boophilus microplus* strains susceptible or resistant to acaricides) in southeastern Mexico. *Vet Parasitol* 2007;146(1-2):158-169.
90. Encalada-Mena LA, López-Arellano ME, Mendoza de Gives P, Liébrano-Hernández E, Vázquez-Prats V, Vera-Ycuspinera G. Primer informe en México sobre la presencia de resistencia a ivermectina en bovinos infectados naturalmente con nematodos gastrointestinales. *Vet Méx* 2008;39:423-428.
91. Alonso-Díaz MA, Arnaud-Ochoa RA, Becerra-Nava R, Torres-Acosta JFJ, Rodríguez-Vivas RI, Quiroz-Romero RH. Frequency of cattle farms with ivermectin resistant gastrointestinal nematodes in Veracruz, Mexico. *Vet Parasitol* 2015;212(3-4):439-443.

92. Kunz SE, Ortiz EM, Fragozo SH. Status of *Haematobia irritans* (Diptera:Muscidae) insecticide resistance in northeastern Mexico. J Med Entomol 1995;32:726-729.
93. Maldonado SE, Apodaca AC, Sumano LH, Bermudez VL, García VZ, Gutierrez OE. Susceptibilidad de *Haematobia irritans* de las zonas norte de Veracruz y centro de Nuevo León, México, a permetrina y diazinón. Vet Mex 2005;36(2)217-227.
94. Hicks R, Gill D, Owens F. Impact of liver flukes on the performance of feedlot steers. Stillwater, Oklahoma Agr Res Sta. 1989.