

dx.doi.org/10.17488/RMIB.41.2.3

E-LOCATION ID: 1043

A Socio-technical Approach to Evaluating an Electronic Medical Record System implemented in the Public Health Services of Aguascalientes

Una Aproximación Socio-técnica a la Evaluación de un Expediente Médico Electrónico implementado en Servicios de Salud Públicos de Aguascalientes

Christian Díaz de León Castañeda^{1,2}

¹Dirección de Cátedras, Consejo Nacional de Ciencia y Tecnología (CONACYT)

²Universidad Michoacana de San Nicolás de Hidalgo (UMSNH)

ABSTRACT

The objective of this study was to explore a strategy for evaluating an Electronic Medical Record (EMR) system implemented in the public health services of Aguascalientes, Mexico. A questionnaire based on DeLone and McLean's Model of Information Systems Success (MISS) was adapted to Spanish and applied with 62 primary care physicians working in health centers of the *Instituto de Servicios de Salud del Estado de Aguascalientes* (ISSEA or the State of Aguascalientes Institute for Health Services). Opportunities for improving EMR systems were also explored from the informants' perspectives. Additionally, the relationships between MISS components were analyzed using Structural Equations Modeling (SEM). Some MISS components and particular items (service quality and overall satisfaction) presented low averages, reflecting opportunities for improving the development and implementation of EMR, such as the need to continuously update information pertaining to diagnostic and medicine catalogs and develop systems that are interoperable between the second and third levels of care. In conclusion, the present study contributes generating evidence on the use of the MISS to evaluating EMR systems in public health services of Mexico. More evidence should be generated in this field in order to promote the continuous improvement of these information systems.

KEYWORDS: Information science; Information technology management; Medical informatics; Implementation science; Primary health care

RESUMEN

El objetivo de este estudio fue explorar una estrategia para la evaluación de un Expediente Clínico Electrónico (ECE) implementado en servicios de salud públicos de Aguascalientes, México. Se adaptó al español un cuestionario basado en el Modelo de Éxito de Sistemas de Información (MISS) de DeLone y McLean y se aplicó a 62 médicos de atención primaria que trabajan en centros de salud del Instituto de Servicios de Salud del Estado de Aguascalientes (ISSEA). Se exploraron también las oportunidades de mejora del ECE desde la perspectiva de los informantes. Además, se analizaron las relaciones entre los componentes del MISS mediante el modelado de ecuaciones estructurales (SEM). Algunos componentes del MISS e ítems particulares mostraron promedios bajos (p.ej., calidad del servicio y satisfacción) que reflejan algunas oportunidades de mejora en el desarrollo e implementación del ECE, como la necesidad de una actualización continua de la información sobre diagnósticos y catálogos de medicamentos; y el desarrollo de sistemas de interoperabilidad con el segundo y tercer nivel de atención. En conclusión, el presente estudio contribuye en la generación de evidencia sobre el uso del MISS para evaluar los sistemas de EMR en servicios de salud públicos de México. Se debe generar más evidencia en este campo para promover la mejora continua de estos sistemas de información.

PALABRAS CLAVE: Ciencia de la información, Administración de las Tecnologías de la Información; Informática médica; Ciencia de la implementación; Atención primaria de salud

Correspondencia

DESTINATARIO: Dr. Christian Díaz de León Castañeda
INSTITUCIÓN: Universidad Michoacana de San Nicolás
de Hidalgo (UMSNH)
DIRECCIÓN: Avenida Francisco J. Múgica S/N, Ciudad
Universitaria, C. P. 58030, Morelia, Michoacán, México
CORREO ELECTRÓNICO: cddeleon@conacyt.mx

Fecha de recepción:

16 de marzo de 2020

Fecha de aceptación:

6 de junio de 2020

INTRODUCTION

Information and Communication Technologies (ICT) are an important resource for improving the performance of health systems. The applications of ICT in health are closely related to the concept of electronic health (e-Health), which is defined as “the cost-effective and secure use of information and communications technologies in support of health and health-related fields, including health-care services, health surveillance, health literature, and health education, knowledge and research” [1]. A similar concept, under the term digital health, has also been recently introduced [2] [3].

The application of ICT in the areas of health surveillance, monitoring, prevention, promotion and care, at various levels or fields of application, has evolved into the concepts of e-Health and digital health [4].

As the principal components of e-health are the following: health information systems (HIS) supported by ICT (public health informatics, clinical informatics, and consumer health informatics); Electronic Medical Records (EMR); Electronic Health Records (EHR); Picture Archiving and Communication Systems (PACS); Clinical Decision Support Systems (CDSS); remote care systems (telehealth and telemedicine); and, mobile systems (mobile health or *m-Health*). It should be noted that the eHealth innovations have been continually growing [5] [6] [7] [8] [9].

Various studies and systematic reviews have found evidence on the effectiveness and efficiency of the components of eHealth [10] [11] [12] [13], on the basis of which, global and regional organizations have issued diagnostics, policies, and recommendations for improving the implementation of ICT in the health systems of developing and developed countries [3] [14] [15] [16] [17] [18] [19] [20] [21]. To this end, various frameworks and recommendations have been developed from different disciplinary perspectives [22] [23] [24] [25] [26] [27].

In order to discuss eHealth in Mexico, it is essential to first describe the country’s *Sistema Nacional de Salud* (SNS or National Health System), which is characterized by its fragmentation into two large public and private sectors. The public sector is subdivided into health systems for people with formal employment and their direct family members (essentially a social security scheme) and health systems for people without formal employment. The social security system is made up the *Instituto Mexicano del Seguro Social* (IMSS or Mexican Social Security Institute), which comprised 39.2% of the sector in 2015; the *Instituto de Seguridad y Servicios Sociales de los Trabajadores del Estado* (ISSSTE or Social Security and Services Institute for State Workers), which comprised 7.7% of the sector; and the health services used by the armed forces (*Secretaría de la Defensa Nacional*, SEDENA or Ministry of Defense, and *Secretaría de Marina*, SEMAR or Ministry of Navy) and employees of *Petróleos Mexicanos* (Pemex), which are provided by their respective institutions and comprised 1.2%. Health services for people without formal employment are mainly provided by the *Secretaría de Salud* (SSA or Ministry of Health) through the *Servicios Estatales de Salud* (SESA or State Health Services), which comprised 49.9% of the sector, although it is currently undergoing a process of incorporation into the newly-formed *Instituto de Salud para el Bienestar* (INSABI or Institute of Health for Welfare). There are also some special programs within the public sector, such as the *IMSS-Bienestar* (IMSS-Welfare) and the *Sistema Nacional para el Desarrollo Integral de la Familia* (DIF Nacional or National System for Integral Family Development) programs, that aim to provide health-care to vulnerable populations. Finally, the private sector comprises a multiplicity of service providers for people with the ability to pay for healthcare [28] [29] [30].

Recent federal administrations have pursued policies promoting the adoption and implementation of ICT, particularly in the public health sector. The last such policy was the *Estrategia Digital Nacional* (EDN or

National Digital Strategy) [31]. However, individual and organizational factors have limited the adoption and implementation of ICT, such as the persistently fragmented structure of the public health system.

Research on the adoption and implementation of ICT in the public health system in Mexico is scarce, which impedes a broad understanding of the problems faced by those making the decisions in this policy area. However, some general overviews have been published [14] [32] [33] [34], as well as some studies related to particular eHealth components, such as telemedicine [35] and mHealth [36] [37], and initiatives undertaken by private institutions [38] [39] [40]. Regarding EMR systems, some studies have analyzed the planning, adoption, and its implementation in public health institutions [32] [41] [42], focusing on identifying the factors that enable or hinder the success of this implementation. Some of these factors are related to technology (i.e., problems with diagnostic catalogs), the user (i.e., the age of some of the physicians providing primary care), and the organization itself (i.e., a lack of training or quality in its provision).

While the afore mentioned studies have focused principally on exploring the advances in and barriers to the implementation of EMR systems, they have not applied models or frameworks taken from behavioral sciences or social psychology [43] [44] as a guide. In this sense, some socio-technical frameworks have been developed to evaluate information systems in organizations, such as DeLone and McLean's Model of Information Systems Success (MISS) [27]. This approach focuses on evaluating information system quality and implementation in an organizational context. This model focuses on evaluating information systems from the users' perspectives, via three main domains: 1) variables for evaluating the quality of the system, the quality of the information provided, and the quality of the service provided to users via the implementation of the system; 2) both system use and user satisfaction; and, 3) the perceived benefits to the health service in which the system was implemented.

The MISS has been used around the world in the evaluation of EMR systems and other eHealth components generating evidence about the validity of this application in different contexts [45] [46] [47] [48] [49] [50] [51] [52] [53]. The objective of the present study was to explore the use of the MISS to evaluate an EMR implemented in the public health services of Aguascalientes, as a case study that could inform its application in other health institutions of Mexico.

MATERIALS AND METHODS

Study design and theoretical model

The case study design [54] applied in the present study used, as a theoretical guide, DeLone and McLean's Model of Information Systems Success (MISS), particularly its most recent update [27].

Study setting

The present study was performed at the *Instituto de Servicios de Salud del Estado de Aguascalientes (ISSEA)*, which is the main health service provider for the population in the state not covered by social security. In 2015, the total population of Aguascalientes was estimated at 1,292,901 inhabitants, of whom an estimated 42.9% do not have social security coverage [55]. The ISSEA comprises the following: 21 urban health centers (*Centros de Salud Urbanos* or CSUs); and, 62 rural health centers (*Centros de Salud Rurales* or CSRs) providing primary care across three sanitary jurisdictions. The ISSEA has six second-level hospitals (385 beds in total) and one third-level hospital (60 beds in total) [56]. A previous published study described with more detail the healthcare provision system of the ISSEA and the general e-Health ecosystem that have been implemented [57].

The EMR system has been progressively implemented in all the primary care clinics and emergency services functioning at the second and third level of care. The advances in EMR development at the ISSEA are shown

in Table 1. The EMR consists of four main modules: 1) Patient registration and payment; 2) Medical consultation; 3) Pharmacy; and, 4) Statistics. The findings presented in Table 1 reveal the significant development of the EMR system [57], with the users of these EMR modules comprising the following, respectively: 1) Administrative and management personnel; 2) Physicians; 3) Pharmacy technicians; and, 4) Administrative personnel (statisticians).

TABLE 1. Development of the EMR in the study setting.

Patient Registration and Payment
<ul style="list-style-type: none"> • Registration of service users. • Identification of user affiliation to other health subsystems and special programs. • User identification via a unique identifier (Clave Única de Registro de Población – CURP, or Unique Population Registry Code). • Administration of fees for service provision • Schedule for medical appointments. • Human resources administration.
Medical Consultation
<ul style="list-style-type: none"> • Integrated padlocks that automatically issue patient-dependent reminders for screening and health prevention activities. • Integrated clinical information requests to promote proper patient management or control (mainly chronic disease patients). • Integrated alerts or recommendations to improve the quality of care (prompts for referrals to specialists once need has been identified based on clinical parameters). • Links to medical diagnostic catalogs. • Provision of information about patient drug coverage (based on their membership of particular programs) • Provision of pharmacy stock information. • Incorporation of clinical practice guidelines and medication information. • Provision of information on recommended dosages in order to make dosage adjustments and promote the reasonable use of medication.
Pharmacy
<ul style="list-style-type: none"> • Provision of information on the essential medication chart (<i>Cuadro Básico de Medicamentos</i>). • Support for the management of stocks of medication and other health supplies.
Statistics
<ul style="list-style-type: none"> • Provision of the necessary forms for maintaining the monthly health service provision reports and compiling epidemiological statistics.

Source: Prepared by the author.

Sampling of medical personnel

In particular, the medical consultation module of the EMR was evaluated from the perspective of medical personnel either working or providing services in ISSEA CSU or CSR primary care health centers. Physicians were selected as informants, as it is, they

who use said module of the EMR. A convenience sample was obtained based on physicians working as medical directors or in charge of personnel in each health center of the state’s sanitary jurisdictions I and III in the period August 2018 to April 2019. In circumstances where the medical director was not available as an informant, medical personnel were included as informants in the sample, in their stead. The inclusion criterion was to have worked or to be currently working with the EMR system.

Data collection

An adaptation of the online questionnaire developed by Canada Health Infoway, which is based on the Benefits Evaluation Framework and the MISS, was applied in the present study as an instrument for evaluating the EMR [46] [58] [59]. Some items from the questionnaire were selected, translated into Spanish and placed on a Google Forms sheet, which was sent to the informants of the study. The variables included in the model were mainly measured using five-level Likert scales, only the use variable was measured through a proposed 10-level scale. A description of the variables and items included in the instrument is shown in Annex 1.

Data analysis

The data collected was analyzed using STATA 15 software, with Structural Equations Modeling (SEM) analysis used to test the theoretical model comprehensively. Adjustments were implemented in order to optimize the model according to suggestions provided by the software. The model was tested using the maximum likelihood method and the Satorra-Bentler method for standard error computation [60].

Ethical considerations

The research protocol was reviewed and approved by the Research Management of the *Centro de Investigación e Innovación en Tecnologías de la Información y Comunicación* (INFOTEC or Center for Research and Innovation in Information and Communication

Technologies). Prior to responding to the survey, the informants were informed about the research objectives and those responsible for the study. All informants gave their consent to participate in the study. The identity of the informants was protected using alphanumeric codes.

RESULTS AND DISCUSSION

Sample characteristics

The sample characteristics are shown in Table 2. A total of 62 physicians answered the online questionnaire, working at a total of 46 health centers, namely 88.5 % of all the primary care health centers, comprising a total of 52 health centers (15 CSUs and 37 CSRs) in sanitary jurisdictions I and III. Moreover, these represented 38.3 % of the 162 physicians working at these health centers, as previously reported by the SSA [56]. This sample comprised mainly qualified general medical practitioners (66.1%), although the sample also included personnel who had yet to qualify and were carrying out their one-year period of social service (30.6%). The majority had worked with the EMR system for more than three years (46.8%), meaning that, therefore, they were qualified physicians.

Measurement of the MISS variables

The results for the measurement of the MISS variables via their respective items are shown in Table 3. As can be seen, means above three were obtained for all items on the five-level Likert scale used in the present study. The lowest means were obtained for two items related to service quality (SQ1 and SQ2, with means of 3.27 and 3.15, respectively), while the highest means were obtained for items related to system quality (SQ5, with a mean of 4.08) and net perceived benefits (NB5 and NB6, with means of 4.05 and 4.11, respectively). It should be noted that the four MISS variables that were measured using Likert items showed high internal consistency values (Coefficient > 0.85).

TABLE 2. Descriptive statistics of the medical staff surveyed (n=62).

Data	n (%)
Sanitary Jurisdiction	
I	40 (64.5)
III	22 (35.5)
Age	
< 20	1 (1.6)
21 to 30	22 (35.5)
31 to 40	10 (16.1)
41 to 50	18 (29.0)
51 to 60	8 (12.9)
> 61	3 (4.8)
Sex	
Female	24 (38.7)
Male	38 (61.3)
Education	
Non-qualified physician (social service)	19 (30.6)
General medical practitioner	41 (66.1)
Specialist physician	1 (1.6)
Master's degree in Public Health	1 (1.6)
Health service type	
Rural (CSR)	39 (62.9)
Urban (CSU)	23 (37.1)
Time working with EMR	
< 1 month	2 (3.23)
1 to 3 months	10 (16.1)
4 to 6 months	6 (16.1)
7 to 12 months	7 (11.3)
1 to 2 years	8 (12.9)
> 3 years	29 (46.8)
Total	62 (100)

NOTES:

Source: Prepared by the author.

Estimations for the MISS

The structural model tested is shown in Figure 1, while the results of the calculation of direct and indirect effects are shown in Table 4. As can be seen, in terms of the dependent variable of *use*, only *system quality* was a significant variable ($\beta = 0.389$). With regard to the dependent variable of *user satisfaction*, the effects of *system quality* ($\beta = 0.907$) and *service quality* ($\beta = 0.331$) were significant, while, in terms of the dependent variable of *net benefits*, only *user satisfaction* was significant ($\beta = 0.728$). The values of the coefficient of determination (R^2) were high for *user satisfaction* and *net benefits* (0.931 and 0.877, respectively), but low for *use* (0.413).

TABLE 3. Measurement of the variables from the Model of Information Systems Success (MISS).

Variable	Item ^a	No ^b		Mean	Standard Deviation	Coefficient α^c
System Quality	SQ1	L5	The EMR is easy to use	3.94	0.14	0.9503
	SQ2	L5	The effort required to complete an action or access information in the EMR is acceptable	3.84	0.13	
	SQ3	L5	The EMR response time is acceptable	3.84	0.13	
	SQ4	L5	The EMR is integrated with my workflow	3.76	0.14	
	SQ5	L5	The access to the EMR is acceptable	4.08	0.12	
	SQ6	L5	The characteristics of the EMR enable me to perform my job well	3.63	0.14	
	SQ7	L5	The EMR is consistent in its performance	3.64	0.13	
	SQ8	L5	In general, the quality of the EMR is excellent	3.42	0.14	
Information Quality	IQ1	L5	The information provided by the EMR is complete	3.71	0.13	0.9410
	IQ2	L5	The information provided by the EMR is timely	3.95	0.11	
	IQ3	L5	The information provided by the EMR is appropriate	3.87	0.12	
	IQ4	L5	The information provided by the EMR is relevant	3.97	0.11	
	IQ5*	L5	The information provided by the EMR is available when needed	3.79	0.13	
	IQ6	L5	The information provided by the EMR enables me to make patient care decisions or recommendations more quickly	3.69	0.14	
Service Quality	SQ1	L5	The current level of EMR training is acceptable	3.27	0.15	0.8584
	SQ2	L5	The level of ongoing support provided for the use of the EMR is acceptable	3.13	0.15	
Use	U1	N10	On a scale of 0 to 10, What amount do you use the EMR for patient care?	8.7	0.27	1
User Satisfaction	US1	L5	Overall, how satisfied are you with the EMR?	3.8	0.13	1
Net Benefits	NB1	L5	The EMR improves my productivity	3.69	0.14	0.9128
	NB2	L5	The EMR improves the quality of care I provide	3.56	0.16	
	NB3	L5	The EMR makes my job easier	3.71	0.15	
	NB4	L5	The EMR improves our capacity for continuous patient care	3.74	0.14	
	NB5*	L5	The EMR improves our ability to share patient information among healthcare staff	4.05	0.12	
	NB6*	L5	The EMR improves the efficiency of requesting laboratory tests, and X-rays, as well as generating prescriptions	4.11	0.11	
	NB7	L5	The EMR improves the quality of my decision-making	3.43	0.15	
	NB8	L5	The EMR reduces fragmentation in the care provided to my patients and facilitates their mobility on the continuum of care	3.53	0.14	
	NB9	L5	The EMR reduces risks to patient safety	3.89	0.13	

NOTES:

^aItem eliminated in order to improve the internal consistency of the respective variable (Coefficient α) and facilitate model adjustment

^bNumber of categories used for the item response: L5 – Likert scale of 5 categories; and, N – numeric scale

^cCoefficient α obtained after model adjustment

Source: Prepared by the author.

Model adjustments were made by means of the software used in the present study, with some items (IQ5, NB5, and NB6) excluded, thus enabling better reliability and model fit. Through the adjustments made, a model was obtained that meets some of the criteria suggested in the literature [60], such as the significance of the Chi² statistical test ($p < 0.000$), $\text{Chi}^2/\text{df} < 2$ (result = 1.47) and the SRMR indicator < 0.1 (result = 0.068). Also, this model was near to meet other criteria such as RMSEA < 0.05 (result = 0.110) and CFI > 0.95 (result = 0.877) indicators.

TABLE 4. Direct, indirect and total effects on the dependent variables obtained in the adjusted model.^a

A Predicts B		Standardized Effects		
A	B	Direct	Indirect	Total
System quality	Use	0.389**	-----	0.389**
Information quality	Use	0.228	-----	0.228
Service quality	Use	0.105	-----	0.105
System quality	User satisfaction	0.907***	-0.168	0.739***
Information quality	User satisfaction	0.064	-0.098	-0.034
Service quality	User satisfaction	0.331*	-0.045	0.286**
Use	User satisfaction	-0.431	-----	-0.431
System quality	Net benefits	-----	0.696***	0.696***
Information quality	Net benefits	-----	0.068	0.068
Service quality	Net benefits	-----	0.251**	0.251**
Use	Net benefits	0.406	-0.314	0.092
User satisfaction	Net benefits	0.728*	-----	0.728*


NOTES:

^aStatistical significance levels: * $p < 0.05$; ** $p < 0.01$; and, *** $p < 0.001$

Source: Prepared by the author.

Improvement opportunities for the EMR

Various improvement opportunities were identified from the physicians' perspectives, principally regarding the medical consultation module (Table 5). Some of the most significant suggestions made by the physicians were: 1) The continuous updating of information, such as diagnostic catalogs, stock information, and



^aThe rectangles represent the variables comprising the MISS, while the arrows show the direct effects of independent variables on the dependent variables of this model (the indirect and total effects are described in Table 4).

^bStatistical significance of the path coefficients: * $p < 0.05$; ** $p < 0.01$; and, *** $p < 0.001$

Source: Prepared by the author.

FIGURE 1. Adjusted MISS.^{a,b}

information pertaining to the co-payment systems for drug catalogs; 2) The implementation of interoperable information systems between the second and third levels of care to improve the availability of information (i.e., results of laboratory tests performed in hospital and counter-referral information); 3) The automation of the monthly reporting of institutional statistics, including the data taken from private sector information systems (Fundación Carlos Slim [38] [61]).

This study used a socio-technical and psychometric approach to evaluate an EMR system by applying the MISS, a comprehensive conceptual framework. The main contribution of the study could be found in the manner in which it applied this model in the primary care physicians and in the context of the study setting.

As an approach of reliability, the data gotten from the instrument adapted for and translated into Spanish obtained good results in terms of the internal consistency (Coefficient α) in the different constructs. Besides, an structural equations model (SEM) analysis was used to study the relationship among the different variables of the MISS allowing to identify the relationship between the variables related to EMR quality (*quality of the system/quality of the information*) and implementation (*quality of the service*) and the closely related *satisfaction/use* variables, as well as the relationship with the *per-*

ceived net benefits reported by medical personnel. However, it was not possible to prove some relationships, mainly those with the use variable, which could be due to the fact that it was measured with a single item, when there may be different modalities and dimensions of the use of the EMR by medical personnel, including the frequency, types, objectives, and quality of use.

In terms of the statements of an international consensus regarding psychometric research reporting [62], this study contributes on generating evidence about the validity of the data gotten from the adapted instrument derived from the MISS, as an approach to evaluating EMR systems in public health services in Mexico. However, it is considered that more studies should be performed in order to improve the adapted instrument and generate robust evidence taking into account the different contexts derived from the country's fragmented health system. Also, the incorporation of the perspectives of physicians working at the second or third levels of care as well as other healthcare professionals could be of great relevance to particular healthcare systems, considering the different levels of EMR implementation in the public health system. In this regard, it should be noted that various published studies have contributed too on generating evidence in this field, using the MISS, and some modified versions, in other countries at different levels of healthcare, from different health workers' perspectives, and focusing on the evaluation of different eHealth systems, such as HIS, EMR and EHR [45] [46] [47] [48] [49] [50], and PACS [51] [52] [53].

Moreover, beyond the focus on developing and validating the MISS or an integrative instrument based on it, the building of indicators from the measurement of MISS variables could also be used as a way to guide and inform the implementation of EMR systems (or other eHealth components) in healthcare institutions. Expanding on this topic, these indicators could be tested and later used in the context of eHealth or digital health policy implementation in Mexico in different contexts.

It should be noted that this has been the focus of studies conducted in other countries [63] [64], some of which have focused their analysis on identifying opportunities to improve the application of these technologies [65] [66].

The following were identified as strengths of the study: 1) The use of MISS, an integrated framework that incorporates various dimensions related to the success of information systems and goes beyond the technology adoption or acceptance models; 2) The use of a structural equation model (SEM) as an analytical methodology that enables an analysis of all the model variables and their relationships (such as their direct and indirect effects); and, 3) The identification of opportunities for improving the EMR system as well as its implementation from the perspective of medical staff, through which the authorities or management can be made aware informed (a formative evaluation approach, which focuses on improving implementation [67] [68] and has been used in other eHealth studies [48]).

As weaknesses of the present study, the following were identified: 1) The case-study design was applied to only one state health service thus limiting the external validity of the study to other state health institutions (SESA, SSA), as these have implemented their own EMR systems; 2) a complete set of responses by the physicians to the questionnaire was not obtained due to the low response rate; and, 3) The need to build a more integrated measurement of the use variable. These weaknesses are identified as further research opportunities.

CONCLUSION

The present study contributes generating evidence on the use of the MISS and the adapted instrument to evaluating EMR systems in public health services of Mexico, although some improvements should be performed in this instrument. Besides, more evidence should be generated in this field in order to promote the continuous improvement of EMR systems, in the context of the e-Health (or Digital health) policy.

Conflicts of interest

The author declares that there is no conflict of interest.

Acknowledgments

The author is extremely grateful to the institutions and individuals that collaborated in carrying out the present study: the *Cátedras CONACyT* (*Consejo Nacional de Ciencia y Tecnología*) program for youth researchers

(project No. 1010 Apropriación Social de Tecnologías de Información y Comunicación para el Desarrollo, proposed by INFOTEC); *Instituto de Servicios de Salud del Estado de Aguascalientes* (ISSEA); the academic staff at the *Centro de Investigación en Matemáticas* (CIMAT or the Center for Mathematics Research), Aguascalientes branch, for the advice provided as part of an SEM course; and, the physicians that participated as informants.

REFERENCES

- [1] World Health Organization. e-Health (WHA58.28). In: Fifty-eighth World Health Assembly (WHA58/2005/REC/1) [Internet]. Geneva: WHO; 2005. p. 108-10. Available from: <http://www.who.int/iris/handle/10665/20398>
- [2] World Health Organization. mHealth, use of appropriate digital technologies for public health. Report by the Director-General. In: Seventy-first World Health Assembly - Provisional agenda item 124 (A71/20) [Internet]. Geneva: WHO; 2018. p. 26. Available from: http://apps.who.int/gb/e/e_wha71.html
- [3] World Health Organization. Digital Health. In: Seventy-first World Health Assembly - Agenda item 124 (A71/A/CONF/1) [Internet]. Geneva: WHO; 2018. p. 21. Available from: http://apps.who.int/gb/e/e_wha71.html
- [4] World Health Organization. WHO guideline: recommendations on digital interventions for health system strengthening [Internet]. Geneva: WHO; 2019. Available from: <https://www.who.int/reproductivehealth/publications/digital-interventions-health-system-strengthening/en/>
- [5] Aceto G, Persico V, Pescapé A. The role of Information and Communication Technologies in healthcare: taxonomies, perspectives, and challenges. *J Netw Comput Appl* [Internet]. 2018;107:125-54. Available from: <https://doi.org/10.1016/j.jnca.2018.02.008>
- [6] Mettler T, Raptis DA. What constitutes the field of health information systems? Fostering a systematic framework and research agenda. *Health Informatics J* [Internet]. 2012;18(2):147-56. Available from: <https://doi.org/10.1177/1460458212452496>
- [7] Barrett D, Liaw S-T, De Lusignan S. Unravelling the tangled taxonomies of health informatics. *J Innov Heal Informatics* [Internet]. 2014;21(3):152-5. Available from: <https://doi.org/10.14236/jhi.v21i3.78>
- [8] Boogerd EA, Arts T, Engelen LJ, van de Belt TH. "What is eHealth": time for an update? *JMIR Res Protoc* [Internet]. 2015;4(1):e29. Available from: <http://dx.doi.org/10.2196/resprot.4065>
- [9] Shaw T, McGregor D, Brunner M, Keep M, Janssen A, Barnet S. What is eHealth (6)? Development of a Conceptual Model for eHealth: Qualitative Study with Key Informants. *J Med Internet Res* [Internet]. 2017;19(10):e324. Available from: <https://dx.doi.org/10.2196%2Fjmir.8106>
- [10] Elbert NJ, Van Os-Medendorp H, Van Renselaar W, Ekeland AG, Hakkaart-Van Roijen L, Raat H, et al. Effectiveness and cost-effectiveness of ehealth interventions in somatic diseases: A systematic review of systematic reviews and meta-analyses. *J Med Internet Res* [Internet]. 2014;16(4):e110. Available from: <https://doi.org/10.2196/jmir.2790>
- [11] Flodgren G, Rachas A, Farmer AJ, Inzitari M, Shepperd S. Interactive telemedicine: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* [Internet]. 2015;(9):CD002098. Available from: <https://doi.org/10.1002/14651858.cd002098.pub2>
- [12] Iribarren SJ, Cato K, Falzon L, Stone PW. What is the economic evidence for mHealth? A systematic review of economic evaluations of mHealth solutions. *PLoS One* [Internet]. 2017;12(2):e0170581. Available from: <https://doi.org/10.1371/journal.pone.0170581>
- [13] Totten AM, Womack DM, Eden KB, McDonagh MS, Griffin JC, Grusing S, et al. Telehealth: mapping the evidence for patient outcomes from systematic reviews [Internet]. Rockville, MD: Agency for Healthcare Research and Quality; 2016. Available from: www.effectivehealthcare.ahrq.gov/reports/final.cfm
- [14] World Health Organization. Atlas of eHealth country profiles: the use of eHealth in support of universal health coverage, based on the findings of the third global survey on eHealth 2015 [Internet]. Geneva: WHO; 2016. Available from: http://www.who.int/goe/publications/atlas_2015/en/
- [15] World Health Organization, International Telecommunication Union. National eHealth Strategy Toolkit [Internet]. Geneva: WHO; 2012. Available from: <https://apps.who.int/iris/handle/10665/75211>
- [16] Organización Panamericana de la Salud. Estrategia y plan de acción sobre eSalud (2012-2017) [Strategy and plan of action on eHealth 2012-2017] [Internet]. 2011. Available from: http://www.paho.org/ict4health/index.php?option=com_content&view=article&id=54:estrategia-y-plan-de-accion-sobre-esalud-2012-2017&Itemid=&lang=es
- [17] Organización Panamericana de la Salud. La eSalud en la región de las américas : derribando las barreras a la implementación. Resultados de la Tercera Encuesta Global de eSalud de la Organización Mundial de la Salud [eHealth in the Americas region: breaking down the barriers to implementation. Results of the Third Global Survey on eHealth of the World Health Organization] [Internet]. Washington, D.C.: OPS; 2016. Available from: <http://iris.paho.org/xmloi/handle/123456789/31287>
- [18] Organization for Economic Cooperation and Development (OECD). Improving Health Sector efficiency: The Role of Information and Communication Technologies. Paris: OECD; 2010. Available from: <https://doi.org/10.1787/2074319x>
- [19] Comisión Económica para América Latina y el Caribe (CEPAL). Salud electrónica en América Latina y el Caribe: avances y desafíos [Electronic health in Latin America and the Caribbean: progress and challenges] [Internet]. Fernández A, Oviedo E, editors. Santiago de Chile: Comisión Económica para América Latina y el Caribe (CEPAL), Naciones Unidas; 2010. Available from: <https://www.cepal.org/es/publicaciones/salud-electronica-en-america-latina-y-el-caribe-avances-y-desafios>
- [20] Palacios J, Flores-Roux E, García Zaballos A. Diagnóstico del sector TIC en México, Conectividad e inclusión social para la mejora de la productividad y el crecimiento económico [Diagnosis of the ICT sector in Mexico, Connectivity and social inclusion for the improvement of productivity and economic growth] [Internet]. Documento de debate IDB-DP-235. Banco Interamericano de Desarrollo (IDB); 2013. Available from: <https://publications.iadb.org/es/diagnostico-del-sector-tic-en-mexico-conectividad-e-inclusion-social-para-la-mejora-de-la>
- [21] World Health Organization. Classification of Digital Health Interventions v 1.0, a shared language to describe the uses of digital technology for health [Internet]. Geneva: WHO; 2018. Available from: <http://www.who.int/iris/handle/10665/260480>
- [22] Díaz de León-Castañeda C. Electronic health (e-Health): a conceptual framework for its implementation in health services facilities. *Gac Med Mex* [Internet]. 2019;155(2):161-8. Available from: <https://doi.org/10.24875/GMM.M19000251>

- [23] Glasgow RE, Phillips SM, Sanchez MA. Implementation science approaches for integrating eHealth research into practice and policy. *Int J Med Inform* [Internet]. 2014 Jul;83(7):e1-11. Available from: <http://dx.doi.org/10.1016/j.ijmedinf.2013.07.002>
- [24] Bergmo T. How to measure costs and benefits of eHealth interventions: An overview of methods and frameworks. *J Med Internet Res* [Internet]. 2015;17(11):e254. Available from: <http://dx.doi.org/10.2196/jmir.4521>
- [25] McNamee P, Murray E, Kelly MP, Bojke L, Chilcott J, Fischer A, et al. Designing and Undertaking a Health Economics Study of Digital Health Interventions. *Am J Prev Med* [Internet]. 2016;51(5):852-60. Available from: <http://dx.doi.org/10.1016/j.amepre.2016.05.007>
- [26] Yusof MM, Papazafeiropoulou A, Paul RJ, Stergioulas LK. Investigating evaluation frameworks for health information systems. *Int J Med Inform* [Internet]. 2008;77(6):377-85. Available from: <https://doi.org/10.1016/j.ijmedinf.2007.08.004>
- [27] Delone W, Mclean E. The DeLone and McLean Model of Information Systems Success: A Ten-Year Update. *J Manag Inf Syst* [Internet]. 2003;19(4):9-30. Available from: <https://doi.org/10.1080/07421222.2003.11045748>
- [28] Gómez-Dantés O, Sesma S, Becerril VM, Knau FM, Arreola H FJ. Sistema de Salud de México [Mexico Health System]. *Salud Publica Mex* [Internet]. 2011;53(supl 2):S220-32. Available from: <http://saludpublica.mx/index.php/spm/article/view/5043/10023>
- [29] Instituto Nacional de Estadística y Geografía (INEGI). Encuesta Intercensal 2015 [Intercensal Survey 2015] [Internet]. 2015. Available from: <http://www.beta.inegi.org.mx/proyectos/enchogares/especiales/intercensal/>
- [30] Diario Oficial de la Federación. DECRETO por el que se reforman, adicionan y derogan diversas disposiciones de la Ley General de Salud y de la Ley de los Institutos Nacionales de Salud [DECREE by which various directions of the General Health Law and the Law of the National Institutes of Health are reformed, added and repealed] [Internet]. 2019 Nov 29; Available from: http://www.dof.gob.mx/nota_detalle.php?codigo=5580430&fecha=29/11/2019
- [31] Gobierno Federal. Estrategia Digital Nacional [National Digital Strategy] [Internet]. 2013. Available from: https://www.gob.mx/cms/uploads/attachment/file/17083/Estrategia_Digital_Nacional.pdf
- [32] Mariscal Avilés J, Gil-García JR, Ramírez-Hernández F. E-Salud en México: antecedentes, objetivos, logros y retos [E-Health in Mexico: background, objectives, achievements and challenges]. *Espac Públicos* [Internet]. 2012;15(34):65-94. Available from: <http://www.redalyc.org/articulo.oa?id=67623463015>
- [33] Gertrudiz N. e-Health: the case of Mexico [e-Health: the case of Mexico]. *Lat Am J Telehealth* [Internet]. 2010;2(2):127-67. Available from: <http://cetec.medicina.ufmg.br/revista/index.php/rlat/article/view/71>
- [34] Organización Panamericana de la Salud. Conversaciones sobre eSalud. Gestión de información, diálogos e intercambio de conocimientos para acercarnos al acceso universal a la salud [Conversations about eHealth. Information management, dialogues and knowledge exchange to get closer to universal access to health] [Internet]. Washington, DC: OPS; 2014. Available from: <http://iris.paho.org/xmlui/handle/123456789/28391>
- [35] González-Rétiz ML, Pacheco-López A. Desarrollo de la telesalud en México [Development of telehealth in Mexico] [Internet]. Santiago de Chile: Comisión Económica para América Latina y el Caribe (CEPAL), Naciones Unidas; 2013. Available from: <http://hdl.handle.net/11362/4055>
- [36] Vélez-Andrade C, Rodríguez-Hernández J. Salud Móvil: Nuevos Horizontes para la Promoción de la Salud. Proyecto m-Salud para la prevención, diagnóstico oportuno y control del sobrepeso/obesidad y diabetes tipo 2 [Mobile Health: New Horizons for Health Promotion. m-Health Project for the prevention, timely diagnosis and control of overweight / obesity and type 2 diabetes]. México City: Fundación Mexicana para la Salud (Funsalud); 2018. Available from: https://funsalud.org.mx/wp-content/uploads/2019/11/Salud-Movil_-Nuevos-Horizontes-para-la-Promocion-de-la-Salud.pdf
- [37] Cabrera NI, Castro P, Demeneghi VP, Fernández L, Morales J, Sainz L, et al. mSalUV: un nuevo sistema de mensajería móvil para el control de la diabetes en México [mSalUV: a new mobile messaging system for diabetes control in Mexico]. *Rev Panam Salud Publica* [Internet]. 2014;35(5-6):371-7. Available from: <https://www.scielosp.org/pdf/rpsp/2014.v35n5-6/371-377>
- [38] Tapia-Conyer R, Gallardo-Rincón H, Saucedo-Martínez R. CASALUD: An innovative health-care system to control and prevent non-communicable diseases in Mexico. *Perspect Public Health* [Internet]. 2015;135(4):180-90. Available from: <https://doi.org/10.1177/1757913913511423>
- [39] Tapia-Conyer R, Saucedo-Martínez R, Mujica-Rosales R, Gallardo-Rincón H, Campos-Rivera PA, Lee E, et al. Enablers and inhibitors of the implementation of the Casalud Model, a Mexican innovative healthcare model for non-communicable disease prevention and control. *Heal Res Policy Syst* [Internet]. 2016;14(1):1-12. Available from: <http://dx.doi.org/10.1186/s12961-016-0125-0>
- [40] Gallardo-Rincón H, Saucedo-Martínez R, Mujica-Rosales R, Lee EM, Israel A, Torres-Beltrán B, et al. Online continuing medical education as a key link for successful noncommunicable disease self-management: The CASALUDTM Model. *Diabetes, Metab Syndr Obes Targets Ther* [Internet]. 2017;10:443-55. Available from: <https://doi.org/10.2147/DMSO.S137891>
- [41] Hernández-Ávila JE, Palacio-Mejía LS, Lara-Esqueda A, Silvestre E, Agudelo-Botero M, Diana ML, et al. Assessing the process of designing and implementing electronic health records in a statewide public health system: The case of Colima, Mexico. *J Am Med Informatics Assoc* [Internet]. 2013;20(2):238-44. Available from: <http://dx.doi.org/10.1136/amiajnl-2012-000907>
- [42] González-Block M, Lopez-Santibañez C, Aguilera Aburto N, Aldama A, Barajas M, Becerril Montekio V, et al. Evaluación y estrategias de portabilidad y convergencia hacia la integración del Sistema Nacional de Salud, resumen ejecutivo [Evaluation and strategies of portability and convergence towards the integration of the National Health System, Executive summary] [Internet]. Cuernavaca: Instituto Nacional de Salud Pública; 2011. Available from: http://pliopencms05.salud.gob.mx:8080/BasesCDI/Archivos/Informes/portabilidad_convergencia.pdf
- [43] Fernández Morales K, McAnally Salas L, Vallejo Casarín A. Apropiación tecnológica: una visión desde los modelos y las teorías que la explican [Technological appropriation: a view from the models and theories that explain it]. *Perspect Educ* [Internet]. 2015;54(2):109-25. Available from: <https://doi.org/10.4151/07189729-Vol.54-Iss.2-Art.331>
- [44] Korpelainen K. Theories of ICT system implementation and adoption - a critical review [Internet]. Helsinki: School of Science and Technology, Aalto University; 2011. Available from: <https://aaltodoc.aalto.fi/handle/123456789/5019>

- [45] Yu P, Qian S. Developing a theoretical model and questionnaire survey instrument to measure the success of electronic health records in residential aged care. *PLoS One* [Internet]. 2018;13(1):e0190749. Available from: <https://doi.org/10.1371/journal.pone.0190749>
- [46] Tharmalingam S, Hagens S, Zelmer J. The value of connected health information: perceptions of electronic health record users in Canada. *BMC Med Inform Decis Mak* [Internet]. 2016;16(1):93. Available from: <http://dx.doi.org/10.1186/s12911-016-0330-3>
- [47] Raymond L, Paré G, Ortiz de Guinea A, Poba-Nzaou P, Trudel M-C, Marsan J, et al. Improving performance in medical practices through the extended use of electronic medical record systems: a survey of Canadian family physicians. *BMC Med Inform Decis Mak* [Internet]. 2015;15(1):27. Available from: <https://doi.org/10.1186/s12911-015-0152-8>
- [48] Bossen C, Jensen LG, Udsen FW. Evaluation of a comprehensive EHR based on the DeLone and McLean model for IS success: Approach, results, and success factors. *Int J Med Inform* [Internet]. 2013;82(10):940-53. Available from: <https://doi.org/10.1016/j.ijmedinf.2013.05.010>
- [49] Messeri P, Khan S, Millery M, Campbell A, Merrill J, Shih S, et al. An Information Systems Model of the Determinants of Electronic Health Record Use. *Appl Clin Inform* [Internet]. 2013;4(2):185-200. Available from: <http://dx.doi.org/10.4338/ACI-2013-01-RA-0005>
- [50] Alharthi H, Youssef A, Radwan S, Al-Muallim S, Zainab A-T. Physician satisfaction with electronic medical records in a major Saudi Government hospital. *J Taibah Univ Med Sci* [Internet]. 2014;9(3):213-8. Available from: <https://doi.org/10.1016/j.jtumed.2014.01.004>
- [51] Tzeng WS, Kuo KM, Lin HW, Chen TY. A Socio-technical assessment of the success of picture archiving and communication systems: The radiology technologist's perspective. *BMC Med Inform Decis Mak* [Internet]. 2013;13(1):1. Available from: <http://dx.doi.org/10.1186/1472-6947-13-109>
- [52] Sicotte C, Paré G, Bini KK, Moreault M-P, Laverdure G. Virtual Organization of Hospital Medical Imaging: A User Satisfaction Survey. *J Digit Imaging* [Internet]. 2010;23(6):689-700. Available from: <https://doi.org/10.1007/s10278-009-9220-x>
- [53] Paré G, Lepanto L, Aubry D, Sicotte C. Toward a multidimensional assessment of picture archiving and communication system success. *Int J Technol Assess Health Care* [Internet]. 2005;21(4):471-9. Available from: <https://doi.org/10.1017/s0266462305050658>
- [54] Yin RK. *Case Study Research: Design and Methods*. 5th ed. Thousand Oaks: SAGE Publications; 2014. 282 p.
- [55] Consejo Nacional de Evaluación de la Política de Desarrollo Social (CONEVAL). Información de Pobreza y Evaluación de las Entidades Federativas [Poverty Information and Evaluation of the Federal Entities] [Internet]. 2019. Available from: <https://www.coneval.org.mx/coordinacion/entidades/Paginas/inicioent.aspx>
- [56] Secretaría de Salud. Portal de datos abiertos de Secretaría de Salud [Open data portal of the Ministry of Health] [Internet]. 2018. Available from: <https://datos.gob.mx/busca/organization/salud>
- [57] Díaz de León Castañeda C, Góngora Ortega J. eSalud en servicios de salud públicos en México: estudio de caso [E-Health in Public Health Services in Mexico: Case Study]. *Región y Soc* [Internet]. 2020;32:e1256. Available from: <https://doi.org/10.22198/rys2020/32/1256>
- [58] Canada Health Infoway. System and use assessment survey (refresh february 2015) [Internet]. 2015. Available from: <https://www.infoway-inforoute.ca/en/component/edocman/resources/toolkits/change-management/national-framework/monitoring-and-evaluation/resources-and-tools/991-benefits-evaluation-survey-process-system-use-assessment-survey?Itemid=101>
- [59] Lau F, Hagens S, Muttitt S. A proposed benefits evaluation framework for health information systems in Canada. *Healthc Q* [Internet]. 2007;10(1):113-8. Available from: <https://www.longwoods.com/content/18657>
- [60] Stata 25 Software. *Structural Equations Modelling Reference Manual* [Internet]. 2019. Available from: <https://www.stata.com/bookstore/structural-equation-modeling-reference-manual/>
- [61] Fundación Carlos Slim. Programas de Innovación [Innovation programs] [Internet]. 2018. Available from: <https://www.salud.carlosslim.org/innovacion/>
- [62] American Educational Research Association (AERA), American Psychological Association (APA), National Council on Measurement in Education (NCME). *Standards for Educational and Psychological Testing*. Washington, DC: AERA, APA, NCME; 2014. 230 p.
- [63] Buabbas AJ, Al-Shamali DA, Sharma P, Haidar S, Al-Shawaf H. Users' Perspectives on a Picture Archiving and Communication System (PACS): An In-Depth Study in a Teaching Hospital in Kuwait. *JMIR Med Informatics* [Internet]. 2016;4(2):e21. Available from: <https://doi.org/10.2196/medinform.5703>
- [64] Otieno GO, Hinako T, Motohiro A, Daisuke K, Keiko N. Measuring effectiveness of electronic medical records systems: Towards building a composite index for benchmarking hospitals. *Int J Med Inform* [Internet]. 2008;77(10):657-69. Available from: <https://doi.org/10.1016/j.ijmedinf.2008.01.002>
- [65] Kuo K, Liu C, Talley PC, Pan S. Strategic Improvement for Quality and Satisfaction of Hospital Information Systems. *J Healthc Eng* [Internet]. 2018:1-14. Available from: <https://doi.org/10.1155/2018/3689618>
- [66] Cohen JF, Coleman E, Kangethe MJ. An importance-performance analysis of hospital information system attributes: A nurses' perspective. *Int J Med Inform* [Internet]. 2016;86:82-90. Available from: <http://dx.doi.org/10.1016/j.ijmedinf.2015.10.010>
- [67] Bauer MS, Damschroder L, Hagedorn H, Smith J, Kilbourne AM. An introduction to implementation science for the non-specialist. *BMC Psychol* [Internet]. 2015;3(1):32. Available from: <https://doi.org/10.1186/s40359-015-0089-9>
- [68] Lau F, Kuziemsy C, Price M, Gardner J. A review on systematic reviews of health information system studies. *J Am Med Informatics Assoc* [Internet]. 2010;17(5):637-45. Available from: <https://doi.org/10.1136/jamia.2010.004838>

ANNEXES

ANNEX 1. Instrument for the Variables Measurement of the Model of Information Systems Success (MISS).

Variable/Items	Strongly disagree	Moderately disagree	Neither agree nor disagree	Moderately agree	Strongly agree
System quality^a					
The EMR is easy to use	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The effort required to complete an action or access information in the EMR is acceptable	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR response time is acceptable	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR is integrated with my workflow	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The access to the EMR is acceptable	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The characteristics of the EMR enable me to perform my job well	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR is consistent in its performance	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
In general, the quality of the EMR is excellent	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Information quality^a					
The information provided by the EMR is complete	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The information provided by the EMR is timely	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The information provided by the EMR is appropriate	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The information provided by the EMR is relevant	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The information provided by the EMR is available when needed	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The information provided by the EMR enables me to make patient care decisions or recommendations more quickly	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Service quality^a					
The current level of EMR training is acceptable	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The level of ongoing support provided for the use of the EMR is acceptable	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
User satisfaction^a					
Overall, how satisfied are you with the EMR?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Perceived net benefits^a					
The EMR improves my productivity	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR improves the quality of care I provide	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR makes my job easier	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR improves our capacity for continuous patient care	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR improves our ability to share patient information among healthcare staff	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR improves the efficiency of requesting laboratory tests, and X-rays, as well as generating prescriptions	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR improves the quality of my decision-making	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR reduces fragmentation in the care provided to my patients and facilitates their mobility on the continuum of care	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
The EMR reduces risks to patient safety	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Use^b					
On a scale of 0 to 10, What amount do you use the EMR for patient care?					
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
1	2	3	4	5	6
7	8	9	10		

NOTES:

^a Variables measured through five-level Likert scales. Items taken or adapted from Canada Health Infoway survey [].

^b This variable was measured from a 10 point scale, proposed in this study.