

# Social determinants for overweight and obesity in a highly marginalized population from Comitán, Chiapas, Mexico

Marisela Vázquez-Durán, MSc,<sup>(1)</sup> María Eugenia Jiménez-Corona, PhD,<sup>(2)</sup> Laura Moreno-Altamirano, PhD,<sup>(3)</sup>  
 Enrique O Graue-Hernández, MSc,<sup>(4)</sup> Noé Guarneros, MSc,<sup>(5)</sup> Leonardo Jiménez-Corona, BIT,<sup>(6)</sup>  
 Aída Jiménez-Corona, PhD.<sup>(1,7)</sup>

Vázquez-Durán M, Jiménez-Corona ME, Moreno-Altamirano L, Graue-Hernández EO, Guarneros N, Jiménez-Corona L, Jiménez-Corona A. Social determinants for overweight and obesity in a highly marginalized population from Comitán, Chiapas, Mexico. *Salud Publica Mex.* 2020;62:477-486. <https://doi.org/10.21149/10691>

## Abstract

**Objective.** We assessed the prevalence of overweight and obesity and its association with some social determinants in a highly marginalized population in Mexico. **Materials and methods.** Cross-sectional study conducted in Comitán, Chiapas, from 2010 to 2012, comprising 1 858 subjects aged  $\geq 20$  years. We evaluated proximal, intermediate, and structural social determinants. **Results.** The prevalence of overweight and obesity was 37.9 and 16.5%, respectively. The probability of overweight and obesity was higher in participants with  $\geq$ primary school, self-reported non-indigenous origin, and medium level of marginalization compared with those with  $<$ primary school, self-reported indigenous origin, and high/very high level of marginalization. **Conclusion.** The probability of overweight and obesity was higher in population with more favorable social conditions, which may be partially explained by changes in the traditional lifestyle with greater access to high energy foods and physical inactivity.

Keywords: prevalence; overweight; obesity; social determinants

Vázquez-Durán M, Jiménez-Corona ME, Moreno-Altamirano L, Graue-Hernández EO, Guarneros N, Jiménez-Corona L, Jiménez-Corona A. Determinantes sociales relacionados con el sobrepeso y obesidad en una población con alto grado de marginación de Comitán, Chiapas, México. *Salud Publica Mex.* 2020;62:477-486. <https://doi.org/10.21149/10691>

## Resumen

**Objetivo.** Estimar la prevalencia de sobrepeso y obesidad y su asociación con determinantes sociales en población con alto grado de marginación. **Material y métodos.** Estudio transversal realizado en Comitán, Chiapas, de 2010 a 2012, que incluyó 1 858 sujetos  $\geq 20$  años de edad. Se evaluaron determinantes sociales proximales, intermedios y estructurales. **Resultados.** La prevalencia de sobrepeso y obesidad fue de 37.9 y 16.5%, respectivamente. La probabilidad de sobrepeso y obesidad fue mayor en sujetos con escolaridad  $\geq$ primaria, en sujetos que se autodefinieron como no indígenas y en sujetos con un grado de marginación medio comparado con individuos con escolaridad  $<$ primaria, con autodefinirse como indígena y tener un grado de marginación alto/muy alto. **Conclusión.** La probabilidad de sobrepeso y obesidad fue mayor en población con condiciones sociales más favorables, parcialmente explicada por cambios en el estilo de vida con mayor acceso a alimentos con alta energía e inactividad física.

Palabras clave: prevalencia; sobrepeso; obesidad; determinantes sociales

- (1) Departamento de Epidemiología Ocular y Salud Visual, Instituto de Oftalmología Conde de Valenciana IAP. Mexico City, Mexico.
- (2) Departamento de Epidemiología, Instituto Nacional de Cardiología Ignacio Chávez, Secretaría de Salud. Mexico City, Mexico.
- (3) Facultad de Medicina, Universidad Nacional Autónoma de México. Mexico City, Mexico.
- (4) Departamento de Córnea y Cirugía Refractiva, Instituto de Oftalmología Conde de Valenciana IAP. Mexico City, Mexico.
- (5) Instituto Nacional de Salud Pública. Cuernavaca, Morelos, Mexico.
- (6) Hospital de la Mujer de Comitán. Comitán de Domínguez, Chiapas, Mexico.
- (7) Dirección General de Epidemiología, Secretaría de Salud. Mexico City, Mexico.

Received on: July 3, 2019 • Accepted on: March 11, 2020 • Published online: August 29, 2020  
 Corresponding author: Aida Jiménez Corona. Departamento de Epidemiología Ocular y Salud Visual, Instituto de Oftalmología Conde de Valenciana IAP. Chimalpopoca 14, col. Obrera, Alcaldía Cuauhtémoc. 06800, Mexico City, Mexico.  
 email: aidaajc@gmail.com

License: CC BY-NC-SA 4.0

According to estimations of the World Health Organization (WHO), in 2016 there were worldwide more than 1 900 million adults aged over 18 years with overweight and obesity (O/O), 650 million corresponding to obesity. It is estimated that by the year 2030, 40% of the world population will be overweight and 20%, obese.<sup>1</sup> Globally, Mexico is ranked as the second country with the largest number of people with O/O, behind the United States of America. Mexico's 2012 National Survey of Health and Nutrition (Ensanut 2012) found that 71.2% of the adult population had O/O; the prevalence was higher in upper (73.5%) compared with lower socioeconomic strata (65.7%) and in urban (72.9%) than in rural areas (65.6%).<sup>2</sup>


The higher prevalence of O/O is the result of environmental and social transformations related to poor supportive policies in public sectors such as health, agriculture, transport, and education, as well as of factors such as food access,<sup>3,4</sup> inequalities of gender and of social, educational, and economic support networks, employment, and access to health services.<sup>5,6</sup> Marmot defines social determinants as the conditions in which the population develops, those conditions being the product of the political and economic context of each region (figure 1).<sup>7</sup> Through these determinants it is possible to examine the unfair distribution of social goods and how avoidable inequalities manifest themselves in the health status.<sup>5</sup>

The State of Chiapas is among the four poorest states in Mexico and is the second state with the highest proportion of indigenous population. Although the prevalence of O/O in Chiapas is lower than that in wealthier states, the prevalence of overweight increased from 39.8% in 2006 to 41% in 2012 in urban areas, whereas it increased from 37.4 to 39.5% in rural areas.<sup>8</sup> Changes on obesity prevalence were less evident, but it seems that this population has a similar trend than populations with better social conditions, so we need specific strategies for them. This study aimed to assess the prevalence of O/O and its association with proximal, intermediate, and structural social determinants in population from the Municipality of Comitán de Domínguez, in the State of Chiapas, Mexico. We hypothesized that more favorable social conditions predispose to higher probability of overweight and obesity.

## Materials and methods

### Study design and subjects

A population-based, cross-sectional study was conducted in the municipality of Comitán de Domínguez, Chiapas, from June 2010 to June 2012. A census track of three urban (Jerusalem, El Cedro and Cerrito de Concepción) and five rural areas (Santa Rosalía, Zaragoza de la Montaña, San



Source: reference 7

**FIGURE 1. SCHEME OF SOCIAL DETERMINANTS FOR OVERWEIGHT AND OBESITY**

José, La Floresta and Yalumá) chosen by random and convenience sampling, respectively, was performed. All eligible indigenous (IND) persons from the rural areas were invited to participate, whereas non-indigenous (NIND) persons were chosen randomly (N=2 500). People were considered of IND origin either by self-report or the speaking of an IND language. Of 1 940 potential participants (response rate 77.6%), 1 858 individuals aged >20 years (885 from IND origin, mainly from the Tzeltal and Tojolabal ethnic groups, and 973 from NIND origin) had complete anthropometric measurements. All eligible individuals were visited at their homes and invited to participate in the study. Those who accepted were examined at the health care center (persons from rural areas) or at the Comitán General Hospital (persons from urban areas). This study was carried out in accordance with the STROBE guidelines (Strengthening the Reporting of Observational studies in Epidemiology) for cross-sectional studies and with the Declaration of Helsinki (2000). The committees of Research, Ethics, and Biosecurity of the National Institute of Public Health (*Instituto Nacional de Salud Pública*, INSP) approved the study protocol. All participants who agreed to participate signed an informed consent form.

### Anthropometric measurements

Anthropometry was performed by trained and standardized personnel (intra- and interclass correlation  $\geq 85$ ). Measurements were made by triplicate with the participants using light clothing and no shoes. Weight, height, waist circumference (WC), mid-upper arm circumference (MUAC), and four skinfolds (bicipital, tricipital, subscapular, and suprailiac) were measured. Body mass index (BMI) was calculated as weight (kg) / height<sup>2</sup> (m) and classified according to the WHO into lightweight (<18.9), normal (18.9 to 24.9), overweight (25.0-29.9), and obesity ( $\geq 30$ ).<sup>9</sup> Abdominal obesity was defined according to the criteria of the International Diabetes Federation for Latin American countries as WC  $\geq 94$  cm in men and  $\geq 88$  cm in women.<sup>10</sup> Body composition was defined as the percentage of fat through the sum of the four skinfolds using the Durnin and Womersley body density equation and by using the Siri's formula.<sup>11,12</sup> The percentage of fat-free mass and fat mass was obtained. The percentage of fat mass was classified by sex (men and women) into normal (12 to 20% and 20 to 30%, respectively), high (21 to 25% and 31 to 33%, respectively), and very high (>25 and >33%, respectively).

### Social determinants of health

Marmot classifies the social determinants into proximal, intermediate and structural.<sup>7</sup> As proximal determinants,

we included age, sex, and self-report of indigenous origin. Also diet information was obtained through a semiquantitative questionnaire on food consumption frequency (FFQ) designed with the methodology of Walter Willett.<sup>13</sup> Both calorie intake (kcal) and percentage of carbohydrates, proteins and lipids included in the diet were calculated. Physical activity was measured using the International Questionnaire for Physical Activity (IPAQ, short version) that measures the frequency (days a week) and intensity (hours and minutes) of the activities carried out by a person; then the metabolic equivalents (METs) were calculated. The level of physical activity (by METs) was classified into inactive, mild, moderate, and vigorous.<sup>14</sup> The comorbidities included were diabetes (fasting glucose  $\geq 126$  or 2-hr post-load glucose  $\geq 200$  mg/dL or previous medical diagnosis) and hypertension (systolic blood pressure >140 mm/Hg or diastolic blood pressure >90 mm/Hg or prior medical diagnosis). As for intermediate determinants, education level was categorized into <primary and  $\geq$ primary school. Occupation was classified into housewife, farmer, trader, and other. The place of residence was classified into urban ( $\geq 2 500$  inhabitants) and rural (<2 500 inhabitants) areas.<sup>15</sup> For structural determinants, residence areas were classified as having low, medium, high, and very high degree of marginalization according to criteria of Mexico's National Council of Population (*Consejo Nacional de Población*, Conapo), which take into account education level, type of housing, type of residence area, overcrowding, and economic wealth.<sup>15</sup> Social exclusion was classified as low, medium, high, and very high in accordance with the criteria of the National Council for the Evaluation of Social Development Policy (*Consejo Nacional de Evaluación de la Política de Desarrollo Social*, Coneval), which consider education level, type of housing, household goods, and affiliation to social security.<sup>16</sup>

### Statistical analysis

Comparisons of some proximal, intermediate, and structural determinants were done by age groups (20-29, 30-39 and  $\geq 40$  years) and sex. Also, comparisons of the distribution and composition of body fat by age, sex, IND origin, area of residence, and marginalization degree were carried out. For categorical variables  $\chi^2$  was used; for continuous variables *t*-student test, analysis of variance (Anova) or Kruskal-Wallis test were used when appropriate. The age- and sex-adjusted prevalence and 95% confidence intervals (95%CI) of overweight and obesity according to some social determinants were estimated through multiple logistic regression. The association between overweight and obesity and determinants such as age (continuous), sex (women/men),

self-reported IND origin (no/yes), education level ( $\geq$ primary and  $<$ primary school), and degree of marginalization (low/medium and high/very high) among other covariables were evaluated using multinomial logistic regression. Finally, models were made stratifying by self-reported IND origin and adjusting for the variables mentioned above. The fit of the models was evaluated through  $\chi^2$  goodness of fit test (Hosmer-Lemeshow test) and by evaluating the influence statistics and outliers. All analyses were performed with Stata/MP 15.1 (Stata Corporation, College Station, TX, USA).

## Results

### Description of the study population

A total of 1 858 individuals participated in the study (644 men and 1 214 women, with a mean age of 42.2 years [ $\pm$ 15.2 years]). By sex, the mean age in men was 43.8 ( $\pm$ 0.6) years and in women, 41.4 ( $\pm$ 0.4) years. 47% of participants self-reported IND origin (46.7% spoke their native language, Tojolabal, in addition to Spanish), 49.7% lived in rural areas, 58.8% lived in localities with a high/very high degree of marginalization and 33.9% in localities with high/very high social exclusion. Differences were found between the percentage of men and women who self-reported IND origin (55.8 and 43.3%, respectively), lived in rural areas (57.8 and 45.9%, respectively), and resided in localities with high/very high marginalization (66.0 and 54.9%, respectively) or high/very high social exclusion (44.1 and 28.5%, respectively).

As for the comparison of proximal determinants by age and sex, although calorie intake was significantly lower in women than in men, the mean percentage of fat intake was higher in women (27.5%) than in men (26.8%) and in both cases it was above the recommended percentage (25%). On the other hand, a higher proportion of women (49.9%) had a lower level of physical activity compared with men (46.6%). Also, significant differences were observed between men and women by level of education, area of residence, and degree of marginalization and social exclusion. In comparisons by age, the percentage of  $\geq$ primary school was higher in the groups of age 20-29 years (87.6%) and 30-39 years (82.9%), whereas in the age group  $\geq$ 40 years 55.5% had  $\geq$ primary and 44.5%  $<$ primary school (table I).

Regarding distribution and composition of body fat, BMI was significantly higher in women than in men and increased with age. It was also higher in those who self-reported NIND origin and those who resided in urban areas and in localities with medium degree of marginalization. The results were similar for WC, MUAC, and waist-to-height ratio. For the percentage of body fat, the

mean was 21.4% in men and 33.6% in women, which was higher than the recommended value ( $<$ 20 and  $<$ 30%, respectively). In both men and women, the percentage of fat mass increased with age. In men aged  $\geq$ 40 years the mean percentage was 23.6% higher than the recommended level ( $<$ 30%). As for women, the mean percentage was 29.5% in the 20-29 years-old group, 32.3% in the 30-39 years-old group, and 36.2% in individuals aged  $\geq$ 40 years. The percentages were greater than the recommended for all age groups ( $<$ 24,  $<$ 27, and  $<$ 30%, respectively) (table II).

### Prevalence of overweight and obesity and social determinants

The age- and sex-adjusted prevalence of O/O was 37.9% (95%CI 35.5-40.3) and 16.5% (95%CI 14.5-18.5), respectively. The prevalence was higher in localities with medium degree of marginalization (45.9% [95%CI 42.1-49.7] and 25.5% [95%CI 22.1-28.9], respectively) than in localities with high/very high marginalization (32.7% [95%CI 29.8-35.5] and 10.6% [95%CI 8.8-12.5], respectively). The results were similar for self-reported IND origin, area of residence, and degree of social exclusion (table III).

In a multinomial regression model, after adjustment for age, physical activity, and lipid intake, the probability of overweight and obesity was higher in women, in persons with  $\geq$ primary school and in those of NIND origin. After stratification by self-report of IND origin, the probability of overweight in the NIND participants was 1.42 (95%CI 1.01-1.99) times higher in women than in men and 2.06 (95%CI 1.36-3.13) times higher in individuals with  $\geq$ primary school than in those with  $<$ primary. The persons of localities with medium degree of marginalization were also more likely to be overweight (OR=2.25, 95%CI 1.61-3.14) compared with those of localities with high/very high marginalization. In the IND population only the medium degree of marginalization (OR=2.79, 95%CI 1.87-4.16) was associated with the probability of overweight.

As for obesity, the probability of being obese was greater in women, in individuals with  $\geq$ primary school and in those of NIND origin. Low physical activity and greater intake of lipids were also associated with obesity. In both IND and NIND participants the probability of obesity was higher in women than in men (OR=2.79, 95%CI 1.84-4.23 and OR=2.64, 95%CI 1.56-4.47, respectively). It was also higher in dwellers of localities with medium degree of marginalization than in those of localities with high/very high marginalization in both IND and NIND population persons (OR=2.83, 95%CI 1.94-4.14 and OR=5.79, 95%CI 3.42-9.78, respectively) (table IV).

**Table I**  
**DESCRIPTION OF THE STUDY POPULATION ACCORDING TO PROXIMAL, INTERMEDIATE, AND STRUCTURAL DETERMINANTS BY AGE AND SEX. THE COMITÁN STUDY, CHIAPAS, MEXICO, 2010-2012**

|                                      | Sex                          |   | Age group                    |                              |                              |
|--------------------------------------|------------------------------|---|------------------------------|------------------------------|------------------------------|
|                                      | Men<br>n =644                | Women<br>n =1 214                       | 20-29 yrs<br>n =426          | 30-39 yrs<br>n =475          | ≥40 yrs<br>n =957            |
| Proximal determinants                |                              |   |                              |                              |                              |
| Self-report of indigenous origin*    |                              |   |                              |                              |                              |
| Yes                                  | 359 (55.8)                   | 526 (43.3)                              | 207 (48.6)                   | 223 (46.9)                   | 455 (47.5)                   |
| No                                   | 285 (44.2)                   | 688 (56.7) <sup>#</sup>                 | 219 (51.4)                   | 252 (53.1)                   | 502 (52.5)                   |
| Speaking of any indigenous language* |                              |   |                              |                              |                              |
| Yes                                  | 177 (49.3)                   | 237 (45.1)                              | 44 (21.3)                    | 88 (39.5)                    | 282 (62.0) <sup>#</sup>      |
| No                                   | 182 (50.7)                   | 289 (54.9)                              | 163 (78.7)                   | 135 (60.5)                   | 173 (38.0)                   |
| Calorie intake (kcal) <sup>‡</sup>   | 1 777.4<br>(1 511.3-2 083.8) | 1 723<br>(1 461.5-2 074.9) <sup>#</sup> | 1 771.0<br>(1 492.1-2 080.0) | 1 739.2<br>(1 479.1-2 080.9) | 1 726.4<br>(1 471.6-2 043.7) |
| Macronutrient intake <sup>§#</sup>   |                              |   |                              |                              |                              |
| Carbohydrates %                      | 56.7 (0.2)                   | 56.4 (0.2)                              | 56.5 (6.3)                   | 56.2 (6.4)                   | 56.7 (6.2)                   |
| Proteins %                           | 17.0 (0.1)                   | 17.0 (0.1)                              | 16.9 (2.2)                   | 17.0 (2.0)                   | 17.1 (2.1)                   |
| Lipids %                             | 26.8 (0.2)                   | 27.5 (0.1) <sup>#</sup>                 | 27.4 (5.1)                   | 27.6 (5.0)                   | 27.0 (4.9)                   |
| METs <sup>‡</sup>                    | 1 638 (819-2 274)            | 996 (699-1 794) <sup>#</sup>            | 1 257 (769-2 019)            | 1 158 (786-2 076)            | 1 116 (720-2 034)            |
| Level of physical activity*          |                              |   |                              |                              |                              |
| Inactive/mild                        | 280 (46.6)                   | 588 (49.9) <sup>#</sup>                 | 177 (43.6)                   | 224 (48.9)                   | 467 (51.0) <sup>#</sup>      |
| Moderate                             | 69 (11.5)                    | 196 (16.6)                              | 75 (18.5)                    | 58 (12.7)                    | 132 (14.4)                   |
| Vigorous                             | 252 (41.9)                   | 394 (33.4)                              | 154 (37.9)                   | 176 (38.4)                   | 316 (34.5)                   |
| Comorbidities*                       |                              |   |                              |                              |                              |
| Diabetes                             | 47 (7.3)                     | 112 (9.2)                               | 10 (2.3)                     | 18 (3.8)                     | 131 (13.7) <sup>#</sup>      |
| Hypertension                         | 124 (19.2)                   | 238 (19.6)                              | 37 (8.7)                     | 54 (11.4)                    | 271 (28.3) <sup>#</sup>      |
| Intermediate determinants            |                              |   |                              |                              |                              |
| Occupation*                          |                              |   |                              |                              |                              |
| Housewife                            | 16 (2.5)                     | 1002 (82.5) <sup>#</sup>                | 244 (57.3)                   | 264 (55.6)                   | 510 (53.3) <sup>#</sup>      |
| Farmer                               | 374 (58.1)                   | 21 (1.7)                                | 62 (14.5)                    | 97 (20.4)                    | 236 (24.7)                   |
| Trader                               | 50 (7.8)                     | 80 (6.6)                                | 27 (6.3)                     | 31 (6.5)                     | 72 (7.5)                     |
| Other                                | 204 (31.7)                   | 111 (9.1)                               | 93 (21.8)                    | 83 (17.5)                    | 139 (14.5)                   |
| Education level*                     |                              |   |                              |                              |                              |
| < Primary                            | 129 (20.0)                   | 431 (35.5) <sup>#</sup>                 | 53 (12.4)                    | 81 (17.0)                    | 426 (44.5) <sup>#</sup>      |
| ≥ Primary                            | 515 (80.0)                   | 783 (64.5)                              | 373 (87.6)                   | 394 (82.9)                   | 531 (55.5)                   |
| Area of residence*                   |                              |   |                              |                              |                              |
| Urban                                | 272 (42.2)                   | 663 (54.6) <sup>#</sup>                 | 191 (44.8)                   | 232 (48.8)                   | 512 (53.5) <sup>#</sup>      |
| Rural                                | 372 (57.8)                   | 551 (45.4)                              | 235 (55.1)                   | 243 (51.2)                   | 445 (46.5)                   |
| Structural determinants              |                              |   |                              |                              |                              |
| Degree of marginalization*           |                              |   |                              |                              |                              |
| Medium                               | 219 (34.0)                   | 547 (45.1) <sup>#</sup>                 | 164 (38.5)                   | 190 (40.0)                   | 412 (43.0)                   |
| High/very high                       | 425 (66.0)                   | 667 (54.9)                              | 262 (61.5)                   | 285 (60.0)                   | 545 (56.9)                   |
| Degree of social exclusion*          |                              |   |                              |                              |                              |
| Medium                               | 360 (55.9)                   | 868 (71.5) <sup>#</sup>                 | 265 (62.2)                   | 314 (66.1)                   | 649 (67.8)                   |
| High/very high                       | 284 (44.1)                   | 346 (28.5)                              | 161 (37.8)                   | 161 (33.9)                   | 308 (32.2)                   |

\* Number of subjects (%).

‡ Median and percentiles 25-75

§ Mean and standard deviation. X<sup>2</sup> was used for comparison of proportions. For comparison of mean between two groups t-Student test was used and one-way ANOVA for more than two groups. For continuous variables with non-normal Kruskal Wallis was used.<sup>#</sup> Significant differences between groups (p<.05).



**Table II**  
**DISTRIBUTION AND COMPOSITION OF BODY FAT ACCORDING TO PROXIMAL, INTERMEDIATE, AND STRUCTURAL DETERMINANTS.**  
**THE COMITÁN STUDY, CHIAPAS, MEXICO, 2010-2012**

|                              | Sex            |                          | Age group            |                      |                          |               |                          | Self-report of indigenous origin |                          |                       | Area of residence        |  |  | Marginalization degree |
|------------------------------|----------------|--------------------------|----------------------|----------------------|--------------------------|---------------|--------------------------|----------------------------------|--------------------------|-----------------------|--------------------------|--|--|------------------------|
|                              | Men<br>n = 644 | Women<br>n = 1214        | 20-29 yrs<br>n = 426 | 30-39 yrs<br>n = 475 | >40 yrs<br>n = 957       | No<br>n = 973 | Yes<br>n = 885           | Urban<br>n = 935                 | Rural<br>n = 923         | Low/medium<br>n = 766 | Highly high<br>n = 1 092 |  |  |                        |
| BMI*                         | 25.4 (4.0)     | 27.0 (4.7) <sup>§</sup>  | 24.8 (3.9)           | 26.9 (4.5)           | 26.9 (4.6) <sup>§</sup>  | 27.3 (4.8)    | 25.4 (4.0) <sup>§</sup>  | 28.1 (4.6)                       | 24.7 (3.7) <sup>§</sup>  | 28.1 (4.6)            | 25.2 (4.0) <sup>§</sup>  |  |  |                        |
| Normal <sup>‡</sup>          | 332 (51.5)     | 471 (38.8) <sup>§</sup>  | 266 (62.4)           | 188 (39.6)           | 349 (36.5) <sup>§</sup>  | 350 (36.0)    | 453 (51.2) <sup>§</sup>  | 251 (26.8)                       | 552 (59.8) <sup>§</sup>  | 203 (26.5)            | 600 (54.9) <sup>§</sup>  |  |  |                        |
| Overweight <sup>‡</sup>      | 242 (37.6)     | 449 (37.0)               | 120 (28.2)           | 175 (36.8)           | 396 (41.4)               | 370 (38.0)    | 321 (36.3)               | 405 (43.3)                       | 286 (31.0)               | 335 (43.7)            | 356 (32.6)               |  |  |                        |
| Obesity <sup>‡</sup>         | 70 (10.9)      | 294 (24.2)               | 40 (9.4)             | 112 (23.6)           | 212 (22.1)               | 253 (26.0)    | 111 (12.5)               | 279 (29.8)                       | 85 (9.2)                 | 228 (29.8)            | 136 (12.4)               |  |  |                        |
| MUJAC* (cm)                  | 29.6 (3.3)     | 29.9 (3.6) <sup>§</sup>  | -                    | -                    | -                        | -             | -                        | -                                | -                        | -                     | -                        |  |  |                        |
| Men                          | -              | -                        | 29.5 (2.9)           | 30.2 (3.2)           | 29.4 (3.5) <sup>§</sup>  | 30.4 (3.5)    | 29.0 (3.0) <sup>§</sup>  | 30.9 (3.6)                       | 28.6 (2.8) <sup>§</sup>  | 31.2 (3.6)            | 28.8 (2.9) <sup>§</sup>  |  |  |                        |
| Women                        | -              | -                        | 28.7 (3.0)           | 30.6 (3.6)           | 30.2 (3.8) <sup>§</sup>  | 30.6 (3.7)    | 29.1 (3.4) <sup>§</sup>  | 31.1 (3.6)                       | 28.6 (3.3) <sup>§</sup>  | 30.9 (3.6)            | 29.1 (3.5) <sup>§</sup>  |  |  |                        |
| WC (cm)*                     | 89.2 (37.2)    | 89.9 (10.9)              | -                    | -                    | -                        | -             | -                        | -                                | -                        | -                     | -                        |  |  |                        |
| Men                          | -              | -                        | 83.2 (10.3)          | 87.2 (9.3)           | 92.6 (49.8) <sup>§</sup> | 90.0 (10.6)   | 88.6 (48.9)              | 95.2 (55.9)                      | 84.8 (8.0)               | 96.5 (62.2)           | 85.5 (8.2)               |  |  |                        |
| Women                        | -              | -                        | 83.5 (8.9)           | 89.9 (10.3)          | 93.0 (10.7) <sup>§</sup> | 91.0 (11.2)   | 88.5 (10.3) <sup>§</sup> | 91.8 (11.0)                      | 87.7 (10.3) <sup>§</sup> | 90.8 (11.0)           | 89.2 (10.7) <sup>§</sup> |  |  |                        |
| Waist-to-height ratio*       | 0.55 (0.05)    | 0.61 (0.00) <sup>§</sup> | -                    | -                    | -                        | -             | -                        | -                                | -                        | -                     | -                        |  |  |                        |
| Men                          | -              | -                        | 0.52 (0.06)          | 0.55 (0.05)          | 0.57 (0.05) <sup>§</sup> | 0.56 (0.00)   | 0.54 (0.00) <sup>§</sup> | 0.57 (0.06)                      | 0.53 (0.05) <sup>§</sup> | 0.58 (0.06)           | 0.54 (0.05) <sup>§</sup> |  |  |                        |
| Women                        | -              | -                        | 0.56 (0.06)          | 0.61 (0.07)          | 0.64 (0.07) <sup>§</sup> | 0.62 (0.07)   | 0.60 (0.07) <sup>§</sup> | 0.62 (0.7)                       | 0.60 (0.07) <sup>§</sup> | 0.62 (0.07)           | 0.61 (0.07)              |  |  |                        |
| Percentage of fat mass*      | 21.4 (0.3)     | 33.6 (0.2) <sup>§</sup>  | -                    | -                    | -                        | -             | -                        | -                                | -                        | -                     | -                        |  |  |                        |
| Men                          | -              | -                        | 16.6 (4.7)           | 20.6 (4.0)           | 23.6 (7.3) <sup>§</sup>  | 22.5 (6.3)    | 20.5 (6.8) <sup>§</sup>  | 23.7 (7.7)                       | 19.6 (5.2) <sup>§</sup>  | 24.2 (8.1)            | 19.9 (5.2) <sup>§</sup>  |  |  |                        |
| Women                        | -              | -                        | 29.5 (6.5)           | 32.3 (4.7)           | 36.2 (5.6) <sup>§</sup>  | 34.5 (5.9)    | 32.5 (6.6) <sup>§</sup>  | 35.2 (5.5)                       | 31.6 (6.6) <sup>§</sup>  | 35.3 (5.6)            | 32.2 (6.4) <sup>§</sup>  |  |  |                        |
| Percentage of fat-free mass* | 78.6 (0.3)     | 66.4 (0.2) <sup>§</sup>  | -                    | -                    | -                        | -             | -                        | -                                | -                        | -                     | -                        |  |  |                        |
| Men                          | -              | -                        | 83.3 (4.7)           | 79.4 (4.0)           | 76.3 (7.3) <sup>§</sup>  | 77.5 (6.3)    | 79.5 (6.9) <sup>§</sup>  | 76.3 (7.7)                       | 80.3 (5.2) <sup>§</sup>  | 75.7 (8.2)            | 80.1 (5.2) <sup>§</sup>  |  |  |                        |
| Women                        | -              | -                        | 70.5 (6.5)           | 67.7 (4.7)           | 63.8 (5.6) <sup>§</sup>  | 65.5 (5.9)    | 67.5 (6.6) <sup>§</sup>  | 64.8 (5.5)                       | 68.3 (6.7) <sup>§</sup>  | 64.7 (5.6)            | 67.8 (6.4) <sup>§</sup>  |  |  |                        |

\* Mean and standard deviation.

<sup>‡</sup> Number of subjects (%).

<sup>§</sup> Significant differences between groups ( $p < 0.05$ ).

MUJAC: Mid-upper arm circumference

X<sup>2</sup> was used for comparison of proportions. For comparison of medians between two groups t-Student test was used for independent samples and ANOVA for more than two groups.

**Table III**  
**AGE- AND SEX-ADJUSTED PREVALENCE OF OVERWEIGHT AND OBESITY ACCORDING TO SOME INTERMEDIATE AND STRUCTURAL DETERMINANTS IN ADULT POPULATION. COMITÁN STUDY, CHIAPAS, MEXICO, 2010-2012**

|                                  | Overweight          |                          |         | Obesity             |                          |         |
|----------------------------------|---------------------|--------------------------|---------|---------------------|--------------------------|---------|
|                                  | Prevalence          | Prevalence ratio (95%CI) | p value | Prevalence          | Prevalence ratio (95%CI) | p value |
| Intermediate determinants        |                     |                          |         |                     |                          |         |
| Self-report of indigenous origin |                     |                          |         |                     |                          |         |
| Yes                              | 36.30 (33.06-39.53) | 1 (Reference)            |         | 10.85 (8.78-12.92)  | 1 (Reference)            |         |
| No                               | 39.47 (36.18-42.76) | 1.09 (0.97-1.23)         | <.001   | 22.07 (19.18-24.96) | 2.03 (1.67-2.56)         | <.001   |
| Level of education               |                     |                          |         |                     |                          |         |
| <primary                         | 29.10 (24.89-33.32) | 1 (Reference)            |         | 13.38 (10.40-16.36) | 1 (Reference)            |         |
| >primary                         | 41.27 (38.42-44.13) | 1.42 (1.19-1.65)         | <.001   | 17.67 (15.38-19.95) | 1.32 (0.99-1.64)         | .016    |
| Area of residence                |                     |                          |         |                     |                          |         |
| Rural                            | 31.03 (27.99-34.06) | 1 (Reference)            |         | 7.93 (6.19-9.66)    | 1 (Reference)            |         |
| Urban                            | 45.30 (41.84-48.75) | 1.46 (1.30-1.67)         | <.001   | 25.69 (22.54-28.83) | 3.24 (2.63-4.17)         | <.001   |
| Structural determinants          |                     |                          |         |                     |                          |         |
| Degree of marginalization        |                     |                          |         |                     |                          |         |
| High/very high                   | 32.70 (29.85-35.55) | 1 (Reference)            |         | 10.65 (8.77-12.52)  | 1 (Reference)            |         |
| Medium                           | 45.91 (42.13-49.70) | 1.40 (1.26-1.59)         | <.001   | 25.51 (22.13-28.90) | 2.39 (2.0-3.03)          | <.001   |
| Degree of social exclusion       |                     |                          |         |                     |                          |         |
| High/very high                   | 29.98 (26.37-33.59) | 1 (Reference)            |         | 6.22 (4.37-8.06)    | 1 (Reference)            |         |
| Medium                           | 42.41 (39.40-45.42) | 1.41 (1.25-1.64)         | <.001   | 22.35 (19.68-25.02) | 3.59 (2.78-5.26)         | <.001   |
| Age-and sex-adjusted prevalence  | 37.90 (35.54-40.27) | -                        | -       | 16.50 (14.55-18.44) | -                        | -       |

## Discussion

Overweight and obesity is a complex condition influenced by the interaction of genetics and environmental risk factors. However, genetic factors only account for 40% of variations in body mass index, so the social environment plays an important role in overweight and obesity. The prevalence of O/O found in this study was 39.5 and 22.1%, respectively, in NIND participants and 36.3 and 10.8%, respectively, in IND participants. When comparing with other studies carried out in Mexico in IND population, we observed a lower prevalence of O/O than that reported in Nahua of Ixtaczoquitlán, Veracruz (41 and 36.5%, respectively),<sup>17</sup> Nahua of the Sierra of Puebla (44.1 and 19.2%, respectively),<sup>18</sup> Maya of Yucatán (11 and 80.3%, respectively),<sup>19</sup> and Triquis of San Juan Copala, Oaxaca (85.5% combined).<sup>20</sup> In a wider scope, in our study, the prevalence of obesity in IND population was lower than that in aboriginal Australian (38.4%) and Canadian population (36.4%), whereas the prevalence of obesity in NIND population was similar to that in non-aboriginal Australian (22.3%) and Canadian population (22.6%).<sup>21,22</sup>

Women are a vulnerable group due to inequalities in education, recreational activities, and food access in comparison with men.<sup>7</sup> In our study, the prevalence of overweight was similar in both sexes, but obesity was twice as high in women as in men. Both prevalence rates were lower than those reported in other populations with high degree of marginalization.<sup>18,20</sup> Women also had a high-fat diet and low physical activity compared with men, with implications for O/O risk. In a study including Mayan Chontal IND population, a higher prevalence of O/O was found in women than in men. In this population physical exercising is prohibited for women. After maternity women assume that having O/O is normal; obesity is considered as an inevitable legacy of parents, not a disease. Thus, weight loss in women is not desirable at any stage of life.<sup>23</sup>

Regarding area of residence, the probability of overweight was similar in both IND and NIND individuals living in localities with medium degree of marginalization compared with those of localities with high/very high marginalization. On the other hand, for localities with medium degree of marginalization, the risk of O/O was higher in IND than in NIND persons

**Table IV**  
**PROXIMAL, INTERMEDIATE, AND STRUCTURAL DETERMINANTS ASSOCIATED WITH OVERWEIGHT AND OBESITY IN ADULT POPULATION. THE COMITÁN STUDY, CHIAPAS, MEXICO, 2010-2012**

|                                   | Whole sample          |         | Indigenous origin      |         |                    |         |
|-----------------------------------|-----------------------|---------|------------------------|---------|--------------------|---------|
|                                   | N=1 858<br>OR (95%CI) | p value | Non-indigenous, n= 973 |         | Indigenous, n= 885 |         |
|                                   |                       |         | OR (95%CI)             | p value | OR (95%CI)         | p value |
| <b>Overweight</b>                 |                       |         |                        |         |                    |         |
| Age (years)                       | 1.02 (1.01-1.03)      | <.001   | 1.03 (1.01-1.04)       | <.001   | 1.01 (1.00-1.03)   | .003    |
| Sex (women)                       | 1.40 (1.12-1.76)      | .003    | 1.42 (1.01-1.99)       | .04     | 1.23 (0.89-1.70)   | .20     |
| Non-indigenous origin             | 1.25 (1.00-1.56)      | .05     | -                      | -       | -                  | -       |
| Schooling >primary                | 1.81 (1.39-2.35)      | <.001   | 2.06 (1.36-3.13)       | .001    | 1.20 (0.84-1.72)   | .31     |
| <b>Level of physical activity</b> |                       |         |                        |         |                    |         |
| Inactive/mild                     | 1 (Reference)         | -       | 1 (Reference)          | -       | 1 (Reference)      | -       |
| Moderate                          | 1.03 (0.76-1.39)      | .85     | 0.94 (0.62-1.44)       | .79     | 1.36 (0.86-2.16)   | .18     |
| Vigorous                          | 0.73 (0.53-1.01)      | .06     | 0.59 (0.37-0.95)       | .03     | 1.06 (0.66-1.70)   | .81     |
| Intake of lipids >25%             | 1.22 (0.97-1.53)      | .08     | 1.09 (0.76-1.58)       | .62     | 0.84 (0.61-1.17)   | .31     |
| Medium degree of marginalization  | -                     | -       | 2.25 (1.61-3.14)       | <.001   | 2.79 (1.87-4.16)   | <.001   |
| <b>Obesity</b>                    |                       |         |                        |         |                    |         |
| Age (years)                       | 1.02 (1.01-1.03)      | <.001   | 1.03 (1.01-1.04)       | .001    | 1.01 (0.99-1.03)   | .16     |
| Sex (women)                       | 2.91 (2.12-3.99)      | <.001   | 2.79 (1.84-4.23)       | <.001   | 2.64 (1.56-4.47)   | <.001   |
| Non-indigenous origin             | 2.27 (1.69-3.03)      | <.001   | -                      | -       | -                  | -       |
| Schooling >primary                | 1.41 (1.03-1.94)      | .03     | 1.28 (0.82-1.99)       | .28     | 0.95 (0.57-1.60)   | .86     |
| <b>Level of physical activity</b> |                       |         |                        |         |                    |         |
| Inactive/mild                     | 1 (Reference)         | -       | 1 (Reference)          | -       | 1 (Reference)      | -       |
| Moderate                          | 0.81 (0.57-1.15)      | .24     | 0.68 (0.43-1.06)       | .09     | 1.67 (0.84-3.29)   | .14     |
| Vigorous                          | 0.68 (0.46-1.01)      | .05     | 0.50 (0.30-0.84)       | .01     | 1.61 (0.79-3.28)   | .19     |
| Intake of lipids >25%             | 1.61 (1.20-2.16)      | .001    | 1.34 (0.88-2.05)       | .17     | 0.87 (0.53-1.42)   | .59     |
| Medium degree of marginalization  | -                     | -       | 2.83 (1.94-4.14)       | <.001   | 5.79 (3.42-9.78)   | <.001   |

Note: A multiple multinomial regression model was done. Biologically relevant variables with  $p < .20$  by simple regression were added to the model.

(OR=2.8 and OR=5.8, respectively). In the present study, nearly 50% of the participants self-reported IND origin. Most of this population are farmers and breed the animals that are part of their diet. Mexico has the largest number of people of IND origin in Latin America, totaling 11 million, of which 2.7 million have limited access to public services, education, and health care.<sup>24</sup> In Chiapas, the percentage of IND population (27.9%) is four times greater than the national average (6.7%).<sup>25</sup> A higher percentage of the IND population lives in rural areas where they have few social, economic, and political opportunities to develop. This coupled with the abandonment of public investment and little subsidy for rural activities has forced the IND population to migrate to big cities. For example, during the decade of 1970s agricultural production accounted for 12% of Mexico's GDP, while by 2014 it had decreased to 3.3%.<sup>26</sup>

In our study, the prevalence of O/O was higher in urban than in rural areas. O/O is spreading rapidly in the urban zones of developing countries. Popkin and colleagues compared the prevalence of O/O in 42 countries of Asia, Middle East, Africa, and Latin America and found a higher prevalence in urban (37.2%) than in rural localities (19%).<sup>27</sup> According to the Ensanut 2000 and the 2016 Ensanut-MC national surveys, O/O prevalence in rural areas increased 7.2 and 7.1%, respectively, whereas in urban locations it decreased 2.6% for overweight and increased 8.1% for obesity.<sup>2</sup>

The high risk of O/O for IND people living in urban areas is linked to changes in lifestyle (food and diet) typical of cities.<sup>28</sup> A study comparing the eating patterns of Pima Indians from Phoenix, Arizona, in the United States of America (urban area residents) and the Pima from Maycoba, Sonora, in Mexico (rural dwellers), found



that the diet in the second group was higher in fiber and lower in fat,<sup>29</sup> whereas the first group had a diet higher in saturated fat and lower in fiber. Thus, the prevalence of obesity was five times higher in the Pima from Arizona (69%) than in the Pima from Sonora (13%).<sup>30</sup>

Populations with lower degree of marginalization are at more risk of O/O compared with those with higher marginalization.<sup>31</sup> In Mexico several studies show that globalization has led to a nutritional transition that has conditioned in part the increase in O/O. In past decades meals consisted of little processed foods and most foods were grown by people for their own consumption.<sup>5</sup> More recently, worldwide advances in communication, infrastructure, and technology have given way to access to a great variety of processed foods with high-calorie content and low nutritional contribution.<sup>27</sup> The analysis of the Food Balance Sheets of the Food and Agriculture Organization (FAO) shows that in Mexico, from 1961 to 2013, the kcal/day consumption per person rose from 2 401 to 3 049 kcal.<sup>32</sup>

Eating patterns are also related to socioeconomic status and area of residence; this however does not guarantee an adequate food choice of the most favored people.<sup>28,29</sup> Unfortunately, in poor rural areas soft-drinks and ultra-processed foods are increasingly consumed because accessibility is growing.<sup>33</sup> Another key factor is physical activity, which has decreased in less marginalized areas as a result of less walking and more use of car or public transport. Insecurity in large cities has also played a role in the decrease of outdoor activities. Likewise, technological tools have replaced physical labor resulting in a greater number of jobs that favor O/O.<sup>33</sup>

Among the strengths of our study was the number of individuals enrolled, which helped us to analyze some social determinants and lifestyle risk factors. Although the nonresponse rate was moderate because of similar reasons in both people with medium marginalization and those with high/very high marginalization, no differences in social determinants between respondents and nonrespondents were found, which reduces the probability of selection bias. Due to the smaller proportion of IND individuals compared with NIND ones, an oversampling of IND persons was done, which could affect the external validity. However, after comparison of people who accepted to participate in the study vs. those who did not accept, no significant differences were observed. On the other hand, because of the definition of IND origin we used, the probability of misclassification bias exists. Yet, self-reported IND origin is more related to social and cultural factors than to biological or genetic aspects, the former being of more interest for the purpose of this study. Also, the cross-sectional design of the study made it difficult to

determine the impact of changes on food intake and physical activity as well as on social determinants related to the prevalence of O/O.

In summary, we consider that public policies aimed to reduce O/O should first be tailored to the particular social environment, ethnicity, socioeconomic status, availability of food, and social and food culture of targeted populations in order to break access barriers and facilitate healthy food choices.

## Acknowledgements

The authors thank the authorities of the General Hospital in Comitán and Health Centers in the participating locations for their support for the study, and the National Institute of Public Health in Mexico for their technical and administrative support. They are also grateful to all those persons who agreed to participate in the study.

## Funding

The Comitán Study was supported by Mexico's National Council of Science and Technology, Conacyt (Grant 87270) and by the National Institute of Public Health.

*Declaration of conflict of interests.* The authors declare that they have no conflict of interests.

## References

1. James PT, Leach R, Kalamara E, Shayeghi M. The Worldwide Obesity Epidemic. *Obes Res* 2001;9(S11):228S-33S. <https://doi.org/10.1038/oby.2001.123>
2. Shamah-Levy T, Campos-Nonato I, Cuevas-Nasu L, Hernández-Barrera L, Morales-Ruán M del C, Rivera-Dommarco J, et al. Overweight and obesity in Mexican vulnerable population. Results of Ensanut 100k. *Salud Publica Mex*. 2019;61(6):852-65. <https://doi.org/10.21149/10585>
3. González-Muniesa P, Martínez-González MA, Hu FB, Després JP, Matsuzawa Y, Loos RJF, et al. Obesity. *Nat Rev Dis Prim*. 2017;3:17034. <https://doi.org/10.1038/nrdp.2017.34>
4. Rivera JA, Barquera S, Campirano F, Campos I, Safdie M, Tovar V. Epidemiological and nutritional transition in Mexico: rapid increase of non-communicable chronic diseases and obesity. *Public Health Nutr*. 2002;5(1a):113-22. <https://doi.org/10.1079/PHN2001282>
5. Devaux M, Sassi F. Social inequalities in obesity and overweight in 11 OECD countries. *Eur J Public Health*. 2013;23(3):464-9. <https://doi.org/10.1093/eurpub/ckr058>
6. Wells JCK, Marphatia AA, Cole TJ, Mccoy D. Associations of economic and gender inequality with global obesity prevalence: Understanding the female excess. *Soc Sci Med*. 2012;75(3):482-90. <https://doi.org/10.1016/j.socscimed.2012.03.029>
7. Marmot M. Social determinants of health inequalities. *Lancet*. 2005;365(9464):1099-104. [https://doi.org/10.1016/S0140-6736\(05\)71146-6](https://doi.org/10.1016/S0140-6736(05)71146-6)
8. Instituto Nacional de Salud Pública. Encuesta Nacional de Salud y Nutrición. Resultados por entidad federativa, Chiapas 2012. Cuernavaca: INSP,

- 2013 [cited April 2019]. Available from: <https://ensanut.insp.mx/encuestas/ensanut2012/doctos/informes/Chiapas-OCT.pdf>
9. Consejo de Salubridad General. Prevención, Diagnóstico y Tratamiento del Sobrepeso y la Obesidad Exógena. Guías de Práctica Clínica. México (actualización 2012). Mexico City: Secretaría de Salud, 2012 [cited May 2019]. Available from: [http://www.cenetec.salud.gob.mx/descargas/gpc/CatalogoMaestro/046\\_GPC\\_ObesidadAdulto/IMSS\\_046\\_08\\_GRR.pdf](http://www.cenetec.salud.gob.mx/descargas/gpc/CatalogoMaestro/046_GPC_ObesidadAdulto/IMSS_046_08_GRR.pdf)
10. Rosas-Guzmán J, González-Chávez A, Aschner P, Bastarrachea R. Epidemiología, Diagnóstico, Control, Prevención y Tratamiento del Síndrome Metabólico en Adultos. *Revista de la ALAD*. 2010;18(1):25-42 [cited April 2019]. Available from: <http://www.revistaalad.com/pdfs/100125-44.pdf>
11. Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr*. 1974;32(1):77-97. <https://doi.org/10.1079/BJN19740060>
12. Martín-Moreno V, Gómez-Gandoy JB, Antoranz-González MJ. Medición de la grasa corporal mediante impedancia bioeléctrica, pliegues cutáneos y ecuaciones a partir de medidas antropométricas. Análisis comparativo. *Rev Esp Salud Publica*. 2001;75(3):221-36 [cited May 2019]. Available from: <http://www.redalyc.org/articulo.oa?id=17075306>
13. Cade J, Thompson R, Burley V, Warm D. Development, validation and utilisation of food-frequency questionnaires a review. *Public Health Nutr*. 2002;5(4):567-87. <https://doi.org/10.1079/PHN2001318>
14. Toloza SC, Gómez-Conesa AA. El Cuestionario Internacional de Actividad Física. Un instrumento adecuado en el seguimiento de la actividad física poblacional. *Revista Iberoamericana de Fisioterapia y Kinesiología*. 2007;1(10):48-52. [https://doi.org/10.1016/S1138-6045\(07\)73665-1](https://doi.org/10.1016/S1138-6045(07)73665-1)
15. Consejo Nacional de Población. Índice de Marginación por Localidad 2010. Mexico City: Conapo, 2012 [cited April 2019]. Available from: [http://www.conapo.gob.mx/en/CONAPO/Indice\\_de\\_Marginacion\\_por\\_Localidad\\_2010](http://www.conapo.gob.mx/en/CONAPO/Indice_de_Marginacion_por_Localidad_2010)
16. Consejo Nacional de Evaluación de la Política de Desarrollo Social. Índice de Rezago Social 2015 [Internet]. Mexico City: Coneval, 2016 [cited April 2019]. Available from: [https://www.coneval.org.mx/Medicion/IRS/Paginas/Indice\\_Rezago\\_Social\\_2015.aspx](https://www.coneval.org.mx/Medicion/IRS/Paginas/Indice_Rezago_Social_2015.aspx)
17. Herrera-Huerta EV, García-Montalvo EA, Méndez-Bolina E, López-López JG, Valenzuela OL. Sobrepeso y obesidad en indígenas nahuas de Ixtaczoquitlán, Veracruz, México. *Rev Peru Med Exp Salud Publica*. 2012;29:345-9 [cited May 2019]. Available from: <http://www.redalyc.org/articulo.oa?id=36325041008>
18. Macín-Pérez G, Gómez-Valdés JA, Ballesteros-Romero M, Canizales-Quinteros S, Acuña-Alonso V. Cambio cultural, estilo de vida, adiposidad y niveles de glucosa en una comunidad totonaca de la Sierra Norte de Puebla. Cuiculco. 2013;20(58):173-98 [cited May 2019]. Available from: [http://www.scielo.org.mx/scielo.php?script=sci\\_arttext&pid=S0185-16592013000300010](http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0185-16592013000300010)
19. Chaves-Asián R, Pasos-Cervera RA. Overweight and obesity in indigenous Mayan communities. *Economy, Culture and Gender. Revista de Estudios Regionales* 2017;109:139-63 [cited May 2019]. Available from: <http://www.revistaestudiosregionales.com/documentos/articulos/pdf-articulo-2523.pdf>
20. Ramos-Rodríguez RM, Sandoval-Mendoza K. Estado nutricional en la marginación y la pobreza de adultos triquis del estado de Oaxaca. *Pan Am J Public Health*. 2007;22(44):260-7 [cited April 2019]. Available from: [http://www.scielo.org.mx/scielo.php?script=sci\\_arttext&pid=S1020-49892007000900006](http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S1020-49892007000900006)
21. Katzmarzyk-Peter T. Obesity and Physical Activity Among Aboriginal Canadians. *Obesity*. 2008;16(1):184-90. <https://doi.org/10.1038/oby.2007.51>
22. Thurber KA, Joshy G, Korda R, Eades SJ, Wade V, Bambrick H, et al. Obesity and its association with sociodemographic factors, health behaviours and health status among Aboriginal and non-Aboriginal adults in New South Wales, Australia. *J Epidemiol Community Health*. 2018;72(6):491-8. <https://doi.org/10.1136/jech-2017-210064>
23. Cruz-Sánchez M, Tuñon-Pablos E, Villaseñor-Farías M, Álvarez-Gordillo GC, Nigh-Nielsen R. Desigualdades de género en sobrepeso y obesidad entre indígenas chontales de Tabasco, México. *Revista de Población y Salud en Mesoamérica*. 2012;9(2):1-20 [cited June 2019]. Available from: <http://www.redalyc.org/articulo.oa?id=44623231003>
24. Instituto Nacional Indigenista. Comisión Nacional para el Desarrollo de los Pueblos Indígenas 1948-2012. Mexico City: Instituto Nacional Indigenista, 2012 [cited June 2019]. Available from: <http://www.cdi.gob.mx/dmdocuments/ini-cdi-1948-2012.pdf>
25. Aguilar-Ortega T. Desigualdad y marginación en Chiapas. *Revista Península*. 2016;11(2):143-59. <https://doi.org/10.1016/j.pnsla.2015.08.011>
26. Moreno-Altamirano L, García-García JJ, Soto-Estrada G, Capraro S, Limón-Cruz D. Epidemiología y determinantes sociales asociados a la obesidad y la diabetes tipo 2 en México. *Rev Med Hosp Gen Mex*. 2014;77(3):114-23. <https://doi.org/10.1016/j.hgmx.2014.07.002>
27. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev*. 2012;70(1):3-21. <https://doi.org/10.1111/j.1753-4887.2011.00456.x>
28. Drewnowski A, Popkin BM. The Nutrition Transition: New Trends in the Global Diet. *Nutr Rev*. 1997;55(2):31-43. <http://doi.org/10.1111/j.1753-4887.1997.tb01593.x>
29. Smith CJ, Nelson RG, Hardy SA, Manahan EM, Bennett PH, Knowler WC. Survey of the diet of Pima Indians using quantitative food frequency assessment and 24-hour recall. *J Am Diet Assoc*. 1996;96(8):778-84. [https://doi.org/10.1016/s0002-8223\(96\)00216-7](https://doi.org/10.1016/s0002-8223(96)00216-7)
30. Esparza-Romero J, Valencia ME, Urquidez-Romero R, Chaudhari LS, Hanson RL, Knowler WC, et al. Environmentally driven increases in type 2 diabetes and obesity in Pima Indians and Non-Pimas in Mexico over a 15-year period: The Maycoba project. *Diabetes Care*. 2015;38(11):2075-82. <http://doi.org/10.2337/dc15-0089>
31. Apolloni A, Marathe A, Pan Z. A longitudinal view of the relationship between social marginalization and obesity. In: Salerno J, Yang SJ, Nau D, Chai SK (eds). *Social Computing, Behavioral-Cultural Modeling and Prediction*. USA: Springer, Berlin, Heidelberg, 2011. [https://doi.org/10.1007/978-3-642-19656-0\\_10](https://doi.org/10.1007/978-3-642-19656-0_10)
32. Soto-Estrada G, Moreno-Altamirano L, García-García JJ, Ochoa-Moreno I, Silberman M. Trends in frequency of type 2 diabetes in Mexico and its relationship to dietary patterns and contextual factors. *Gac Sanit*. 2018;32(3):283-90. <https://doi.org/10.1016/j.gaceta.2017.08.001>
33. Ewing R, Meakins G, Hamidi S, Nelson AC. Relationship between urban sprawl and physical activity, obesity, and morbidity-Update and refinement. *Health Place*. 2014;26:118-26. <https://doi.org/10.1016/j.healthplace.2013.12.008>