

Julio A. Anduro-Jordan¹

id 0000-0001-8900-622X

Ignacio E. Maldonado-Mendoza²

id 0000-0001-9952-1508

Alejandro Miguel Figueroa-López¹

id 0000-0003-2437-6523

María del Refugio Torres-Vitela³

id 0000-0002-9934-5943

Olga N. Campas-Baypoli¹

id 0000-0003-2240-0884

Dalia Isabel Sánchez-Machado¹

id 0000-0001-5574-9500

Ernesto Uriel Cantú-Soto^{1*}

id 0000-0003-3916-5892

¹Instituto Tecnológico de Sonora,
Departamento de Biotecnología
y Ciencias Alimentarias,
Obregón, México

²Instituto Politécnico Nacional,
Centro Interdisciplinario de Investigación
para el Desarrollo Integral Regional,
Unidad Sinaloa, Guasave,
Sinaloa, México.

³Universidad de Guadalajara,
Centro Universitario de Ciencias Exactas e Ingenierías,
Departamento de Farmacobiología,
Guadalajara, México.

***Corresponding author:**

Email address:

ernesto.cantu@itson.edu.mx

Submitted: 2019-12-21

Accepted: 2022-01-14

Published: 2022-10-21

© Copyright 2022

Julio A. Anduro-Jordan *et al.*open access 

Distributed under Creative Commons CC-BY 4.0

Supplementary material


STEC non-0157 strains in meat from Southern Sonora, Mexico and their antibiotic resistance

Cite this as: Anduro-Jordan JA., Maldonado-Mendoza IE, Figueroa-López AM, Torres-Vitela M del R, Campas-Baypoli ON, Sánchez-Machado DI, Cantú Soto EU. STEC non-0157 strains in meat from Southern Sonora, Mexico and their antibiotic resistance. Veterinaria México OA. 2022;9. doi: 10.22201/fmvz.24486760e.2022.811.855

Supplementary material

Table S1. Sampling sites, kind and number of samples, frequency and period of sampling

Establishment	Meat cuts	Sampling period
TIF slaughterhouse	52 Pork samples (17 loins, 17 ribs, and 21 pulps)	Between October to November 2015 were programed seven visits to the TIF slaughterhouse with a weekly frequency; was not possible to choose a fixed day for the sampling
TIF slaughterhouse	55 beef samples (17 loins, 17 ribs, and 21 pulps) and 33 pork samples (10 loins, 10 ribs, and 13 pulps)	Between October to december 2015 were programed eleven visit to the non-TIF slaughterhouse with a weekly frequency was not possible to choose a fixed day for the sampling
Butchery	50 ground beef samples	Between March to December 2016 were programed visit to the 12 butcherries in urban area of Ciudad Obregon, Sonora with a weekly frequency without a fixed day in the week. It is possible that some butcherries were sampling once a time

**Figure S1.** Detection of *E. coli* O157:H7 pathogenicity genes using multiplex PCR. Lanes 1 and 8: molecular marker 1 Kb plus DNA Ladder; lane 2: negative control (*E. coli* ATCC 25922), lane 3: single PCR for gene *eaeA* (384 bp); lane 4: single PCR for gene *stx2* (255 bp); lane 5: single PCR for gene *stx1* (180 bp); lane 6: single PCR for gene *rfb* (1 000 bp); lane 7: multiplex PCR reaction with *E. coli* O157:H7 DNA. PCR reactions were run on a 2% agarose gel and visualized with ethidium bromide.

Supplementary material




Figure S2. Amplification by multiplex PCR of pathogenicity genes (*stx1*, *stx2*, *eaeA*, *rfb*) from *E. coli* presumptive isolates. Lane 1: negative control (ATCC 25922); lane 2: molecular marker 1Kb plus DNA ladder, lane 3: positive control *E. coli* O157:H7, lanes 4, 5, and 6: *E. coli* JA31 isolates (beef loin), JA49 and JA50 (beef pulp).




Figure S3. End-point PCR to detect the presence of gene *fliCH4* in *E. coli* presumptive isolates from Ciudad Obregon, Sonora. Lane 1: molecular marker 1Kb plus DNA ladder; lane 2: negative control *E. coli* (ATCC 25922); lane 3: positive control JA31 (*fliCH4*, beef loin); lane 4: JA107 (beef ribs); lane 5: JA113 (beef ribs); lane 6: JA116 (pork pulp). Isolates JA49 (beef pulp); JA50 (beef pulp) did not show gene *fliCH4*.