

Functional gluten-free beverage elaborated from whole quinoa and defatted chia extruded flours: antioxidant and antihypertensive potentials

Bebida funcional libre de gluten elaborada con harinas extrudidas de quinoa integral y chíá desgrasada: potencial antioxidante y antihipertensivo

Luis Martín Sánchez-Magaña¹, Cuauhtémoc Reyes-Moreno^{1,2}, José Antonio Garzón-Tiznado^{1,2}, Jorge Milán-Carrillo^{1,2}, Mario Armando Gómez-Favela³, Edith Oliva Cuevas-Rodríguez^{1,2}, Ramona Julieta Espinoza-Moreno³, Roberto Gutiérrez-Dorado^{1,2*}

¹Programa de Posgrado en Ciencia y Tecnología de Alimentos.

²Programa de Posgrado Integral en Biotecnología, Facultad de Ciencias Químico Biológicas (FCQB), Universidad Autónoma de Sinaloa (UAS), Sinaloa, México.

³Ingeniería Agroindustrial, Universidad Politécnica del Mar y la Sierra (UPMyS).

Tel. +52 (667) 7157641. robe399@hotmail.com

*Corresponding author.

Abstract

The current market demands gluten-free (GF) foods with better nutritional/nutraceutical value. In this study, a functional GF beverage (GFB) from a flours mixture of optimized extruded quinoa (OEQ) (70%) and extruded defatted chia (EDC) (30%) was developed. To obtain OEQ, a central composed experimental design with two factors (extrusion temperature [TE] = 60 °C-140 °C and screw speed [VT] = 60 rpm-240 rpm) and three responses (antioxidant activity [AoxA], total phenolic compounds [TPC], and *in vitro* protein digestibility [IVPD]) was used. The optimal extrusion conditions for OEQ were 138 °C/138 rpm, and for EDC they were 147 °C/237 rpm (obtained in a previous work). The mixture had calculated protein efficiency ratio (C-PER) = 2.28 and angiotensin-converting enzyme (ACE) inhibitory potential (IC₅₀) = 0.271 mg/ml. 200 ml of functional GFB contains 5.61 g of good quality protein, 6.13 g of dietary fiber, 88 kcal, and AoxA = 5091 µmol TE. The functional GFB is a healthy alternative for celiac people.

Keywords: Quinoa; chia; optimization; functional gluten-free beverage; antioxidant; antihypertensive.

Resumen

El mercado actual demanda alimentos libres de gluten (LG) con mejor valor nutricional/nutraceutico. En este estudio, se desarrolló una bebida LG (BLG) funcional a partir de una mezcla de harinas de quinoa extrudida optimizada (QEO) (70%) y chíá desgrasada extrudida (CDE) (30%). Para obtener QEO se usó un diseño experimental central compuesto con dos factores (temperatura de extrusión [TE] = 60 °C-140 °C y velocidad de tornillo [VT] = 60 rpm-240 rpm) y tres respuestas (actividad antioxidante [AAox], compuestos fenólicos totales [CFT] y digestibilidad proteínica *in vitro* [DPIV]). Las condiciones óptimas de extrusión para QEO fueron 138 °C/138 rpm y para CDE fueron: 147 °C/237 rpm (obtenidas en trabajo previo). La mezcla tuvo relación de eficiencia proteínica calculada (C-PER) = 2.28 y potencial inhibitorio de enzima convertidora de angiotensina (ECA) (IC₅₀) = 0.271 mg/ml. 200 ml de BLG funcional contiene 5.61 g proteína de buena calidad, 6.13 g fibra dietaria, 88 kcal y AAox = 5091 µmol ET. La BLG funcional es una alternativa saludable para personas celiacas.

Palabras clave: Quinoa; chíá; optimización; bebida funcional libre de gluten; antioxidante; antihipertensivo.

Recibido: 07 de diciembre de 2021

Aceptado: 04 de mayo de 2022

Publicado: 29 de junio de 2022

Cómo citar: Sánchez-Magaña, L.M., Reyes-Moreno, C., Garzón-Tiznado, J.A., Milán-Carrillo, J., Gómez-Favela, M. A., Cuevas-Rodríguez, E. O., Espinoza-Moreno, R. J., & Gutiérrez-Dorado, R. (2022). Functional gluten-free beverage elaborated from whole quinoa and defatted chia extruded flours: antioxidant and antihypertensive potentials. *Acta Universitaria* 32, e3413. doi: http://doi.org/10.15174/au.2022.3413

Introduction

The growing consumer interest in “healthy living” has boosted the market for more natural and functional food claims; additionally, producing foods that contain a high content of phytochemicals and high protein quality that are beneficial to health and have a relatively low cost has been considered (Medina *et al.*, 2020). The population of Americans suffering from celiac, gluten intolerance, and wheat allergy is 1 in every 14 people. Also, many choose gluten-free (GF) diets nowadays because of the perception that it is a healthier option for them. Therefore, in the last decade, the GF market in the US and all over the world has seen a significant growth. Globally, GF product sales reached 4.63 billion USD in 2017 and are expected to reach 6.47 billion USD by 2023, a projected compound annual growth rate of 7.6% (Woomer & Adedeji, 2021). Celiac disease (CD), or gluten-sensitive enteropathy, is a generalized autoimmune disease characterized by chronic inflammation and atrophy of the small intestine’s mucosa by exposure to dietary gluten that affects genetically predisposed individuals (Kamel *et al.*, 2020; Lebwohl & Rubio-Tapia, 2021; Wieser *et al.*, 2021). Gluten refers to a family of proteins known as prolamins (primarily glutenin and gliadin) that constitute the storage protein in the starchy endosperm of many cereal grains such as wheat, barley, and rye (Niland & Cash, 2018). The treatment for CD is a strict gluten-free (GF) diet for life, excluding dietary wheat, rye, barley, and hybrids like Kamut and triticale. A strict GF diet improves histological lesions, blood biochemistry, clinical manifestations, and CD-related complications (Sharma *et al.*, 2020).

GF food products existing in the market today are more expensive than gluten-containing food products. Also, these are known to have lower nutritional value than the critical needs of celiac patients. It is necessary to develop cheap, nutritionally rich gluten-free food products to promote healthy living in people who have the CD (Gómez-Favela *et al.*, 2021). Some of the GF and nutritious grains include cereals (corn, rice), pseudo-cereals (buckwheat, amaranth, chia, quinoa), millets (pearl millet, finger millet, Kodo millet, foxtail millet, pearl millet), and pulses (soybean, common bean, pea, chickpea, Bengal gram), which can be used for the preparation of gluten-free functional foods (Gómez-Favela *et al.*, 2021; Kumar, 2018).

Quinoa (*Chenopodium quinoa* Willd.) is a crop native to the Andean region. It is an annual dicotyledonous herb belonging to the genus *Chenopodium*, which has been cultivated for 5000–7000 years (Tan *et al.*, 2020). Due to its increasing popularity, the grain has been recently introduced as a cereal crop in non-indigenous regions such as Europe, North America, Australia, China, and Japan. Quinoa is a member of the family Amaranthaceae rather than Poaceae (Gramineae), but because the seeds are mostly milled into flour and used as a cereal crop, it is referred to as a pseudo-cereal (Li *et al.*, 2021). Quinoa seeds reveal high levels of essential fatty acids, vitamins, minerals, dietary fibers (8%-13%), and proteins (12%-23%) with a balanced amino acid spectrum, with high contents of lysine and methionine (Angeli *et al.*, 2020; Aziz *et al.*, 2018; Montemurro *et al.*, 2019; Pellegrini *et al.*, 2018). In addition to this rich nutritional composition, these seeds have been found to contain a large variety of bioactive compounds (phenolic compounds, phytosterols, squalene, phytoecdysteroids, carotenoids), which have proved protective against a variety of diseases, particularly cancer and inflammatory diseases, and they may reduce the risk of cardiovascular diseases, considering quinoa seeds a functional food (Montemurro *et al.*, 2019; Tang & Tsao, 2017). Moreover, quinoa is considered gluten-free and suitable for celiac patients and people who have a wheat allergy (Caratini & Rosentrater, 2019; Demir & Bilgicli, 2021).

Chia (*Salvia hispanica* L.) seeds were used together with bean, corn, and amaranth by Precolumbian Mesoamerican cultures (Mayas and Aztecs) to prepare foods and medicines. Chia was the second main crop after beans. In the Aztecs communities, chia was used for food, cosmetics, and religious rituals (Hrnčić *et al.*, 2020). In Mexico, chia seeds are currently used for their nutritional and medicinal properties, i.e., endurance for athletes, appetite suppressor, blood glucose control, weight loss agent, and intestinal regulation (Martínez-Cruz & Paredes-López, 2014). Chia seed has approximately 15%-24% protein, 26%-41% carbohydrates, and 25%-40% fat. It has been studied principally due to its oil quality content, possessing almost 55%-60% linolenic acid (ω -3), 18%-20% linoleic acid (ω -6), 6% monounsaturated fat (ω -9), and 10% saturated fat. On the other hand, the seed has highly soluble and insoluble dietary fiber, over 35% of the total weight, and a rich source of the B vitamins and minerals. It also contains six more-fold calcium, 11 more-fold phosphorus, and four more-fold potassium than 100 g of milk; besides this, it possesses magnesium, iron, zinc, and copper (León-López *et al.*, 2019; Masood, 2022; Orón-Tamayo *et al.*, 2017). On top of that, it has many natural antioxidants, such as phenolic compounds (chlorogenic and caffeic acids, and flavonols [quercetin, kaempferol, myricetin]), which protect against some adverse conditions (Hrnčić *et al.*, 2020). Another important characteristic of this seed is that it does not contain gluten. Persons with celiac disease can consume it, thus, making seeds a safe ingredient for gluten-free diets (Menga *et al.*, 2017). Recently, Rubavathi *et al.* (2020) reported that chia seeds extracts possess anti-obesity activity.

Extrusion process is an adequate technology to generate functional foods. This process increases the digestibility and biological value of proteins and maintains or increases the content of bioactive compounds and nutraceutical properties of foods, due to the use of high-temperature, high-pressure, and high shear conditions in a short period. These accelerate process conditions, used during the extrusion, cause important molecular transformation and chemical reactions (modification in starch, proteins, dietary fiber and lipids, inactivation of enzymes and microorganisms, and formation of color polymers and volatile flavor components) on the food materials. In addition, this process is a highly efficient alternative technology that minimizes energy and water consumption, and does not produce polluting effluents (León-López *et al.*, 2019; Ortiz-Cruz *et al.*, 2020; Ramos-Enríquez *et al.*, 2018; Reyes-Moreno *et al.*, 2018).

In the functional food market, the functional drinks market is one of the fastest-growing segments (Tolun & Altintas, 2019). Consumer demand for "healthy" foods and beverages is seen as a driving force behind the functional foods sector (Kim & Adhikari, 2020). An expert report on science from the Institute of Food Technologists (IFT) defined functional food and food components as products that provide a health benefit beyond basic nutrition (for the intended population). The functional drinks are considered as a most active and effective sector because of (1) the possibility to cover consumer demands for appearance, container contents, shape, and size; (2) the ease of distribution and better storage for low temperatures and shelf-stable products; and (3) the opportunity to incorporate bioactive compounds (peptides, antioxidants, omega three fatty acids, dietary fiber, prebiotics) and nutrients (good quality protein, with a good balance of EAA, lipids -essential fatty acids-, vitamins, minerals) (Argüelles-López *et al.*, 2018).

In Mexico, 76.6% of adults, 36.3% of adolescents, and 33.2% of children suffer from overweight and obesity, and 25.5% of adults suffer from arterial hypertension, as reported by Encuesta Nacional de Salud y Nutrición de Medio Camino (ENSANUTMC) (Instituto Nacional de Salud Pública [INSP], 2016). In the initiation and progression of hypertension, oxidative stress is a significant factor. An essential factor in blood pressure regulation is the angiotensin-converting enzyme (ACE). ACE inactivates Bradykinin (vasodilator) and catalyzes the conversion of angiotensin I into angiotensin II (vasoconstrictor). One of the strategies for combating hypertension is searching for ACE inhibitors from natural products such as peptides, triterpenes, and phenolic compounds (Argüelles-López *et al.*, 2018). Some researchers (Argüelles-López *et al.*, 2018; Masood, 2022; Montoya-Rodríguez *et al.*, 2015; Pellegrini, *et al.*, 2018) have reported that phenolic compounds derived from pseudo-cereals (quinoa and amaranth) and chia have antioxidant and antihypertensive effects.

In this context, public policies must be implemented to reverse the obesogenic environment, characterized by increased access to high-calorie food and beverages, a sedentary lifestyle, and constant promotion of non-healthy product consumption.

In this sense, the objectives of this work were: (1) to optimize the extrusion process of whole quinoa seeds to find the best combination extrusion temperature (ET)/screw speed (SS) for producing optimized extruded quinoa (OEQ) flour with high antioxidant activity (AoxA), total phenolic content (TPC), and *in vitro* protein digestibility (IVPD); and (2) to develop a functional gluten-free beverage (GFB) high nutritional/nutraceutical value from a mixture of optimized extruded quinoa and extruded defatted chia (OEQ, EDC) flours.

Materials and Methods

Materials

Commercial varieties of quinoa (*Chenopodium quinoa* Willd) and chia (*Salvia hispanica* L.) seeds were purchased in a local market (Mercado Rafael Buelna, Culiacán, Sinaloa, México). Seeds were cleaned and kept in tightly sealed containers at 4 °C until used.

Production of extruded quinoa (EQ) flours

The EQ flours were prepared as suggested by León-López *et al.* (2019). Response surface methodology (RSM) was applied to find the best combination of process variables (Extrusion temperature [ET = 60 °C-140 °C]/screw speed [SS = 60 rpm-240 rpm]) for producing optimized extruded quinoa (OEQ) flour with maximum values of AoxA, TPC, and IVPD. The central composite experimental design generated 13 treatments (Table 1). Portions of 250 g of whole quinoa seeds were mixed with lime (0.21%) and water (28% humidity), and they were allowed to rest (4 °C-8 °C/12 h) and tempered (25 °C/1 h). After tempered, the portions of quinoa flours were fed to the extruder and underwent various extrusion temperature/screw speed (Table 1). The extrudates coming from each of the 13 treatments were cooled, ground, packed in plastic bags, and stored (4 °C). AAox, TPC, and DPIV were evaluated to the 13 different EQ flours.

Optimization of the extrusion process to obtain optimized extruded quinoa (OEQ) flour

Predictive models of second order were generated for each response variable from the data experimentally obtained from the extrusion process for whole quinoa seeds (Table 1). The models were adjusted, and, subsequently, the non-significant terms ($p > 0.1$) were eliminated from the second-order polynomials, and new prediction models were calculated for each response variable (Yolmeh & Jafari, 2017). The program Design-Expert version 7.00 was used to carry out these analyses.

The RSM's conventional graphical method was applied to obtain maximum values (optimization) of AoxA, TPC, and IVPD in optimized extruded quinoa (OEQ) flour. The prediction models were used to graphically represent the system (Figure 1a, 1b, 1c). The overlay of each one of the response variables contour plots allowed to obtain a unique contour plot (Figure 1d) for observation and selection of the best (optimum) combination ET/SS for producing OEQ flour (Figure 1d) with high AoxA, TPC, and IVPD. Once the optimized extrusion conditions were obtained, OEQ flour was prepared; it was packed in plastic bags and stored at 4 °C. After that, OEQ flour was analyzed for chemical composition, nutritional and nutraceutical properties.

Antioxidant activity and total phenolic content (AoxA, TPC)

The ORAC assay carried out by dilution of free and bound phenolic extracts were diluted in 75 mM phosphate buffer (pH 7.4). Aliquots of 25 µL of diluted extracts were mixed with 150 µL of fluorescein (0.1 mM) and 25 µL of the peroxy radical AAPH (200 mM). After 30 minutes, fluorescence (485 nm for excitation and 538 nm for emission) was measured (37 °C) at 2 min intervals for 60 min, using a Synergy Microplate Reader (Synergy™ HT Multi-Detection, BioTek, Inc., Winooski, VT) (León-López *et al.*, 2019). The results ORAC assay were expressed as micromol of Trolox equivalent (TE) per 100 g of dry weight sample. The TPC of free and bound extracts were determined using 20 ml of appropriate dilutions of extracts, and they oxidized with 180 µL of Folin-Ciocalteu reagent (Gámez-Valdez *et al.*, 2021). The resulting blue color's absorbance was measured after 20 min, at 750 nm, using the Synergy Microplate Reader and expressed the TPC in mg gallic acid equivalents (GAE) / 100 g DW. All measurements were made in triplicate.

In vitro protein digestibility (IVPD)

The IVPD was evaluated according to Salas-López *et al.* (2018), using a modified multienzyme system. Fifty milliliters of a multienzyme solution were prepared (80 mg of pancreatic trypsin [15 600 U/mg], 155 mg of bovine pancreatic chymotrypsin [83.9 U/mg], 65 mg of porcine intestinal peptidase [102 U/g] and distilled water). Also, a protein-water suspension was prepared (6.25 g of protein/L). The pH of multienzyme solution and protein suspension was adjusted to 8.0 using 0.1 N HCl and 0.1 N NaOH solutions. Also, 5 ml of multienzyme solution were mixed with 50 ml of the protein suspension employing constant stirring in a water bath at 37 °C. The extent of hydrolysis was determined by pH-drop after a 10 min incubation period (X), and the percentage of protein digestibility (IVPD) was calculated by the regression equation:

$$\text{IVPD} = 210.469 - 18.10 X$$

where X is the pH of protein suspension at 10 min of digestion with the multi-enzymatic system.

Preparation of extruded defatted chia (EDC) flour

Production of EDC flour was carried out according to the methodology proposed by León-López *et al.* (2019). For this, 500 g lots of partially defatted chia seeds (lipids content = 5%-6%, DW) were placed in a domestic blender at low velocity, the grits obtained were ground in the same blender to obtain grits that passed through a 40-US mesh (0.425 mm) screen. The extrusion cooking was carried out on a single screw laboratory extruder, model 20 DN (CW Brabender Instruments, Inc, NJ, USA), at extrusion temperature (ET) and screw speed (SS) of 147 °C and 237 rpm, respectively. Extrudates were cooled and dried at environmental conditions (25 °C, RH = 65%), milled to pass through an 80-US mesh (0.180 mm) screen, packed in plastic bags, and stored at 4 °C. EDC flour was analyzed for chemical composition, nutritional and nutraceutical properties.

Functional gluten-free beverage (GFB) preparation. Sensorial evaluation

The GFB was developed from the mixture: 70% OEQ flour + 30% EDC flour. The mixture (25 g) was added with 2.5 g of sweetener Stevia™ (no-calorie), vanillin powder (1 g), and water (170 ml). The suspension was stirred with a domestic shaker (medium velocity), refrigerated (8 °C–10 °C), and sensory evaluated for acceptability. The sensorial evaluation was carried out by 80 non-trained panelists using a labeled affective magnitude (LAM) scale. These values were transformed to a 0-100 scale (Cardello & Schutz, 2004).

Chemical composition

The official methods (Association of Official Analytical Chemists [AOAC], 2012) were used to determine protein (micro-Kjeldahl method 960.52), lipids (920.39), and moisture (method 925.09B) contents. Soluble and insoluble dietary fiber (SDF/IDF) were evaluated according to the enzymatic-gravimetric method for total dietary fiber (TDF) (method 985.29) (AOAC, 2012).

Nutritional properties: essential amino acid (EAA), chemical score (CS), and calculated protein efficiency ratio (C-PER)

The EAA composition was determined using an analytical scale (4.6mm × 250mm) hypersil ODS C18 column kept at 38 °C and connected to an HPLC system (León-López *et al.*, 2019). The HPLC system (GBC, Dandenong, Australia) was equipped with a fluorescence detector >LC 5100 set at 270 nm (excitation) and 316 nm (emission). We used an ultraviolet detector to detect tryptophan at 280 nm.

The chemical score (CS) was calculated as $CS = (\text{Content of the most limiting EAA}/\text{REAA}) \times 100$; EAA = Essential amino acid and REAA = Recommended EAA requirements for three-year-old children and older, adolescents, and adults (Food and Agriculture Organization of the United Nations [FAO], 2013).

C-PER was calculated as described by Satterlee *et al.* (1982) and summarized by the AOAC (2012). This procedure is based on the IVPD and the EAA composition of the optimized mixture. All determinations were made in triplicate.

ACE inhibitory potential

The angiotensin-converting enzyme (ACE) inhibitory activity in free and bound phenolic extracts was determined using the Dojindo ACE Kit-WST test kit (Dojindo Laboratories, Kumamoto, Japan). Briefly, 20 μL of sample solution was added to a sample well, and 20 μL of deionized water was added to blank 1 and blank 2 wells. 20 μL of substrate buffer was added to each well. 20 μL of deionized water were added to blank 2 well, while 20 μL of enzyme working solution (ACE 0.2 U/ml, aminoacylase 172 kU/ml, and 3HBMGGG 3.4 mg/ml) was added to each sample and blank 1 well. All wells containing above solutions were incubated at 37 °C for 1 h. Afterwards, 200 μL of the indicator working solution (0.333 mM NAD⁺, 0.333 mM WST⁻¹, 0.1 mM EDTA, 0.633 U ml⁻¹ diaphorase, and 0.700 U ml⁻¹ 3-hydroxybutyrate dehydrogenase) was added to each well with further incubation for 10 min. The yield of 3HB was monitored indirectly through formazan concentration, which was measured at 450 nm using a Microplate Reader (SynergyTM HT Multi-Detection, BioTek, Inc., Winooski, VT, USA). ACE inhibition was calculated from the equation:

$$\text{ACE inhibition (\%)} = [(A_{\text{blank 1}} - A_{\text{sample}}) / (A_{\text{blank 1}} - A_{\text{blank 2}})] \times 100$$

where A_{blank1} represented positive control (without ACE inhibition), A_{blank2} was the absorbance of the reagent blank (negative control), and A_{sample} was the absorbance of the sample (extract). The concentration of extract inducing 50% inhibition (IC_{50}) of ACE was determined with non-linear regression using the Prism v5 software (GraphPad Prism) (Gámez-Valdez *et al.*, 2021).

Statistical analyses

The experimental results on the chemical composition, nutritional properties, antioxidant activity, phenolic compounds content, and ACE inhibitory potential (IC_{50}) of flours were subjected to analysis of variance using the statistical software MINITAB 17.0; the multiple range Duncan's test at $p \leq 0.05$ was used to compare differences among means.

Results

Prediction models for antioxidant activity (AoxA), total phenolic content (TPC), and *in vitro* protein digestibility (IVPD) for extruded quinoa (EQ) flours

The experimental values for AoxA, TPC, and IVPD of EQ varied from 13 230 $\mu\text{mol TE}/100\text{g sample (DW)}$ to 16 470 $\mu\text{mol TE}/100\text{g sample (DW)}$, from 475 mg GAE/ 100g sample (DW) to 640 mg GAE/ 100g sample (DW), and from 78.2% to 87.2 %, respectively (Table 1). The analysis of variance (Anova) showed that AoxA, TPC, and IVPD depend significantly ($p < 0.1$) on ET and SS linear and quadratic terms. The prediction models, using decoded variables, for the response variables were:

$$Y_{\text{AoxA}} = +15196.60 + 924.14 (\text{ET}) + 301.80 (\text{SS}) - 221.86 (\text{ET})^2 + 309.64 (\text{SS})^2$$

$$Y_{\text{TPC}} = +540.97 + 58.34 (\text{ET}) + 10.98 (\text{SS}) + 24.02 (\text{SS})^2$$

$$Y_{\text{IVPD}} = + 85.74 + 3.50 (\text{ET}) - 0.52 (\text{SS}) - 0.96 (\text{ET})^2 - 1.76 (\text{SS})^2$$

The regression models explained 94.7%, 96.4%, and 91.7% of the total variability ($p < 0.001$) in AoxA, TPC, and IVPD. The lack of fit was not significant ($p > 0.05$). The relative dispersion of the experimental points from the predictions of the models (CV) was $< 10\%$. These values indicated that the experimental models were adequate and reproducible. In general, AoxA and TPC of extruded quinoa (EQ) flours increased with ET until reaching maximum values around 140 °C and 240 rpm (Figure 1a, 1b), while the IVPD of EQ increased with ET showed maximum values at high ET (140 °C) and intermediate SS (150 rpm) (Figure 1c). The potential of pseudo-cereals seeds as functional food has been related to their antioxidant activity and other nutraceutical properties (antihypertensive, antidiabetic, anti-inflammatory, antiatherosclerotic) associated with phenolic compounds and bioactive peptides (Martínez-López *et al.*, 2020; Orona-Tamayo *et al.*, 2018; Tan *et al.*, 2020).

Table 1. Experimental design¹ used to obtain different combinations of extrusion temperature/screw speed for producing extruded quinoa (EQ) flours, and experimental results for response variables.

Assay	Process variables		Response variables		
	Extrusion temperature (°C)	Screw speed (rpm)	AoxA ²	TPC ³	IVPD ⁴
1	71.72	86.36	14 150	485.4	80.5
2	128.28	86.36	15 730	609.9	86.3
3	71.72	213.64	14 891	503.7	79.7
4	128.28	213.64	16 470	640.0	87.2
5	60.00	150.00	13 230	475.0	78.2
6	140.00	150.00	16 224	620.6	88.6
7	100.00	60.00	15 460	580.3	83.3
8	100.00	240.00	16 120	608.2	80.3
9	100.00	150.00	15 318	539.7	86.6
10	100.00	150.00	15 386	550.1	84.6
11	100.00	150.00	15 251	543.3	85.3
12	100.00	150.00	15 072	530.2	86.4
13	100.00	150.00	14 956	538.4	85.8

¹ Central composite experimental design with two factor and five variation levels (13 assays); ² AoxA = Antioxidant activity ($\mu\text{mol Trolox equivalents [TE]}/100 \text{ g, DW}$); ³ TPC = Total phenolic content (mg Galic acid equivalents [GAE]/100 g, DW); ⁴ IVPD = *in vitro* protein digestibility (%).

Source: Author's own elaboration.

Optimization of the whole quinoa seeds extrusion process

Figure 1d, which was obtained from the superimposition of response variables contour plots (Figure 1a, 1b, 1c), was utilized to determine the best combination of process variables for the production of optimized extruded quinoa (OEQ) flour. The central point of the optimization region in Figure 1d corresponds to the optimum combination of process variables (ET = 138 °C/SS = 138 rpm) for producing OEQ flour with the highest AoxA, TPC, and IVPD values. The predicted values of AoxA, TPC, and IVPD, using the predictive models of each response variable and the optimal conditions of extrusion, were 15 992 $\mu\text{mol TE}/100\text{ g}$ sample (DW), 618.4 mg GAE/100 sample (DW), and 88.7%, respectively (Figure 1d). The experimental values of AoxA, TPC, and IVPD of OEQ flour (Table 4) were similar to the predicted values, meaning that the extrusion process's optimal conditions were appropriate and reproducible.




Figure 1. Contour plots showing the effect of extrusion temperature (ET) and screw speed (SS) on (A) Antioxidant activity, (B) Total phenolic content, and (C) *in vitro* protein digestibility. (D) Region of the best combination of extrusion process variables for producing optimized extruded whole quinoa flour (OEQ)

Source: Author's own elaboration.

Antioxidant activity (AoxA) and total phenolic content (TPC) of optimized extruded quinoa (OEQ) flour

The extrusion of quinoa seeds increased ($p < 0.05$) AoxA (12.95%) and bound (44.81%) and total (29.09%) phenolic compounds when compared with unprocessed whole quinoa seeds (Table 2). These results show that the bound phenolics were the primary contributors (64%-72%) to the antioxidant activity in unprocessed and optimized extruded quinoa flours (Table 2).

Table 2. Phenolic content and antioxidant activity of quinoa and chia flours.

Property	Unprocessed whole	OEQ ¹	Unprocessed	EDC ¹
	quinoa flour	flour	defatted chia flour	flour
Phenolic compounds²				
Free	174.5 ± 15.2A	176.1 ± 14.2A	248.94 ± 2.97B	302.44 ± 4.81A
Bound	312.6 ± 10.1B	452.7 ± 10.2A	252.32 ± 4.19A	339.40 ± 3.61B
Total	487.1 ± 17.6B	628.8 ± 14.6A	501.26 ± 12.70B	641.84 ± 14.2A
Antioxidant activity³				
Free phytochemicals	3570 ± 198 ^A	3673 ± 183 ^A	8456 ± 680 ^B	10 781 ± 898 ^A
Bound phytochemicals	10 710 ± 710 ^B	12 258 ± 872 ^A	14 268 ± 1102 ^B	15 986 ± 975 ^A
Total phytochemicals	14 280 ± 927 ^B	16 129 ± 933 ^A	22 724 ± 1132 ^B	26 767 ± 1076 ^A

^{A-B} Means with the same letter, in first and second columns, are not different (Duncan, $p > 0.05$ - no significant difference); ^{A-B} Means with the same letter, in third and fourth columns, are not different (Duncan, $p > 0.05$ - no significant difference); ¹ OEQ = Optimized extruded whole quinoa, EDC = Extruded defatted chia; ² mg Gallic acid equivalents (GAE)/100 g sample (DW); ³ μmol Trolox equivalents (TE) / 100 g sample (DW).
Source: Author's own elaboration.

Antioxidant activity (AoxA) and total phenolic content (TPC) of extruded defatted chia (EDC) flour

The extrusion cooking of defatted chia seeds increased ($p < 0.05$) AoxA in 17.80%, when compared with unprocessed defatted chia seed (26 767 μmol TE/100 sample, DW vs. 22 724 μmol TE/100 sample, DW) (Table 2). The AoxA of free, bound, and total phenolic compounds increased ($p < 0.05$) by 21.49%, 34.51%, and 28.04%, respectively, after the optimized extrusion cooking process (Table 2).

Chemical composition, essential amino acid (EAA) content, *in vitro* protein digestibility (IVPD), and C-PER of optimized extruded quinoa (OEQ) and extruded defatted chia (EDC) flours

The extrusion process caused slight changes on chemical composition of quinoa and chia seeds during production of OEQ and EDC flours (Table 3). Milán-Carrillo *et al.* (2012), Argüelles-López *et al.* (2018), and León-López *et al.* (2019) reported similar results to the extrusion of the pseudo-cereals amaranth and chia.

Table 3 shows the essential amino acid content of unprocessed and processed flours. In general, the essential amino acid content of unprocessed and extruded quinoa/chia flours were higher than those suggested by FAO (2013) for the essential amino acid requirement for children (3 years old and older), adolescents, and adults. Lys was the only one presented as limiting amino acid in unprocessed and extruded defatted chia flours. This tendency is similar to literature results of essential amino acid content in amaranth (Argüelles-López *et al.*, 2018) and chia (León-López *et al.*, 2019) seeds. The extrusion process caused only a slight decrease ($p < 0.05$) in the content of some EAA (Met+Cys, Phe+Tyr, and Trp) in EDC flour. Unprocessed and extruded quinoa/chia flours present a high EAA chemical score (91.66-100.00). On the other hand, the extrusion process increased IVPD of OEQ and EDC flours in 20% and 8%, respectively. Results in Table 3 also indicated that C-PER increased during the OEQ and EDC flours preparation.

ACE Inhibitory potential of optimized extruded quinoa (OEQ) and extruded defatted chia (EDC) flours

The phenolic extracts of OEQ and EDC flours had better ACE inhibitory potential (IC_{50} = Concentration of phenolic extracts in mg extract/ml suspension, required to produce an inhibition of 50% of the activity of ACE) than phenolic extracts from unprocessed quinoa (0.150 mg extract/ml vs. 0.907 mg extract/ml) and defatted chia (0.210 mg extract/ml vs. 0.871 mg extract/ml) flours, respectively (Table 3).

Table 3. Chemical composition and nutritional/nutraceutical properties of quinoa and chia flours.

Property	Unprocessed whole quinoa flour	OEQ ¹ flour	Unprocessed defatted chia flour	EDC ¹ flour	Standard FAO ²
Chemical composition (% DW)					
Protein	15.95 ± 0.07 ^A	15.99 ± 0.07 ^A	31.09 ± 0.37 ^A	30.66 ± 0.29 ^A	---
Lipids	6.02 ± 0.07 ^A	5.63 ± 0.07 ^B	5.27 ± 0.06 ^A	5.12 ± 0.07 ^B	---
Minerals	3.50 ± 0.07 ^A	3.54 ± 0.07 ^A	6.88 ± 0.07 ^A	6.91 ± 0.08 ^A	---
Dietary fiber					---
Soluble	2.46 ± 0.05 ^B	3.22 ± 0.03 ^A	6.71 ± 0.09 ^B	7.36 ± 0.07 ^A	---
Insoluble	8.03 ± 0.04 ^A	7.34 ± 0.07 ^B	41.21 ± 0.13 ^B	43.01 ± 0.13 ^A	---
Total	10.49 ± 0.08 ^A	10.56 ± 0.07 ^A	47.92 ± 0.21 ^B	50.37 ± 0.25 ^A	---
Carbohydrate	64.04 ± 0.19 ^B	64.28 ± 0.22 ^B	8.84 ± 0.03 ^A	6.94 ± 0.03 ^B	---
Nutritional					
EAA3 (g/100 g protein)					
His	2.82 ± 0.02 ^A	2.76 ± 0.02 ^A	1.99 ± 0.04 ^A	2.01 ± 0.03 ^A	1.60
Ile	4.57 ± 0.03 ^A	4.50 ± 0.04 ^A	3.37 ± 0.03 ^A	3.29 ± 0.04 ^A	3.00
Leu	7.03 ± 0.04 ^A	6.88 ± 0.03 ^A	6.17 ± 0.03 ^A	6.08 ± 0.05 ^A	6.10
Lys	5.83 ± 0.03 ^A	5.72 ± 0.03 ^A	4.48 ± 0.04 ^A	4.40 ± 0.05 ^A	4.80
Met+Cys	4.60 ± 0.05 ^A	4.59 ± 0.02 ^A	2.69 ± 0.03 ^A	2.48 ± 0.02 ^B	2.30
Phe+Tyr	7.17 ± 0.03 ^A	7.01 ± 0.05 ^A	5.18 ± 0.02 ^A	5.02 ± 0.03 ^B	4.10
Thr	3.59 ± 0.04 ^A	3.61 ± 0.04 ^A	2.58 ± 0.04 ^A	2.51 ± 0.04 ^A	2.50
Trp	0.91 ± 0.01 ^A	0.94 ± 0.02 ^A	1.63 ± 0.02 ^A	1.54 ± 0.05 ^B	0.66
Val	4.38 ± 0.04 ^A	4.41 ± 0.03 ^A	4.23 ± 0.02 ^A	4.25 ± 0.04 ^A	4.00
Total	40.90	40.42	32.32	31.58	29.06
Chemical score	100	100	93.33	91.66	---
Limiting EAA	None	None	Lys	Lys	---
IVPD4 (%)	68.2 ± 0.17 ^B	82.1 ± 0.12 ^A	77.9 ± 0.13 ^B	84.05 ± 0.18 ^A	---
C-PER5	1.94	2.37	1.69	2.15	---
Nutraceutical					
AoxA	14 280 ± 927 ^B	16 129 ± 933 ^A	22 724 ± 1132 ^B	26 767 ± 1076 ^A	---
ACE7 Inhibitory potential (IC50) ⁸	0.907	0.150	0.871	0.210	---

^{A-B} Means with the same letter, in first and second columns, are not different (Duncan, $p > 0.05$ - no significant difference); ^{A-B} Means with the same letter, in third and fourth columns, are not different (Duncan, $p > 0.05$ - no significant difference); ¹ OEQ = Optimized extruded quinoa, EDC = Extruded defatted chia; ² EAA requirements for children, adolescents and adults (3 years old and older) according to FAO (2013); ³ EAA = Essential amino acid, ⁴ IVPD = *in vitro* protein digestibility, ⁵ C-PER = Calculated protein efficiency ratio, ⁶ μmol Trolox equivalents (TE)/100 g sample (DW); ⁷ angiotensin converting enzyme; ⁸ mg extract/ml suspension.

Source: Author's own elaboration.

Chemical composition and nutritional/nutraceutical properties of the mixture of flours (70% OEQ + 30% EDC)

The mixture contains 20.39% proteins, 5.48% lipids, and 22.50% total dietary fiber (TDF) on a dry basis. The EAA content in the mixture was greater than that recommended (FAO, 2013) for three-year-old children, adolescents, and adults (39.52 g/100 g protein vs. 29.06 g/100 g protein); their chemical score was 100. It did not present limiting EAA (Table 4). The mixture presented IVPD and calculated protein efficiency ratio (C-PER) of 81.12% and 2.28, respectively (Table 4). A mixture of nixtamalized maize (26.7%) and extruded chickpea (73.3%) flours had IVPD and C-PER of 54.6% and 1.86, respectively (Alarcón-Valdez *et al.*, 2005).

The mixture had an antioxidant activity of 18 943 $\mu\text{mol TE}/100\text{ g (DW)}$ and an antihypertensive potential, expressed as IC_{50} , of 0.271 mg extract/ml suspension (Table 4). This IC_{50} value was lesser than those reported for soybean protein hydrolysates (80 $\mu\text{g}/\text{ml}$ -360 $\mu\text{g}/\text{ml}$) (Pyo & Lee, 2007) and for protein fractions of amaranth (albumins: 0.35 mg/ml; globulins: 15 mg/ml) (Tovar-Pérez *et al.*, 2009).

Table 4. Chemical composition and nutritional/nutraceutical properties of the mixture.

Property	Mixture (70% OEQ +30% EDC) ¹	Standard FAO ²
Chemical composition (% DW)		
Protein	20.39	---
Lipids	5.48	---
Dietary fiber		
Soluble	4.46	---
Insoluble	18.04	---
Total	22.50	---
Nutritional		
EAA³ (g/100 g protein)		
His	4.24	1.60
Ile	4.14	3.00
Leu	6.64	6.10
Lys	5.40	4.80
Met+Cys	3.96	2.30
Phe+Tyr	6.41	4.10
Thr	3.28	2.50
Trp	1.09	0.66
Val	4.36	4.00
Total	39.52	29.06
Chemical score	100	---
Limiting EAA	None	---
In vitro protein digestibility (%)	81.12	---
C-PER ⁴	2.28	---
Nutraceutical		
Antioxidant activity ⁵	18 943	---
ACE ⁶ Inhibitory potential (IC_{50}) ⁷	0.271	---

¹ OEQ = Optimized extruded quinoa flour, EDC = Extruded defatted chia flour; ² EAA requirements for children, adolescents, and adults (3 years old and older) according to FAO (2013); ³ EAA = Essential amino acid, ⁴ C-PER = Calculated protein efficiency ratio, ⁵ $\mu\text{mol Trolox equivalents (TE)}/100\text{ g sample (DW)}$; ⁶ Angiotensin converting enzyme; ⁷ mg extract/ml suspension.

Source: Author's own elaboration.

Formulation, nutrimental composition, energy content, and antioxidant activity of the functional gluten-free beverage (GFB) are prepared from the mixture

The formulation of the GFB (main ingredient: mixture of flours [70% OEQ + 30% EDC]) was based on the preparation of traditional drinks consumed in Mexico made from grains (e.g., rice, barley, maize). Sensory evaluations were carried out to define the quantities of the ingredients. The formulation originated a beverage containing 12.6% of the main ingredient (mixture) and a protein content > 1.5%, with quality similar to casein. Stevia™ (Non-caloric vegetable sweetener) was used to comply with the Ministry of Health recommendation: "A 200 ml portions of the beverage (food) should not contain more than 140 kcal".

A portion of 200 ml of GFB prepared with 25 g of the mixture of flours (70% OEQ + 30% EDC) contains 5.61 g good protein quality, 1.50 g lipids, 12.92 g carbohydrates, and dynamic content of 88 kcal (Table 5). This portion of beverage covers between 35% and 23.3% of the daily protein requirements for children of 1-3 and 4-8 years of age. Functional beverage (200 ml) had the total hydrophilic antioxidant activity of 5091 µmol TE, which contributes 100%-169% of the recommended daily consumption of antioxidants (from 3000 µmol TE to 5000 µmol TE) (United States Department of Agriculture [USDA], 2010). Untrained panelists assessed with an average value of 83 the acceptability of the GFB (satisfaction level between "I like it very much" and "I really like it"). It is expected that this acceptability of beverages will allow for adequate consumption and, thus, can provide benefits to consumers' health.

Table 5. Nutrimental and energy contents, antioxidant activity and acceptability of a portion of 200 ml of the functional gluten free beverage (GFB).

Property	Functional gluten free beverage (GFB) ¹
Nutrient (g)	
Protein	5.61
Lipids	1.50
Dietary fiber	
Soluble	1.25
Insoluble	4.78
Total	6.13
Carbohydrates	12.92
Energy (kcal)	88
AoxA ² (µmol TE)	5091
Acceptability ³	83

¹ Main ingredient: Mixture = 70% optimized extruded whole quinoa flour + 30% extruded defatted chia flour; ² AoxA = Antioxidant activity; ³ A value between "I like it very much" and "I really like it" on a hedonic scale of 11 points.

Source: Author's own elaboration.

Discussion

Antioxidant activity (AoxA) and total phenolic content (TPC) of optimized extruded quinoa (OEQ) flour

These results coincide with the reported by Milán-Carrillo *et al.* (2012), Argüelles-López *et al.* (2018), and Song *et al.* (2020). They observed that the extrusion of a pseudo-cereal (amaranth, chia, and quinoa) increased AoxA and TPC when compared with crude grains, which improves its functionality and therapeutic potential. This behavior could be attributed to (1) the breaking of conjugated phytochemicals and release free of phytochemicals (Dewanto *et al.*, 2002), (2) the prevention of enzymatic oxidation, and (3) the darker colors of the extruded flours indicating the formation of Maillard reaction products having antioxidant properties (Fares & Menga, 2012).

Bioactive phytochemicals exist in free, soluble-conjugated, and bound forms. Bound phytochemicals, mostly in cell wall materials, are difficult to digest in the upper intestine and may be digested by bacteria in the colon to provide health benefits and reduce colorectal cancer incidence. Dietary polyphenols significantly impact cancer stem cells (CSC), which may be applied in cancer prevention and treatment (Liskova *et al.*, 2019). Colorectal cancer (CRC) is a major cancer type and a leading cause of death. Current medical treatments are not sufficient due to the lack of effective therapy, chemoresistance, adverse side effects, and disease recurrence. Recently, epidemiologic observations have highlighted the association between the ingestion of foods enriched with phytochemicals and nutrients and the lower CRC risk. According to preclinical studies, dietary phytochemicals exert chemopreventive effects on CRC by regulating different markers and signaling pathways. Additionally, gut microbiota plays a strong effect on CRC onset and progression (Afrin *et al.*, 2020).

Antioxidant activity (AoxA) and total phenolic content (TPC) of extruded defatted chia (EDC) flour

Gómez-Favela *et al.* (2017) and León-Murillo *et al.* (2021) reported similar values of AoxA to free, bound, and total phenolic compounds in chia seeds. León-Murillo *et al.* (2021) and Argüelles-López *et al.* (2018) reported that the application of the extrusion process to pseudo-cereals such as chia and amaranth, at optimal conditions, increased the AoxA and bound and free phenolic content when compared to unprocessed seeds.

Antioxidant activity increase could also be a result of the release of antioxidant phenolic compounds during extrusion process. Prevention of oxidation could be caused by phenolic compounds in the extruded product, enzymatic inactivation during the processing and due to the presence of Maillard reaction products with antioxidant activity, generated during extrusion of raw material that contains amino acids and reducing sugars (Argüelles-López, *et al.*, 2018). Félix-Medina *et al.* (2021) stated that the extrusion process conditions (humidity, temperature, screw speed) have a crucial impact on the levels of bioactive compounds (phenolic acids, flavonoids) by releasing, by degrading, or by composition. The increase in the AoxA might be due to the products formed during the Maillard reaction. These non-enzymatically formed products have affinities for unstable chemical radicals, thus inflating the value of AoxA. However, in some cases, the level of bioactive compounds in the extruded products may have elevated certain phenolic acid levels due to their release from the cell wall matrix (Félix-Medina *et al.*, 2020; Ramos-Enriquez *et al.*, 2018)

Chemical composition, essential amino acid (EAA) content, *in vitro* protein digestibility (IVPD), and C-PER of optimized extruded quinoa (OEQ) and extruded defatted chia (EDC) flours

According to literature reports, extrusion is a benevolent process with the proximal chemical composition and essential amino acid content of grains such as cereals and pseudo-cereals because it is a high-temperature, short-time (HTST) process (Argüelles-López *et al.*, 2018; Gámez-Valdez *et al.*, 2021; León-Murillo *et al.*, 2021; León-López *et al.*, 2019; Reyes-Moreno *et al.*, 2018).

During food processing by extrusion, both an IVPD increase and an IVPD decrease may occur. This increase of IVPD occurs due to the destruction of labile anti-nutritional factors (lectins, trypsin and chymotrypsin inhibitors) and protein denaturation, as a result of the applied conditions (shearing forces, temperature, and humidity) during processing (Argüelles-López *et al.*, 2018; Félix-Medina *et al.*, 2021; León-López *et al.*, 2019; Reyes-Moreno *et al.*, 2018). The decrease of IVPD in extrusion occurs because the components of dietary fiber, phenolic compounds, and the products of the Maillard reaction (reaction carried out between reducing sugars and amino acids of the proteins) can bind to protein, partially unavailing it for its digestion (Reyes-Moreno *et al.*, 2018). However, the increase of IVPD during food processing by extrusion is the most reported phenomenon in literature (León-López *et al.*, 2019), which agrees with the results obtained in the present investigation.

The nutritional parameter C-PER of unprocessed and extruded flours was calculated using the IVPD and the EAA composition of the samples, so its increase was mainly due to the increase in IVPD, since the EAA content only slightly decreased in EDC flour, while in OEQ flour this EAA content did not undergo significant changes ($p > 0.05$) (Félix-Medina *et al.*, 2021; León-López *et al.*, 2019).

ACE inhibitory potential of optimized extruded quinoa (OEQ) and extruded defatted chia (EDC) flours

The IC₅₀ values to ACE inhibitory potential obtained in this research are in concordance with reported results for phenolic extracts of extruded blue maize and extruded defatted chia flour (0.51 extract mg/ml and 0.38 extract mg/ml, respectively) and extruded amaranth flour (0.42 mg/ml) (Argüelles-López *et al.*, 2018; Gámez-Valdez *et al.*, 2021; León-Murillo *et al.*, 2021). During the extrusion of whole quinoa seeds and defatted chia seeds, the release of phytochemicals with biological activity (e.g. ACE inhibitory activity) occurs; in this study, an increase in the content of bound and total phenolic compounds was observed during the production of OEQ and EDC by extrusion (Table 2). Also, during the gastrointestinal digestion other bioactive compounds with ACE inhibition capacity are released. The phytochemicals with ACE inhibitory potential released during both processes above mentioned have the potential to reduce the content of angiotensin II (vasoconstrictor) and increase the levels of Bradykinin (vasodilator) in hypertensive people, resulting in a decrease in their blood pressure. Phenolic compounds (phenolic acids, flavonoids, tannins, stilbenes) can inhibit the *in vitro* ACE activity. In the literature, some hypotheses on the mechanisms of inhibition of phenolic compounds on ACE have been posed: competitive inhibition, by the structure of phenols that alter their function by agglutination; noncompetitive inhibition, both substrate and inhibitor bind to the enzyme simultaneously and reversibly; metal sequestration, ACE is a Zn (II)-dependent metalloproteinase, and phenolic compounds can chelate non-specific metals exhibiting an ACE inhibitory effect; and the interactions between phenols and the disulfide bridges (oxidized cysteines) reside on the surface of the ACE, causing slight modifications in the structure of the ACE (Al Shukor *et al.*, 2013; Gámez-Valdez *et al.*, 2021; León-Murillo *et al.*, 2021). Also, peptides released during enzymatic digestion has shown ACE inhibitory activity. Montoya-Rodríguez *et al.* (2015) reported that extruded amaranth flours were sources of peptides with potential biological activity, such as ACE-inhibitor. The extrusion process had a higher impact on the peptide profile, producing more peptides with lower molecular mass and biological activity. Previous studies have shown that the amino acid composition of a peptide has a significant effect on its ACE inhibitory activity, and it has been reported that ACE prefers to bind to a polypeptide with a high content of hydrophobic amino acids. Hydrophobic peptides can coordinate to the Zn (II) ion. This interaction may have caused distortion of the tetrahedrally coordinated Zn (II), which further resulted in the loss of ACE activity (Sun *et al.*, 2019).

In the literature, it has also been reported that Maillard reaction products (MRP) exhibit ACE inhibitory activity *in vitro*. The *in vitro* ACE inhibitory activity of melanoidins (end products of the Maillard reaction) from food is similar to that reported for well-known antihypertensive peptides. Bounded melanoidin compounds (BMC), e.g., polyphenols, and low molecular weight MRP, which are non-covalently bound to the melanoidin backbone, also contribute to ACE inhibitory activity. On the other hand, it has also been reported that the *in vitro* ACE inhibitory activity of melanoidins is related to the structure of melanoidin, and that the melanoidin chromophores are not involved. Melanoidins are low molecular weight pigments or high molecular weight compounds with low molecular weight chromophore (Wang *et al.*, 2011).

Chemical composition and nutritional/nutraceutical properties of the mixture (70% OEQ + 30% EDC)

TDF is an important functional component in the human diet. It possesses recognized health properties and can reduce the risk of specific chronic diseases, such as diabetes type-2 and coronary heart disease. It has been proven that TDF is an ingredient of commercial functional foods designed for consumers who are concerned about elevated circulating blood glucose and cholesterol levels (Yegin *et al.*, 2020).

According to León-López *et al.* (2019), the C-PER technique predicted the same differences observed in the rat bioassay (PER); both techniques C-PER and PER can detect significant differences among samples of different origin. Therefore, it is recommended to use the *in vitro* techniques as fast, accurate indicators of protein digestibilities and PER.

Formulation, nutrimental composition, energy content, and antioxidant activity of the functional gluten-free beverage (GFB) are prepared from the mixture

The Mexican norm NMX-F-439-1983 for food and non-alcoholic beverages (Secretaría de Comercio y Fomento Industrial, 1983) defines a beverage as nutritional when it contains at least 1.5% protein or hydrolyzed proteins with a quality equivalent to that of casein; beverages must contain between 10% and 25% main ingredient. These may contain about 2% ethanol, carbon dioxide, flavoring agents, juices, fruit pulp, sweeteners, vegetables or legumes, and other additives authorized by the Mexican Ministry of Health.

Conclusions

The extrusion process is an excellent technological strategy that increased the nutritional value, antioxidant activity, total phenolic content, and inhibitory ACE potential of whole quinoa and chia seeds. Extruded defatted chia flour added to extruded whole quinoa flour enhanced nutritional, antioxidant, and antihypertensive potential of the mixture used to produce a functional gluten-free beverage. Therefore, this functional beverage can be used for health promotion and as an alternative to beverages with high caloric content and low nutritional value, which predominate in a market of consumers with significant trends or incidences of overweight/obesity and chronic degenerative diseases such as hypertension. Additionally, the novel functional gluten-free beverage is an excellent alternative to promote healthy living in people with celiac disease.

Acknowledgements

To the Mexican Consejo Nacional de Ciencia y Tecnología (Conacyt), Convocatoria 2019 - Ciencia de Frontera (Grupal) for the funds supporting this research (ID Number: 263352). To Programa de Fomento y Apoyo a Proyectos de Investigación (Profapi) under the Universidad Autónoma de Sinaloa, for supplementary funds.

Conflicts of interest

The authors declare no conflict of interest.

References

- Afrin, S., Giampieri, F., Gasparrini, M., Forbes-Hernández, T. Y., Cianciosi, D., Reboredo-Rodríguez, P., Zhang, J., Manna, P. P., Daglia, M., Atanasov, A. G., & Battino, M. (2020). Dietary phytochemicals in colorectal cancer prevention and treatment: A focus on the molecular mechanisms involved. *Biotechnology Advances*, *38*, 107322. doi: <https://doi.org/10.1016/j.biotechadv.2018.11.011>
- Alarcón-Valdez, C., Milán-Carrillo, J., Cárdenas-Valenzuela, O. G., Mora-Escobedo, R., Bello-Pérez, L. A., & Reyes-Moreno, C. (2005). Infant food from quality protein maize and chickpea: Optimization for preparing and nutritional properties. *International Journal of Food Science and Nutrition*, *56*(4), 273-285. doi: <https://doi.org/10.1080/09637480500146804>.
- Al Shukor, N., Van Camp, J., Gonzales, G. B., Staljanssens, D., Struijs, K., Zotti, M. J., Raes, K., & Smagghe, G. (2013). Angiotensin-converting enzyme inhibitory effects by plant phenolic compounds: A study of structure activity relationships. *Journal of Agricultural and Food Chemistry*, *61*(48), 11832-11839. doi: <https://doi.org/10.1021/jf404641v>
- Angeli, V., Silva, P. M., Massuela, D. C., Khan, M. W., Hamar, A., Khajehei, F., Graeff-Hönninger, S., & Piatti, C. (2020). Quinoa (*Chenopodium quinoa* Willd.): An overview of the potentials of the "golden grain" and socio-economic and environmental aspects of its cultivation and marketization. *Foods*, *9*(29), 216. doi: <https://doi.org/10.3390/foods9020216>
- Argüelles-López, O. D., Reyes-Moreno, C., Gutiérrez-Dorado, R., Sánchez-Osuna, M. F., López-Cervantes, J., Cuevas-Rodríguez, E. O., Milán-Carrillo, J., Perales-Sánchez, J. X. K. (2018). Functional beverages elaborated from amaranth and chia flours processed by germination and extrusion. *Biotecnia*, *20*(3), 135-145. <https://biotecnia.unison.mx/index.php/biotecnia/article/view/721/289>
- Association of Official Analytical Chemists (AOAC). (2012). Official methods of analysis of the Association of Official Analytical Chemists. Methods: 925.09B, 920.39 C, 960.52, and 985.29. AOAC.
- Aziz, A., Akram, N. A., & Ashraf, M. (2018). Influence of natural and synthetic vitamin C (ascorbic acid) on primary and secondary metabolites and associated metabolism in quinoa (*Chenopodium quinoa* Willd.) plants under water deficit regimes. *Plant Physiology and Biochemistry*, *123*, 192-203. doi: <https://doi.org/10.1016/j.plaphy.2017.12.004>
- Caratini, C., & Rosentrater, K. A. (2019). Developing and testing gluten-free spaghetti using quinoa. *Journal of Food Research*, *8*(5), 20-32. doi: <https://doi.org/10.5539/jfr.v8n5p20>
- Cardello, A. V., & Schutz, H. G. (2004). Research note numerical scale-point locations for constructing the LAM (labeled affective magnitude) scale. *Journal of Sensory Studies*, *19*(4), 341-346. doi: <https://doi.org/10.1111/j.1745-459X.2004.tb00152.x>
- Demir, B., & Bilgili, N. (2021). Utilization of quinoa flour (*Chenopodium quinoa* Willd.) in gluten-free pasta formulation: Effects on nutritional and sensory properties. *Food Science and Technology International*, *27*(3), 242-250. doi: <https://doi.org/10.1177/1082013220940092>
- Dewanto, V., Wu, X., & Liu, R. H. (2002). Processed sweet corn has higher antioxidant activity. *Journal of Agricultural and Food Chemistry*, *50*(17), 4959-4964. doi: <https://doi.org/10.1021/jf0255937>
- Fares, C., & Menga, V. (2012). Effects of toasting on the carbohydrate profile and antioxidant properties of chickpea (*Cicer arietinum* L.) flour added to durum wheat pasta. *Food Chemistry*, *131*(4), 1140-1148. doi: <https://doi.org/10.1016/j.foodchem.2011.09.080>
- Félix-Medina, J. V., Montes-Ávila, J., Reyes-Moreno, C., Perales-Sánchez, J. X. K., Gómez-Favela, M. A., Aguilar-Palazuelos, E., & Gutiérrez-Dorado, R. (2020). Second-generation snacks with high nutritional and antioxidant value produced by an optimized extrusion process from corn/common bean flours mixtures. *LWT – Food Science and Technology*, *124*, 109172, doi: <https://doi.org/10.1016/j.lwt.2020.109172>
- Félix-Medina, J. V., Gutiérrez-Dorado, R., López-Valenzuela, J. A., López-Ángulo, G., Quintero-Soto, M. F., Perales-Sánchez, J. X. K., & Montes-Ávila, J. (2021). Nutritional, antioxidant and phytochemical characterization of healthy ready-to-eat expanded snack produced from maize/common bean mixture by extrusion. *LWT – Food Science and Technology*, *142*, 111053, doi: <https://doi.org/10.1016/j.lwt.2021.111053>.
- Food and Agriculture Organization of the United Nations (FAO). (2013). Findings and Recommendations of the 2011 FAO Expert Consultation on Protein Quality Evaluation in Human Nutrition. En FAO Food and Nutrition Paper 92, *Dietary protein quality evaluation in human nutrition: Report of an FAO expert consultation* (pp. 29). FAO. <http://www.fao.org/3/a-i3124e.pdf>

- Gómez-Valdez, L. C., Gutiérrez-Dorado, R., Gómez-Aldapa, C. A., Perales-Sánchez, J. X. K., Milán-Carrillo, J., Cuevas-Rodríguez, E. O., Mora-Rochín, S., & Reyes-Moreno, C. (2021). Effect of the extruded amaranth flour addition on the nutritional, nutraceutical and sensory quality of tortillas produced from extruded creole blue maize flour. *Biotechnia*, 23(2), 103-112. <https://biotecnia.unison.mx/index.php/biotechnia/article/download/1385/561>
- Gómez-Favela, M. A., Gutiérrez-Dorado, R., Cuevas-Rodríguez, E. O., Canizalez-Román, V. A., León-Sicaños, C. R., Milán-Carrillo, J., & Reyes-Moreno, C. (2017). Improvement of chia seeds with antioxidant activity, GABA, essential amino acids, and dietary fiber by controlled germination bioprocess. *Plant Foods for Human Nutrition*, 72, 345-352. doi: <https://doi.org/10.1007/s11130-017-0631-4>
- Gómez-Favela, M. A., Reyes-Moreno, C., Milán-Carrillo, J., Partida-Preciado, R. A., Espinoza-Moreno, R. J., Preciado-Ortiz, R., & Gutiérrez-Dorado, R. (2021). Gluten-free healthy snack with high nutritional and nutraceutical value elaborated from a mixture of extruded underutilized grains (quality protein maize/tepyary bean). *Acta Universitaria*, 31, 1-18. doi: <https://doi.org/10.15174/au.2021>
- Hrncic, M. K., Ivanovski, M., Cör, D., & Knez, Z. (2020). Chia seeds (*Salvia hispanica* L.): An overview - Phytochemical profile, isolation methods, and application. *Molecules*, 25(1), 11. doi: <https://doi.org/10.3390/molecules25010011>
- Instituto Nacional de Salud Pública (INSP). (2016). *Encuesta Nacional de Salud y Nutrición de Medio Camino*. INSP.
- Kamel, M. A., Aly, M. H., & Abd-El-Khalek, M. H. (2020). Preparation of gluten-free corn crackers supplemented with some legume flours. *Journal of Advanced Research in Food Science and Nutrition*, 3(1), 42-49. doi: <https://doi.org/10.24321/2582.3892.202006>
- Kim, J., & Adhikari, K. (2020). Current trends in kombucha: Marketing perspectives and the need for improved sensory research. *Beverages*, 6(1), 15. doi: <https://doi.org/10.3390/beverages6010015>
- Kumar, K. (2018). Role of gluten-free functional foods in the management of celiac disease. *EC Nutrition*, 13, 742-744. https://www.researchgate.net/publication/328858424_Role_of_GlutenFree_Functional_Foods_in_the_Management_of_Celiac_Disease
- Lebwohl, B., & Rubio-Tapia, A. (2021). Epidemiology, presentation, and diagnosis of celiac disease. *Gastroenterology*, 160(1), 63-75. doi: <https://doi.org/10.1053/j.gastro.2020.06.098>
- León-López, L., Reyes-Moreno, C., Ley-Osuna, A. H., Perales-Sánchez, J. X. K., Milán-Carrillo, J., Cuevas-Rodríguez, E. O., & Gutiérrez-Dorado, R. (2019). Improvement of nutritional and nutraceutical value of nixtamalized maize tortillas by addition of extruded chia flour. *Biotechnia*, 21(3), 56-66. <https://biotecnia.unison.mx/index.php/biotechnia/article/view/1012/342>
- León-Murillo, J. R., Gutiérrez-Dorado, R., Reynoso-Camacho, R., Milán-Carrillo, J., Perales-Sánchez, J. X. K., Cuevas-Rodríguez, E. O., & Reyes-Moreno, C. (2021). Tortillas made with extruded flours of blue maize and chia seeds as a nutritious and nutraceutical food option. *Agrociencia*, 55, 487-506. <https://agrociencia-colpos.mx/index.php/agrociencia/article/view/2555/2081>
- Li, L., Lietz, G., & Seal, C. J. (2021). Phenolic, apparent antioxidant and nutritional composition of quinoa (*Chenopodium quinoa* Willd.) seeds. *International Journal of Food Science & Technology*, 56(7), 3245-3254. <https://ifst.onlinelibrary.wiley.com/doi/pdf/10.1111/ijfs.14962>
- Liskova, A., Kubatka, P., Samec, M., Zubor, P., Mlyncek, M., Bielik, T., Samuel, S. M., Zulli, A., Kwon, T. K., & Büsselberg, D. (2019). Dietary phytochemicals targeting cancer stem cells. *Molecules*, 24(5), 899. doi: <https://doi.org/10.3390/molecules24050899>
- Martínez-Cruz, O., & Paredes-López, O. (2014). Phytochemical profile and nutraceutical potential of chia seeds (*Salvia hispanica* L.) by ultrahigh performance liquid chromatography. *Journal of Chromatography A*, 1346, 43-48. doi: <https://doi.org/10.1016/j.chroma.2014.04.007>
- Martínez-López, A., Millán-Linares, M. C., Rodríguez-Martin, N. M., Millán, F., & Montserrat-de la Paz, S. (2020). Nutraceutical value of kiwicha (*Amaranthus caudatus* L.). *Journal of Functional Foods*, 65, 103735. doi: <https://doi.org/10.1016/j.jff.2019.103735>
- Masood, M. A. B. (2022). Chia seeds as potential nutritional and functional ingredients: A review of their applications for various food industries. *Journal of Nutrition Food Science and Technology*, 4(1), 1-14. <https://unisciencepub.com/storage/2022/02/Chia-Seeds-as-Potential-Nutritional-and-Functional-Ingredients.pdf>

- Medina, O. D., Lopes, R. C., Vieira-Queiroz, V. A., Ribeiro, M., Duarte, H. S., & Ribeiro, F. A. (2020). Mixed sorghum and quinoa flour improves protein quality and increases antioxidant capacity *in vivo*. *LWT - Food Science and Technology*, 129, 1-8. doi: <https://doi.org/10.1016/j.lwt.2020.109597>
- Menga, V., Amato, M., Phillips, T. D., Angelino, D., Morreale, F., & Fares, C. (2017). Gluten-free pasta incorporating chia (*Salvia hispanica* L.) as thickening agent: An approach to naturally improve the nutritional profile and the *in vitro* carbohydrate digestibility. *Food Chemistry*, 221, 1954–1961. doi: <https://doi.org/10.1016/j.foodchem.2016.11.151>
- Milán-Carrillo, J., Montoya-Rodríguez, A., Gutiérrez-Dorado, R., Perales-Sánchez, J. X. K., & Reyes-Moreno, C. (2012). Optimization of extrusion process for producing high antioxidant instant amaranth (*Amaranthus hypochondriacus* L.) flour using response surface methodology. *Applied Mathematics*, 3, 1516-1525. doi: <https://doi.org/10.4236/am.2012.330211>
- Montemurro, M., Pontonio, E., & Rizzello, C. G. (2019). Quinoa flour as an ingredient to enhance the nutritional and functional features of cereal-based foods. In V. R. Preedy & R. R. Watson (eds.), *Flour and breads and their fortification in health and disease prevention* (pp. 453-464). Academic Press. doi: <https://doi.org/10.1016/B978-0-12-814639-2.00036-8>
- Montoya-Rodríguez, A., Milán-Carrillo, J., Reyes-Moreno, C., & González, E. (2015). Characterization of peptides found in unprocessed and extruded amaranth (*Amaranthus hypochondriacus*) pepsin/pancreatin hydrolysates. *International Journal of Molecular Sciences*, 16(4), 8536-8554. doi: <https://doi.org/10.3390/ijms16048536>
- Niland, B., & Cash, B. D. (2018). Health benefits and adverse effects of a gluten-free diet in non-celiac disease patients. *Gastroenterology & Hepatology*, 14(2), 82-91. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5866307/>
- Orona-Tamayo, D., Valverde, M. E., & Paredes-López, O. (2017). Chia - The new golden seed for the 21st century: Nutraceutical properties and technological uses. In S. R. Nadathur, J. P. D. Wanasundara & L. Scanlin (eds.), *Sustainable protein sources* (pp. 265-281). Elsevier. doi: <https://doi.org/10.1016/B978-0-12-802778-3.00017-2>
- Orona-Tamayo, D., Valverde, M. E., & Paredes-López, O. (2018). Bioactive peptides from selected Latin American food crops - A nutraceutical and molecular approach. *Critical Reviews in Food Science and Nutrition*, 59(12), 1949-1975. doi: <https://doi.org/10.1080/10408398.2018.1434480>
- Ortiz-Cruz, R. A., Ramírez-Wong, B., Ledesma-Osuna, A. I., Torres-Chávez, P. I., Sánchez-Machado, D. I., Montañón-Leyva, B., López-Cervantes, J., & Gutiérrez-Dorado, R. (2020). Effect of extrusion processing conditions on the phenolic compound content and antioxidant capacity of sorghum (*Sorghum bicolor* L.) Moench) Bran. *Plant Foods for Human Nutrition*, 75, 252-257. doi: <https://doi.org/10.1007/s11130-020-00810-6>
- Pellegrini, M., Lucas-Gonzales, R., Ricci, A., Fontecha, J., Fernández-López, J., Pérez-Álvarez, J. A., & Viuda-Martos, M. (2018). Chemical, fatty acid, polyphenolic profile, techno-functional and antioxidant properties of flours obtained from quinoa (*Chenopodium quinoa* Willd) seeds. *Industrial Crops and Products*, 111, 38-46. doi: <https://doi.org/10.1016/j.indcrop.2017.10.006>
- Pyo, Y. H., & Lee, T. C. (2007). The potential antioxidant capacity and Angiotensin-I converting enzyme inhibitory activity of Monascus-fermented soybean extracts: Evaluation of Monascus-fermented soybean extracts as multifunctional food additives. *Journal of Food Science*, 72(3), S218-S223. doi: <https://doi.org/10.1111/j.1750-3841.2007.00312.x>
- Ramos-Enríquez, J. R., Ramírez-Wong, B., Robles-Sánchez, R. M., Robles-Zepeda, R. E., González-Aguilar, G. A., & Gutiérrez-Dorado, R. (2018). Effect of extrusion conditions and the optimization of phenolic compound content and antioxidant activity of wheat bran using response surface methodology. *Plant Foods for Human Nutrition*, 73, 228–234. doi: <https://doi.org/10.1007/s11130-018-0679-9>
- Reyes-Moreno, C., Reyes-Fernández, P. C., Cuevas-Rodríguez, E. O., Milán-Carrillo, J., & Mora-Rochín, S. (2018). Changes in nutritional properties and bioactive compounds in cereals during extrusion cooking. In S. Z. Qamar (ed.), *Extrusion of metals, polymers, and food products*. IntechOpen. doi: <http://dx.doi.org/10.5772/intechopen.68753>
- Rubavathi, S., Ayyappadasan, G., Sangeetha, N., Harini, T., Saranya, D., & Harshapradha, P. (2020). Studies on antioxidant and anti-obesity activity of *Salvia hispanica* (Chia) seeds extracts. *Journal of Drug Delivery and Therapeutics*, 10(3), 98-106. doi: <https://doi.org/10.22270/jddt.v10i3-s.4169>

- Salas-López, F., Gutiérrez-Dorado, R., Milán-Carrillo, J., Cuevas-Rodríguez, E. O., Canizalez-Roman, V. A., León-Sicairos, C. R., & Reyes-Moreno, C. (2018). Nutritional and antioxidant potential of a desert underutilized legume-tepary bean (*Phaseolus acutifolius*). Optimization of germination bioprocess. *Food Science and Technology*, 38(1), 254-262. doi: <https://doi.org/10.1590/fst.25316>
- Satterlee, L. D., Kendrick, J. G., Marshall, H. F., Jewell, D. K., Ali, R. A., Heckman, M. M., Steinke, H. F., Larson, P., Phillips, D., Sarwar, G., & Slump, P. (1982). *In vitro* assay for predicting protein efficiency ratio as measured by rat bioassay: Collaborative study. *Journal of the Association of Official Analytical Chemists*, 65(4), 798-809. doi: <https://doi.org/10.1093/jaoac/65.4.798>
- Secretaría de Comercio y Fomento Industrial (1983). Norma Mexicana NMX-F-439-1983. Alimentos - Bebidas No Alcohólicas - Bebidas y Refrescos - Clasificación y Definiciones. https://caisatech.net/uploads/XXI_2_MXD_C10_NMX-F-439-1983_R0_11FEB1983.pdf
- Sharma, N., Bhatia, S., Chunduri, V., Kaur, S., Sharma, S., Kapoor, P., Kumari, A., & Garg, M. (2020). Pathogenesis of celiac disease and other gluten related disorders in wheat and strategies for mitigating them. *Frontiers in Nutrition*, 7. doi: <https://doi.org/10.3389/fnut.2020.00006>
- Song, J., Shao, Y., Chen, X., & Li, X. (2020). Release of characteristic phenolics quinoa based on extrusion technique. *Food Chemistry*, 374, 128780. doi: <https://doi.org/10.1016/j.foodchem.2020.128780>
- Sun, S., Xu, X., Sun, X., Zhang, X., Chen, X., & Xu, N. (2019). Preparation and identification of ACE inhibitory peptides from the marine macroalga *Ulva intestinalis*. *Marine Drugs*, 17(3), 179. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6471128/>
- Tan, M., Chang, S., Liu, J., Li, H., Xu, P., Wang, P., Wang, X., Zhao, M., Zhao, B., Wang, L., & Zhao, Q. (2020). Physicochemical properties, antioxidant and antidiabetic activities of polysaccharides from Quinoa (*Chenopodium quinoa* Willd.) seeds. *Molecules*, 25(17), 3840. doi: <https://doi.org/10.3390/molecules25173840>
- Tang, Y., & Tsao, R. (2017). Phytochemicals in quinoa and amaranth grains and their antioxidant, anti-inflammatory, and potential health beneficial effects: A review. *Molecular Nutrition & Food Research*, 61(7), 1600767. doi: <https://doi.org/10.1002/mnfr.201600767>
- Tolun, A., & Altintas, Z. (2019). Medicinal properties and functional components of beverages. En A. M. Grumezescu & A. M. Holban (eds.), *Functional and medicinal beverages* (pp. 235-284). Woodhead Publishing. doi: <https://doi.org/10.1016/B978-0-12-816397-9.00007-8>
- Tovar-Pérez, E. G., Guerrero-Legarreta, I., Farrés-González, A., & Soriano-Santos, J. (2009). Angiotensin I-converting enzyme-inhibitory peptide fractions from albumin 1 and globulin as obtained of amaranth grain. *Food Chemistry*, 116(2), 437-444. doi: <https://doi.org/10.1016/j.foodchem.2009.02.062>
- United States Department of Agriculture [USDA]. (2010). *Antioxidants and health*. ACES publications.
- Wang, H., Qian, H., & Yao, W. (2011). Melanoidins produced by the Maillard reaction: Structure and biological activity. *Food Chemistry*, 128(3), 573-584. doi: <https://doi.org/10.1016/j.foodchem.2011.03.075>
- Wieser, H., Ruiz-Carnicer, A. R., Segura, V., Comino, I., & Sousa, C. (2021). Challenges of monitoring the gluten-free diet adherence in the management and follow-up of patients with celiac disease. *Nutrients*, 13(7), 2274. doi: <https://doi.org/10.3390/nu13072274>
- Woomer, J. S., & Adedeji, A. A. (2021). Current applications of gluten-free grains—a review. *Critical Reviews in Food Science and Nutrition*, 61(1), 14-24. doi: <https://doi.org/10.1080/10408398.2020.1713724>
- Yegin, S., Kopec, A., Kitts, D. D., & Zawistowski, J. (2020). Dietary fiber: A functional food ingredient with physiological benefits. En H. G. Preuss & D. Bagchi (eds.), *Dietary sugar, salt and fat in human health* (pp. 531-555). Academic Press. doi: <https://doi.org/10.1016/B978-0-12-816918-6.00024-X>
- Yolmeh, M., & Jafari, S. M. (2017). Applications of response surface methodology in the food industry processes. *Food and Bioprocess Technology*, 10, 413-433. doi: <https://doi.org/10.1007/s11947-016-1855-2>