

The Collection of Benefic and Entomophagous Insects (CIBE) of the Universidad Autónoma de Nuevo León (UANL), México: A collection focused on parasitoids Hymenoptera

La Colección de Insectos Benéficos y Entomófagos (CIBE) de la Universidad Autónoma de Nuevo León (UANL), México: Una colección enfocada a Hymenoptera parasitoides



Acta Zoológica Mexicana (nueva serie)

¹ IRAM P. RODRÍGUEZ-SÁNCHEZ, ² KENZY I. PEÑA-CARRILLO, ¹ MAYRA A. GÓMEZ-GOVEA, ³ MANUEL DE LUNA, ^{1*} MARÍA DE LOURDES RAMÍREZ-AHUJA

*Corresponding author:

^{ID} María de Lourdes Ramírez-Ahuja
lulu.ahuja@hotmail.com

How to cite:

Rodríguez-Sánchez, I. P., Peña-Carrillo, K.I., Gómez-Govea, M. A., de Luna, M., Ramírez-Ahuja, M.L. (2024) The Collection of Benefic and Entomophagous Insects, (CIBE) of the Universidad Autónoma de Nuevo León (UANL), México: A collection focused on parasitic Hymenoptera. *Acta Zoológica Mexicana (nueva serie)*, 40, 1–13.

10.21829/azm.2024.4012715
elocation-id: e4012715

Received: 27 August 2024

Accepted: 20 November 2024

Published: 31 December 2024

¹ Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de Fisiología Molecular y Estructural, Av. Universidad s/n Cd. Universitaria, San Nicolás de los Garza, N.L. 66455, México.

² INIFAP, Campo Experimental General Terán, km 31 carretera Montemorelos-China, 67400, General Terán, N.L. México.

³ Universidad Autónoma de Nuevo León, Facultad de Ciencias Forestales, Laboratorio de Entomología Forestal, Linares, N.L. 67700 México.

Responsible editor: Magdalena Cruz Rosales

ABSTRACT. The Collection of Benefic and Entomophagous Insects, CIBE is an entomological collection held at the Facultad de Ciencias Biológicas (FCB) of the Universidad Autónoma de Nuevo León (UANL) which specializes in parasitoid wasps (order Hymenoptera). This collection holds 105,000 specimens of the superfamilies Ceraphronoidea, Chalcidoidea, Cynipoidea, Evanoidea, Ichneumonoidea,

Megalyroidea, Mymarommatoidea, Platygastroidea, Proctotrupoidea, Stephanoidea and Trigonalioidea. Among the best represented families are Braconidae with 123 genera and 23 species, Ichneumonidae with 52 genera and 59 species, Encyrtidae with 100 genera and 29 species, and Eulophidae with 53 genera and 65 species. In the CIBE there are ten paratypes of the species: *Triaspis eugenii* Wharton & López-Martínez 2000, *Retusigaster albopilosus* 2003 Mercado, *Retusigaster noguerai* 2003 Mercado (Braconidae), *Itopectis gonzalezi* Kasparyan 2007, *Itopectis nigrithorax* Kasparyan 2007, *Itopectis specularis* Kasparyan 2007, *Iseropus hylesiae* Kasparyan 2006 (Ichneumonidae), *Aranobroter rayorae* LaSalle 1990, *Baryscapus uetzi* LaSalle 1990 (Eulophidae), and *Spalangia imitator* Gibson 2009 (Pteromalidae).

Key words: Scientific collections; parasitoid wasps; biologic control

RESUMEN. La Colección de Insectos Benéficos y Entomófagos, CIBE, es una colección entomológica que se encuentra en la Facultad de Ciencias Biológicas (FCB) de la Universidad Autónoma de Nuevo León (UANL) y que se especializa en avispa parasitoides (orden Hymenoptera). Esta colección alberga 105,000 ejemplares de las superfamilias Ceraphronoidea, Chalcidoidea, Cynipoidea, Evanoidea, Ichneumonoidea, Megalyroidea, Mymarommatoidea, Platygastroidea, Proctotrupoidea, Stephanoidea y Trigonalioidea. Entre las familias mejor representadas se encuentran Braconidae con 123 géneros y 23 especies, Ichneumonidae con 52 géneros y 59 especies, Encyrtidae con 100 géneros y 29 especies y Eulophidae con 53 géneros y 65 especies. En CIBE existen diez paratipos de las especies: *Triaspis eugenii* Wharton & López-Martínez 2000, *Retusigaster albopilosus* 2003 Mercado, *Retusigaster noguerai* 2003 Mercado (Braconidae), *Itopectis gonzalezi* Kasparyan 2007, *Itopectis nigrithorax* Kasparyan 2007, *Itopectis specularis* Kasparyan 2007, *Iseropus hylesiae* Kasparyan 2006 (Ichneumonidae), *Aranobroter rayorae* LaSalle 1990, *Baryscapus uetzi* LaSalle 1990 (Eulophidae) y *Spalangia imitator* Gibson 2009 (Pteromalidae).

Palabras clave: Colecciones científicas; avispa parasitoides; control biológico

INTRODUCTION

Biological collections represent the natural heritage of a country or region (Suárez-Mayorga & Vivas, 2003, Montaña *et al.*, 2012, Trujillo-Trujillo *et al.*, 2014); and their increase, through the collection of specimens, supports the knowledge of the biodiversity (Suárez-Mayorga & Vivas, 2003). Biological collections allow us to document the richness and distribution of species, which is essential for any study on taxonomy, systematics, ecology and biogeography, and essential for the conservation of species (Andrade *et al.*, 2013). In Mexico, there are several entomological collections, the portal of the National Commission for the Knowledge and Use of Biodiversity (CONABIO) reports at least 38 entomological collections. One of the largest entomological collections in the country is the Colección Nacional de Insectos (CNIN) del Instituto de Biología de la Universidad Nacional Autónoma de México (UNAM), which has approximately five million dry-mounted specimens (CONABIO, 2023). It includes the orders: Coleoptera, Dermaptera, Diptera, Embioptera, Hemiptera-Heteroptera, Homoptera, Hymenoptera, Lepidoptera, Odonata, Orthoptera, Psocoptera, Thysanoptera and Trichoptera. Another important collection is the Entomological Collection of the Institute of Ecology (IEXA) of Xalapa, Veracruz, which is made up of 419,535 specimens (INECOL, 2024).

The CIBE is the acronym for the Colección de Insectos Benéficos y Entomófagos, it was started in 1986 by Dr. Alejandro González Hernández with the support of projects from the Ministry of Public Education, the National Council of Science and Technology, the National Commission for Knowledge and Use of Biodiversity, University of California (UC MEXUS), Texas A&M University-CONACYT and the Scientific and Technological Research Support Program (PAICYT) of the Autonomous University of Nuevo León (CONABIO, 2023). It is a collection specialized in parasitoid wasps (Hymenoptera), particularly, those in the superfamilies Ceraphronoidea, Chalcidoidea, Cynipoidea, Evanioidea, Ichnemionoidea, Megalyroidea, Mymarommatoidea, Platygastroidea, Proctotrupoidea, Stephanoidea and Trigonalioidea (Huber, 2017). These superfamilies are of great importance in many fields because they are parasitoids of other insects, that is, they need a host to develop (Fig. 1). Several of these families include species that are considered beneficial because they act as biological control of insect pests of importance to agriculture. In fact, some countries mass-breed and release these organisms for integrated pest management (Simaz *et al.*, 2023).




Figure 1. Life cycle of the wasp *Tamarixia triozae* (Chalcidoidea: Eulophidae) parasitoid of *Bactericera cockerelli* (Hemiptera: Triozoidae) or the potato psyllid. The adult wasp oviposits on its host; later, larvae emerge from the eggs and feed on its hemolymph until completing their life cycle, which results in the death of the host.

The importance of the CIBE lies in the documentation, recording and preservation of the biodiversity of Mexican parasitoid wasps; in addition, it constitutes a reference collection for both national and international specialists. The collection has type specimens and provides parasitoid identification services to research institutions, private industry, as well as undergraduate and

graduate students. According to our knowledge, to date, the CIBE is the largest and best kept collection of parasitoid wasps in Mexico.

MATERIALS AND METHODS

The dry and slide-mounted specimens (as well as those kept in ethanol) deposited in the CIBE were revisited and counted. The dry-mounted material is stored in metal drawers (Fig. 2A–B) and arranged in entomological boxes with airtight glass lids (Fig. 2C). Each specimen contains a label with the locality data, a second label with the host data, and a third one with the taxonomic determination. The material that is in slides is found in plastic boxes and each slide includes collection data, determinant (who identified it) and the CIBE reference key (Fig. 3). The material preserved in ethanol (75%) is stored in refrigerators at a temperature of 4°C, the specimens are separated by families in 2 mL microtubes. Each tube has an internal label with the CIBE key and an external one with the replica of said key (Fig. 4).




Figure 2. A, B) Metal drawers where the entomological boxes of dry-mounted material are stored. C) Inside the drawers, the insects are arranged in entomological boxes with airtight glass lids.



Figure 3. A) Slides in plastic boxes. B) Detail of labeling of the individual slides and the arrangement of the specimens.



Figure 4. A) Refrigerators for preserving insects in alcohol. B) Plastic microtubes with specimens.

The curatorial processes of the dry-mounted specimens follow the process of Borror & DeLong (1971). Which basically consists of dehydrating the specimens in alcohol at increasing concentrations (70%, 80%, 90% and 96%), then they are placed in amyl acetate for 30 minutes and then on paper for drying. Finally, the specimens are mounted in triangles and labeled. The specimens prepared in permanent mounting on slides follow the standardized techniques for micro Hymenoptera recommended by Castro (1996) and Noyes (1982). It basically consists of permanent preparations in Canada balsam.

RESULTS

Currently the collection has more than 105,000 specimens. The collection represents 40 families of parasitoid wasps, which have been collected in practically all states of the country. Although there is also material from countries such as Argentina [*Tamarixia radiata* (Waterston)], Brazil [*Tamarixia radiata*], Colombia [*Diaphorencyrtus aligharensis* (Shafee, Alam and Argarwal) and *T. radiata*], Ecuador (*Euplectrus* Westwood), USA (*Diaphorencyrtus aligarhensis*, *Euplectrus* spp., *T. dahlsteni* Zuparko, *T. radiata*, and *T. schina* Zuparko), Pakistan (*T. radiata*), Trinidad and Tobago (*Ixodiphagus* Howard) and Uruguay [*Melittobia acasta* (Walker) and *Sceliphron asiaticum* (Linnaeus)].




Figure 5. Genera, families and superfamilies of Hymenoptera. The numbers next to the family represent the genera that it has in CIBE.

The collection is organized into superfamilies, in the first is the superfamily Ichneumonoidea with its families Braconidae and Ichneumonidae, Braconidae is organized by genera and species; these are arranged in alphabetical order. In another drawer the superfamily

Iseropus hylesiae Kasparyan (Ichneumonidae), *Aranobroter rayorae* LaSalle, *Baryscapus uetzi* LaSalle (Eulophidae), and *Spalangia imitator* Gibson (Pteromalidae).

Superfamily: Ichneumonoidea

Family: Braconidae

***Triaspis eugenii* Wharton & López-Martínez, 2000**

Paratypes. Mexico, Nayarit, Baladero, 02-II-1997. E. Mariscal (1♀; 1♂); Host: No hosts. Additional data: Malaise trap, Drawer 1, box II (CIBE). They do not have geographic location and altitude information.

***Retusigaster albopilosus* Mercado, 2003**

Paratypes. Mexico, Yucatan, Xmatkuil, 04-08-VI-1996. F. Leon (1♀); Mexico, Yucatan, Xmatkuil, 18-28-VI-1996. H. Delfin & F. Leon (1♂); Host: No hosts. Additional data: Malaise Trap, Drawer 1, box II (CIBE). They do not have geographic location and altitude information.

***Retusigaster noguera* Mercado, 2003**

Paratype. Mexico, Jalisco, Biología Chamela Station, 04-05-VII-1993. Wharton & Sharkey (1♀; 1♂). Additional data: Malaise trap, Drawer 1, box II (CIBE). They do not have geographic location and altitude information.

Superfamily: Ichneumonoidea

Family: Ichneumonidae

***Itoplectis gonzalezi* Kasparyan, 2007**

Paratype. Mexico, Chiapas, San Cristobal, Reserva Cerro El Huitepec, 02-14-VIII-1997. A. González-Hernández (1♂); Host: No hosts. Additional data: Malaise Trap, 7900 msnm, 16° 46'06'' N, 92° 41'04'' W. Drawer 1, box III (CIBE 97-072).

***Itoplectis nigrithorax* Kasparyan, 2007**

Paratype. Mexico, Chiapas, San Cristobal, El Huitepec Hill Reserve, 02-14-VIII-1997. A. González-Hernández (1♀); Host: No hosts. Additional data: Malaise Trap, 7900 msnm, 16° 46'06'' N, 92° 41'04'' W Drawer 1, box III (CIBE 97-072).

***Itoplectis specularis* Kasparyan, 2007**

Paratype. Mexico, Chiapas, San Cristobal, El Huitepec Hill Reserve, 02-14-VIII-1997. A. González-Hernández (1♀); Host: No hosts. Additional data: Malaise Trap, 7900 msnm, 16° 46'06'' N, 92° 41'04'' W Drawer 1, box III (CIBE 97-072).

***Iseropus hylesiae* Kasparyan, 2006**

Paratype. Mexico, Chiapas, San Cristobal, El Huitepec Hill Reserve, 17-II-05. Y. Balam-Ballote (1♀); Host: Ex nidos de *Hylesia* sp (Lepidoptera: Saturniidae), sobre *Arbutus xalapensis* (Ericaceae). Box # 14. Additional data: Drawer 1, box III. They do not have geographic location and altitude information.

Superfamily: Chalcidoidea

Family: Eulophidae

***Aranobroter rayorae* LaSalle, 1990**

Paratype. Mexico, Veracruz, Fortin de las Flores, 1000m, 27-VIII-1983. G. W. Uetz (4♀); Host: From eggs sacs *Metepeira incrassata* (Araneidae). Additional data: Drawer 3, box VIII. They do not have geographic location and altitude information.

***Baryscapus uetzi* LaSalle, 1990**

Paratype. Mexico, Veracruz, Fortin de las Flores, 1000m, -XI-1988. R. S. Rayer (1♀); Host: From eggs sacs *Metepeira incrassata* (Araneidae). Additional data: Drawer 3, box VIII. They do not have geographic location and altitude information.

Superfamily: Chalcidoidea

Family: Pteromalidae

***Spalangia imitator* Gibson, 2009**

Paratypes. Mexico, Michoacan, La Posta, Tarimbaro, 02-X-1987. S. Diaz F. (1♀); Tamaulipas, Cd. Victoria, Valle del Libramiento 20-VII-1987. C. Hernández (1♀); Tamaulipas, Cd. Victoria, Cañon del Novillo, 22-V-1987. A. González (1♂); Host: No data. Additional data: Drawer 4, box I. They do not have geographic location and altitude information.

Publications. Since its creation to date, the CIBE has served as a reference for the making of taxonomic catalogs and identification tools such as: Illustrated catalog of the parasitic Hymenoptera of Mexico (González-Hernández, 1997); illustrated catalog of Mexican Braconidae (Hymenoptera: Ichneumonoidea) (González-Hernández *et al.*, 2003 a,b); Identification and distribution of the genera of Encyrtidae (Hymenoptera: Chalcidoidea) in Mexico (González-Hernández & Woolley, 2001). State studies have been carried out such as the Ichneumonoidea Wasps work in Veracruz (González-Hernández *et al.*, 2011). Taxonomic keys such as Mymaridae, Scelionidae have been published to recognize the species of the different families and, in addition, CIBE has contributed to the training of specialists in various families of Hymenoptera such as the family Aphelinidae, Bethylidae, Encyrtidae, Mymaridae, Trichogrammatidae, Scelionidae, Signiphoridae, Braconidae, Ichneumonidae, etc.

Regarding the publication of scientific articles, since its creation, CIBE has served as a source of information and reference for the publication of various articles, focused on the biological control of a wide variety of pests of agricultural importance. For example, "Species of *Trichogramma* Westwood (Hymenoptera: Trichogrammatidae) present in breeding centers in Mexico" (García-González *et al.*, 2005). According to Van Lenteren & Bueno (2003), in our country, *Trichogramma* is released in 1.5 million hectares annually, so having reference specimens is of utmost importance to guarantee the certainty of the species that are reproducing.

On the other hand, research has also been carried out based on the records of the collection in order to provide candidates for use as biological control agents against pests of possible invasion in our country. As an example, given the high risk of invasion by the stink bug *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae) which invaded the United States of America. In Mexico, studies were carried out to search for and determine parasitoids associated with stink bugs of the Pentatomidae family, which could represent potential biocontrol agents against this species of polyphagous bug (Ramírez-Ahuja *et al.*, 2019).

In 2009, when *Diaphorina citri* (Kuwayama) (Hemiptera: Liviidae) invaded the country, its associated native parasitoids were studied, and the genetic variability of wild populations of *Tamarixia radiata* (its main biocontrol agent) was also characterized. This study was carried out in order to search for races or populations of *T. radiata* that are more aggressive against *D. citri* (Peña-Carrillo *et al.*, 2014; Peña-Carrillo *et al.*, 2015).

During 2018, work was carried out on the determination of the parasitoids and hyperparasitoids of the yellow sorghum aphid *Melanaphis sacchari* (Zehntner) (Hemiptera: Aphididae) in the sorghum crop (Payán-Arzapalo *et al.*, 2018; García-González *et al.*, 2018). Since *M. sacchari* was introduced into the country in 2013, this pest has spread to practically all sorghum-producing states, causing losses of 30% to 100% in sorghum cultivation (Rodríguez del Bosque & Terán, 2015). Accordingly, in areas where the yellow sorghum aphid occurs in Mexico, the species *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae) is released for its control. Although *C. carnea* is the only species that is released against *M. sacchari*, at least eight other species of Chrysopidae reproduce in Mexico which represent potential biocontrol agents for said pest (Arredondo-Bernal, 2019). Chrysopidae egg collections were recently carried out in Sinaloa, where two species of *Telenomus* Haliday that parasitize Neuroptera eggs were found (Ramírez-Ahuja *et al.*, 2020). Therefore, the activity of the parasitoid on Chrysopidae populations must be considered in biological control and integrated pest management programs.

In another work, the parasitoid species associated with the American leaf miner, *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae), were identified in Baja California Sur during 2017 (Holguín-Peña *et al.*, 2019). In another study, the parasitoid *Ganaspis brasiliensis* Ihering (Hymenoptera: Figitidae) was reported for the first time in the country, in Michoacán, parasitizing *Drosophila suzukii* Matsumura (Diptera: Drosophilidae), one of the main pests of strawberries in Mexico. Additionally, in this study, other parasitoid species such as *Leptopilina bouvardi* (Barbotin, Carton, & Kelner-Pillault) (Hymenoptera: Figitidae), *Pachycrepoideus vindemmiae* (Rondani) (Hymenoptera: Pteromalidae) and *Trichopria drosophilae* (Perkins) (Hymenoptera: Diapriidae) were reported attacking this fly species (Toledo-Hernández *et al.*, 2021).

Among the CIBE collection, the parasitoid wasps from the "El Edén" reserve are also registered at the family level, which has contributed to the list of the Diversity of Chalcididae (Hymenoptera: Chalcidoidea) in protected natural areas of the state of Yucatán (Chay-Hernández *et al.*, 2009).

Perspectives. Currently, the CIBE has included the molecular identification of species through the use of the barcoding technique or barcode of life. To date, of the species deposited in the collection, only 5% have a barcode. In the future, it is planned to have all the species in the collection identified morphologically, and for each one to have its barcoding sequences registered in databases such as GenBank and BoldSystems (http://www.boldsystems.org/index.php/MAS_Management_UserConsole) for universal availability (Boldsystem, 2024). The possibility of digitizing the content of the collection is also being considered, since previously there was a server on which photographs and data of the species were available.

Due to the importance of native bees, a new line of research has been created at CIBE focused on the study of these species. Stingless bees of the Meliponini tribe are native bees in Mexico and are classified as threatened due to habitat loss, changes in land use, indiscriminate use of insecticides, among other factors. This new line of research consists of determining the profile of amino acids and acylcarnitines (metabolomes), identifying the bacteria found in the midgut of these species (microbiome) and knowing the miRNome of these species. With these objectives, the aim is to know and contribute to the preservation of native bees.

If we want our collections to be considered a permanent source of information and national heritage, they must be properly cared for and maintained. With the care, management and conservation of specimens from biological collections and their associated data, a wide variety of

research can be carried out that leads us to know the biodiversity of the planet. The richness of a scientific collection is found in the information it contains.

ACKNOWLEDGMENTS. We would like to thank all the students, professors and researchers who in some ways have contributed at the CIBE. To the institutions and researchers who have donated specimens to us for preservation. The corresponding author greatly thanks Dr. Alejandro González Hernández for all his teachings and for introducing her to the wonderful world of parasitoids. Both MLRA and MdL thank CONAHCYT for their postdoctoral research grants.

LITERATURE CITED

- Arredondo-Bernal, H. C. (2019) Directorio de Laboratorios Reproductores y Comercializadores de Agentes de Control Biológico. Available online:
<https://sivico.cnf.gob.mx/ContenidoPublico/10%20Material%20de%20Consulta/Literatura/Especies%20ex%C3%B3ticas%20invasoras/DIRECTORIODELABORATORIOS.pdf>
(Accessed on 20 october 2023).
- Andrade, M. G., Henao E. R., Triviño, P. (2013) Técnicas y procesamiento para la recolección, preservación y montaje de mariposas en estudios de biodiversidad y conservación (Lepidóptera: Hesperoidea – Papilionoidea). *Revista de la Academia Colombiana de Ciencias*, 37 (144), 311–325.
- BOLDSYSTEMS. (2024) Código de barras de la vida Boldsystems. Available online:
http://www.boldsystems.org/index.php/Login/page?destination=MAS_Management_User Console (Accessed on 18 october 2023).
- Borror, D. J., DeLong, D. M. (1971) An introduction to the study of insects (No. Ed. 3). New York, USA.
- Castro, L. (1996) Notas sobre la recogida y preparación de Himenópteros. *Boletín de la Sociedad Entomológica Aragonesa*, 15 (1), 47–52.
- Chay-Hernández, D., Delfin H., Meléndez-Ramírez V., González-Hernández A. (2009) Diversity of Chalcididae (Hymenoptera: Chalcidoidea) in protected natural areas of the state of Yucatán. *Entomología Mexicana*, 8, 176–181.
- CONABIO (2023) Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México. Available online:
<https://www.biodiversidad.gob.mx/fichas-conabio-war/resources/coleccion/61#:~:text=Descripci%C3%B3n%20general%3A%20La%20Colecci%C3%B3n%20IEXA,Ley%20General%20de%20Bienes%20Nacionales> (Accessed on 04 july 2023).
- García-González, F., González-Hernández, A., España-Luna, M. P. (2005) Especies de *Trichogramma* Westwood (Hymenoptera: Trichogrammatidae) presentes en centros reproductores de México. *Acta Zoológica Mexicana (n.s.)*, 21 (3), 125–135.
- García-González, F., Ramírez-Delgado, M., Cortez-Madrigal, H., Ramírez-Ahuja, M. L. (2018) New reports of parasitoids and hyperparasitoids of *Melanaphis sacchari* (Zehntner) in Mexico. *Southwestern Entomologist*, 43 (3), 787–790.
<https://doi.org/10.3958/059.043.0324>
- Gibson, G. A. (2009) Revision of new world spalangiinae (Hymenoptera: Pteromalidae). *Zootaxa*, 2259 (1), 1-159.
<https://doi.org/10.11646/zootaxa.2259.1.1>

- González-Hernández A. (1997) Catálogo ilustrado de Hymenoptera: Parasítica de Mexico. CONABIO-Universidad Autónoma de Nuevo León, pp. 40
- González-Hernández, A., Lomelí-Flores, J. R., Ruíz-Cancino, E. (2011) Avispas Ichneumonoidea (Insecta: Hymenoptera). CONABIO. La Biodiversidad en Veracruz. Estudio de estado, pp. 441–448.
- González-Hernández, A., Wharton, R. A., Sánchez García, J. A., López-Martínez, V., Lomelí-Flores, R., Figueroa, I. (2003a) Catálogo ilustrado de Braconidae (Hymenoptera: Ichneumonoidea) en México. Available online:
<http://www.conabio.gob.mx/institucion/proyectos/resultados/FichapubAP008.pdf>
- González-Hernández, A., Wharton, R.A., Sánchez-García, J.A., López-Martínez, V., Lomelí-Flores, J. R., Figueroa-De La Rosa, J. I., Delfín González, H. (2003b) Catálogo ilustrado de Braconidae (Hymenoptera: Ichneumonoidea) de Mexico. Universidad Autónoma de Nuevo León-CONABIO – CONACyT. CD Interactivo.
- González-Hernández, A., Woolley, J. B. (2001) Identificación y distribución de los géneros de Encyrtidae de México. Available online:
<http://www.conabio.gob.mx/institucion/proyectos/resultados/FichapubU039.pdf>
- Holguín-Peña, R. J., Ramírez-Ahuja, M. L., Medina-Hernández, D., Torres, R. E. P., & Servín-Villegas, R. (2019) Parasitoid wasps associated with *Liriomyza trifolii* mortality in pepper at Baja California Sur, Mexico. *Southwestern Entomologist*, 44 (4), 867–875.
<https://doi.org/10.3958/059.044.0402>
- Huber, J. T. (2017) Biodiversity of Hymenoptera. In *Insect biodiversity: science and society*. pp. 419–461. Robert G. Foottit and Peter H. Adler (Eds).
<https://doi.org/10.1002/9781118945568.ch12>
- INECOL (2024) Instituto de Ecología A. C. Available online:
<https://www.inecol.mx/index.php/biblioteca/coleccion/coleccion-entomologica-iexa>
(Accessed on 16 november 2024).
- Kasparyan, D. R. (2006) A new species of *Iseropus* Foerster from Mexico (Hymenoptera: Ichneumonidae). *Zoosystematica Rossica*, 14 (2), 210.
- Kasparyan, D. R., Êàññäðŷŷŷ, Ä. Ð. (2007) Review of Mexican species of the genus *Itopectis* Foerst. (Hymenoptera: Ichneumonidae: Pimplinae) with description of four new species. *Russian Entomological Journal*, 16(1), 109–114.
- LaSalle, J. (1990) Tetrastichinae (Hymenoptera: Eulophidae) associated with spider egg sacs. *Journal of Natural History*, 24 (6), 1377–1389.
- Mercado, I., Wharton, R. A. (2003) Mexican cardiochiline genera (Hymenoptera: Braconidae), including a preliminary assessment of species-groups in *Toxoneuron* Say and *Retusigaster* Dangerfield, Austin and Whitfield. *Journal of Natural History* 37(7): 845–902.
<https://doi.org/10.1080/00222930110097167>
- Montaño, M., Meza, A. M., Dias, L. G. (2012) La colección entomológica CEBUC y su potencial como colección de referencia de insectos acuáticos. *Boletín científico. Centro de Museos. Museo de Historia Natural*, 16(2), 173–184.
- Noyes, J. S. (1982) Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History*, 16(1), 315–334.
- Payán-Arzapalo, M. A., Catzim, C. E. A., Luque, R. G., Liera, J. E. G., Juárez, M. G. Y., Ramírez-Ahuja, M. L. (2018) Parasitism and Hyperparasitism in *Melanaphis sacchari* Zehntner in sorghum in Mexico. *Southwestern Entomologist*, 43(2), 433–437.
<https://doi.org/10.3958/059.043.0214>

- Peña-Carrillo, K. I., González-Hernández, A., Lozano-Contreras, M. G., Jasso-Argumedo, J., López-Arroyo, J. I. (2014) Parasitoides de *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) y caracterización de haplotipos de *Tamarixia radiata* (Waterston) (Hymenoptera: Eulophidae) en Yucatán, México. *Vedalia*, 15, 16–24.
- Peña-Carrillo, K. I., González-Hernández, A., López-Arroyo, J. I., Mercado-Hernández, R., Favela-Lara, S. (2015) Morphological and genetic variation in Mexican wild populations of *Tamarixia radiata* (Hymenoptera: Eulophidae). *Florida Entomologist*, 98 (4), 1093–1100.
- Ramírez-Ahuja, M. L., López-Arroyo, J. I., Rodríguez-Sánchez, I. P., González-Hernández, A. (2019) Hymenoptera parasitoids associated with stink bugs in Mexico. *Southwestern Entomologist*, 44 (1), 35–49.
<https://doi.org/10.3958/059.044.0104>
- Ramírez-Ahuja, M. L., Garza-González, E., Talamas, E. J., Gómez-Govea, M. A., Rodríguez-Pérez, M. A., Zambrano-Robledo, P., Rodríguez-Sánchez, I. P. (2020) Parasitoids of Chrysopidae eggs in Sinaloa Mexico. *Insects*, 11 (12), 849.
<https://doi.org/10.3390/insects11120849>
- Rodríguez-del-Bosque, L. A., Terán, A. P. (2015) *Melanaphis sacchari* (Hemiptera: Aphididae): A new sorghum insect pest in Mexico. *Southwestern Entomologist*, 40 (2), 433–434.
<https://doi.org/10.3958/059.040.0217>
- Simaz, O., Michaelson, J., Wilson, J. K., Talamas, E., Gut, L., Pote, J., Szűcs, M. (2023) Field releases of the exotic parasitoid *Trissolcus japonicus* (Hymenoptera: Scelionidae) and survey of native parasitoids attacking *Halyomorpha halys* (Hemiptera: Pentatomidae) in Michigan. *Environmental Entomology*, 52 (6), 998–1007.
<https://doi.org/10.1093/ee/nvad102>
- Suárez-Mayorga, A.M., A.J. Vivas. (2003) Guía para la evaluación de software para la administración de colecciones biológicas, versión preliminar. Instituto de investigación de recursos biológicos Alexander von Humboldt. pp 28, Bogotá, Colombia.
- Toledo-Hernández, R. A., Martínez, F., Ramírez-Ahuja, M. L., Sánchez, A., Rodríguez, D., Driskell, A., Buffington, M. (2021) The description of an efficient trap for monitoring *Drosophila suzukii* parasitoids in organic soft fruit crops, and a new record of *Ganaspis brasiliensis* (Ihering) (Hymenoptera: Figitidae) from Michoacan, Mexico. *Proceedings of the Entomological Society of Washington*, 123 (1), 230–243.
<https://doi.org/10.4289/0013-8797.123.1.230>
- Trujillo-Trujillo, E., Triviño, P. A. V., Fajardo, L. V. S. (2014) Clasificación, manejo y conservación de colecciones biológicas: una mirada a su importancia para la biodiversidad. *Momentos de Ciencia*, 11 (2), 97–106.
- Van Lenteren, J. C., Bueno, V. H. (2003) Augmentative biological control of arthropods in Latin America. *BioControl*, 48, 123–139.
<https://doi.org/10.1023/A:1022645210394>
- Wharton, R. A., Lopez-Martinez, V. (2000) A new species of *Triaspis* Haliday (Hymenoptera: Braconidae) parasitic on the pepper weevil, *Anthonomus eugenii* Cano (Coleoptera: Curculionidae). *Proceedings-Entomological Society of Washington*, 102 (4), 794–923.