



Evaluation of antimicrobial susceptibility and genetic profiles (ERIC-PCR) of *Enterococcus* species isolated from chicken viscera

Evaluación de la susceptibilidad a antibióticos y perfiles genéticos (ERIC-PCR) de especies *Enterococcus* aislados de vísceras de pollo

Arantxa Almada-Corral¹, Maritza Alejandra Cordero-Ortiz¹, Lizbeth Dolores Ballesteros-Herrera¹, Siria de Jesús Calderón-Montoya¹, Enrique Bolado-Martínez¹, Reyna Isabel Sánchez-Mariñez¹, Alfonso García-Galaz², Iracema del Carmen Rodríguez-Hernandez¹, Mario Onofre Cortez-Rocha³ and Maritza Lizeth Álvarez-Ainza^{1*}

¹ Laboratory of Microbiology, Department of Chemical and Biological Sciences, University of Sonora. Boulevard Luis Encinas y Rosales s/n, C.P. 83000, Hermosillo, Sonora, Mexico.

² Coordination of Food Science. Research Center in Food and Development A.C., Carretera a la Victoria Km 0.6, P.O. Box 1735, Hermosillo, Sonora, Mexico.

³ Laboratory of Microbiology, Department of Food Research and Graduate Program, University of Sonora. Boulevard Luis Encinas y Rosales s/n, C.P. 83000, Hermosillo, Sonora, Mexico.

ABSTRACT

The genus *Enterococcus* can be found in water, soil, and food, and above all, they are part of the animal (including human) intestinal microbiota. Food prevalence is mainly due to their resistance to adverse environmental conditions. Seventy-three isolates from chicken viscera were confirmed to belong to the *Enterococcus* genus by biochemical tests and carbohydrate fermentation. Phenotypic characterization of the species indicates that *E. casseliflavus* was the predominant specie. Antimicrobial resistance to amikacin (42 %), kanamycin (38 %), streptomycin (55 %) and erythromycin (33 %) in the isolates was notorious. The analysis obtained from ERIC-PCR profiles shows a high genetic variability among the isolates. In addition, a relationship between the antimicrobial and ERIC-PCR profiles was observed among isolates. These results indicate the presence of multi-resistant *Enterococcus* in commercial chicken viscera with high genetic variability, which could be a potential nosocomial bacterium or transfer this resistance to another pathogenic species causing human diseases.

Keywords: *Enterococcus*; antimicrobials; identification; resistance; food safety.

RESUMEN

El género *Enterococcus* puede ser encontrado en agua, suelo y en general en todo ambiente, son parte de la microbiota intestinal de animales, incluyendo a humanos. La prevalencia en alimentos se debe principalmente a la resistencia que presentan a las condiciones adversas del ambiente. Setenta y tres aislados de vísceras de pollo fueron confirmados como pertenecientes al género a través de pruebas bioquímicas y fermentación de carbohidratos. La caracterización bioquímica indicó que la especie predominante fue *E. casseliflavus*. La resistencia de los aislados a antimicrobianos fue notoria para amikacina (42 %), kanamicina (38 %), estreptomicina (55 %), y eritromicina (33 %). El análisis obtenido por ERIC-PCR mostró una alta variabilidad genética entre los aislados. Así también,

los aislados mostraron relación entre los perfiles obtenidos con la susceptibilidad a antimicrobianos y por ERIC-PCR. Estos resultados muestran la multi-resistencia de *Enterococcus* en vísceras de pollo expedidos comercialmente, con una alta variabilidad genética. Además, estos *Enterococcus* pueden ser potencialmente patógenos en individuos como una bacteria nosocomial y además transferir la resistencia en especies que son estrictamente patógenas y causan enfermedades al humano.

Palabras clave: *Enterococcus*; antimicrobianos; identificación; resistencia; seguridad alimentaria.

INTRODUCTION

Enterococcus genus are Gram-positive cocci, generally grouped in pairs or short chains. They are facultative anaerobes, catalase-negative, and have Lancefield group D oligosaccharides. Their primary habitat is water, soil, and food; above all, they are part of animal (including) intestinal microbiota (Ortega-González, 2010). The *Enterococcus* genus include species associated with human diseases whose prevalence in food is mainly due to their resistance to adverse environmental conditions (Marguet *et al.*, 2008; Chajęcka-Wierzchowska *et al.*, 2017). *Enterococcus faecalis* and *E. faecium* are the two main species involved in human infections such as bacteremia, infective endocarditis, wound infections, and urinary tract infections, among others (Alzahrani *et al.*, 2022; Hammerum 2012; Hayes *et al.*, 2004), and they are resistant to antimicrobials. *Enterococcus* is inherently resistant to several first-line antimicrobial agents; they show low-level resistance to β -lactams and aminoglycosides (Alzahrani *et al.*, 2022). Therefore, controlling bacterium infections and avoiding cross-contamination with food is of utmost importance (Hammerum, 2012).

Since the detection of vancomycin-resistant *Enterococcus* and streptogramin-resistant *E. faecium* outside hospitals, molecular typing methods have been implemented to compare enterococci from different sources. The genetic

*Autor para correspondencia: Maritza Lizeth Álvarez Ainza
Correo electrónico: maritza.alvarez@unison.mx

Recibido: 10 de octubre de 2022

Aceptado: 9 de noviembre de 2022

profiles obtained gave a clearer picture of the isolates involved and a comparison of the clonal transfer of plasmids (Hammerum, 2012).

Molecular techniques are preferable to phenotypic ones due to their better reproducibility, typing power, versatility, and discrimination. They are widely used as a complement to conventional epidemiological investigations, for the study of nosocomial infections of bacterial etiology as well as in the epidemiological study of different bacterial isolates distribution, based on genetic profiles (Corrales and López-Cánovas, 2016). One of the molecular techniques proven useful to precisely distinguish between *E. faecalis* and *E. hirae* isolates, and to reveal the genetic diversity of *E. faecalis* isolates from birds is the amplification of intergenic repetitive consensus sequences (ERIC-PCR). In addition, it has shown great discriminating power between cheese *E. faecium* isolates, compared to other techniques such as pulsed-field gel electrophoresis (PFGE). Therefore, it is useful to obtain a bacterial genetic distribution and epidemiology (Blanco *et al.*, 2017). Also, in other studies, ERIC-PCR has shown a good differentiation power for *E. coli* molecular typing (Ardakani and Ranjbar, 2016; Ramakrishnan *et al.*, 2022).

Thus, the aim of this study was to determine the antibiotic susceptibility and the genetic profiles using ERIC-PCR of *Enterococcus* spp. previously isolated from commercial chicken viscera in order to observe the relationship of these isolates.

MATERIALS AND METHODS

Strain reactivation

Enterococcus samples isolated from chicken viscera purchased in retail stores from Hermosillo, Sonora, México, were conserved by deep freezing (-80 °C). Vials were defrosted at 25 °C, placed into BHI broth, and incubated at 36 °C for 24 - 48 h for reactivation. Initially, every chicken viscera was evaluated according to the Official Mexican Norm NOM-210-SSA1-2014 (DOF, 2015) at the University of Sonora Microbiology laboratory, for the obtention of *Enterococcus*. All the positive and confirmed *Enterococcus* isolates were used in this study.

Phenotypic characterization

Biochemical tests were performed on the previous isolates to observe the typical characteristics of the genus *Enterococcus*. The tests carried out were Gram staining, catalase, growth in salty BHI broth, and Bile Esculin Medium (BEM) (Devriese *et al.*, 1993). Also, the carbohydrate fermentation test was carried out according to the recommendations of García-Galaz *et al.* (2004). Sorbose, lactose, adonitol, melezitose, melibiose, raffinose, arabinose, glucose, galactose, sorbitol, sucrose, trehalose, inulin, xylose and mannitol were used.

Antimicrobial resistance determination

The disk diffusion method was used according to the recommendation of the Clinical & Laboratory Standards

Institute (CLSI, 2020) to evaluate the isolates resistance to the following antimicrobials: ampicillin (10 µg), streptomycin (10 µg), gentamicin (10 µg), amikacin (30 µg), kanamycin (30 µg), amoxicillin/clavulanic acid (20 +10 µg), erythromycin (15 µg), ciprofloxacin (5 µg), norfloxacin (10 µg), nitrofurantoin (100 µg), imipenem (10 µg), tetracycline (30 µg) and vancomycin (30 µg). *Staphylococcus aureus* (ATCC 25923), *E. faecalis* (ATCC 29212) and *E. faecalis* (ATCC 51299) were used as control strains. The assay was performed in Mueller-Hinton Agar (Difco), and the inoculum of each bacterium was obtained by the absorbance 665 nm (0.08 - 01) using a spectrometer (Thermo Spectronic 40001/4).

Genomic characterization by ERIC-PCR

DNA extraction. Once the isolates were phenotypically characterized, their genomic DNA was extracted according to Wright *et al.* (2017) with some modifications using lysozyme (24000 kU/mL, Sigma, Canada) and mutanolysin (40 kU/mL). The DNA integrity was checked through a 1.0 % agarose gel electrophoresis and developed with GelRed (Biotum, U.S.A.) under ultraviolet light. Subsequently, the bacterial genetic material was quantified using an AquaMate Vis spectrophotometer (Thermo Scientific).

ERIC-PCR. Amplification of the DNA extracted from each isolate was carried out by the ERIC-PCR technique. The reaction mixture contained 200 µM of dNTP's, 0.4 µM of ERIC-1 (5'-ATGTAAGCTCCTGGGGATTAC-3'), 0.4 µM ERIC-2 (5'-AAGTAAGTGACTGGGTGAGCG-3'), 2 mM of MgCl₂ (Promega, USA), 5 µL of 5X buffer with dye (Promega, USA), and one U of Taq DNA polymerase (Promega, USA). Then, sterile deionized water was added to complete a final volume of 20 µL and gently homogenized. Finally, 5 µL of previously extracted DNA (containing 100 ng) were added to each microtube with the reaction mixture (Modified from Cabral *et al.*, 2012; Durmaz *et al.*, 2015). After obtaining the ERIC-PCR results, the profiles were visualized or checked through a 1.5 % agarose gel electrophoresis and developed with GelRed (Biotum, U.S.A.) under ultraviolet light, and the image captured by the photodocumentator WiseDOc (WGD-305, Korea). The fragments size was estimated and compared with the 100-bp molecular ladder.

Statistical analysis

The results of the carbohydrate fermentation test and the ERIC-PCR profiles were analyzed with the BioNumerics version 6.5 software (Applied Maths, Belgium), using Pearson or UPGMA (Unweighted Pair Group Method with Arithmetic Mean) to make the clusters and the DICE coefficient (80 %), which provides a percentage of similarity among the *Enterococcus* isolates. Furthermore, the ERIC-PCR results were correlated with the determination of antibiotic resistance through the disk diffusion test (Durmaz *et al.*, 2015).

RESULTS AND DISCUSSION

A total of 73 isolates were confirmed to belong to the *Enterococcus* genus. The carbohydrate fermentation results

gave the presumptive phenotypic characterization of the species, and by means of the BioNumerics program output, it was possible to form isolates groups, a dendrogram (Figure 1). It allowed us to relate and discriminate the species with more than 60 % similarity. According to the prevalence (%) for each *Enterococcus* species, *E. casseliflavus* and *E. faecalis* showed the highest prevalence, corresponding to 53 % (39 isolates) and 20.5 % (15 isolates), respectively. The rest of the species showed lower than or equal to 6 %, as shown in Table 1.




Figure 1. Dendrogram of *Enterococcus* species isolated from chicken viscera, according to their biochemical characteristics.

Figure 1. Dendrograma de las especies de *Enterococcus* aisladas de vísceras de pollo, de acuerdo con sus características bioquímicas.

Table 1. *Enterococcus* species isolated from chicken viscera identified by biochemical characteristics.

Tabla 1. Especies de *Enterococcus* aisladas de vísceras de pollo, de acuerdo con sus características bioquímicas.

Identified species	Total isolated	% Isolated
<i>E. casseliflavus</i>	49	53 %
<i>E. faecalis</i>	21	23 %
<i>E. raffinosus</i>	8	9 %
<i>E. faecium</i>	5	5 %
<i>E. solitarius</i>	3	3 %
<i>E. mundtii</i>	3	3 %
<i>E. malodoratus</i>	2	2 %
<i>E. cecorum</i>	1	1 %
<i>E. durans</i>	1	1 %
Total	73	100 %

In chickens, it has been observed that an age-dependent succession of enterococcal species colonizes the intestines. Initially, they are colonized by *E. faecalis*, then displaced mainly by *E. faecium*, and afterwards, these species are replaced by *E. cecorum* in adult chickens (Aarestrup *et al.*, 2002; Lebreton *et al.*, 2014). Nevertheless, Lebreton *et al.* (2014) indicate that *E. faecalis* is also found in young chickens. For this reason, the *E. faecalis* species found in our study could be due to human contamination during handling the product since they were not young chickens.

Our results differ from those reported by Hidano *et al.* (2015) and Alzahrani *et al.* (2022), who isolated *Enterococcus* species from chicken product samples. They found that *E. faecalis* (75 and 50 %) dominated as the high-prevalence species followed by *E. faecium* (16 and 33 %). This contrasts with our results since *E. faecalis* appeared with high frequency, whereas *E. faecium* had a prevalence near 5 %.

On the other hand, *E. cecorum* was initially described as part of the normal microbiota in poultry, which coincides with the findings in this work. Even if it was found in low proportion, it is recognized as an important pathogen in feedlots. In the case of humans, it has had sporadic participation in infections such as septicemia, endocarditis and osteomyelitis (Hayes *et al.*, 2004).

Gallegos and Guerrero (2004) performed a similar study in which *Enterococcus* was isolated from chicken viscera. They found that 23 % of isolates belonged to *E. casseliflavus*, also reporting *E. cecorum* (22.8 %), *E. gallinarum* (19.4 %), *E. sulfureus* (10.1 %), *Enterococcus* spp. (8.4 %), *E. avium* (6.7 %), *E. malodoratus* (5.9 %), *E. faecalis* (1.6 %) and *E. hirae* (0.8 %). These results partially agree with the data obtained in our study. The diversity of species is similar, *E. casseliflavus* was predominated in both cases, but there is dissimilarity in the other species prevalence, which might be due to the raising poultry practices, that can modify the *Enterococci* species in the microbiota.

Resistance to Antibiotics by Disk Diffusion

Results of the resistance profiles determined by the disk diffusion test are presented in Table 2. The antimicrobial resistance demonstrated by all the isolates to antibiotics was notorious for amikacin (42 % of the isolates), kanamycin (38 %), streptomycin (55 %) in addition to erythromycin (33 %). These results indicate that the susceptibility to the antimicrobials is low and a large number of the isolated showed intermediate susceptibility. Also, it is noteworthy the representative number of isolated resistance to vancomycin (22 %), which is interesting since these isolates were obtained from chicken viscera, not from a hospital source, where phenotypes with resistance to this antibiotic are commonly found. The results in this work are similar to those reported by Alzahrani *et al.* (2022), where resistance to tetracycline (55.6 %), erythromycin (31.1 %), ciprofloxacin (21.1 %) ampicillin (30 %) and nitrofurantoin (17.8 %) were the most frequent, except the last two antibiotics which in our work, showed low resistance. In theory, the chicken viscera isolated have not been exposed to antibiotics. However, it is possible that the human handling of the product in the supermarket was the contamination source.

Moreover, it is acknowledged that different types of antibiotics are used in poultry raising, mainly those that belong to macrolides (erythromycin), trimethoprim, fluoroquinolones and tetracyclines, either in a therapeutic or sub-therapeutic way. For this reason, the resistance shown may be due to the chicken viscera samples obtained. Poultry may have been exposed to antibiotics belonging to any of these groups. However, it should be noted that enterococci have intrinsic resistance to macrolides (Novais *et al.*, 2013; Daniel *et al.*, 2015; Woolhouse *et al.*, 2015).

The results of the disk diffusion test shows that the isolates are multidrug-resistant. However, one isolate showed resistance to eight antibiotics and was characterized as *Enterococcus faecium*. Its high percentage of resistance to vancomycin is worrisome because it is a naturally multi-resistant bacterium, in which daptomycin and linezolid resistance also begins to appear. In some cases, this has led to a total absence of effective antibiotics (Alós, 2015).

The multidrug resistance among *Enterococcus* species may be due to their intrinsic resistance to aminoglycosides and β -lactams, or because they can easily acquire resistance from the environment where extrinsic resistance is found.

Table 2. *Enterococcus* spp. susceptibility to antibiotics analyzed by the disc diffusion method.
Table 2. Susceptibilidad a antibióticos en *Enterococcus* spp. analizados a través del método de disco difusión.

Drug group	Antibiotic	% Resistant (n)	% Intermediate (n)	% Susceptible (n)
1. Aminoglycosides	AMK	42 % (31)	14 % (10)	44 % (32)
	KAN	38 % (28)	35 % (26)	26 % (19)
2. Aminopenicillins	STR	55 % (40)	20 % (15)	25 % (18)
	AMC	8 % (6)	0 % (0)	92 % (67)
3. Carbapenems	AMP	2 % (3)	1 % (1)	96 % (70)
	IMP	3 % (2)	0 % (0)	97 % (71)
4. Fluoroquinolones	CIP	18 % (17)	27 % (20)	55 % (40)
	NOR	14 % (10)	9 % (7)	77 % (56)
5. Glycopeptides	VAN	22 % (16)	21 % (15)	57 % (42)
6. Macrolides	ERI	33 % (24)	38 % (38)	15 % (11)
7. Nitrofurans	NIT	10 % (14)	9 % (7)	84 % (61)
8. Tetracyclines	TCY	22 % (16)	4 % (3)	74 % (54)

AMK: Amikacin; KAN: Kanamycin; STR: Streptomycin; AMC: Amoxicillin/clavulanic acid; AMP: Ampicillin; IMP: Imipenem; CIP: Ciprofloxacin; NOR: Norfloxacin; VAN: Vancomycin; ERI: Erythromycin; NIT: Nitrofurantoin; TCY: Tetracycline. The bacterium susceptibility (Resistant, Intermediate and Susceptible) was determined based on CLSI recommendations and the disk fabricant of each antimicrobial, using control strains.

AMK: Amikacina; KAN: Kanamicina; STR: Estreptomicina; AMC: Amoxicilina/ácido clavulánico; AMP: Ampicilina; IMP: Imipenem; CIP: Ciprofloxacino; NOR: Norfloxacino; VAN: Vanomicina; ERI: Eritromicina; NIT: Nitrofurantoina; TCY: Tetraciclina. La susceptibilidad de las bacterias (Resistente, Intermedio y Susceptible) fue determinada basada en las recomendaciones del CLSI y las del fabricante de cada antimicrobiano, utilizando las cepas control.

Due to the massive use of antibiotics, a highly significant increase in the prevalence of resistance has been observed worldwide (Cercenado, 2011; Alós, 2015).

The appearance of MDR (Multidrug-resistant) bacteria (in this case *Enterococcus*) in livestock and poultry industry is a public health concern due to the reported transmission of these bacteria to humans. The spread of MDR bacteria occurs by consuming contaminated food, direct contact with farmers and veterinarians or indirectly by handling animal waste, contaminated soil, water, or surfaces. Resistance determinants, carried by MDR strains, can also be transmitted to other commensal isolates in the host and cause future complications. In addition, infections caused by MDR isolates have been associated with extended hospital stays, high morbidity and mortality rates (Daniel *et al.*, 2015; Price *et al.*, 2018; Begonovic *et al.*, 2018).

It is known that the microbiota plays a fundamental role in the regulation of the organism, and that the type of microbiota depends on various factors, including diet. Currently, selective microbiota associated with the consumption of contaminated products with resistant bacteria has been observed, which is also of great concern. Additionally, bacteria such as enterococci can easily acquire and transfer resistance. For this reason, it is essential to know the profiles of resistance and proper of use antibiotics, whether in human, veterinary, or animal husbandry medicine.

ERIC-PCR

All the isolates showed around 2 - 11 bands in the ERIC-PCR test. Most of them resulted in 3 (23.29 %), 4 (28.77 %), or 6 (13.70 %) bands. The ERIC segment size was 100 to 4,361 bp, coinciding with those reported by Wijetunge *et al.* (2012) who characterized *E. cecorum* species isolated from diseased poultry, and obtained between 2-7 bands with a size of 200-5000 bp, similar to those obtained in our study.

The band analysis from the ERIC-PCR products electrophoretic gels, through a dendrogram obtained by UPGMA analysis using the DICE coefficient with a similarity value of 80 %, is shown in Figure 2. As can be seen, the analysis shows a high genetic variability among the isolates, which coincides with the findings of Wijetunge *et al.* (2012) and Blanco *et al.* (2017).

Most of the clusters had around 20 and 60 % similarity in the dendrogram analysis. Based on this, they were classified into 10 groups (I to X). Group I includes two *E. casseliflavus* isolates with 100 % similarity and one group of an isolate with ≥ 80 % similarity. Group II contains two isolates with 100 % similarity (one *E. casseliflavus*, and the other *E. solitarius*), and two groups with ≥ 80 % similarity. Group III comprises two groups with ≥ 80 % similarity, each with two isolates. Groups IV and V have only one group with ≥ 80 % similarity containing two isolates each. Groups VI to X contain isolates that do not have a significant percentage of similarity, which is why they are considered different. This analysis shows the genomic diversity among the isolated *Enterococcus*, even among the specie characteristics determined by biochemical




Figure 2. Dendrogram obtained from ERIC-PCR analysis of 73 *Enterococcus* species isolated from chicken viscera.

Figure 2. Dendrograma obtenido del análisis por ERIC-PCR de 73 especies de *Enterococcus* aisladas de vísceras de pollo.

tests and carbohydrate fermentation.

Figure 3 shows the dendrogram that includes the analysis of the ERIC-PCR genomic profiles and antibiotic susceptibility tests. In this analysis, the two *E. casseliflavus* isolates with 100% similarity in group I coincide in presenting resistance to kanamycin, erythromycin and streptomycin. On the other hand, they have $\geq 80\%$ similarity with another *E. casseliflavus* isolate, sharing resistance to kanamycin and erythromycin. In group II are two other isolates (*E. casseliflavus* and *E. solitarius*) with 100% similarity. This resistance profile coincides with amikacin, kanamycin, erythromycin, vancomycin, and streptomycin. Group III presents two similar isolates (*E. casseliflavus* and *E. raffinosus*) with resistance to kanamycin, erythromycin and streptomycin; and two other isolates (*E. mundtii* and *E. faecalis*) that have resistance to erythromycin, tetracycline, vancomycin and streptomycin.




Figure 3. Dendrogram obtained from the ERIC-PCR and antimicrobial profiles analysis of 73 *Enterococcus* species isolated from chicken viscera. AMK: Amikacin; KAN: Kanamycin; STR: Streptomycin; AMC: Amoxicillin/clavulanic acid; AMP: Ampicillin; IMP: Imipenem; CIP: Ciprofloxacin; NOR: Norfloxacin; VAN: Vancomycin; ERI: Erythromycin; NIT: Nitrofurantoin; TCY: Tetracycline.

Figura 3. Dendrograma obtenido del análisis por ERIC-PCR y de perfiles de susceptibilidad a antimicrobianos de 73 especies de *Enterococcus* aisladas de vísceras de pollo. AMK: Amikacina; KAN: Kanamicina; STR: Estreptomicina; AMC: Amoxicilina/ácido clavulánico; AMP: Ampicilina; IMP: Imipenem; CIP: Ciprofloxacino; NOR: Norfloxacino; VAN: Vancomicina; ERI: Eritromicina; NIT: Nitrofurantoina; TCY: Tetraciclina.

In group IV, there are two isolates (*E. casseliflavus* and *E. faecalis*) with resistance to amikacin, kanamycin, erythromycin, ciprofloxacin and streptomycin. Group V contains two similar isolates (*E. faecalis* and *E. casseliflavus*) with resistance to kanamycin and norfloxacin.

The similarity percentage used for the dendrogram analysis was $\geq 80\%$. Fifteen isolates had a resistance percentage higher than or equal to 80%, showing a great diversity

of strains among the isolates, even of the same species according to the phenotypic characterization. As mentioned above, this was perhaps due to the product handling in the different supermarkets where poultry viscera were obtained. Also, the chicken samples could come from different farms and/or companies.

These results agree with those of Bedendo and Pignataro (2000), where identical profiles were observed through molecular techniques in *E. malodoratus* and *E. hirae* samples. They stated that this may indicate that conducting studies only at the phenotypic level, such as biochemical tests or susceptibility to antibiotics, does not provide accurate results about the species. Also, that among different species, it is possible to obtain an identical band, thus both, phenotypic and genotypic studies, complement each other. The molecular techniques used to typify bacterial strains are known to be statistically significant in discriminatory power and reproducibility, standardization, and interpretation, making them indispensable in any epidemiological study (Palomino-Camargo and González-Muñoz, 2014).

The isolates related through the ERIC-PCR profile also showed a relationship with their antibiotic resistance profile. This has been previously reported in enterococcal isolates from hospitalized patients, which were related to their ERIC band profile and antibiogram. A similar resistance pattern among isolates classified in the same group could indicate that they came from a common site and have both chromosomal genes and resistance mechanisms (Zalipour et al., 2019).

CONCLUSIONS

This study shows the prevalence of *E. casseliflavus* isolates in chicken viscera. A high percentage of antimicrobial resistance was observed to aminoglycoside and macrolide antibiotics; also, the isolates had a worrying resistance percentage to vancomycin. The ERIC-PCR analysis showed high genetic variability among the isolates and a correlation between the antimicrobial and ERIC-PCR profiles.

ACKNOWLEDGEMENTS

The authors would like to thank to Gutiérrez-Franco, L. E. for proofreading.

CONFLICT OF INTEREST

There is no conflict of interest.

REFERENCES

- Aarestrup, F.M., Butaye, P. and Witte, W. 2002 Non-human reservoirs of enterococci. In The Enterococci: Pathogenesis, Molecular Biology and Antibiotic Resistance. Washington DC: ASM Press. <https://doi.org/10.1128/9781555817923.ch2>
- Alós, J.I. 2015 Resistencia bacteriana a los antibióticos: una crisis global. Enfermedades Infecciosas y Microbiología Clínica. 33(10): 692-699. <https://doi.org/10.1016/j.eimc.2014.10.004>
- Ardakani, M.A. and Ranjbar, E. 2016. Molecular typing of uropathogenic *E. coli* strains by the ERIC-PCR method. Electronic Physician. 8(4): 2291-2296. <http://dx.doi.org/2291>

Alzahrani, O.M., Fayed, M., Alswat, A.S., Alkafafy, M., Mahmoud S., Al-Marry, T., Almuslem, A., Ashfaq, H. and Yusuf, S. 2022. Antimicrobial resistance, biofilm formation, and virulence genes in *Enterococcus* species from small backyard chicken flocks. *Antibiotics* 11(3): 320. <https://doi.org/10.3390/antibiotics11030380>

Bedendo, J. and Pignatari, A. C. 2000. Typing of *Enterococcus faecium* by polymerase chain reaction and pulsed field gel electrophoresis. *Brazilian Journal of Medical Biological Research.* 33(11): 1269-1274. <https://doi.org/10.1590/S0100-879X2000001100002>

Beganovic, M., Luther, M.K., Rice, L.B., Arias C.A., Rybak M.J. and LaPlante K.L. 2018. A re-view of combination antimicrobial therapy for *Enterococcus faecalis* bloodstream infections and infective endocarditis. *Clinical Infectious Diseases.* 67(2): 303-309. <https://doi.org/10.1093/cid/ciy064>

Blanco, A.E., Barz, M., Cavero D., Icken, W., Sharifi, A.R., Voss, M., Buxadé, C. and Preisinger, R. 2017. Characterization of *Enterococcus faecalis* isolates by chicken embryo lethality assay and ERIC-PCR. *Avian Pathology.* 47: 23-32. <https://doi.org/10.1080/03079457.2017.1359404>

Cabral, A.B., Melo, R. de C., Viera, M.A. and Souza, A.C. 2012. Multidrug resistance genes, including *bla_{KPC}* and *bla_{CTX-M-2'}* among *Klebsiella pneumoniae* isolated in Recife, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical.* 45(5): 572-578. <https://doi.org/10.1590/S0037-86822012000500007>

Cercenado, E. 2011. *Enterococcus*: resistencias fenotípicas y genotípicas y epidemiología en España. *Enfermedades Infecciosas y Microbiología Clínica.* 29: 59-65. [https://doi.org/10.1016/S0213-005X\(11\)70045-3](https://doi.org/10.1016/S0213-005X(11)70045-3)

Chajęcka-Wierzchowska, W., Zadernowska, A. and Łaniewska-Trockenheim, Ł. 2017. Virulence factors of *Enterococcus* spp. presented in food. *LWT.* 75: 670-676. <https://doi.org/10.1016/j.lwt.2016.10.026>

CLSI. 2020. Performance Standards for Antimicrobial Susceptibility Testing (M100). 32nd Edition. Wayne, P.A.: Clinical and Laboratory Standards Institute; 2020.

Corrales, F. and López-Cánovas, L. 2016. Las infecciones nosocomiales en Cuba y su control mediante las técnicas moleculares de tipificación de microorganismos. *Revista CENIC. Ciencias Biológicas.* 47: 27-32.

Daniel, D. S., Lee, S. M., Dykes, G. A. and Rahman, S. 2015. Public health risks of multiple-drug-resistant *Enterococcus* spp. in Southeast Asia. *Applied and Environmental Microbiology.* 81(18): 6090-6097. <https://doi.org/10.1128/AEM.01741-15>

Devriese, L.A., Pot, B., and Collins, M.D. 1993. Phenotypic identification of the genus *Enterococcus* and differentiation of phylogenetically distinct enterococcal species and species groups. *Journal of Applied Bacteriology.* 75(5): 399-408. <https://doi.org/10.1111/j.1365-2672.1993.tb02794.x>

DOF. 2015. Norma Oficial Mexicana NOM-210-SSA1-2014, Productos y servicios. Métodos de prueba microbiológico. Determinación de microorganismos indicadores. Determinación de microorganismos patógenos. 2015. Disponible: http://dof.gob.mx/nota_detalle.php?codigo=5398468&fecha=26/06/2015

Durmaz, S., Bal, E., Gunaydin, M., Yula, E. and Percin, D. 2015. Detection of β-lactamase genes, ERIC-PCR typing and phylogenetic groups of ESBL producing quinolone resistant clinical *Escherichia coli* isolates. *Biomedical Research.* 26(1): 43-50.

Gallegos, J.C. and Guerrero, J. L. 2004. Resistencia a antibióticos y de alto nivel a aminoglucósidos en *Enterococcus*, aislados de vísceras de pollo de supermercados de Hermosillo Sonora (Bachelor's Theses). Universidad de Sonora. Hermosillo, Sonora, México.

García-Galaz, A., Pérez-Morales, R., Díaz-Cinco, M. and Acedo-Feliz, E. 2004. Resistance of *Enterococcus* strains isolated from pigs to gastrointestinal tract and antagonistic effect against *Escherichia coli* K88. *Revista Latinoamericana de Microbiología.* 46(1-2), 5-11.

Hammerum A.M. 2012 Enterococci of animal origin and their significance for public health. *Clinical Microbiology and Infection.* 18(7): 619-625. <https://doi.org/10.1111/j.1469-0691.2012.03829.x>

Hayes, J.R., English, L.L., Carr, L.E., Wagner, D.D. and Joseph, S. W. 2004 Multiple-antibiotic resistance of *Enterococcus* spp. isolated from commercial poultry production environments. *Applied Environmental Microbiology.* 70(10), 6005-6011. <https://doi.org/10.1128/AEM.70.10.6005-6011.2004>

Hidano, A., Takehisa, Y., Yoko, H., Norihiko, M., Sota, K., Takeshi, N., and Toshiyuki, T. 2015. Unraveling antimicrobial resistance genes and phenotype patterns among *Enterococcus faecalis* isolated from retail chicken products in Japan. *PLoS ONE.* 10(3):1-15. <https://doi.org/10.1371/journal.pone.0121189>

Lebreton, F., Willems, R. J., and Gilmore, M. S. 2014. *Enterococcus* diversity, origins in nature, and gut colonization. In *Enterococci: from commensals to leading causes of drug resistant infection* [Internet]. Boston: Massachusetts Eye and Ear Infirmary; 2014-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK190427/>

Marguet, E. R., Vallejo, M., and Olivera, N. L. 2008. Factores de virulencia de cepas de *Enterococcus* aisladas de quesos ovinos. *Acta Bioquímica Clínica Latinoamericana.* 42(4): 543-548.

Novais, C., Freitas, A.R., Silveira, E., Antunes, P., Silva, R., Coque, T.M. and Peixe, L. 2013. Spread of multidrug-resistant *Enterococcus* to animals and humans: an underestimated role for the pig farm environment. *Journal of Antimicrobial Chemotherapy.* 68(12): 2746-2754. <https://doi.org/10.1093/jac/dkt289>

Ortega-González, L. M. 2010. Enterococos: actualización. *Revista Habanera Ciencia Médica.* 9(4): 507-515.

Ramakrishnan V., Marialouis, X.A., Al-Ansari, M.M., Al-Humaid L., Santhanam A., and Obulisanmy K. 2022. Multilocus sequence typing and ERIC-PCR fingerprinting of virulent clinical isolates of uropathogenic multidrug resistant *Escherichia coli*. *Journal of King Saus University- Science.* 34: 101874. <https://doi.org/10.1016/j.jksus.2022.101874>

Palomino-Camargo, C., and González-Muñoz, Y. 2014. Técnicas moleculares para la detección e identificación de patógenos en alimentos: ventajas y limitaciones. *Revista Peruana de Medicina Experimental y Salud Pública.* 31: 535-546.

Price, V.J., McBride, S.W., Duerkop, B.A. and Palmer, K.L. 2018. CRISPR-Cas blocks antibiotic resistance plasmid transfer between *Enterococcus faecalis* strains in the gastrointestinal tract. *bioRxiv* 312751. <https://doi.org/10.1101/312751>

Woolhouse, M., Ward, M., Van Bunnik, B. and Farrar, J. 2015. Antimicrobial resistance in humans, livestock and the wider environment. *Philosophical Transactions of the Royal Society B.* 370:20140083. <https://doi.org/10.1098/rstb.2014.0083>

Wijetunge, D.S., Dunn, P., Wallner-Pendleton, E., Lintner, V., Lu, H. and Kariyawasam, S. 2012 Fingerprinting of poultry isolates of *Enterococcus cecorum* using three molecular typing methods. *Journal of Veterinary Diagnostic Investigation*. 24: 1166-1171. <https://doi.org/10.1177/1040638712463563>

Wright, M. H., Adelskov, J. and Greene, A. C. 2017. Bacterial DNA Extraction using individual enzymes and Phenol/Chloroform separation. *Journal of Microbiology and Biology Education*. 18(2): 1-3. <https://doi.org/10.1128/jmbe.v18i2.1348>

Zalipour, M., Esfahani, B. N., and Havaei, S. A. 2019. Phenotypic and genotypic characterization of glycopeptide, aminoglycoside and macrolide resistance among clinical isolates of *Enterococcus faecalis*: a multicenter based study. *BMC Research Notes*. 12(1): 292. <https://doi.org/10.1186/s13104-019-4339-4>