




Boletín Médico del Hospital Infantil de México

www.elsevier.es/bmhim



RESEARCH ARTICLE

Proteomic changes in a childhood acute lymphoblastic leukemia cell line during the adaptation to vincristine



Ana Laura Guzmán-Ortiz^{a,b}, Gerardo Aparicio-Ozores^b, Ricardo Valle-Rios^{a,c},
Oscar Medina-Contreras^a, Genaro Patiño-López^a, Héctor Quezada^{a,*}

^a Laboratorio de Investigación en Inmunología y Proteómica, Hospital Infantil de México Federico Gómez, Mexico City, Mexico

^b Departamento de Microbiología, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Mexico City, Mexico

^c División de Investigación, Facultad de Medicina, Universidad Nacional Autónoma de México (UNAM), Mexico City, Mexico

Received 11 February 2017; accepted 28 March 2017

Available online 23 May 2017

KEYWORDS

Leukemia;
Chemoresistance;
Vincristine

Abstract

Introduction: Relapse occurs in approximately 20% of Mexican patients with childhood acute lymphoblastic leukemia (ALL). In this group, chemoresistance may be one of the biggest challenges. An overview of complex cellular processes like drug tolerance can be achieved with proteomic studies.

Methods: The B-lineage pediatric ALL cell line CCRF-SB was gradually exposed to the chemotherapeutic vincristine until proliferation was observed at 6 nM, control cells were cultured in the absence of vincristine. The proteome from each group was analyzed by nanoHPLC coupled to an ESI-ion trap mass spectrometer. The identified proteins were grouped into over-represented functional categories with the PANTHER classification system.

Results: We found 135 proteins exclusively expressed in the presence of vincristine. The most represented functional categories were: Toll receptor signaling pathway, Ras Pathway, B and T cell activation, CCKR signaling map, cytokine-mediated signaling pathway, and oxidative phosphorylation.

Conclusions: Our study indicates that signal transduction and mitochondrial ATP production are essential during adaptation of leukemic cells to vincristine, these processes represent potential therapeutic targets.

© 2017 Hospital Infantil de México Federico Gómez. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: hquezada@himfg.edu.mx (H. Quezada).

PALABRAS CLAVE

Leucemia;
Quimiorresistencia;
Vincristina

Cambios en el proteoma de una línea celular de leucemia linfoblástica aguda infantil durante la adaptación a vincristina

Resumen

Introducción: Aproximadamente el 20% de los pacientes mexicanos con leucemia linfoblástica aguda (LLA) infantil presentan recaídas. En este grupo, la quimiorresistencia es uno de los principales desafíos. Los estudios proteómicos pueden dar un panorama general de procesos celulares complejos como la tolerancia a fármacos.

Métodos: La línea celular de LLA de linaje B, CCRF-SB, fue expuesta de manera gradual al fármaco quimioterapéutico vincristina hasta observar proliferación celular en presencia de 6 nM, como control se cultivaron células en ausencia del fármaco. Se analizó el proteoma de cada grupo mediante nanoHPLC acoplado a un espectrómetro de masas de tipo trampa de iones. Las proteínas identificadas se agruparon en categorías funcionales sobre-representadas con el sistema de clasificación PANTHER.

Resultados: Encontramos 135 proteínas expresadas exclusivamente en presencia de vincristina. Las categorías funcionales más representadas fueron la señalización asociada a los receptores tipo Toll, señalización dependiente de Ras, activación de células B y T, mapa de señalización CCKR, señalización mediada por citoquinas y la fosforilación oxidativa.

Conclusiones: Nuestro estudio indica que la transducción de señales y la producción de ATP mitocondrial son procesos esenciales durante la adaptación de células leucémicas a vincristina por lo que estos procesos representan potenciales blancos terapéuticos.

© 2017 Hospital Infantil de México Federico Gómez. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Childhood acute lymphoblastic leukemia (ALL) is the most common type of cancer and the second leading cause of mortality among Mexican children.¹ Overall cure rates of newly diagnosed ALL patients is around 80%, while chemoresistance is one of the main challenges in the relapsed population.^{2–4} Although much progress has been achieved in the detailed characterization of the molecular processes directly involved in drug tolerance of cancer cells,^{5,6} more research is needed to develop effective therapeutic strategies to resensitize chemoresistant cells.⁶

Vincristine is a vinca alkaloid, which interacts with tubulin disrupting microtubule polymerization and favoring the cell cycle arrest in the M phase which is followed by induction of apoptosis.⁷ It is used in several stages of the ALL treatment.³ It has been reported that the P-glycoprotein MDR1 actively pumps vincristine outside the cell reducing its therapeutic effect.⁸ Inactivation of intracellular vincristine by the myeloperoxidase has also been reported to contribute to resistance in some types of leukemia.⁹ Moreover, resistant leukemia, and other types of cancer cells, commonly show deregulated apoptosis^{4,6,10} and signaling pathways involved in survival.^{5,6,11} The complex state of resistance is the result of the concerted action of multiple interacting genes, proteins, and metabolites; and this scenario is well suited for characterization with the omic technologies. Proteomics studies provide an overview of the changes in the relative abundance of the proteins of a cell or tissue under different conditions.¹² The characterization of proteins, as the final executors of cellular activities, may lead to identification of putative therapeutic targets and a better understanding of the pathological states.¹³ In the present work, we

describe the changes in the proteome of a B-ALL cell line after adaptation to vincristine. Our results allowed the identification of targetable signaling and metabolic steps which may represent potential targets to resensitize leukemic cells to vincristine.

2. Methods

2.1. Growth conditions

The B-lineage pediatric ALL cell line CCRF-SB (ATCC CCL-120) was grown in RPMI-1640 with 10% (v/v) fetal bovine serum, 100 U/ml penicillin, 0.1 mg/ml streptomycin, 1% sodium pyruvate at 37 °C under an atmosphere with 5% CO₂.

2.2. Vincristine exposition

Cell viability was estimated with the MTT assay in 96 well plates.¹⁴ To determine the IC₅₀, vincristine was added to 7 × 10⁴ cells in 100 µL of media per well at 0, 1, 2, 3, 4, 5, 6, 8, 10, 20 and 40 nM in triplicate for 48 hours (Figure 1A). Gradual exposition was made as follows: 3 × 10⁶ cells were cultured in 5 ml of media in the presence of 1 nM vincristine for five days, and then were exposed to 2 nM for the next five days, 3.5 nM for five days, 4.5 nM for three days, and 6 nM for four days (Figure 1B). In every step viable cells were enriched by centrifugation at 800 rpm 5 min, only the pelleted cells were transferred to the subsequent drug concentration. After 22 days of gradual exposition, cells proliferated in the presence of 6 nM vincristine, and cell viability was higher than 70%. Control cells were subjected to the same manipulations but cultured in the absence of vincristine.




Figure 1 The CCRF-SB cell line is sensitive to vincristine. A. To obtain the IC₅₀ (10 nM), cells were exposed to growing concentrations of vincristine for 48 h. B. Cell viability (□) and vincristine concentration (—) during the gradual adaptation protocol. The arrow shows the time point at which cells were harvested for proteomic studies.

2.3. Proteomic methods

Cells were washed with cold PBS and resuspended in 500 μ L lysis buffer (4% SDS, 100 mM DTT, 100 mM TRIS pH 8.6 and a protease/phosphatase inhibitor cocktail, Thermo Scientific). Complete cell disruption was achieved by 20 cycles of 2 seconds sonication at 50% amplitude in a sonics 130-Watt Ultrasonic Processor and 5 seconds ice. After reduction (30 min at 40 °C) and alkylation (200 mM iodoacetamide, 30 min at room temperature in the darkness), 50 μ g of protein were precipitated overnight with nine volumes of ethanol at -20 °C and washed twice with 90% ethanol. The dried pellet was dissolved in 50 mM guanidinium chloride, 20 mM TRIS and digested with mass spectrometry grade trypsin (1:50, Promega) overnight at 37 °C. The resulting peptides were desalted (sep-pak C18, cartridges, Waters), dried and kept at -80 °C until used.

Just before injection, peptides were dissolved in 50 μ L of 0.1% formic acid. 5 μ L of each sample were injected in triplicate to an Ultimate 3000 nanoHPLC system with a 1 cm Acclaim PepMap100 C18 pre-column and a 50 cm Acclaim PepMap100 C18 column. Peptides were eluted with

a non-linear gradient from 5% to 90% acetonitrile with 0.1% formic acid in 240 min. The HPLC system was coupled to the ESI-Ion Trap mass spectrometer Amazon speed (Bruker Daltonics) operated in positive mode, 400-1400 m/z, the 20 most abundant ions were fragmented every 50 msec, single charged ions were excluded. The ion list was constructed with the Data Analysis software (Bruker Daltonics) and used for subsequent protein identification with the Mascot search engine using the SwissProt human database, two missed trypsin cleavages, carbamidomethylation as fixed modification, methionine oxidation as variable modification, and a peptide mass tolerance of 0.6 Da. Only those proteins identified in the three replicates were considered as valid identifications. Venn diagrams were drawn with the website from the University of Southern Mississippi (<http://genevenn.sourceforge.net/>) and functional categories were analyzed with the statistical overrepresentation test of the PANTHER classification system¹⁵ (<http://pantherdb.org>) in which the analyzed list contained those proteins detected exclusively in the presence of vincristine whereas the reference list contained the proteins detected in both groups (with and without vincristine), the Bonferroni correction for multiple testing was not used. Only the functional categories with a fold enrichment value equal or higher than three were considered for further analysis.

3. Results

3.1. After gradual adaptation, the CCRF-SB cell line grew in the presence of 6 nM vincristine

Sudden vincristine exposition for 48 h resulted in IC₅₀ of 10 nM (Figure 1A), which is similar to previous reports using other types of sensitive cell lines.^{9,16,17} However, gradual exposition allowed growth and cell duplication at 6 nM. The adaptation procedure depicted in Figure 1B resulted to be highly reproducible and allowed that cell density doubled during the last four days (from about 6x10⁵ to 1.5 x10⁶ cells/ml) keeping cell viability above 70%. These adapted cells were used for proteomic studies.

3.2. Approximately 15% of the proteome is differentially expressed after adaptation to vincristine

The proteomic approach used resulted in 849 and 858 valid identifications in the control and adapted cells respectively (Figure 2). As expected, most of the proteins (approximately 85%) were present in both conditions. However, 126 proteins were detected exclusively in the absence of vincristine and 135 in the presence of this drug. This result indicated that during the adaptation process, significant changes in the proteome took place.

To identify the cellular processes that mainly contributed to drug tolerance, the statistical overrepresentation test was made using the PANTHER classification system¹⁵. The 135 proteins identified exclusively in the presence of vincristine were compared with the 723 proteins common to both groups (Figure 2). This resulted in the identification of 17 overrepresented pathways and functions (Figure 3)

Table 1 Details of the proteins assigned to the overrepresented pathways in adapted cells.

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
<i>Signaling proteins</i>					
MP2K1	Dual specificity mitogen-activated protein kinase 1	Q02750	Kinase	Toll receptor signaling pathway, Ras pathway, B-cell activation, CCKR signaling map, T-cell activation	Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway
MK14	Mitogen-activated protein kinase 14	Q16539	Non-receptor serine/threonine protein kinase	Toll receptor signaling pathway, p53 pathway feedback loops 2, p38 MAPK pathway, Ras pathway, B-cell activation, CCKR signaling map	Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK14 is one of the four p38 MAPKs which play an important role in the cascades of cellular responses evoked by extracellular stimuli such as proinflammatory cytokines or physical stress leading to direct activation of transcription factors
RB1	Retinoblastoma-associated protein	P06400	Transcription factor	p53 pathway feedback loops	Key regulator of entry into cell division that acts as a tumor suppressor. Promotes G0-G1 transition when phosphorylated by CDK3/cyclin-C
RAC1	Ras-related C3 botulinum toxin substrate 1	P63000	Small GTPase	p38 MAPK pathway, Ras pathway, B-cell activation, CCKR signaling map, T-cell activation	Plasma membrane-associated small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state it binds to a variety of effector proteins to regulate cellular responses such as secretory processes, phagocytosis of apoptotic cells, epithelial cell polarization and growth-factor induced formation of membrane ruffles
VAV	Proto-oncogene vav	P15498	Guanyl-nucleotide exchange factor, signaling molecule	B-cell activation, T-cell activation	It couples tyrosine kinase signals with the activation of the Rho/Rac GTPases, thus leading to cell differentiation and/or proliferation
CASP3	Caspase-3	P42574	Cysteine protease, protease inhibitor	CCKR signaling map, cell differentiation	Involved in the activation cascade of caspases responsible for apoptosis

Table 1 (Continued)

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
ZAP70	Tyrosine-protein kinase ZAP-70	P43403	Non-receptor tyrosine protein kinase	T-cell activation, cell differentiation	Tyrosine kinase that plays an essential role in regulation of the adaptive immune response, regulates motility, adhesion and cytokine expression of mature T-cells, as well as thymocyte development. It also contributes to the development and activation of primary B-lymphocytes
VRK1	Serine/threonine-protein kinase VRK1	Q99986	Non-receptor serine/threonine protein kinase		Serine/threonine kinase involved in Golgi disassembly during the cell cycle, following phosphorylation by PLK3 during mitosis, required to induce Golgi fragmentation
IMA4	Importin subunit alpha-4	O00505	Transfer/carrier protein	Cytokine-mediated signaling pathway, cell surface receptor signaling pathway	Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1
IMA3	Importin subunit alpha-3	O00629	Transfer/carrier protein	Cytokine-mediated signaling pathway, cell surface receptor signaling pathway	Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1
RAP1A	Ras-related protein Rap-1A	P62834	Small GTPase	Cell surface receptor signaling pathway	Induces morphological reversion of a cell line transformed by a Ras oncogene, counteracts the mitogenic function of Ras, at least partly because it can interact with Ras GAPs and RAF in a competitive manner
PIPSL	Putative PIP5K1A and PSMD4-like protein	A2A3N6	Kinase	Cell surface receptor signaling pathway, phospholipid metabolic process	Negligible PIP5 kinase activity. Binds to ubiquitinated proteins
STK25	Serine/threonine-protein kinase 25	O00506	Kinase	Response to environmental stress, cell differentiation	Oxidant stress-activated serine/threonine kinase that may play a role in the response to environmental stress. Targets to the Golgi apparatus where it appears to regulate protein transport events, cell adhesion, and polarity complexes important for cell migration

Table 1 (Continued)

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
<i>Modulators of signaling proteins</i>					
ZYX	Zyxin	Q15942	Cell junction protein, kinase modulator	Nervous system development	Adhesion plaque protein. Binds alpha-actinin and the CRP protein. Important for targeting TES and ENA/VASP family members to focal adhesions and for the formation of actin-rich structures. It may be a component of a signal transduction pathway that mediates adhesion-stimulated changes in gene expression. A cell-adhesion and mechanotransducer protein
AT2A2	Sarcoplasmic/endoplasmic reticulum calcium ATPase 2	P16615	Cation transporter, ion channel	Homeostatic process	This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen
CK5P3	CDK5 regulatory subunit-associated protein 3	Q96JB5	Probable tumor suppressor	Regulation of catalytic activity	Probable tumor suppressor initially identified as a CDK5R1 interactor controlling cell proliferation. Negatively regulates NF-kappa-B-mediated gene transcription through the control of RELA phosphorylation. Also, it regulates mitotic G2/M transition checkpoint and mitotic G2 DNA damage checkpoint
OXSRI	Serine/threonine-protein kinase OSR1	O95747	Kinase	Regulation of catalytic activity	Regulates downstream kinases in response to environmental stress. It may also have a function in regulating the actin cytoskeleton
<i>Production of mitochondrial ATP</i>					
ATPK	ATP synthase subunit f, mitochondrial	P56134	ATP synthase	Oxidative phosphorylation	Mitochondrial membrane ATP synthase (F1F0 ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane, which is generated by electron transport complexes of the respiratory chain

Table 1 (Continued)

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
NDUS8	NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial	O00217	Dehydrogenase, reductase	Oxidative phosphorylation	Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain
SDHB	Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial	P21912	Dehydrogenase	Oxidative phosphorylation	Iron-sulfur protein (IP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q)
QCR1	Cytochrome b-c1 complex subunit 1, mitochondrial	P31930	Metalloprotease	Oxidative phosphorylation	This is a component of the ubiquinol-cytochrome c reductase complex (complex III or cytochrome b-c1 complex), which is part of the mitochondrial respiratory chain. This protein may mediate formation of the complex between cytochromes c and c1
COX41	Cytochrome c oxidase subunit four isoform 1, mitochondrial	P13073	Oxidase	Oxidative phosphorylation	This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport
ETFA	Electron transfer flavoprotein subunit alpha, mitochondrial	P13804	Dehydrogenase	Homeostatic process	The electron transfer flavoprotein serves as a specific electron acceptor for several dehydrogenases, including five acyl-CoA dehydrogenases, glutaryl-CoA, and sarcosine dehydrogenase. It transfers the electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase)

Table 1 (Continued)

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
<i>Cytoskeletal dynamics</i>					
MAP4	Microtubule-associated protein 4	P27816	–	Cell differentiation	Promotes microtubule assembly
<i>Others</i>					
APOA1	Apolipoprotein A-I	P02647	–	Phospholipid metabolic process, steroid metabolic process, cell differentiation	Participates in the reverse transport of cholesterol from tissues to the liver for excretion by promoting cholesterol efflux from tissues and by acting as a cofactor for the lecithin cholesterol acyltransferase (LCAT). As part of the SPAP complex activates spermatozoa motility
DCXR	L-xylulose reductase	Q7Z4W1	Dehydrogenase, reductase	Steroid metabolic process	Catalyzes the NADPH-dependent reduction of several pentoses, tetroses, trioses, alpha-dicarbonyl compounds and L-xylulose
SPRE	Sepiapterin reductase	P35270	Dehydrogenase, reductase	Steroid metabolic process	Catalyzes the final one or two reductions in tetrahydrobiopterin biosynthesis to form 5,6,7,8-tetrahydrobiopterin
RAD50	DNA repair protein RAD50	Q92878	Oxidase, transferase	Homeostatic process	Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis
TFR1	Transferrin receptor protein 1	P02786	Receptor	Ion transport	Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system. Positively regulates T and B cell proliferation through iron uptake

Table 1 (Continued)

Protein code	Protein name	Uniprot code	Protein class	Pathways	Protein function
THTM	3-mercaptopyruvate sulfurtransferase	P25325	Transfer/carrier protein, transferase	Ion transport	Transfer of a sulfur ion to cyanide or other thiol compounds. In combination with cysteine aminotransferase (CAT), contributes to the catabolism of cysteine and is an important producer of hydrogen sulfide in the brain, retina and vascular endothelial cells. Hydrogen sulfide H ₂ S is an important synaptic modulator, signaling molecule, smooth muscle contractor and neuroprotectant. Its production by the 3MST/CAT pathway is regulated by calcium ions




Figure 2 Venn diagram of valid protein identifications with and without gradual exposition of the CCRF-SB cell line to 6 nM vincristine.

to which 30 different proteins were assigned (Table 1). Interestingly, 43% of the proteins in Table 1 and 53% of the pathways in Figure 3 were directly involved in signal transduction, whereas 13% of the proteins were involved in regulation of signaling and 20% in oxidative phosphorylation.

4. Discussion

This study addressed the question of which are the cellular processes that sensitive leukemic cells induced to achieve tolerance to vincristine. To this end, the B-ALL cell line CCRF-SB was gradually exposed until cell proliferation was




Figure 3 Functional categories overrepresented in adapted cells. The analysis was made with the statistical overrepresentation test of the PANTHER classification system as described in the Proteomic methods section.

observed in the presence of 6 nM vincristine, and the corresponding proteomic profile was compared to that of cells grown in the absence of the chemotherapeutic drug.

Chemoresistance may be intrinsic or acquired.¹⁸ The ability to tolerate high concentrations of chemotherapeutics of an intrinsically resistant cancer cell is not developed as a result of an exposition to the drugs; instead it is the result of genetic abnormalities the cell carries before exposition.¹⁸ By contrast, acquired chemoresistance is developed after

the cancer cell is exposed to the drug and may be the result of molecular evolution of resistant clones.¹⁹ Experimental settings to study acquired chemoresistance include the comparison of matched paired samples at diagnosis and after relapse²⁰ or the comparison of sensitive cell lines and resistant sublines which are obtained after prolonged exposure (several months) to the drug.²¹ It is likely that resistant clones may have evolved during the gradual adaptation protocol depicted in Figure 1B. However, the low growth rate, the short time of exposure (22 days) and the fact that it was highly reproducible when it was assayed from frozen stocks of sensitive cells suggest that the observed tolerance to vincristine was the result of induction of cellular processes to counteract the cytotoxic effect of the drug rather than the result of events involving mutation, selection, and evolution of resistant clones.

The cytotoxic effects of drugs are dependent on the concentration and the time of exposure. In many studies, cells have been exposed to relatively high concentrations for short times (24 or 48 h). In such experimental setting, at concentrations close to the IC₅₀, cells are not growing actively but remain viable and quiescent. Our results showed that even at concentration as low as 1 nM for five days, a reduction in cell viability was observed (Figure 1B). The gradual adaptation protocol used allowed cell division even after four days of exposition to 6 nM. By contrast, in cells exposed to 6 nM vincristine for 48 h without previous adaptation, no cellular division was observed (data not shown). In this regard, our results may be complementary to previous studies and may reflect some aspects of leukemic cells *in vivo* during advanced stages of therapy.

Successful adaptation was also reflected in the absence of overrepresented categories typically associated with cell death (Figure 3).

Interestingly, 43% of the proteins included in the overrepresented categories of adapted cells (namely MP2K1, MK14, RB1, RAC1, VAV, CASP3, ZAP70, VRK1, IMA4, IMA3, RAP1A, PIPSL, and STK25) were involved in signaling pathways like Toll receptor signaling pathway, Ras pathway, B- and T-cell activation, CCKR signaling map, p53 pathway feedback loops, p38 MAPK pathway, cytokine-mediated signaling pathway, cell surface receptor signaling pathway (Table 1); some of them have been reported to be important for growth and proliferation of lymphocytes and chemoresistance.^{22–24} For example, the MP2K1, MK14 and RAC1 proteins positively regulate the Ras pathway, and the MP2K1 protein is an essential component of the MAP kinase pathway. These signal transduction pathways have been associated with tumorigenesis and chemoresistance.^{22,25,26} Inhibition of the MP2K1 protein with PD-0325901, BAY86-9766, trametinib, selumetinib, pimasertib or GDC-0973 has been studied in advanced clinical trials.²² The MK14 protein has been reported to be closely associated with drug resistance in leukemia,²⁷ and the RAC1 protein has been proposed as a therapeutic target for gefitinib-resistant non-small-cell lung cancer and pancreatic cancer cells,^{28–30} and its overexpression has been associated with poor prognosis in non-small-cell lung cancer.³⁰ Deregulation of the VAV protein has been reported in neuroblastoma, melanoma, pancreatic, lung and breast cancers, and B-cell chronic lymphocytic leukemia.³¹ Furthermore, the proteins ZYX, AT2A2, CK5P3 and OXSR1, which were assigned to categories like *Nervous*

system development, *Homeostatic process and Regulation of catalytic activity* (Table 1), are modulators of signaling proteins.^{32,33} By contrast, in the control cells, any pathway resulted to be overrepresented when the statistical overrepresentation test was made comparing the 126 proteins detected exclusively in the absence of vincristine with the proteins detected in both groups (with and without vincristine). This difference may indicate that some signaling proteins and pathways may be crucial for adaptation to vincristine. In this regard, some drugs have been developed to target signaling pathways and to overcome drug tolerance.^{22,25,34} Further research is necessary to explore this possibility during the gradual adaptation to vincristine.

The cell-adhesion and mechanotransducer protein ZYX was assigned to the *Nervous system development* category (Table 1). However, it has been reported to participate in signal transduction pathways leading to cell migration, proliferation, and tumorigenesis.³² The ZYX protein is upregulated in human breast cancer and positively correlates with metastasis. It has been proposed as a potential therapeutic target in breast cancer and as an early detection biomarker in non-small cell lung cancer.^{32,35}

Verrills et al. reported the proteomic changes of the T-lineage ALL cell line CCRF-CEM after exposition to 2, 4 and 8 nM vincristine for 24 h using two-dimensional polyacrylamide electrophoresis and MALDI-TOF mass spectrometry.¹⁷ They found that vincristine-induced changes in the expression of 39 proteins and that a resistant subline differentially expressed 42 proteins mainly involved in cytoskeleton metabolism.¹⁷ Other differentially expressed proteins resulted to be regulators of apoptosis, chaperones, gene regulators and ribosomal proteins. The involvement of alterations in the metabolism of actin and tubulin during the cellular response to vincristine was confirmed in a subsequent study by the same group using primary cells engrafted into NOD/SCID mice and intraperitoneal vincristine (0.5 mg/kg every seven days).³⁶ Results shown in Figure 3 and Table 1 were based in comparisons of those proteins identified exclusively in the adapted cells versus the shared proteins (Figure 2). As cytoskeletal metabolism is very active in every cell, many cytoskeletal regulating proteins were among the 723 shared proteins; this may have prevented the overrepresentation of cytoskeletal metabolism in the adapted cells. To detect changes in the relative abundance of the shared proteins, a quantitative analysis is required. However, the microtubule-associated protein 4 (MAP4) was assigned to the overrepresented pathway *Cell differentiation* (Table 1). This protein promotes microtubule assembly and may reflect a more active cytoskeletal metabolism in the adapted cells compared to the control cells. Further research is necessary to explore this possibility.

After the combined effect of signaling pathways, *Oxidative phosphorylation* had the second highest percentage of proteins among the overrepresented pathways (Table 1). This suggested that a higher ATP production is needed to deal with the presence of vincristine. In this regard, modulation of the metabolic peculiarities of cancer cells has been proposed as a promising therapeutic strategy.^{37,38} The inhibitors metformin, rotenone, α -tocopheryl succinate, benzylisothiocyanate, oligomycin and resveratrol have been used to target mitochondrial energetic metabolism.³⁸ Our results

suggest that these compounds may be good candidates to overcome tolerance to vincristine.

Detection of the TFR1 and THTM proteins resulted in the overrepresentation of the *Ion transport* category in the adapted cells (Table 1 and Figure 3). The protein TFR1 may have contributed to B-cell growth through iron uptake.³⁹ Interestingly, this protein has been implicated in resistance to tamoxifen in a subgroup of ER+/luminal-like breast cancer.⁴⁰ Calcium is translocated from the cytosol to the sarcoplasmic reticulum lumen by the AT2A2 protein (assigned to the *Homeostatic process* category in Table 1). Inhibition of this protein has been proposed as a promising strategy to overcome multidrug-resistance in leukemic cells.³³ Moreover, calcium signaling modulates the activity of the THTM protein which participates in the production of hydrogen sulfide (Table 1). Interestingly, this compound has been reported to be involved in chemoresistance of lung adenocarcinoma cells, and inhibition of hydrogen sulfide-producing enzymes has been proposed as a strategy to sensitize resistant cells.⁴¹

DNA repair protein RAD50 was assigned to the overrepresented category *Homeostatic process* (Table 1). As part of the MRN complex, this protein has been proposed as a predictor for poor prognosis and chemoresistance in gastric cancer.⁴²

Unexpectedly, p-glycoprotein was not detected in our proteomic analysis of adapted cells. This may be due to the low relative abundance of this protein compared to other cellular proteins. To achieve higher sensitivity, peptide mixture can be fractionated in a previous HPLC run before the nanoHPLC-ESI-MS analysis.

Although monoculture of cell lines is a useful model for in vitro studies, the influence of the bone marrow microenvironment in chemoresistance⁴³ cannot be studied. In this regard, the co-culture of bone marrow stromal cells and leukemic cells⁴⁴ will certainly give more information of the involved mechanisms.

As a general rule, findings from the omic analyses must be confirmed with traditional techniques like Western blot, flow cytometry, ELISA or RT-PCR. Confirmation would be necessary only for those few genes of interest in a particular study. Although we used stringent criteria for protein identification in the MASCOT search engine, only those proteins identified in the three analytical replicates were considered as valid identifications, and only pathways with a fold enrichment value higher or equal than three were considered as overrepresented. Our results must be validated if they are to be used in further studies aimed to deepen into the molecular mechanisms involved in tolerance to vincristine.

Ethical responsibilities

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article

Funding

Federal funding HIM/2015/053 SSA1183, HIM/2016/026 SSA1222 for HQ and HIM/2014/003 SSA1129 for GPL. CONACYT grant 589666 for ALGO.

Conflict of interest

The authors declare no conflict of interest of any nature.

References

1. Rivera R, Cárdenas R, Olaya A, Shalkow J, Pérez M, Pérez OA, et al. El niño de población abierta con cáncer en México. *An Med.* 2015;60:91–7.
2. Ko RH, Ji L, Barnette P, Bostrom B, Hutchinson R, Raetz E, et al. Outcome of patients treated for relapsed or refractory acute lymphoblastic leukemia: a therapeutic advances in childhood leukemia consortium study. *J Clin Oncol.* 2010;28:648–54.
3. Pui CH, Yang JJ, Hunger SP, Pieters R, Schrappe M, Biondi A, et al. Childhood acute lymphoblastic leukemia: Progress through collaboration. *J Clin Oncol.* 2015;33:2938–48.
4. Nguyen K, Devidas M, Cheng SC, La M, Raetz EA, Carroll WL, et al. Factors influencing survival after relapse from acute lymphoblastic leukemia: a Children's Oncology Group study. *Leukemia.* 2008;22:2142–50.
5. Holohan C, Van Schaeybroeck S, Longley DB, Johnston PG. Cancer drug resistance: an evolving paradigm. *Nat Rev Cancer.* 2013;13:714–26.
6. Housman G, Byler S, Heerboth S, Lapinska K, Longacre M, Snyder N, et al. Drug resistance in cancer: an overview. *Cancers (Basel).* 2014;6:1769–92.
7. Risinger AL, Giles FJ, Mooberry SL. Microtubule dynamics as a target in oncology. *Cancer Treat Rev.* 2009;35:255–61.
8. Huang RS, Murry DJ, Foster DR. Role of xenobiotic efflux transporters in resistance to vincristine. *Biomed Pharmacother.* 2008;62:59–64.
9. Schlaifer D, Cooper MR, Attal M, Sartor AO, Trepel JB, Laurent G, et al. Myeloperoxidase: an enzyme involved in intrinsic vincristine resistance in human myeloblastic leukemia. *Blood.* 1993;81:482–9.
10. Fraser M, Leung B, Jahani-Asl A, Yan X, Thompson WE, Tsang BK. Chemoresistance in human ovarian cancer: the role of apoptotic regulators. *Rep Biol Endocrinol.* 2003;1:66.
11. Lee YY, Kim HP, Kang MJ, Cho BK, Han SW, Kim TY, et al. Phosphoproteomic analysis identifies activated MET-axis PI3K/AKT and MAPK/ERK in lapatinib-resistant cancer cell line. *Exp Mol Med.* 2013;45:e64.
12. Anderson NL, Anderson NG. Proteome and proteomics: new technologies, new concepts, and new words. *Electrophoresis.* 1998;19:1853–61.
13. Johnston HE, Carter MJ, Cox KL, Dunscombe M, Manousopoulou A, Townsend PA, et al. Integrated cellular and plasma proteomics of contrasting B-cell cancers reveals common, unique and systemic signatures. *Mol Cell Proteomics.* 2017;16:386–406.
14. Gerlier D, Thomasset N. Use of MTT colorimetric assay to measure cell activation. *J Immunol Methods.* 1986;94:57–63.
15. Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, et al. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res.* 2003;13:2129–41.

16. Cocker HA, Tiffin N, Pritchard-Jones K, Pinkerton CR, Kelland LR. In vitro prevention of the emergence of multidrug resistance in a pediatric rhabdomyosarcoma cell line. *Clin Cancer Res.* 2001;7:3193–8.
17. Verrills NM, Walsh BJ, Cobon GS, Hains PG, Kavallaris M. Proteome analysis of vinca alkaloid response and resistance in acute lymphoblastic leukemia reveals novel cytoskeletal alterations. *J Biol Chem.* 2003;278:45082–93.
18. Lippert TH, Ruoff HJ, Volm M. Intrinsic and acquired drug resistance in malignant tumors. The main reason for therapeutic failure. *Arzneimittelforschung.* 2008;58:261–4.
19. Giaccone G, Pinedo HM. Drug Resistance. *Oncologist.* 1996;1:82–7.
20. Kim HK, Choi IJ, Kim CG, Kim HS, Oshima A, Michalowski A, et al. A gene expression signature of acquired chemoresistance to cisplatin and fluorouracil combination chemotherapy in gastric cancer patients. *PLoS ONE.* 2011;6:e16694.
21. Rumjanek VM, Trindade GS, Wagner-Souza K, de-Oliveira MC, Marques-Santos LF, Maia RC, et al. Multidrug resistance in tumour cells: characterization of the multidrug resistant cell line K562-Lucena 1. *An Acad Bras Cienc.* 2001;73:57–69.
22. Neuzillet C, Tijeras-Raballand A, de Mestier L, Cros J, Faivre S, Raymond E. MEK in cancer and cancer therapy. *Pharmacol Ther.* 2014;141:160–71.
23. Zhang J, Mullighan CG, Harvey RC, Wu G, Chen X, Edmonson M, et al. Key pathways are frequently mutated in high-risk childhood acute lymphoblastic leukemia: a report from the Children's Oncology Group. *Blood.* 2011;118:3080–7.
24. Rak JW, Coomber B, Yu JL. Oncogenes and Tumor Suppressor Genes in Therapeutic Resistance, the Role of Evolving Interrelationships between Cancer Cells and Host Tissues. In: Teicher EA, editor. *Cancer Drug Resistance*. Totowa, NJ: Ed. Humana Press; 2006. p. 67–103.
25. McCubrey JA, Steelman LS, Chappell WH, Abrams SL, Wong EW, Chang F, et al. Roles of the Raf/MEK/ERK pathway in cell growth, malignant transformation and drug resistance. *Biochim Biophys Acta.* 2007;1773:1263–84.
26. Cox AD, Der CJ. Ras history: The saga continues. *Small GTPases.* 2010;1:2–27.
27. Gao F, Liu WJ. Advance in the study on p38 MAPK mediated drug resistance in leukemia. *Eur Rev Med Pharmacol Sci.* 2016;20:1064–70.
28. Kaneto N, Yokoyama S, Hayakawa Y, Kato S, Sakurai H, Saiki I. RAC1 inhibition as a therapeutic target for gefitinib-resistant non-small-cell lung cancer. *Cancer Sci.* 2014;105:788–94.
29. Yan Y, Hein AL, Etekpó A, Burchett KM, Lin C, Enke CA, et al. Inhibition of RAC1 GTPase sensitizes pancreatic cancer cells to gamma-irradiation. *Oncotarget.* 2014;5:10251–70.
30. Zhou Y, Liao Q, Han Y, Chen J, Liu Z, Ling H, et al. Rac1 overexpression is correlated with epithelial mesenchymal transition and predicts poor prognosis in non-small cell lung cancer. *J Cancer.* 2016;7:2100–9.
31. Katzav S. Vav1: A Dr. Jekyll and Mr. Hyde protein-good for the hematopoietic system, bad for cancer. *Oncotarget.* 2015;6:28731–42.
32. Ma B, Cheng H, Gao R, Mu C, Chen L, Wu S, et al. Zyxin-Siah2-Lats2 axis mediates cooperation between Hippo and TGF- β signalling pathways. *Nat Commun.* 2016;7:11123.
33. Bleeker NP, Cornea RL, Thomas DD, Xing C. A novel SERCA inhibitor demonstrates synergy with classic SERCA inhibitors and targets multidrug-resistant AML. *Mol Pharm.* 2013;10:4358–66.
34. Tasian SK, Teachey DT, Rheingold SR. Targeting the PI3K/mTOR pathway in pediatric hematologic malignancies. *Front Oncol.* 2014;4:108.
35. Kim YJ, Sertamo K, Pierrard MA, Mesmin C, Kim SY, Schlessner M, et al. Verification of the biomarker candidates for non-small-cell lung cancer using a targeted proteomics approach. *J Proteome Res.* 2015;14:1412–9.
36. Verrills NM, Liem NL, Liaw TY, Hood BD, Lock RB, Kavallaris M. Proteomic analysis reveals a novel role for the actin cytoskeleton in vincristine resistant childhood leukemia an in vivo study. *Proteomics.* 2006;6:1681–94.
37. Vander Heiden MG. Targeting cancer metabolism: a therapeutic window opens. *Nat Rev Drug Discov.* 2011;10:671–84.
38. Zhang Y, Yang JM. Altered energy metabolism in cancer: a unique opportunity for therapeutic intervention. *Cancer Biol Ther.* 2013;14:81–9.
39. Lo B. The requirement of iron transport for lymphocyte function. *Nat Genet.* 2016;48:10–1.
40. Habashy HO, Powe DG, Staka CM, Rakha EA, Ball G, Green AR, et al. Transferrin receptor (CD71) is a marker of poor prognosis in breast cancer and can predict response to tamoxifen. *Breast Cancer Res Treat.* 2010;119:283–93.
41. Szczesny B, Marcatti M, Zatarain JR, Druzhyna N, Wiktorowicz JE, Nagy P, et al. Inhibition of hydrogen sulfide biosynthesis sensitizes lung adenocarcinoma to chemotherapeutic drugs by inhibiting mitochondrial DNA repair and suppressing cellular bioenergetics. *Sci Rep.* 2016;6:36125.
42. Altan B, Yokobori T, Ide M, Bai T, Yanoma T, Kimura A, et al. High expression of MRE11-RAD50-NBS1 is associated with poor prognosis and chemoresistance in gastric cancer. *Anticancer Res.* 2016;36:5237–47.
43. Castells M, Thibault B, Delord JP, Couderc B. Implication of tumor microenvironment in chemoresistance: tumor-associated stromal cells protect tumor cells from cell death. *Int J Mol Sci.* 2012;13:9545–71.
44. Purroy N, Abrisqueta P, Carabia J, Carpio C, Palacio C, Bosch F, et al. Co-culture of primary CLL cells with bone marrow mesenchymal cells, CD40 ligand and CpG ODN promotes proliferation of chemoresistant CLL cells phenotypically comparable to those proliferating in vivo. *Oncotarget.* 2015;6:7632–43.