

Fossil freshwater gastropods from northern Mexico – A case of a “silent” local extirpation, with the description of a new species

Alexander Czaja, Alan P. Covich, José Luis Estrada-Rodríguez, Ulises Romero-Méndez, Jorge Saenz-Mata, Iris Gabriela Meza-Sánchez, Verónica Ávila-Rodríguez, Jorge Luis Becerra-López, David Ramiro Aguillón-Gutiérrez, José Gamaliel Castañeda-Gaytán

Alexander Czaja
 José Luis Estrada-Rodríguez
 Ulises Romero-Méndez
 Jorge Saenz-Mata
 Iris Gabriela Meza-Sánchez
 Verónica Ávila-Rodríguez
 Jorge Luis Becerra-López
 David Ramiro Aguillón-Gutiérrez
 José Gamaliel Castañeda-Gaytán
 drjlestrada@ujed.mx

Facultad de Ciencias Biológicas, Universidad Juárez del Estado de Durango, 35010 Gómez Palacio, Durango, Mexico.

Alan P. Covich
 Institute of Ecology, Odum School of Ecology,
 University of Georgia, Athens, GA 30602-
 2202, USA.

ABSTRACT

The decline of freshwater species diversity is a worldwide phenomenon but it is especially pronounced in arid regions such as in the Chihuahuan Desert of Coahuila and Durango, Northern Mexico. There are few historical data on the distribution but no data on extinctions of Mexican freshwater gastropods. In such cases, paleontological data can be very helpful and are often the only way to document past distributions and local extinction or extirpation events. This new paleoecological approach, called Conservation Paleobiology, can provide important perspectives in favor of conservation of recent analogue ecosystems. The study site contains subfossil (Late Holocene) deposits of the Laguna District, a small area in Coahuila and Durango. Here we describe a new species and report several new records of subfossil cochliopid, hydrobid, planorbisid and neritid gastropods. Most of these species (and some genera) have not been known as fossil from Mexico and all of them disappeared in very recent times. Together with data from our former paleomalacological studies, we present herein an example of a “silent” local extirpation in northern Mexico that began in the Middle Holocene. This loss of species apparently increased rapidly in the second half of the 20th century. Of 32 species of freshwater snails present in the area of study through the Holocene only four (12.5%) are still extant. At least 24 (75.0%) of them disappeared from the area in the 20th century due to habitat loss caused by anthropogenic activities. The results show that especially hydrobiid species with small geographic ranges (local endemics) are highly sensitive and vulnerable to such extirpation events. In this study area the extirpation of freshwater snails happened silently (unrecorded) while at the same time some two hundred kilometers farther north, one of the greatest recent hotspots of gastropods diversity in North America remained in the Cuatrociénegas Basin. The same imperilment that led to local extirpation in the area of study still threatens the freshwater snail communities of the Cuatrociénegas valley.

Keywords: Paleobiology, extirpation, gastropods, freshwater, loss of habitat.

RESUMEN

El declive de la diversidad de especies dulceacuicolas es un fenómeno mundial, pero es especialmente pronunciado en las regiones áridas como en el Desierto Chihuahuense de Coahuila y Durango, norte-centro de México. Existen pocos datos históricos sobre la distribución y ninguna información sobre extinciones de gasterópodos dulceacuicolas mexicanos. En tales casos, los datos paleontológicos pueden ser muy útiles y, a menudo, son la única forma de documentar las distribuciones pasadas y los eventos locales de extinción u extirpación. Este nuevo enfoque paleoecológico, llamado Paleobiología de Conservación, puede proporcionar perspectivas importantes para los esfuerzos de conservación de ecosistemas análogos recientes. El sitio de estudio contiene depósitos fósiles (Holoceno Tardío) del Distrito Laguna, un área pequeña en Coahuila y Durango. De estos depósitos fósiles describimos una nueva especie y reportamos varios registros nuevos de gasterópodos cochliópodos, hidróbidos, planórbidos y neritidos. La mayoría de estas especies y algunos géneros no se conocen como fósiles de México y todos ellos desaparecieron en tiempos muy recientes. En conjunto con datos de nuestros estudios paleomalacológicos anteriores, presentamos aquí un ejemplo de una extirpación local “silenciosa” en el norte de México la cual comenzó en el Holoceno Medio. Esta pérdida de especies aumentó rápidamente en la segunda mitad del siglo XX. De las 32 especies de caracoles dulceacuicolas presentes en el área de estudio durante el Holoceno, solo 4 (12.5%) aún existen. Al menos 24 (75.0%) de ellos desaparecieron del área en el siglo XX debido a la pérdida de hábitat causada por actividades antropogénicas. Nuestros resultados muestran que especialmente los hidróbidos con intervalos geográficos pequeños (especies endémicas locales) son altamente sensibles y vulnerables a tales acontecimientos de extirpación. La extirpación de los caracoles dulceacuicolas en el área de estudio ocurrió silenciosamente (sin registros), mientras que al mismo tiempo unos doscientos kilómetros más al norte, uno de los mayores focos de diversidad de gasterópodos en América del Norte permaneció en la cuenca de Cuatrociénegas. Las mismas amenazas que llevaron a la extirpación local en nuestra área de estudio aún amenazan a las comunidades de caracoles dulceacuicolas del valle de Cuatrociénegas.

Palabras clave: Paleobiología, extirpación, gasterópodos, dulceacuicolas, pérdida de hábitat.

BOL. SOC. GEOL. MEX. 2019
 VOL. 71 NO. 3
 P. 609 – 629
<http://dx.doi.org/10.18268/BSGM2019v71n3a2>

Manuscript received: August 29, 2018.
 Corrected manuscript received: September 25, 2018.
 Manuscript accepted: October 2, 2018.

1. Introduction

The extinction of freshwater snails due to habitat loss is of wide interest and occurs especially among endemic species with restricted distributions, such as those in karst landscapes where lakes, groundwater-fed springs, and other habitats are threatened (e.g., Strong *et al.* 2008, Régnier *et al.* 2009). The distribution of extant freshwater gastropods of northern Mexico is poorly known and freshwater habitats are rapidly being altered by water diversions and dam construction. Only a portion of living species have been studied ecologically, anatomically or using molecular methods. Recent advances have focused mainly on systematics and biogeography of newly collected extant cochliopid and hydrobiid snails or species previously known only from shell morphology (Hershler *et al.*, 2002, 2011a, 2011b, 2014a, 2014b). Without these data our systematic research affiliation of a part of the subfossil material would be much more difficult, or, as in some cases of shell convergences, almost impossible.

The study area is located in the central part of the Laguna District, a small karst region and it comprises the adjoining portions of western Coahuila and eastern Durango states, northern Mexico (Figure 1). The region, a large endorheic basin, was one of the largest paleo-lakes in Latin America during Late Pleistocene/Early Holocene (Butzer *et al.*, 2008; Czaja *et al.*, 2014b). Although this lake had become significantly smaller during the Holocene, large lakes were still present until the early 1900s (Ortega-Guerrero, 2003; Butzer *et al.*, 2008). In the first half of the 20th century the region became an important center of agriculture and industry in Mexico. This economic process was accompanied by habitat perturbations such as damming and channelization of the two rivers, alteration of springs, and especially by excessive groundwater pumping. This aquifer depletion, described in detail by Wolfe (2013), finally eliminated the habitats provided by the remaining inland lakes and springs in the area of study.

From this semi-desert and karst region of south-

west Coahuila and northeast Durango, we report subfossil shells belonging to cochliopid, hydrobiid, planorbisid and neritid freshwater gastropod genera. The aim of the present study is to describe a new species, report new records and show their original biogeographic distribution prior to the loss of their habitats in recent times. To achieve that, we follow the “near-time” approach *sensu* Dietl and Flessa (2011) which utilizes the fossil record to understand the present-day conditions. In Mexico, this new interdisciplinary approach was applied so far only once (Figuerola-Rangel *et al.*, 2012) but its potential has already been pointed out in detail by Guerrero-Arenas and Jiménez-Hidalgo (2015).

2. Material and methods

Shells of *Juturnia gracilis* n. sp., *J. coahuilae* (Taylor, 1966), *Pseudotryonia pasajae* (Hershler, Liu and Landye, 2011), *Promenetus exacuus* (Say, 1821) and *Tryonia* cf. *seemani* (Frauenfeld, 1863) were collected in present-day superficial sediments of dried-out springs near to the town of Viesca, Coahuila (Figure 1, site 2). All springs were still active in 1959 and dried out in the following years (Blásquez, 1959; Wolfe, 2013). Shells of *Biomphalaria havanensis* (Pfeiffer, 1839) were collected at west of Viesca springs in superficial sediments of a former wetland area of these springs (Figure 1, site 3). Shells of both species of *Probythinella* Thiele, 1929, *Cochliopina riograndensis* (Pilsbry and Ferriss, 1906) and *Dilatata dilatata* (Gould, 1841) were collected in fluvial deposits of the Aguanaval River and in lacustrine sub-recent deposits of the paleo-lake Viesca (Figure 1, sites 5 and 4). A single shell of *Vitta* cf. *usnea* (Röding, 1798) was collected in a gray silt horizon with abundant planorbisid snails of the former Paleo-Lake Irritila (Figure 1, site 1). These sediments with *Vitta* were not dated directly but the same horizon with abundant snails from a neighboring site was dated by Butzer *et al.* (2008) as Holocene playa deposits with radiometric calibrated ^{14}C ages between 3320 ± 100 at the top of the section and not older than ~7130 years BP at the bottom.




Figure 1 Map of the study area with localization of the sampled sites of the Laguna District of Coahuila and Durango. 1 = Holocene playa deposits of the Paleo-Lake Irritila (Laguna Mayrán). 2 = Springs of Viesca. 3 = Former wetland area of the springs of Viesca. 4 = Lacustrine sub-recent deposits of the Paleo-Lake Viesca. 5 = Sub-recent fluvial deposits of the Aguanaval River. Dotted squares = Area of study.

For a comparison of the subfossil material, empty shells of the extant species *Pseudotryonia pasajae* were collected in September 2016 from an unnamed spring in San Pedro de Ocuila, Durango, which is the type locality for this species. Empty shells of the extant *Juturnia coahuilae* from Poza Churince in Cuatrociénegas were collected in summer 2017 directly from sediments of this spring. Only the outflows of the dams of the Nazas and Aguanaval rivers and the small spring La Peña still contain permanent water in the area of study. Table 1 contains the coordinates of the localities that are monitored by the authors since 2009. These three sites were last sampled in summer 2017. The snails were hand collected from stones, sand and aquatic vegetation, 4 kg sediment per site was screened through sieves.

To visualize morphological shell differences between subfossil and living populations, a discriminant analysis was carried out. This was done

using the software PAST (Paleontological Statistics) v 3.14 (Hammer *et al.*, 2001). This technique has frequently been used to differentiate closely similar shells of other hydrobiid and cochliopid gastropods (Czaja *et al.*, 2017b). We used following shell morphometric dataset (excluding ratios): Total number of whorls, shell height, shell width, aperture height, aperture width, height of body whorl and width of body whorl. The mean, standard deviation and sample size are given in text (shell measurements).

Sediments of all localities were screened through two sieves with 0.5 mm and 0.3 mm mesh size. Some specimens (especially the apertures of the shells) were cleaned using hydrogen peroxide to remove sediments. All shells were photographed with a Zeiss AxioCam ERc5s-camera attached to a Zeiss Stemi 2000-C microscope. Morphometric variables of the shells were measured using the same microscope. Shell whorls were counted

Table 1. The coordinates of three localities that still contain permanent water in the study area (outflows of the dams of the Nazas and Aguanaval rivers and the small spring La Peña).

| | Nazas River | Aguanaval River | La Peña spring |
|--------|---------------------------|---------------------------|---------------------------|
| Site 1 | 25°27'56" N, 103°42'13" W | 25°07'49" N, 103°22'22" W | 25°27'02" N, 102°35'10" W |
| Site 2 | 25°27'55" N, 103°42'12" W | 25°13'14" N, 103°26'58" W | |
| Site 3 | 25°39'30" N, 103°22'51" W | 25°01'36" N, 103°16'09" W | |

according to the method of Pilsbry (1939). The material is housed in the Malacological Collection of the Faculty of Biological Science of the Juárez State University of Durango (UJED). Paratypes of the new species were deposited in the National Collection of Mollusks, National Autonomous University of Mexico (CNMO).

Abbreviations used in the text: WN, total number of whorls; SH, shell height; SW, shell width; AH, aperture height; AW, aperture width; HBW, height of body whorl; WBW, width of body whorl; sd, standard deviation; UJMC = University Juárez Malacological Collection.

3. Results

3.1. SYSTEMATICS

We emphasize that in the present paper we apply the morphological species concept widely used for paleobiological and paleontological studies. However, our morphospecies definitions are also based on insight from modern species concepts of living species that includes biogeographic considerations and population genetic aspects. A combination of neo- and paleobiological approach is mutually supportive in these determinations.

Superfamily Neritacea Lamarck, 1809

Family Neritidae Lamarck, 1809

Subfamily Neritinae Poey, 1852

Genus *Vitta* Mörch, 1852

Type species: *Nerita virginea* Linnaeus, 1758

Vitta cf. *usnea* (Röding, 1798)

(Figure 2a to 2c)

Description: Shell dextral, ovoid, with flat spire, height 7.82 mm, width 6.79 mm, with 3.00 whorls, whorls rapidly increasing in size, bodywhorl constitutes most of shell, surface smooth and porcelainous; sutures faintly impressed, with numerous microscopic axial growth lines and no spiral sculpturing, color pattern visible in the area of the sutures consisting of dark, wavy lines, which are closely spaced on a whitish background; aperture ovate, the outer lip is thickened and without teeth, parietal shelf straight with 12 (or 13?) irregular teeth (Figure 2c), teeth absent from the top and the bottom of the inner lip; numerous thin, parallel, brown to black axial lines as remains of the original color patterning (thin, parallel axial lines) visible only near the sutures (Figure 2d).

Shell Measurements: SH = 7.82 mm, SW = 6.79 mm, AH = 5.49 mm, AW = 3.67 mm, WN = 3.00.

Material Examined: 1 specimen (UJMC 360).

Distribution: Florida to Texas, Mexico (gulf coast), West Indies (Eichhorst, 2016).

Remarks: Because it is a single specimen without color and pattern on the shell, our material can only be identified tentatively. Further, the specimen shows the teeth on the columellar lip much more defined than on modern *Vitta usnea*. The outer shell layer on the parietal shelf that leads up to these teeth probably has been broken off, exposing more of the teeth than is normally visible. The species is reported as *Neritina reclinata* (Say, 1822) in much of the recent literature, but we follow Eichhorst (2016) and placed this species in the genus *Vitta* because it lacks the characteristics




Figure 2 *Vitta cf. usnea* (Röding, 1798) and *Cochliopina riograndensis* (Pilsbry & Ferriss, 1906). a-e) *V. cf. usnea*, UJMC 360. d) Black axial lines as remains of the original color patterning. e) Parietal shelf with irregular teeth. f-h) *Cochliopina riograndensis*. f) Specimen UJMC361a. g) Specimen UJMC 361b with cords on the bodywhorl. h) Specimen UJMC 361c, protoconch view.

of *Neritina*. *V. usnea*, as it occurs in fresh/brackish water, on rocks, plant stems, or other hard surfaces and was found living in freshwater 160 km inland in Florida (Eichhorst, 2016). However, its Late Holocene presence in the study area almost 1000 km from the coast is notable. Although the species has a wide ecological range, occurrences in freshwater at long distances from the coast habitats are not known in its present-day distribution. Last record in the area: Middle/Late Holocene (radiometric calibrated ^{14}C ages between 3320 ± 100 and ~ 7130 years BP, Butzer *et al.*, 2008).

Superfamily Truncatelloidea Gray, 1840

Family Cochliopidae Tryon, 1866

Genus *Cochliopina* Morrison, 1946

Type species: *Cochliopa riograndensis* Pilsbry & Ferriss, 1906 (by original designation)

Cochliopina riograndensis (Pilsbry and Ferriss, 1906)
(Figure 2f to 2h)

Description: Shell small, broadly heliciform, height 1.39–2.02 mm, width 1.91–3.01 mm, openly umbilicate; apical whorl with pitted structure; teleoconch nearly smooth or frequently with spiral threads or (two) cords on the bodywhorls, one thread at the shoulder usually more prominent (Figure 2g); whorls moderately convex with deep sutures; the aperture is roundly ovate and angled adapically; peristome thin, inner lip partly fused to the penultimate whorl.

Shell Measurements: (mean, with sd in parentheses; N = 10) SH = 1.65 (0.28) mm, SW = 2.41 (0.31) mm, AH = 1.15 (0.08) mm, AW = 1.02 (0.08) mm, WN = 3.40 (0.21).

Material Examined: 10 specimens (UJMC 361a-j).

Distribution: Streams in Val Verde County, Texas and Northwest Mexico (Coahuila, San Luis Potosí, Tamaulipas) (Thompson, 2011).

Remarks: The subfossil shells from Durango and Coahuila correspond morphologically in all details to those from the type locality in Texas. Pilsbry and Ferriss (1906) noticed in the original description of *C. riograndensis* a great similarity to shells of the genus *Valvata* (Müller, 1773), but they

emphasized that species of the last genus have rounded apertures. According to Riedel (1993), the protoconchs of *Valvata* exhibit as well typical valvatid sculpture in form of numerous spiral lirae which are radially interconnected. Both shell features are not present in our material so we place the shells into the genus *Cochliopina*.

Genus *Juturnia* Hershler, Liu and Stockwell, 2002
Type species: *Durangonella coahuilae* Taylor, 1966 (by original designation)

Juturnia gracilis Czaja, Covich, Estrada-Rodríguez and Romero-Méndez sp. nov.
(Figure 3a to 3g)

Diagnosis: Shell medium-sized, turritiform, fragile, light tan or transparent, teleoconch of 8.00 to $11\frac{1}{2}$ strongly convex whorls; teleoconch without sculpture, only with very fine and closely spaced growth lines; aperture ovate, inner and outer lip thin; umbilicus rimate to broadly open.

Description: Shell turritiform, medium-sized, 4.82–7.32 mm shell length and 1.29–1.78 mm shell wide, with 8.00 to $11\frac{1}{2}$ strongly convex and smooth whorls; teleoconch sculpture of closely spaced growth lines, from light tan to beige-white or transparent, in some specimens from light tan to brown periostracum preserved; protoconch of 0.7–1.0 whorl, smooth (Figure 3g); aperture ovate, inner lip thin to slightly thickened, broadly adnate to completely separate from parietal wall, outer lip thin or slightly thickened, the peristome is complete; umbilicus rimate to broadly open.

Shell Measurements: (mean, with sd in parentheses; N = 15) SH = 5.75 (0.61) mm, SW = 1.49 (0.13) mm, AH = 1.11 (0.11) mm, AW = 0.95 (0.08) mm, HBW = 2.05 (0.21) mm, WBW = 1.29 (0.12) mm, WN = 10.07 (0.62).

Measurements of Holotype: SH = 5.83 mm, SW = 1.47 mm, HBW = 1.97 mm, WBW = 1.30 mm, AH = 1.07 mm, AW = 1.01 mm, WN = $10\frac{1}{4}$.

Differential Diagnosis: Of the four extant species of the genus *Juturnia*, the new species only resembles *J. coahuilae* but differs clearly from this by having larger shells with considerably more whorls (*J. coahuilae* is possibly the described freshwater




Figure 3 *Juturnia gracilis* new species, *Juturnia coahuilae* (Taylor, 1966) Hershler, Liu and Stockwell, *Probythinella emarginata* (Küster) and *Probythinella protera* Pilsbry. a, b, g) *J. gracilis* new species, Holotype, UJMC-362. g) Protoconch view. c, d) Paratype 1, UJMC-363. e, f) Paratype 2, UJMC-364. h-j) *Juturnia coahuilae* (Taylor, 1966) Hershler, Liu and Stockwell from spring Tunel 7, Viesca, Coahuila, Mexico, UJMC-366a. j) Protoconch view. k-m) *J. coahuilae* from Poza Churince, Cuatrociénegas, Coahuila, Mexico, a present-day specimen. m) Protoconch view. n-p) *Probythinella emarginata*, UJMC-367a. p) Protoconch view. q-t) *Probythinella protera*. q, r) Specimen UJMC-368a. s) Specimen UJMC-368b. t) Protoconch view, UJMC-368a.

snail with most whorls in North America). Reliable diagnostic distinctions between these two species are as well the whorl height/whorl width ratio that is 0.21–0.29 in *J. gracilis* n. sp. and 0.30–0.40 in *J. coahuilae*. As a result of this different ratio, *J. gracilis* n. sp. has a narrower general form than *J. coahuilae* (Figure 3a to 3f). Also, the discriminant analysis (Figure 4), based on seven standard shell morphometric data, confirms that the subfossil shells of *J. gracilis* n. sp. described herein are morphologically clearly distinct to shells of the extant endemic species from the Cuatrociénegas Basin. A combination between the detailed morphological description and morphometric based discriminant analyses were efficient in distinguishing among the recent and subfossil species.

Type Material: Holotype (Figures 3a, 3b, 3g), UJMC-362. Paratype 1 (Figures 3c, 3d), UJMC-363, Paratype 2 (Figures 3e, 3f), UJMC-364. (leg. Alexander Czaja and José Luis Estrada-Rodríguez, 25/vi/2016). All from type locality.

Type Locality: Mexico, Coahuila state, Viesca, Tunel 4 Spring (25° 20' 20" N, 102° 52' 32" W, 1099 m a.s.l.).

Stratum Typicum: Spring sediments (layer c. 5 cm below the present-day surface). Latest Holocene, sub-recent.

Other Material Examined: 12 specimens from the same locality (UJMC-365a-l).

Etymology: The species was named after the slender (lat. *gracilis*) general shape of its shells.

Distribution: *J. gracilis* n. sp. is endemic to the Túnel 4 spring at Viesca, Coahuila, Mexico.

Remarks: Hershler *et al.* (2002) mentioned that shells of *Juturnia* broadly overlap in size, shape, and sculptural features with those of several other epigeal cochliopines of southwestern North America such as *Eremopyrgus*, *Littoridinops*, *Pseudotryonia* and *Tryonia*. However, such narrow and turri-form shells with numerous whorls (more than 7) are reported only from extant members of *Juturnia* and *Tryonia*. Both genera are difficult to separate




Figure 4 Graphic depiction of the discriminant analysis of morphometric variables of three populations of *Juturnia*. Circles = *Juturnia gracilis* new species, subfossil population from Spring Tunel 4, Viesca, Coahuila. Square = *J. coahuila* (Taylor, 1966), a subfossil population from Spring Tunel 7, Viesca, Coahuila. Triangles = *J. coahuila*, a present-day population from Poza Churince, Cuatrociénegas, Coahuila.

by shells and it seems that conchologically only the well-developed growth lines on *Juturnia* shells is the unique useful differentiating feature between both genera. Of the three living species of the genus, *J. gracilis* n. sp. closely resembles *J. coahuilae* (Taylor, 1966), an extant endemic species from the Cuatrociénegas Basin. In neighboring springs and spring-outflows in Viesca subfossil shells of *J. coahuilae* are very common (see below). This geographical proximity and the similarity of the shell structure of *J. coahuilae* and *J. gracilis* n. sp. reflect undoubtedly a narrow phylogenetic relationship between both species. We suppose that the subfossil *J. gracilis* n. sp. has evolved from *J. coahuilae* in a separated spring (allopatric evolution) but a confirmation of this hypothesis requires further study of the vertical section of the spring deposits. According to Hershler *et al.* (2002), all four extant species of *Juturnia* are associated with saline environments. This can also be assumed for the former springs at Viesca since the groundwater, which once fed the springs with water, had (and has) high levels of salinity (Ortega-Guerrero, 2003).

Juturnia coahuilae (D. W. Taylor, 1966)

(Figure 3h to 3j)

Durangonella coahuilae Taylor, 1966

Durangonella coahuilae Hershler, 1985, 81, fig. 32

Description: Morphologically, the subfossil shells are identical to those described by Taylor (1966) and Hershler (1985) from the type locality of this species.

Shell Measurements: (mean, with sd in parentheses; N = 15) SH = 3.55 (0.34), SW = 1.27 (0.10) mm, AH = 0.95 (0.13) mm, AW = 0.76 (0.04) mm, HBW = 1.80 (0.13) mm, WBW = 1.19 (0.09) mm, WN = 6.12 (0.21).

Material Examined: 15 specimens (UJMC 366a-o).

Distribution: The present-day species is endemic to the spring complex of the Cuatrociénegas basin, Coahuila.

Remarks: The subfossil *J. coahuilae* from Viesca corresponds in its conchological features in almost

all details to the recent species from Cuatrociénegas (Figures 3h to 3m). The unique difference between the subfossil and extant species is the slightly narrower form of the subfossil species, which is also reflected in the discriminant analysis (Figure 4). However, these small morphological differences are, in view of the temporal and spatial distance of both populations, not surprising and we consider them within the variation of the extant species from Cuatrociénegas. The present record extends the (fossil) range of the genus *Juturnia* c. 300 km to the south.

Genus *Probythinella* Thiele, 1928

Type species: *Paludina emarginata* Küster, 1852

Probythinella emarginata (Küster, 1852)

(Figure 3n to 3p)

Description: Shell small, sub-globose to narrow-conic or sub-cylindrical, height ranging from 3.24–4.69 mm, width 2.24–3.18 mm; with $4\frac{3}{4}$ – $5\frac{1}{4}$ whorls; whorls inflated, slightly convex to well rounded, sometimes slightly shouldered below suture, suture impressed; apex blunt or depressed, often tilted; body whorl rarely abruptly descending toward end of growth; teleoconch sculptured with strong, regularly spaced collabral striations extending from the suture to the base of whorl; first whorl sometimes planorboid, depressed; aperture ovate, sometimes weakly angled above, peristome usually narrowly adnate, or slightly separated, outer and inner lips thin; shell clear-white, some specimens transparent.

Shell Measurements: (mean, with sd in parentheses; N = 10) SH = 4.11 (0.63) mm, SW = 2.75 (0.41) mm, AH = 1.87 (0.18) mm, AW = 1.45 (0.20) mm, HBW = 3.05 (0.34) mm, WBW = 2.43 (0.30) mm, WN = 4.78 (0.25).

Material Examined: 10 specimens (UJMC 367a-j).

Distribution: Eastern North America in lotic and lentic inland habitats throughout the Mississippi River basin; drainages of Great Lakes, Hudson Bay, and Mackenzie River to the north (Hershler, 1996).

Remarks: The genus *Probythinella* includes only two extant species: *P. emarginata* and *P. protera*, both distributed along the eastern part of North America. Both species are well characterized by the extremely blunt apex of their shells (Figures 3p, 3t), a synapomorphy of this small group of North American Hydrobiidae (Hershler, 1996). Our records spread out the (fossil) range of *P. emarginata* c. 300 km to the south.

Probythinella protera Pilsbry, 1953

Type species: *Amnicola emarginata* Küster, 1852
(Figures 3q to 3t)

Description: Shell medium in size, ovate-conic to pupiform, height ranging from 2.54–2.92 mm, width 1.51–1.82 mm; with $4\frac{1}{4}$ – $4\frac{1}{2}$ slightly rounded whorls (accuracy to a $\frac{1}{4}$ whorl), first whorl almost planorbiform; umbilicus absent; apex blunt, sometimes depressed; aperture small, ovate and inclined approximately 40° away from the coiling axes, inner lip thickened and adnate to the parietal wall, outer lip strongly thickened.

Shell Measurements: (mean, with sd in parentheses; N = 10) SH = 2.69 (0.13) mm, SW = 1.69 (0.10) mm, AH = 1.16 (0.04) mm, AW = 0.99 (0.07) mm, HBW = 2.13 (0.12) mm, WBW = 1.64 (0.10) mm, WN = 5.50 (0.12).

Material Examined: 10 specimens (UJMC 368a-j).

Distribution: Endemic to United States, brackish waters along the eastern Gulf of Mexico coast (Hershler, 1996).

Remarks: Hershler (1996) mentioned that the recent distributions of both *Probythinella* species appear to be contiguous, but non-overlapping. That does not apply to our area, where both species occurred during the Holocene in lotic and lentic habitats as sympatric species. *P. protera* occurs in present-day only in coastal regions but our findings confirm that this species was, during the Holocene, also abundant in inland sites.

Genus *Pseudotryonia* Hershler, 2001

Type species: *Tryonia alamosae* Taylor, 1987 (by original designation)

Pseudotryonia pasajae Hershler, Liu and Landye, 2011

(Figure 5a to 5d)

Pseudotryonia pasajae Hershler *et al.*, 2011: 14, 16, figures. 2E–F, I–J, M–O, 3E–H, 4D–E.

Pseudotryonia pasajae Hershler *et al.*, 2014: 59–60.

Description: Morphologically, the shells are identical to those described by Hershler *et al.* (2011b, p. 16) from the type locality of this species in Durango.

Shell Measurements: (mean, with sd in parentheses; N = 11) SH = 2.21 (0.50) mm, SW = 1.38 (0.22) mm, AH = 0.98 (0.14) mm, AW = 0.77 (0.08) mm, HBW = 1.64 (0.26) mm, WBW = 1.23 (0.19) mm, WN = 4.23 (0.54).

Material Examined: 11 specimens (UJMC 369a-k).

Distribution: *P. pasajae* occurs living only in two springs northwest of Cuencamé de Cenicerós, Rio Nazas basin, Durango, Mexico (Hershler *et al.*, 2014a).

Remarks: The results of the discriminant analysis did not reveal significant morphological differences among the subfossil shells from Viesca and shells of the endemic extant species *P. pasajae* from an unnamed spring at San Pedro de Ocuila, Durango, near the type locality (Figure 6). Almost all morphometric characters are within the phenotypic variation of the recent species. Small differences between the subfossil and extant populations remain only in the size of the shells, but these differences are even smaller than differences among the two known populations described by Hershler *et al.* (2014a) from the type locality and they are also visible in the discriminant analysis (Figure 6). The same author mentioned that *P. pasajae* shows sexual dimorphism in form of shells size differences, which is also observed in our material. Since all other shell characteristics of both forms are similar, we conclude that they may reflect the sexual dimorphism of the species. The subfossil shells were found in spring El Molino, Viesca, Coahuila, c. 100 km NE from the type locality of this species in Durango.

Genus *Tryonia* Stimpson, 1865

Type species: *Tryonia clathrata* Stimpson, 1865 (by original designation)

Tryonia cf. *seemani* (Frauenfeld, 1863)

(Figure 5i to 5k)

Hydrobia seemani Frauenfeld, 1863: 1025. Frauenfeld, 1865: 525, pl. 8.

Durangonella seemani (Morrison, 1945): Morrison, 1945, 19, pl. 3, Fig. 1

Description: Shell turritiform, medium-sized, height 5.99 mm, with 5½ rounded whorls. Teleoconch whorls highly convex, evenly rounded with strong impressed sutures. Sculpture of fine growth lines; aperture pyriform, parietal lip complete; umbilicus absent; periostracum tan.

Shell Measurements: SH = 5.99 mm, SW = 2.54 mm, AH = 2.09 mm, AW = 1.56 mm, HBW = 3.64 mm, WBW = 2.25 mm, WN = 5½.

Material Examined: 1 specimen (UJMC-370).

Distribution: Endemic to a few springs near Durango City, Durango State (Hershler *et al.*, 2011a).

Remarks: Our shell closely resembles shells from Durango described originally as *Hydrobia seemani* by Frauenfeld (1863). This species was later redescribed and placed as type species in the new erected genus *Durangonella* by Morrison (1945). Hershler *et al.* (2002) finally examined the rehydrated material and transferred this species to the genus *Tryonia* (Figure 5m). Our single high spired and strongly convex-whorled specimen is




Figure 5 *Pseudotryonia pasajae* Hershler, Liu and Landye, 2011 and *Tryonia* cf. *seemani* (Frauenfeld, 1863). a-d) *P. pasajae*, specimen UJMC-369a. c) Male (?) specimen, UJMC-369b. d) Protoconch view, UJMC-369a. e-h) *P. pasajae*, a present-day specimen from San Pedro de Ocuila, Durango, Mexico. h) Protoconch view. i-m) *T. seemani*. i-k) *T. cf. seemani*, subfossil specimen from Viesca, Coahuila, Mexico, UJMC-370. k) Protoconch view. l) *T. seemani* from Durango, syntype (BMNH 20001099) illustrated by Hershler *et al.* (2002, Fig. 1b). m) *T. seemani*, a present-day specimen from an unnamed spring on west side of Nombre de Dios, Durango, Mexico (Hershler *et al.*, 2011a, Fig. 12c).

morphologically near identical to the syntypes of the species from Durango illustrated by Hershler et al. (2002) (Figure 5l) and can therefore be considered as conspecific. A convergence is unlikely, however, since it is a single specimen we identify the subfossil shell from Viesca only tentatively. The single specimen was found in superficial sediments of the Juan Guerra Spring, Viesca.

Family Planorbidae H. and A. Adams, 1855

Genus *Biomphalaria* Preston, 1910

Type species: *Biomphalaria smithii* Preston, 1910

Biomphalaria havanensis (Pfeiffer, 1839)

(Figure 7a to 7c)

Biomphalaria obstructa (Morelet 1849): Test. Noviss.

I: 16.

Biomphalaria “temascalensis” Rangel-Ruiz 1987: 25–34; pl. 1.

Description: Shells medium to large (maximum diameter 13.25 mm), flattened, discoidal; with $4\frac{3}{4}$ – $5\frac{3}{4}$ rounded whorls that increase moderately

in diameter; spire flattened; sculpture of fine lines of growth pronounced on the bodywhorl, whorls separated by deep sutures; aperture ovate, the peristome is thin, without apertural lamellae.

Shell Measurements: (mean, with sd in parentheses; N = 8) SH = 3.41 (0.20) mm, SW (= shell diameter) = 12.10 (0.83) mm, WN = 5.19 (0.35).

Material Examined: 8 specimens (UJMC 371a-h).

Distribution: The species occurs in southeastern USA, West Indies, Mexico and Central America. South American distribution remains to be determined (Thompson, 2011). According to Naranjo-García (2003), *B. havanensis* is the most common species of the genus in Mexico and inhabits mainly in areas along the Atlantic and Pacific coasts.

Remarks: The taxonomic history and systematics of the recent members of the genus *Biomphalaria* are long and confusing (DeJong et al. 2001). The subfossil shells from Viesca resemble shells of two extant species reported frequently from United States and Mexico: *B. havanensis* and *B. obstructa*.




Figure 6 Graphic depiction of the discriminant analysis of morphometric variables of three populations of *Pseudotryonia*. Square = *P. pasajae*, a subfossil population from Spring Tunel 4, Viesca, Coahuila. Circles = *P. pasajae*, a present-day population from an unnamed spring at San Pedro de Ocuila, Durango. Triangles = *P. pasajae*, a present-day population from el Tanque, Durango, type locality (data from Hershler et al., 2011b).

Both species were considered for a long time as different species (DeJong *et al.*, 2001; Thompson, 2011) but recent molecular studies show that *B. obstructa* (as well as *B. tamascalensis* Rangel-Ruiz from Oaxaca, Mexico) is a junior synonym of *B. havanensis* (Aguilar-Silva *et al.*, 2014; Rosser *et al.*, 2016).

Genus *Dilatata* Clessin, 1885

Type species: *Planorbis dilatatus* Gould, 1841

Dilatata dilatata (Gould, 1841)

(Figure 7d to 7f)

Description: Shell small, ultradextral, flattened, discoidal (flat spiral), with few rapidly enlarging whorls; body whorl with less well-developed carina (keel), placed just above the center of the body;

with 3 whorls (accuracy to a $\frac{1}{4}$ whorl), whorls rapidly increasing in size, sculpture of fine lines of growth; umbilicus deep; aperture large, expanded; lips slightly thickened.

Shell Measurements: Specimen 1: SH = 1.63 mm, SW (= shell diameter) = 3.23 mm, AH = 1.28 mm, AW = 1.59 mm, WN = 3. Specimen 2: SH = 1.22 mm, SW (= shell diameter) = 2.68 mm, AH = 1.11 mm, AW = 1.31 mm, WN = 2.75.

Material Examined: 2 specimens (UJMC 372a-b).

Distribution: *Dilatata dilatata* has a broad present-day distribution across United States from Florida and Texas to Canada (Baker, 1945; Burch, 1989). In Mexico in Zacatecas and Puebla (Albrecht *et al.*, 2007; Thompson, 2011 as *Micrometetus dilatatus*).




Figure 7 *Biomphalaria havanensis* (Pfeiffer, 1839), *Dilatata dilatata* (Gould, 1841) and *Prometetus exacuus* (Say, 1821). a-c) *B. havanensis*, fossil specimen UJMC-371a from a former wetland area of Viesca, Coahuila, Mexico. d-f) *D. dilatata*, subfossil specimen UJMC-372a from fluvial deposits of the Aguanaval River, Durango, Mexico. g-j) *P. exacuus* subfossil specimens from spring el Molino, Viesca, Coahuila, Mexico. g, h, j) Specimen UJMC-373a. i) Specimen UJMC-373b.

Remarks: The two subfossil specimens correspond in their shell morphology in all details to the recent material of this species. It is the first fossil record of this genus from Mexico.

Genus *Promenetus* Baker, 1935

Type species: *Planorbis exacuus* Say, 1821

Promenetus exacuus (Say, 1821)

(Figure 7g to 7j)

Menetus exacuus (Say, 1821). Baker, 1928, Wisconsin Geol. Nat. History Survey Bull. 70, pt. 1, p. 361, pi. 23, fig. 1-5.

Promenetus exacuus exacuus (Say, 1821): Baker, 1945, Molluscan family Planorbidae, p. 182.

Promenetus exacuus megas (W.H. Dall, 1905): Baker, Molluscan family Planorbidae, p. 188.

Description: Shells medium (maximum diameter 8.43 mm), flattened, discoidal (flat spiral), ultradextral, with carinate periphery (sharply keeled); keel close to or above the center of the body whorl; with $2\frac{1}{4}$ – $3\frac{1}{2}$ whorls, whorls rapidly increasing in size; spire flattened; sculpture of fine lines of growth and fine spiral lines; umbilicus deep; aperture triangular or ovate, outer and inner lip thin to slightly thickened: some specimens with shell repair scars.

Shell Measurements: (mean, with sd in parentheses; N = 10) SH = 1.91 (0.07) mm, SW (= shell diameter) = 6.96 (0.59) mm, WN = 2.60 (0.36).

Material Examined: 10 specimens (UJMC 373a-j).

Distribution: *P. exacuus* has a broad present-day distribution across Canada and United States (including Alaska) but does not reach Mexico.

Remarks: The prominent keel on the body whorl is the main characteristic of the shell of this species. Although there are no anatomical or radular differences, some authors such as Clarke (1981) and Prescott and Curteanu (2004) distinguish two subspecies of the Keeled *Promenetus*: *P. e. exacuus* and *P. e. megas*. However, we follow Burch (1989) and Turgeon *et al.* (1998) and do not recognize these subspecies because we considered that small differences in shell size alone are insufficient for a distinction. The southern limit of occurrence of living *P. exacuus* is Tarrant

County, Texas (Pratt, 1983), and our present record spread out the (fossil) range of the genus c. 1000 km to the south. It is the first fossil record of this genus from Mexico.

4. Discussion

There are few data about extinctions of freshwater gastropods in Mexico but Johnson *et al.* (2013) suggested that similar or greater extinctions likely occurred in Mexico and North America. The largest single modern extinction event of freshwater snails in North America took place in Coosa River, Alabama, United States, where within 50 years 33 species went extinct (Johnson *et al.*, 2013). Our results from northern Mexico show a similar rapid loss of local freshwater snail species as in United States. This general decline of mollusks diversity began in the early to middle Holocene (Czaja *et al.*, 2014a, 2014b), but then increased rapidly in the 20th century. The estimated times of local gastropod extirpation in the area (Table 2) are based on species described in the present study as well several other subfossil species reported recently from the same area (Czaja *et al.*, 2014a, 2014b, 2015, 2017a, 2017b, 2017c). Of the 32 species present through the Holocene in Laguna District only four (12.5%) are still living here. A quarter (25%) is totally extinct, half of them (50%) still live in Mexico and three species (9%) occur only in United States and Canada. This means the loss of 88% of the number of freshwater gastropod species in the area of study (Figure 8). The area is now a very dry region and belongs to the most populated regions in Mexico. Only the outflows of the dams of the Nazas and Aguanaval rivers and the small spring La Peña contain permanent waters. These sites are monitored by the authors since 2009 and contain only four gastropod species (Table 2). Snails with small geographic ranges were more vulnerable to such extinction events than species with a wider distribution. Of the 28 species that disappeared from the area during the Holocene, 14 (43%) were endemics (Table 2). Extirpation of all endemic species in this area apparently depended

Table 2. List of gastropod species of the study area and estimated time prior of the local extirpation or extinction based on species described in the present study as well subfossil species reported previously from the same area. * = Endemic species, bold = Species described in present paper.

| Family | Species | Holocene Records | 20th Century Records | Present-day Occurrence in area |
|--------------|---|------------------|----------------------|--------------------------------|
| Assimineidae | <i>Assiminea cienegensis</i> * | X | X | |
| Cochliopidae | <i>Coahuilix hubbsi</i> * | X | | |
| Cochliopidae | <i>Coahuilix landyei</i> * | X | | |
| Cochliopidae | <i>Coahuilix parrasense</i> * | X | X | |
| Cochliopidae | <i>Cochliopina riograndensis</i> | X | X | |
| Cochliopidae | <i>Juturnia coahuilae</i> * | X | X | |
| Cochliopidae | <i>Juturnia gracilis</i> * | X | X | |
| Cochliopidae | <i>Mexipyrgeus viescaensis</i> * | X | X | |
| Cochliopidae | <i>Paludiscala thompsoni</i> * | X | X | |
| Cochliopidae | <i>Phreatoceras taylori</i> | X | ? | |
| Cochliopidae | <i>Tryonia cf. seemani</i> | X | X | |
| Cochliopidae | <i>Tryonia hershleri</i> * | X | | |
| Cochliopidae | <i>Tryonia paleocircumstriata</i> * | X | X | |
| Hydrobiidae | <i>Cincinnatia integra</i> | X | X | |
| Hydrobiidae | <i>Probythinella emarginata</i> | X | X | |
| Hydrobiidae | <i>Probythinella protera</i> | X | X | |
| Hydrobiidae | <i>Pseudotryonia pasajae</i> * | X | X | |
| Hydrobiidae | <i>Pyrgulopsis manantiali</i> * | X | | |
| Hydrobiidae | <i>Pyrgulopsis paleoacarinatus</i> * | X | X | |
| Hydrobiidae | <i>Pyrgulopsis paleominckleyi</i> * | X | X | |
| Lymnaeidae | <i>Galba obrussa</i> | X | X | |
| Neritidae | <i>Vitta cf. usnea</i> | X | | |
| Physidae | <i>Physella acuta</i> | X | X | X |
| Physidae | <i>Physella pomilia</i> | X | | |
| Planorbidae | <i>Biomphalaria havenensis</i> | X | X | |
| Planorbidae | <i>Ferrissia californica</i> | X | X | X |
| Planorbidae | <i>Gyraulus parvus</i> | X | X | X |
| Planorbidae | <i>Hebetancylus excentricus</i> | X | X | |
| Planorbidae | <i>Helisoma anceps</i> | X | X | X |
| Planorbidae | <i>Dilatata dilatata</i> | X | X | |
| Planorbidae | <i>Planorbella duryi</i> | X | | |
| Planorbidae | <i>Promenetus exacuus</i> | X | X | |




Figure 8 Decline in species richness of freshwater gastropods in the area of study during the Holocene.

on, among others factors, their limited dispersal ability, which is especially true for the phreatic species of *Coahuilix*, *Paludiscala* and *Phreatoceras*. Similar patterns of extinctions of endemics were also reported from other sites (Hershler *et al.*, 2014b). However, also taxa with relatively wide recent distribution such as *Promenetus*, *Dilatata*, *Cincinnatia*, *Biomphalaria* and both species of *Probythinella* did not survive in this area. The genera *Promenetus* and *Probythinella* apparently disappeared completely from Mexico. Only *Physella acuta* (Draparnaud, 1805), *Ferrissia californica* (Rowell, 1863), *Helisoma anceps* (Menke, 1830) and *Gyraulus parvus* (Say, 1817) survived. Not surprisingly, all of them are known as very tolerant species with wide geographic ranges. It seems that of the 28 disappeared species only *Planorbella duryi* (Wetherby, 1879), *Tryonia hershleri* Czaja and Estrada-Rodriguez, 2015, *Physella pomilia* (Conrad, 1834) and *Vitta usnea* (Röding, 1798) disappeared from the area of study before human activities impacted the region.

The families most affected by the local extirpation were the Cochliopidae (12) and Hydrobiidae (7 species) and none of both currently have members in the area. These families belong to the most vulnerable freshwater gastropod families in North America, and they depend on groundwater inflows. Almost all endemic species of Cuatrociénegas also belong to these hydrobiid families.

The environmental impact was especially severe in the complex of 15 springs of Viesca in the central part of the Laguna District that dried up completely in the second half of the last century. The exact time of the drying is difficult to determine but in 1959 all 15 springs were still active (Blásquez, 1959). In 1961, some freshwater habitat must have existed around Viesca because type specimens of the Viesca Mud Turtle (*Kinosternon hirtipes megacephalum* Iverson) were collected (Legler and Vogt, 2013). Rhodin *et al.* (2011) reported 1970 as the approximate extinction date of the Viesca Mud Turtle and this could also be time of extirpation of the freshwater snails. That most snails from Viesca were still living in the 20th century is established by their common occurrence with *Melanoides tuberculata* (Müller, 1774) in the superficial sediments of the springs. This exotic species was introduced to North America in the 1930's (Rader *et al.*, 2003). The loss of aquatic habitats in Viesca occurred after decades of extensive overexploitation of groundwater pumping principally for irrigated agricultural production. In addition, the construction of two large dams further altered the groundwater hydrology. Wolfe (2013) has described how groundwater resources in Laguna District (especially at Viesca) were depleted from the 1920s to the 1960s. In this period, the number of water pumps increased from 12 to almost 3000 and, as a result, the groundwater level dropped by almost 100 meters. A few years later the lakes disappeared and also all 15 springs at Viesca had dried out (Figure 9). Similar local extinction events documented by Hershler *et al.* (2014b) from various sites in arid western North America coincide in time with the environment change in Laguna District. Also, there were especially hydrobiid snails affected in these arid regions.

Assiminea cienegensis, *J. coahuilae*, *T. seemani*, *P. pasajae* and *Pyrgulopsis manantiali* (Hershler, 1985) are still living near the area of study but all five species are now endemic to very restricted areas, including single springs. The subfossil records show that they had a wider distribution; hence, they are currently not only endemics but also relicts.



Figure 9 Spring Tunel 1 (Bilbao), Viesca, Coahuila, Mexico, that dried up in the second half of the last century (May 2018).

Although *Promenetus exacuous* and *Probythinella emarginata* are extirpated in Mexico, they have a widespread distribution in eastern United States and Canada and their populations are currently stable (Johnson *et al.*, 2013). However, both species occur in semi-arid region of United States in only a very few sites. *P. protera*, a species common in late Holocene Rivers and lakes in the area of study, shows currently a very limited distribution in United States (Hershler, 1996). However, contrary to the more widespread *P. emarginata*, this species is not included in the IUCN Red List and does also not appear in the list of endangered freshwater species of Canada and the United States published by Johnson *et al.* (2013).

It cannot be ruled out that invasive species such as *Melanooides tuberculata* are at least partly responsible for the loss of gastropod diversity in the Laguna District. This species has long been suspected of displacing native populations (Thompson, 2004; Almeida *et al.*, 2018). Although extirpation and

invasion coincide temporally and spatially in the area of study, a causal relationship such as interspecific competition or increased parasitism is unclear. The current sampling demonstrates that the superficial sediments contain only a few specimens of *Melanooides* and its relative abundance never exceeds 0.1%. Today, however, *Melanooides* is now by far the most abundant species in the localities that still contain water in the study area.

5. Conclusions

The local extirpation event described here from the Laguna District happened silently and unrecognized only some two hundred kilometers away from one of the currently greatest hotspots of gastropods diversity in North America, the Cuatrociénegas Basin. Unfortunately, the same human activities, especially groundwater pumping, that caused loss of aquatic habitat in the study area

continue to endanger the ecosystems of Cuatrociénegas. One of the mottos of the Conservation Paleobiology concept created by Dietl and Flessa (2011) was “Using the Past to Manage for the Future”. We hope that the large number of extirpated species reported here will strengthen the efforts to conserve threatened freshwater habitats in Mexico, especially in Cuatrociénegas basin.

Acknowledgements

We thank the Faculty of Biological Science of the Juárez University of the State of Durango for support us during this research. We thank M.C. Juan Carlos Ibarra Flores of the Área de Protección de Flora y Fauna Cuatrociénegas for permission to collect soil samples of the Poza Churince in Cuatrociénegas. Dr. Thomas A. Neubauer (Justus Liebig University Giessen, Germany) and an anonymous reviewer provided numerous constructive comments.

References

- Albrecht, C., Kuhn, K., Streit, B., 2007, A molecular phylogeny of Planorbioidea (Gastropoda, Pulmonata): insights from enhanced taxon sampling: *Zoologica Scripta*, 36 (1), 27–39. <https://doi.org/10.1111/j.1463-6409.2006.00258.x>
- Almeida, P.R.S., Nascimento Filho, S.L., Viana, G.F.S., 2018, Effects of invasive species snails in continental aquatic bodies of Pernambuco semiarid: *Acta Limnologica Brasiliensis*, 30, 103. <https://doi.org/10.1590/s2179-975x10616>
- Aguiar-Silva, C., C. Mendonça, P. da Cunha Kellis Pinheiro, S. Mesquita, Carvalho O., Caldeira, R., 2014, Evaluation and updating of the Medical Malacology Collection (Fiocruz-CMM) using molecular taxonomy: *SpringerPlus*, 3, 1–6. <https://doi.org/10.1186/2193-1801-3-446>
- Baker, F.C., 1945, The molluscan family Planorbidae: University Illinois Press, Urbana xxxvi + pp. 530.
- Blásquez, L.L., 1959, Hidrogeología de las regiones desérticas de México: *Anales del Instituto de Geología*, 15, pp. 172.
- Burch, J. B., 1989, North American Freshwater Snails: Malacological Publications, Hamburg, Michigan, pp. 365.
- Butzer, K.W., Abbott, J.T., Frederick, C.D., Lehman, P.H., Cordova, C.E., Oswald, J.F., 2008, Soil-geomorphology and “wet” cycles in the Holocene record of North-Central Mexico: *Geomorphology*, 101, 237–267. <https://doi.org/10.1016/j.geomorph.2008.06.005>
- Clarke, A.H., 1981, The Freshwater Molluscs of Canada: National Museum of Natural Sciences, National Museums of Canada, Ottawa, Canada, pp. 446.
- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., 2014a, Freshwater mollusks of the Valley of Sobaco, Coahuila, Northeastern Mexico – a subfossil ecosystem similar to Cuatrociénegas: *Boletín de la Sociedad Geológica Mexicana*, 66 (3), 459–469. <https://doi.org/10.18268/bsgm2014v66n3a4>
- Czaja, A., Palacios-Fest, M.R., Estrada-Rodríguez, J.L., Romero-Méndez, U., Alba-Ávila, J.A., 2014b, Inland Dunes mollusks fauna from the Paleolake Irritila in the Comarca Lagunera, Coahuila, Northern Mexico: *Boletín de la Sociedad Geológica Mexicana*, 66 (3), 541–551. <https://doi.org/10.18268/bsgm2014v66n3a9>
- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., 2015, A new species of the genus Mexipyrurgus Taylor, 1966 (Caenogastropoda: Truncatelloidea: Cochliopidae) from late Holocene spring deposits in Viesca, Coahuila, Mexico: *The Nautilus*, 129, 163–168.
- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., Orona-Espino, A., 2017a, Two new subfossil species of springsnails (Hydrobiidae: Pyrgulopsis) from north Mexico and their relation with extant species from Cuatrociénegas: *Boletín de la Sociedad Geológica Mexicana*, 69 (1), 199–208. <https://doi.org/10.18268/bsgm2017v69n1a9>

- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., Estrada-Arellano, J.R., González-Zamora, A., 2017b, Primer registro fósil del gasterópodo Cincinnatia (Hydrobiidae: Nymphophilinae) en México: Revista Mexicana de Biodiversidad, 88, 912–917. <https://doi.org/10.1016/j.rmb.2017.10.025>
- Czaja, A., Estrada-Rodríguez, J.L., Romero-Méndez, U., Ávila-Rodríguez, V., Meza-Sánchez, I.G., Covich, A.P., 2017c, New species and records of phreatic snails (Caenogastropoda: Cochliopidae) from the Holocene of Coahuila, Mexico: Archiv für Molluskenkunde, 146 (2), 227–232. <https://doi.org/10.1127/arch.moll/146/227-232>
- DeJong, R.J., Morgan, J.A.T., Paraense, W.L., Pointier, J.P., Amarista, M., Aych-Kumi, P.F.K. et al. 2001, Evolutionary relationships and biogeography of Biomphalaria (Gastropoda: Planorbidae) with implications regarding its role as host of the human bloodfluke, Schistosoma mansoni: Molecular Biology and Evolution, 18 (12), 2225–2239. <https://doi.org/10.1093/oxfordjournals.molbev.a003769>
- Dietl, G.P. and Flessa, K.W., 2011, Conservation paleobiology: putting the dead to work: Trends in Ecology and Evolution, 26 (1), 30–37. <https://doi.org/10.1016/j.tree.2010.09.010>
- Eichhorst, T.E., 2016, Neritidae of the world. Vol. 2. Harxheim: Conchbooks, 696–1366.
- Figueroa-Rangel, B.L., Willis, K.J. and Olvera-Vargas, M., 2012, Late-Holocene successional dynamics in a transitional forest of west-central Mexico: The Holocene, 22(2), 143–153. <https://doi.org/10.1177/0959683611414929>
- Frauenfeld, G.R. von, 1863, Vorläufige Aufzählung der Arten der Gattungen Hydrobia Htm. und Amnicola Gld. Hldm. in der kaiserlichen und in Cuming's Sammlung: Verhandlungen der Kaiserlich-Königlichen Zoologisch-botanischen Gesellschaft in Wien, 13, 1017–1032.
- Frauenfeld, G.R. von, 1865, Zoologische Miscellen: Verhandlungen der Kaiserlich-Königlichen Zoologisch-botanischen Gesellschaft in Wien, 15, 525–536.
- Guerrero-Arenas, R., Jiménez-Hidalgo, E., 2015, El registro fósil y la conservación de la biodiversidad actual: Ciencia y Mar, 23, 67–75.
- Hammer, R., Harper, D.A.T., Ryan P.D., 2001, PAST: Paleontological statistics software package for education and data analysis. http://palaeo-electronica.org/2001_1/past/issue1_01.htm. [Accessed 23. July. 2019].
- Hershler, R., 1985, Systematic revision of the Hydrobiidae (Gastropoda: Rissoacea) of the Cuatro Ciénegas Basin, Coahuila, Mexico: Malacologia, 26 (1–2), 31–123.
- Hershler, R., 1996, Review of the North American aquatic snail genus Probythinella (Rissooidea: Hydrobiidae): Invertebrate Biology, 115 (2), 120–144. <https://doi.org/10.2307/3227043>
- Hershler, R., 2001, Systematics of the North and Central American aquatic snail genus Tryonia (Rissooidea: Hydrobiidae): Smithsonian Contributions to Zoology, 612, 1–53. <https://doi.org/10.5479/si.00810282.612>
- Hershler, R., Liu, H.-P., Stockwell, C.A., 2002, A new genus and species of aquatic gastropods (Rissooidea: Hydrobiidae) from the North American Southwest: phylogenetic relationships and biogeography: Proceedings of the Biological Society of Washington, 115 (1), 171–188.
- Hershler, R., Liu, H.-P., Landye, J.J., 2011a, New species and records of springsnails (Caenogastropoda: Cochliopidae: Tryonia) from the Chihuahuan Desert (Mexico and United States), an imperiled biodiversity hotspot: Zootaxa, 3001, 1–32. <https://doi.org/10.11646/zootaxa.3001.1.1>
- Hershler, R., Liu, H.-P., Landye, J.J., 2011b, Two new genera and four new species of freshwater cochliopid gastropods (Rissooidea) from northeastern Mexico: Journal of Molluscan Studies, 77 (1), 8–23. <https://doi.org/10.1093/mollus/eyq033>

- Hershler, R., Landye, J.J., Liu, H.-P., De la Maza-Benignos, M., Ornelas, P., Carson E.W., 2014a, New species and records of Chihuahuan Desert springsnails, with a new combination for *Tryonia brunei*: Western North American Naturalist, 74 (1), 47–65. <https://doi.org/10.3398/064.074.0105>
- Hershler, R., Liu, H.-P., Howard, J., 2014b, Springsnails: a new conservation focus in western North America. *BioScience*, 64 (8), 693–700. <https://doi.org/10.1093/biosci/biu100>
- Johnson, P.D., Bogan, A.E., Brown, K.M., Burkhead, N.M., Cordeiro, J.R., Garner, J.T., Hartfield, P.D., Lepitzki, D.A.W., Mackie, G.L., Pip, E., Tarpley, T.A., Tiemann, J.S., Whelan, N.V., Strong, E.E., 2013, Conservation status of freshwater gastropods of Canada and the United States: *Fisheries*, 38 (6), 247–282. <https://doi.org/10.1080/03632415.2013.785396>
- Legler, J.M., Vogt, R.C., 2013, *The Turtles of Mexico: Land and Freshwater Forms*: University of California Press, 416 pp.
- Morrison, J.P.E., 1945, *Durangonella*, a new hydrobiine genus from Mexico, with threenew species: *The Nautilus*, 59, 18–23.
- Naranjo-García, E., 2003, Moluscos continentales de México: dulceacuícolas: *Revista de Biología Tropical*, 51 (3), 495–505.
- Ortega-Guerrero, A., 2003, Origin and geochemical evolution of groundwater in a closed-basin clayey aquitard, Northern Mexico: *Journal of Hydrology*, 284 (1), 26–44. [https://doi.org/10.1016/S0022-1694\(03\)00239-7](https://doi.org/10.1016/S0022-1694(03)00239-7)
- Pilsbry, H.A., 1939, *Land Mollusca of North America*: Academy of Natural Sciences of Philadelphia, Monograph 3, pp. 547.
- Pilsbry, H.A., Ferriss, J.H., 1906, *Mollusca of the southwestern states, II*: *Proceedings of the Academy of Natural Sciences of Philadelphia*, 58, 123–175.
- Pratt, W.L., 1983, *Living Promenetus exacuus* (Pulmonata: Planorbidae) from North Central Texas: *The Nautilus*, 97 (2), 73–74.
- Prescott, D.R.C., Curteanu, M.M., 2004, *Survey of Aquatic Gastropods in the Central Parkland Subregion of Alberta*: Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 92, Edmonton, AB, pp. 50. <https://doi.org/10.5962/bhl.title.114093>
- Rader, R.B., Belk, M.C., Keleher, M.J., 2003, The introduction of an invasive snail (*Melanoides tuberculata*) to spring ecosystems of the Bonneville Basin, Utah: *Journal of Freshwater Ecology*, 18 (4), 647–657. <https://doi.org/10.1080/02705060.2003.9664007>
- Régner, C., Fontaine, B., Bouchet, P., 2009, Not knowing, not recording, not listing: numerous unnoticed mollusk extinctions: *Conservation Biology*, 23 (5), 1214–1221. <https://doi.org/10.1111/j.1523-1739.2009.01245.x>
- Rhodin, A.G.J., Walde, A.D., Horne, B.D., van Dijk, P.P., Blank, T., Hudson, R., 2011, *Turtles in Trouble: The World's 25+ Most Endangered Tortoises and Freshwater Turtles*, Lunenburg, MA: IUCN/SSC Tortoise and Freshwater Turtle Specialist Group, Turtle Conservation Fund, Turtle Survival Alliance, Turtle Conservancy, Chelonian Research Foundation, Conservation International, Wildlife Conservation Society, and San Diego Zoo Global, pp. 54.
- Riedel, F., 1993, Early ontogenetic shell formation in some freshwater gastropods and taxonomic implications of the protoconch: *Limnologia*, 23, 349–368.
- Rosser, T.G., Alberson, N.R., Khoo, L.H., Woodyard, E.T., Wise, D.J., Pote, L.M. and Griffin, M.J., 2016, *Biomphalaria havanensis* is a Natural First Intermediate Host for the Trematode *Bolbophorus damnificus* in Commercial Catfish Production in Mississippi: *North American Journal of Aquaculture*, 78 (3), 189–192. <https://doi.org/10.1080/15222055.2016.1150922>

- Strong, E.E., Gargominy, O., Ponder, W.F., Bouchet, P., 2008, Global diversity of gastropods (Gastropoda; Mollusca) in freshwater: *Hydrobiologia*, 595 (1), 149–166. <https://doi.org/10.1007/s10750-007-9012-6>
- Taylor, D.W., 1966, A remarkable snail fauna from Coahuila, Mexico: *The Veliger*, 9, 152–228.
- Thompson, F.G., 2004, An identification manual for the freshwater snails of Florida <http://www.flmnh.ufl.edu/malacology/fl-snail/snails1.htm> [Accessed 14 May 2018].
- Thompson, F.G., 2011, The land and freshwater snails of Mexico and Central America: *Bulletin of the Florida Museum of Natural History*, 50, 1–299.
- Turgeon, D.D., Quinn, J.F. Jr., Bogan, A.E., Coan, E.V., Hochberg, F.G., Lyons, W.G., Mikkelsen, P.M., Neves, R.J., Roper, C.F.E., Rosenberg, G., Roth, B., Scheltema, A., Thompson, F.G., Vecchione, M., Williams, J.D., 1998, Common and scientific names of aquatic invertebrates from the United States and Canada: mollusks, 2nd edition: American Fisheries Society, Special Publication 26. American Fisheries Society, Bethesda, Maryland, pp. 526.
- Wolfe, M., 2013, The historical dynamics of Mexico's groundwater crisis in La Laguna: Knowledge, resources, and profit, 1930s–1960s: *Mexican Studies/Estudios Mexicanos*, 29 (1), 3–35. <https://doi.org/10.1525/msem.2013.29.1.3>