

## Revision of corals (Late Jurassic to Early Cretaceous, Northern Mexico) previously established by Imlay and Wells

*Revisión de los corales (Jurásico tardío a Cretácico temprano, norte de México) previamente establecidos por Imlay y Wells*

Hannes Löser<sup>1,\*</sup>

<sup>1</sup> Estación Regional del Noroeste, Instituto de Geología, Universidad Nacional Autónoma de México. Av. Luis Donaldo Colosio y Madrid S/N. Hermosillo 83110, Sonora, México.

\* Corresponding author: (H. Löser)  
hannes@paleotax.de

### ABSTRACT

New Mesozoic (Kimmeridgian to Albian) corals from northern Mexico (Mexican states Coahuila, Durango and Sonora), that were established by Imlay and Wells between 1940 and 1946, are revised on the basis of the type material kept at the Museum of Paleontology of the University of Michigan in Ann Arbor (Mich., USA). From the six new species, only one may remain in use, and the other five species are considered to be synonyms of older taxa.

**Keywords:** Scleractinia, Jurassic, Cretaceous, Mexico.

### RESUMEN

Nuevos corales del Mesozoico fueron establecidos por Imlay y Wells entre 1940 y 1946 desde del norte de México (estados Coahuila, Durango y Sonora). Estos corales se revisan en base al material tipo que se encuentra en el Museo Paleontológico de la Universidad de Michigan en An Arbor (Mich., EE.UU.). De las seis especies nuevas, solo una queda en uso; las otras cinco son sinónimos de taxones más antiguos.

**Palabras clave:** Scleractinia, Jurásico, Cretácico, México.

#### How to cite this article:

Löser, H., 2024, Revision of corals (Late Jurassic to Early Cretaceous, Northern Mexico) previously established by Imlay and Wells: Boletín de la Sociedad Geológica Mexicana, 76 (3), A190424. <http://dx.doi.org/10.18268/BSGM2024v76n3a190424>

Manuscript received: March 14, 2024.  
Corrected manuscript received: April 14, 2024.  
Manuscript accepted: April 20, 2024.

Peer Reviewing under the responsibility of Universidad Nacional Autónoma de México.

This is an open access article under the CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

## 1. Introduction

The investigation of Mesozoic corals in Mexico began with the work of Felix (1891). We know of nearly 90 publications that have at least some citations of Mesozoic coral species, and nearly 60 of those have descriptions and/or illustrations of fossil material (Löser, 1994 and later supplements). Half of the systematic taxonomic investigations were undertaken in the past 25 years. There are two older works that were realized by American workers, Imlay (1940) and Wells (1946). Imlay (1940) erected a new coral species from Durango. Wells (1946) published a small collection of Upper Jurassic and Lower Cretaceous corals collected by Kellum and Imlay in Coahuila, Durango, and

Sonora, for which he established five new species. In the present paper, the coral material is examined again, provided with morphometric data, and set into a modern taxonomic context.

## 2. Study area

The material comes from the Mexican states Coahuila, Durango, and Sonora (Figure 1). The geology for each of these areas is described by Imlay (1939, 1940), Kellum (1936), and Kelly (1936). Exact localities or sample locations are unknown and geographic coordinates are, therefore, not given.



Figure 1 Study area with the approximate position of the localities.

## 2.1. COAHUILA

Barril Viejo, SE of Cuatro Ciénegas. Barril Viejo shale, Lower Hauterivian. See Imlay (1940).

Outlier of the Sierra de Venado. Cuchillo Fm, Upper Aptian to Lower Albian with *Dufrenoya* cf. *justinae*. See Kellum (1936).

Los Vagos, Potrero de Oballos. Barril Viejo shale, Lower Hauterivian. See Imlay (1940). The place name probably refers to the small village of Obayos NW of Cuatro Ciénegas.

Sierra de Acatita, at the head of the canyon east of the Acatica ranch. Lower Cuchillo fm, Upper Aptian to Lower Albian. See Kellum (1936).

## 2.2. DURANGO

Outcrop about 4 km south of the Las Cuevas Ranch. La Casita Fm, Kimmerdigian. The geology was described by Kellum (1936) and Imlay (1939). Las Cuevas lies about 13 km SW from the city centre of Torreón.

Outcrop 3.2 km SW of Las Cuevas. Carbonara fm, middle Valanginian. For details see Imlay (1940).

Outcrop at the NW end of the Sierra de España (or Hispana), about 1.6 km NE of La Goma station. Carbonara fm, middle Valanginian. See Kellum (1936) and Imlay (1940). La Goma lies about 23 km SW of Torreón.

## 2.2. SONORA

Sierra el Tigre, 400 m west of the King ranch, Cañón Sta. Rosa. Mural Fm, Upper Aptian to Lower Albian.

## 3. Material and methods

The studied material is almost entirely kept at the University of Michigan Museum of Paleontology (UMMP) in Ann Arbor (Mich., USA). One specimen (23343) could not be found. Most specimens are still complete, without polished or thin sections. Some thin sections are available. All specimens

were photographed and measured. Their generic taxonomy was updated and the species were determined based on the morphometric data.

It was not possible to prepare the coral specimens, such as cutting, polishing, or taking thin sections. Available thin sections were scanned by passing light through them using a flatbed scanner with an optical resolution of 6,400 dpi. Specimens were scanned with an optical resolution of 4,800 dpi. The images were then transferred to greyscale bitmaps and their quality was amended by histogram contrast manipulation (contrast stretching) where possible.

Corallite dimensions and the septal counts were systematically measured using the scanned thin sections. To achieve statistical significance, in colonial corals the largest number of possible measurements was taken. This number was mainly determined by the size and quality of the thin sections, and the size of the single corallites in relation to the size of the thin sections.

For each type of measurement (such as, for instance, the corallite diameter or septal counts), the following values were obtained (all measurements in millimetres):

n number of measurements

min–max lowest and highest measured values

$\mu$  arithmetic mean (average)

s standard deviation

cv coefficient of variation according to K. Pearson

$\mu \pm s$  first interval

Thin sections were measured and values were calculated using the Palaeontological Database System PaleoTax, module PaleoTax/Measure (<https://www.paleotax.de/measure>); for details on the mathematical background, see Löser (2012). Characters that were visible on the fossils were compared against characters on specimens in worldwide fossil coral collections, and an associated image database (29,000 specimens, 17,000 illustrated), located in the Estación Regional de Noroeste (ERNO), Sonora, Mexico. Data storage and processing were carried out using the PaleoTax database programme (Löser, 2004).

## 4. Systematic description

The classification follows Löser (2016). The following abbreviations are used to describe the dimensions of the corals: c, outer diameter of the solitary coral; ccd, distance between corallite centres; clmax, large lumen; clmin, small lumen; septa, number of septa in the adult corallite. The abbreviations used in the synonymy lists follow Matthews (1973): \*: earliest valid publication of the species name; v: the specimen was observed; p: the material described in this citation belongs only partly to the present species.

Superfamily Actinastreoidea Alloiteau, 1952

**Remarks:** For the morphology of the superfamily refer to Samaniego-Pesqueira *et al.* (2023).

Family Actinastreae Alloiteau, 1952

Genus *Stelidioseria* Tomes, 1893

**Type species:** *Stelidioseria gibbosa* Tomes, 1893

**Remarks:** For the morphology of the superfamily refer to Samaniego-Pesqueira *et al.* (2023).

*Stelidioseria icaunensis* (Orbigny, 1850)

Figure 2:1-2

### Synonymy:

- \*v 1850 *Prionastraea icaunensis* Orbigny, (2), p. 93
- v 1871 *Astrocoenia Kunthi* – Bölsche, p. 56, pl. 12, fig. 7
- v 1897 *Astrocoenia urgoniensis* – Koby, p. 58, pl. 15, figs. 5-8
- v 1924 *Astrocoenia hexamera* n.sp. – Fritzsche, p. 318, pl. 3, fig. 7
- v 1933 *Astrocoenia budaensis* n.sp. – Wells, p. 78, pl. 6, fig. 3
- v 1933 *Siderastraea tuckerae* n.sp. – Wells, p. 104, pl. 9, figs. 6, 7
- v 1946 *Astrocoenia kellumi* Wells, n. sp. – Wells, p. 2, pl. 1, fig. 1
- v 2020 *Stelidioseria icaunensis* (Orbigny, 1850) – Löser, Mendicoa and Fernández Mendiola, p. 221, figs. 3a-c

**Material:** UMMP 15969; 1 thin section.

### Dimensions: (UMMP 15969)

	n	min-max	$\mu$	s	cv	$\mu \pm s$
clmin	20	1.34-1.91	1.55	0.16	10.1	1.39-1.70
clmax	20	1.72-2.46	2.09	0.23	10.8	1.87-2.32
ccd	20	1.55-2.10	1.84	0.18	9.7	1.66-2.02
septa	24					

**Remarks:** Systematic measurements of the thin section of the holotype of *Astrocoenia kellumi* shows clearly synonymy of this species with *Stelidioseria icaunensis*.

**Occurrence:** Mexico, Durango, Rancho Las Cuevas Rancho (Kimmeridgian).

**Other occurrences:** Kimmeridgian to middle Cenomanian, worldwide.

Superfamily Cladocoroidea Orbigny, 1851

**Remarks:** For the morphology of the superfamily refer to Löser *et al.* (2023).

Family Columastreae Alloiteau, 1952

**Description:** Plocoid colonies. Septal symmetry regular radial and mostly hexamerous. Pali present in some genera. Columella varies: styloform, styloform and double, lamellar, or absent. Coenosteum with costae.

Genus *Eocolumastrea* Löser and Zell, 2015

**Type species:** *Columnocoenia bucovinensis* Morycowa, 1971

**Description:** Plocoid coral with septa in a regular hexamerous or decamerous symmetry. Columella lamellar or small and styloform. Irregular pali at the first septal cycle, not very pronounced. Coenosteum narrow.

**Remarks:** The genus received a more detailed study by Garberoglio *et al.* (2021).

*Eocolumastrea octaviae* (Prever, 1909)

Figure 2: 3-4

### Synonymy:

- \*v 1909 *Ulastraea Octaviae* Prever, p. 91, pl. 5, fig. 5
- v 1946 *Stephanocoenia guadalupe minor* Wells, n.var. – Wells, p. 3, pl. 1, figs. 2-4
- v 2021 *Eocolumastrea octaviae* (Prever, 1909) – Gar-

beroglio, Löser and Lazo, p. 9, fig. 8 [= here more detailed synonymy]

**Material:** UMMP 15993, 23339; 2 thin sections.

**Dimensions:** (UMMP 23339)

	n	min-max	$\mu$	s	cv	$\mu \pm s$
clmin	25	1.32-1.78	1.53	0.13	8.7	1.40-1.66
clmax	25	1.65-2.04	1.84	0.09	4.7	1.75-1.92
ccd	30	1.51-2.19	1.87	0.18	9.6	1.69-2.05
septa	24					

**Remarks:** The present type specimen fits well in the dimensions of *E. octaviae*.

**Occurrences:** Mexico, Coahuila, Sierra de Acatita, Acatita ranch (upper Aptian). Mexico, Sonora, Sierra El Tigre, Rancho de King, Cañón Sta. Rosa (upper Aptian to lower Albian).

**Other occurrences:** Valanginian to lower Cenomanian, world-wide.

Superfamily Cyclolitoidea Milne Edwards and Haime, 1849

**Remarks:** For the morphology of the superfamily refer to Löser *et al.* (2023).

Family Latomeandridae Fromentel, 1861

**Remarks:** For the morphology of the family refer to Löser *et al.* (2023).

Genus *Thalamocaeniopsis* Alloiteau, 1954

**Type species:** *Thalamocaeniopsis ouenzensis* Alloiteau, 1954

**Description:** Cerioid colony. Corallite outline polygonal with centres slightly depressed. Symmetry of septa irregular. Synapticulae occasional, mainly in the space between calices. No costae. Columella consists of isolated trabeculae or one more solid element. Wall subcompact, made of synapticulae. Budding extracalicular.

**Remarks:** After having been established, the genus was poorly cited in the literature, probably because the first description was poorly illustrated. *Thalamocaeniopsis* is a common Lower Cretaceous and Cenomanian coral genus with numerous species.

*Thalamocaeniopsis neocomiensis* (Tomes, 1885)

Figure 2: 5-6

**Synonymy:**

\*v 1885 *Isastraea neocomiensis* Tomes, p. 547

v 1926 *Isastraea* cf. *geometrica* Koby – Dietrich, p. 79, pl. 7, fig. 3

v 1946 *Isastrea whitneyi* Wells 1932 – Wells, p. 3, pl. 2, figs. 1-3

1985 *Latiastrea whitneyi* (Wells, 1932) – Sikharulidze, p. 54, pl. 25, fig. 5

**Material:** UMMP 15786; 1 thin section.

**Dimensions:** (UMMP 15786)

	n	min-max	$\mu$	s	cv	$\mu \pm s$
clmin	25	2.43-3.46	2.96	0.28	9.3	2.68-3.24
clmax	25	3.85-5.34	4.53	0.45	10.0	4.08-4.98
ccd	25	3.09-4.00	3.54	0.28	8.0	3.25-3.82
septa	25	38-55	46.08	5.04	10.9	41-51

**Remarks:** This specimen was originally assigned to *Isastrea whitneyi* Wells, 1932. *Isastrea* does not occur in the Cretaceous and *Isastrea whitneyi* belongs to *Thalamocaeniopsis* as the present specimen. The present specimen differs from *T. whitneyi* by slightly larger dimensions and higher septal counts. *T. whitneyi* is a junior synonym of *T. fleurbaeusi* (Orbigny, 1850).


**Occurrence:** Mexico, Coahuila, Outlier of Sierra del Venado (upper Aptian to lower Albian).

**Other occurrence:** Hauterivian of the Eastern Tethys (Georgia), Lower Aptian of the European Boreal (UK) and the Southern Tethys (Tanzania), Lower Albian of the Western Atlantic (Mexico), and the Western Tethys (Spain).

Informal *Holocoenia* superfamily

**Remarks:** For the morphology of the informal group refer to Garberoglio *et al.* (2022).

**Remarks:** In older literature, the genus *Holocoenia* Milne Edwards and Haime, 1851 (and its synonyms *Paretallonia* Sikharulidze, 1972 and *Stereocoenia* Alloiteau, 1952) were assigned to the family Thamnasteriidae Reuss, 1864 (Löser, 2009). The Thamnasteriidae (now within the superfamily Rhizangioidea Orbigny, 1851) are characterised by medium-sized trabeculae and interconnected



**Figure 2** 1-2. *Stelidioseris icaunensis* (Orbigny, 1850), Holotype of *Astrocoenia kellumi*, (UMMP 15969). 1, transversal thin section. 2, transversal thin section, detail.  
 3-4. *Eocolumastrea octaviae* (Prever, 1909), Holotype of *Stephanocoenia guadalupae minor*, (UMMP 23339). 3, transversal thin section. 4, transversal thin section, detail.  
 5-6. *Thalamocaeniopsis neocomiensis* (Tomes, 1885), (UMMP 15786). 5, transversal thin section. 6, transversal thin section, detail.  
 7-8. *Holocoenia micrantha* (Roemer, 1841), Holotype of *Astrocoenia hispaniensis*, (UMMP 19345). 7, transversal thin section. 8, longitudinal thin section.  
 Scale bar 1 mm.

septa, whereas *Holocoenia* is characterised by smaller trabeculae and free septa. A new family (and superfamily) should be established for *Holocoenia* and the related genus *Etallonasteria* Roniewicz, 1987, but the type material of its type species (*Astrea micrantha* Roemer, 1841) is not available and the type locality (Germany, Niedersachsen, Berklingen; uppermost Valanginian to lowermost Hauterivian) does not yield anymore fossil corals, and comparable collection material is scarce. The genus was revised by Löser (2009) based on a small topotypical specimen.

Genus *Holocoenia* Milne Edwards and Haime, 1851

**Type species:** *Astrea micrantha* Roemer, 1841

**Description:** Cerioid colony with small corallites (diameter below 2 mm). Septa confluent, two septal generations differing in septal length. The number of septa vary between 18 and 22, and may differ within one colony. Thick styliform columella. Incomplete wall made of synapticalae. Endotheca absent.

*Holocoenia micrantha* (Roemer, 1841)

Figure 2: 7-8

**Synonymy:**

- \* 1841 *Astraea micrantha* Roemer, p. 113, pl. 16, fig. 27
- v 1940 *Astrocoenia hispaniensis* Imlay, n. sp. – Imlay, p. 138, pl. 1, figs. 21, 22
- v 1964 *Stereocaenia triboleti* (Koby, 1896) – Morycowa, p. 77, pl. 23, figs. 1-3
- v 1997 *Paretallonia bendukidzeae* Sikharulidze, 1972 – Baron-Szabo, p. 76, pl. 11, figs. 1, 3, 4
- v 2009 *Holocoenia micrantha* (Roemer, 1841) – Löser, p. 96, figs. 1, 2a-c
- v 2019 *Holocoenia micrantha* (Roemer, 1841) – Löser, Arias and Vilas, p. 280, figs. 13.1-3

**Material:** UMMP 19345; 1 thin section.

**Dimensions:** (UMMP 19345)

	n	min-max	$\mu$	s	cv	$\mu \pm s$
clmin	6	1.29-1.55	1.42	0.10	7.0	1.32-1.52
clmax	8	1.38-1.98	1.65	0.20	12.2	1.45-1.85

ccd	15	1.33-1.85	1.54	0.18	11.6	1.36-1.72
septa	20					

**Remarks:** The holotype of *Astrocoenia hispaniensis* belongs to the comparably rare genus *Holocoenia*. *Astrocoenia* Milne Edwards and Haime, 1848 is a Palaeogene genus but was widely used in the 19th and early 20th centuries for material that is now assigned to *Actinastrea* Orbigny, 1849 and *Stelidioseris* Tomes, 1893. *Holocoenia* shows a certain affinity to *Stelidioseris*, but clearly differs due to the wall made of synapticalae, which is clearly visible in the present specimen.

**Occurrence:** Mexico, Durango, Las Cuevas (Valanginian).

**Other occurrence:** Upper Valanginian to lower Hauterivian of the North Africa (Algeria) and the European Boreal (Germany, France), Hauterivian to Barremian of the Western Tethys (Spain, France), lower Barremian of the Western Tethys (France), upper Barremian to lower Aptian of the Central Tethys (Germany), Barremian to lower Aptian of the Central Tethys (Poland).

Superfamily Misistelloidea Eliášová, 1976

**Remarks:** For the morphology of the superfamily refer to Löser *et al.* (2023).

Family Rayasmiliidae Löser, 2022

**Description:** Solitary and phaceloid corals. The septa are always free. A lamellar columella is present in some genera. One or two septa may be connected to the columella.

Genus *Rayasmilia* Löser, 2022

**Type species:** *Rayasmilia salvata* Löser, 2022

**Description:** Solitary turbinate coral. Septa compact, not connected to each other, in a regular radial symmetry. Septa can be connected to the columella. Wall absent. Epitheca present. Endotheca well developed. Columella lamellar.

*Rayasmilia bangoimensis* (Liao and Xia, 1985)

Figure 3: 1-2

**Synonymy:**

- vp 1933 *Pleurosmilia saxi-fisi* Wells, p. 62, pl. 2, fig.

17, pl. 5, figs. 1, 2

vp 1946 *Axosmia mexicana* Wells, n.sp. – Wells, p. 6, pl. 1, figs. 6-11

\*v 1985 *Axosmia bangoinensis* (sp. nov.) – Liao and Xia, p. 146, pl. 8, figs. 6-9

v 1994 *Axosmia bangoinensis* Liao et Xia – Liao and Xia, p. 80, 223, pl. 8, figs. 11, 12

2002 *Aulophyllia belbekensis* Kusmicheva, sp. nov. – Kusmicheva, p. 178, pl. 27, fig. 2

v 2013a *Plesiosmia hennigi* Dietrich, 1926 – Löser, p. 104, fig. 5.10

v 2013b *Plesiosmia hennigi* (Dietrich, 1926) – Löser, fig. 3, fig. 1

**Material:** UMMP 23350#2; 1 thin section.

**Dimensions:** (23350#2)

c 11.8×15.6

septa 46

**Remarks:** Wells doubtfully included this specimen into *Axosmia mexicana*, but excluded it from the type series.

**Occurrence:** Mexico, Coahuila, Barril Viejo, Cuatrociénegas, (lower Hauterivian).

**Other occurrences:** Berriasian of the Central Tethys (Ukraine), Berriasian to Valanginian of the Southern Tethys (China), Valanginian to Aptian of the Western Atlantic (Mexico), lower Hauterivian of the European Boreal (France), lower Aptian of the Western Atlantic (Mexico), of the Central Tethys (Greece), and of the Western Tethys (Spain), upper Aptian of the Central Tethys (Greece), lower Cenomanian of the Western Atlantic (USA), upper Cenomanian of the Western Tethys (France).

*Rayasmia fromenteli* (Angelis d'Ossat, 1905)

Figure 3: 5-6

**Synonymy:**

\*v 1905 *Peplosmia Fromenteli* Angelis d'Ossat, p. 242, pl. 17, figs. 6 a-g

v 1926 *Pleurosmia hennigi* n.sp. – Dietrich, p. 87, pl. 7, figs. 6 a-c

vp 1946 *Axosmia mexicana* Wells, n.sp. – Wells, p. 6, pl. 1, figs. 6-11

1977 *Rhipidosmia tauridae* – Babaev and Krasnov, p. 147, text-fig. 9, pl. 43, fig. 3

v 1991 *Axosmia viai* n. sp. – Reig Oriol, p. 8, pl. 1, fig. 6, pl. 3, figs. 7, 8

v 2008 *Plesiosmia neocomiensis* (de Fromentel, 1867) – Löser, p. 55, pl. 4, figs. 2-3

**Material:** UMMP 23349; 1 thin section.

**Dimensions:** (23349)

c 20.5×24.9

septa 50

**Remarks:** This paratype of *Axosmia mexicana* can clearly be assigned to *Rayasmia*. *Axosmia* is a poorly defined genus and should no longer be used. The lectotype of the type species (*Caryophyllia extincorium* Michelin, 1841; Muséum National d'Histoire Naturelle Paris M00053) is a small and poorly defined specimen that does not reveal any details about the septal microstructure, the endotheca, or possible pali and/or columella. The genus is used (for instance, as per Vasseur and Lathuilière 2021) in a sense that was arbitrarily defined by Beauvais (1986) and Roniewicz (2008), but not based on proper type material.

**Occurrence:** Mexico, Coahuila, Los Vagos, Potrero de Oballos (lower Hauterivian).

**Other occurrence:** Upper Tithonian to Berriasian of the Central Tethys (Ukraine), upper Valanginian to lower Aptian of the Southern Tethys (Tanzania), lower Aptian of the Western Tethys (Spain), upper Aptian to lower Albian of the Western Tethys (Spain).

Genus *Trochophyllia* Alloiteau, 1952

**Type species:** *Montlivaltia melania* Fromentel, 1861


**Description:** Cylindric or turbinat solitary coral with circular or slightly elliptical outline. No columella, no pali. Endotheca made of large dissepiments. No wall, just a thin epitheca that is often not present. There is microstructural evidence for the position of the genus within the informal group.

*Trochophyllia communis* (Prever, 1909)

Figure 3: 7-9

**Synonymy:**

\*v 1909 *Trochosmia communis* Prever, p. 106, text-



**Figure 3** 1-2. *Rayasmilia bangoinensis* (Liao and Xia, 1985), (UMMP 23350#2). 1, transversal thin section. 2, transversal thin section, detail.

3-4. *Trochophyllia* sp., (UMMP 23350#1). 3, transversal thin section. 4, transversal thin section, detail.

5-6. *Rayasmilia fromenteli* (d'Angelis d'Ossat, 1905), Paratype of *Axosmilia mexicana*, (UMMP 23349). 5, transversal thin section. 6, transversal thin section, detail.

7-9. *Trochophyllia communis* (Prever, 1909), Holotype of *Axosmilia mexicana*, (UMMP 23348). 7, corallite surface. 8, (UMMP 23346#1), transversal thin section. 9, transversal thin section, detail.

Scale bar 1 mm.

figs. 8-10, pl. 10, fig. 4

v 1930 *Montlivaultia ignorata* nom. mut. – Oppenheim, p. 278

vp 1946 *Axosmilia mexicana* Wells, n.sp. – Wells, p. 6, pl. 1, figs. 6-11

vp 1946 *Montlivaultia burckhardti* Wells, n.sp. – Wells, p. 5, pl. 2, figs. 7-14

**Material:** UMMP 23346#1, 23348; 1 thin section.

**Dimensions:** (23346#1)

c 14.4×17.6

septa 78

(23348)

c 14.7×20.5

septa 74

**Remarks:** The figured holotype of *Axosmilia mexicana* is an unsectioned solitary coral. Fortunately, a thin section of a none-type specimen of *Montlivaultia burckhardti* from the same locality is available, that shows the same morphology and measurements, and certainly is conspecific.

**Occurrence:** Mexico, Coahuila, Los Vagos, Potrero de Oballos (lower Hauterivian).

**Other occurrences:** Lower Aptian of the Central Tethys (Italy), Aptian of the Eastern Tethys (Iran), lower Albian to lower Cenomanian of the Western Tethys (Spain), Coniacian to Santonian of the Central Tethys (Austria).

*Trochophyllia* sp.

Figure 3: 3-4

**Synonymy:**

v 1889 *Trochosmilia* aff. *inflexa* Reuss – Toula, p. 84, pl. 6, fig. 4

vp 1946 *Axosmilia mexicana* Wells, n.sp. – Wells, p. 6, pl. 1, figs. 6-11

**Material:** (UMMP 23350); 1 thin section.

**Dimensions:** (UMMP 23350)

c 12.3×13.5

septa 25

**Remarks:** The specimen 23350 was assigned by Wells to *Axosmilia mexicana*, but it is not a type specimen.

**Occurrence:** Mexico, Coahuila, Barril Viejo, Cuatrociénegas (lower Hauterivian).

**Other occurrence:** Barremian of the Central Tethys (Bulgaria).

*Rayasmiliidae* indet. *coahuilensis* Wells, 1946

Figure 4.1-2

**Synonymy:**

\*v 1946 *Montlivaultia coahuilensis* Wells, p. 4, pl. 1, fig. 5, pl. 2, figs. 4-6

**Material:** UMMP 23340.

**Dimensions:** (23340)

c 36.4 × 38.3

septa 100

**Description:** Solitary cupolate coral. Septa thin, compact, not connected to each other, in a subregular radial symmetry. Septal generations differ by length and thickness. No synapticalae. Endotheca probably by dissepiments. Wall absent. Columella unknown.

**Remarks:** The holotype is a complete specimen, without any polished surface or thin section. The septa are thin and show very little ornamentation. The specimen does, therefore, not belong to *Montlivaultia*. The differing thickness of septa is a clear indication of a position in the *Rayasmiliidae*. Its systematic position cannot be precisely defined without having information about the fine skeletal structure.

**Occurrence:** Mexico, Coahuila, Los Vagos, Potrero de Oballos (lower Hauterivian).

Superfamily *Montlivaultioidea* Felix, 1900

**Remarks:** For the morphology of the superfamily refer to Löser *et al.* (2023).

Family *Montlivaultiidae* Felix, 1900


**Description:** The ornamentation of septal lateral faces is more pronounced in this family. Generally no columella.

Genus *Montlivaultia* Lamouroux, 1821

**Type species:** *Montlivaultia caryophyllata* Lamouroux, 1821

**Description:** Solitary turbinate to cylindrical coral. Without columella.

**Remarks:** *Montlivaultia* is a generic name that has



**Figure 4** 1-2. *Rayasmiliidae* indet. *coahuilensis* (Wells, 1946), Holotype of *Montlivaltia coahuilensis*, (UMMP 23340). 1, coral surface. 2, coral surface, detail. 3-4. *Montlivaltia icaunensis* Orbigny, 1850, Syntype of *Montlivaltia burckhardti*, (UMMP 23342). 3, coral surface. 4, coral surface, with details of the microstructure. 5-6. *Montlivaltia subturbinata* Beauvais and M'Rabet, 1977, (UMMP 23345). 5, transversal thin section. 6, transversal thin section, detail. 7-8. *Styliina bucheti* de Fromentel, 1856, (UMMP 23347). 7, transversal thin section. 8, transversal thin section, detail. Scale bar 1 mm.

been misused for nearly 200 years in been applied to any Mesozoic solitary coral. As a result, the genus currently accounts for a ridiculously high number of species, with an overall count of 384 species (Felix, 1925; Lathuilière, 1989; Löser, 2000; Marchal, 1991); that is, 40 in the Triassic, 238 in the Jurassic, 75 in the Cretaceous, and 40 in the Palaeogene. With so many species, it is impossible to assign any of the present material to any existing species; hence, it became a bad strategy to establish new species for our own material. For example, of the 75 Cretaceous species assigned to the genus, only 24 may belong to *Montlivaltia*, based on the studied type material. On first inspection, most of the remaining species belong to the similar genus *Rayasmilia* Löser, 2022, that differs completely in its septal microstructure. When comparing the true 24 Cretaceous *Montlivaltia* species critically against each other, 15 species may remain. The genus became extinct during the Cenomanian (Löser, 2016), so all Palaeogene material needs to be assigned to other genera.

*Montlivaltia icaunensis* Orbigny, 1850

Figure 4: 3-4

**Synonymy:**

- \*v 1850 *Montlivaltia icaunensis* Orbigny, (2), p. 90
- v 1863 *Montlivaultia icaunensis* – Fromentel, p. 315, pl. 42, figs. 2, 2 a-b, pl. 78, figs. 3, 3 a, pl. 81, figs. 1, 1 a, 2, 2 a
- vp 1946 *Montlivaltia burckhardti* Wells, n.sp. – Wells, p. 5, pl. 2, figs. 7-14
- v 2003 *Montlivaltia* sp. – Baron-Szabo, Hamedani and Senowbari-Daryan, p. 204, pl. 36, fig. 8, pl. 39, fig. 6

**Material:** UMMP 23342.

**Dimensions:** (23342)

c 25 × 28.4mm

septa 94

**Remarks:** Wells mentioned several specimens and indicated two cotypes, that probably correspond to syntypes. Syntype 23343 could no longer be found. Measurements and illustration are, therefore, based on syntype 23342. The specimen is poorly preserved and damaged. No polished

surface or thin section are available. Nevertheless, the ornamentation of the upper margin of the septal and the outline clearly indicate the typical montlivaltiid microstructure. *Montlivaltia burckhardti* Wells, 1946 is here considered as a junior synonym of *Montlivaltia icaunensis* because it presents the same measurements and septal counts.

**Occurrence:** Mexico, Coahuila, Barril Viejo, Cuatrociénegas, (Lower Hauterivian).

**Other occurrences:** Kimmeridgian to Hauterivian of the European Boreal (Germany, France), Barremian of the Central Tethys (Bulgaria), upper Aptian to Albian of the Eastern Tethys (Iran).

*Montlivaltia subturbinata* Beauvais and M'Rabet, 1977

Figure 4: 5-6

**Synonymy:**

v 1882 *Trochosmilia* spec. – Toula, p. 30, pl. 2, fig. 19

vp 1946 *Montlivaltia burckhardti* Wells, n.sp. – Wells, p. 5, pl. 2, figs. 7-14

\* 1977 *Montlivaltia subturbinata* Beauvais and M'Rabet, p. 111, pl. 2, fig. 6

1977 *Montlivaltia kairouanensis* nov. sp. – Beauvais and M'Rabet, p. 114, pl. 3, fig. 5

**Material:** UMMP 23345; 1 thin section.

**Dimensions:** (23345)

c 30.5×33.25

septal 152

**Remarks:** The specimen was named *Montlivaltia burckhardti* by Wells but it is not a type specimen.

**Occurrence:** Mexico, Coahuila, Los Vagos, Potrero de Oballos, (Lower Hauterivian).

**Other occurrence:** Lower Kimmeridgian of the Western Tethys (Spain), upper Kimmeridgian of the European Boreal (Germany), Kimmeridgian to upper Berriasian of the North Africa (Tunisia), Barremian of the Central Tethys (Bulgaria).

Superfamily Styliinoidea Orbigny, 1851

**Remarks:** For the morphology of the superfamily refer to Löser *et al.* (2023).

Family Stylinidae Orbigny, 1851

**Description:** Colonial (phaceloid, plocoid) corals. Septal symmetry radial, in varying systems, bilateral in one genus. Lonsdaleoid septa absent. Columella well-developed, styliform or lamellar.

Genus *Stylina* Lamarck, 1816

**Type species:** *Stylina insignis* Fromentel, 1861

**Description:** Plocoid colony. Corallite outline circular. Symmetry of septa regular radial. Costae non-confluent. Columella styliform. Endotheca consists of thin tabulae. Wall compact, septothecal. Coenosteum broad, consisting of costae and exothecal dissepiments.

*Stylina bucheti* Fromentel, 1856

Figure 4: 7-8

**Synonymy:**

\* 1856 *Stylina bucheti* Fromentel, p. 857

v 1889 *Placocoenia Kaulbarsi* nov. spec. – Toula, p. 82, pl. 5, fig. 12

v 1946 *Placocoenia* n.sp. – Wells, p. 5

1977 *Stylina carthagensis* nov. sp. – Beauvais and M'Rabet, p. 106, text-figs. 1, pl. 1, fig. 1

v 1998 *Stylina pyrenaica* Alloiteau 1946/47 – Schöllhorn, p. 77

**Material:** UMMP 23347; 1 thin section.

**Dimensions:** (23347)

clmin 3.3-3.4

clmax 3.6-4.4

septal 24

**Remarks:** The specimen is poorly preserved, even the genus *Stylina* cannot be clearly confirmed.

**Occurrence:** Mexico, Durango, Las Cuevas, Cuesta del Carbonera, (Valanginian).

**Other occurrence:** Tithonian of the Central Tethys (France), Kimmeridgian of the European Boreal (Germany), Berriasian of the North Africa (Tunisia), Aptian of the Central Tethys (Bulgaria, Greece), upper Aptian of the Western Tethys (Spain).

**Table 1. The coral specimens from the Museum of Paleontology of the University of Michigan in Ann Arbor with their revised taxonomy. Bold names in the second column refer to the type status.**

Specimen UM	Published as	Figured	Type status	Here assigned to
15786	Wells (1946): <i>Isastrea whitneyi</i> Wells 1932 - p. 3, pl. 2: 1-3	pl. 2: 2		<i>Thalamocaeniopsis neocomiensis</i>
15969	Wells (1946): <b><i>Astrocoenia kellumi</i></b> Wells, n. sp. - p. 2, pl. 1: 1	pl. 1: 1	holotype	<i>Stelidioseris icaunensis</i>
15993	Wells (1946): <i>Stephanocoenia guadalupe minor</i> Wells, n.var. - p. 3, pl. 1: 2-4	pl. 1: 4		<i>Eocolumastrea octaviae</i>
19345	Imlay (1940): <b><i>Astrocoenia hispaniensis</i></b> Imlay, n. sp. p. 138, pl. 1: 21, 22	pl. 1: 21, 22	holotype	<i>Holocoenia micrantha</i>
22150	Wells (1942b): <b><i>Thamnasteria imlayi</i></b> Wells, n. sp. - p. 127, pl. 21: 1-3	pl. 21: 1, 2	holotype	(unassigned)
22156	Wells (1942b): <b><i>Stylina arkansasensis</i></b> Wells, n. sp. - p. 128, pl. 21: 4	pl. 21: 4	holotype	(unassigned)
23339	Wells (1946): <b><i>Stephanocoenia guadalupe minor</i></b> Wells, n.var. - p. 3, pl. 1: 2-4	pl. 1: 2, 3	holotype	<i>Eocolumastrea octaviae</i>
23340	Wells (1946): <b><i>Montlivaltia coahuilensis</i></b> Wells, n.sp. - p. 4, pl. 1: 5, pl. 2: 4-6	pl. 2: 4, 5	holotype	Rayasmiliidae indet. <i>coahuilensis</i> Wells 1946
23341	Wells (1946): <i>Montlivaltia coahuilensis</i> Wells, n.sp. - p. 4, pl. 1: 5, pl. 2: 4-6	pl. 1: 5		(unassigned)
23342	Wells (1946): <b><i>Montlivaltia burckhardti</i></b> Wells, n.sp. - p. 5, pl. 2: 7-14	pl. 2: 7, 8	syntype	<i>Montlivaltia icaunensis</i>
23343	Wells (1946): <b><i>Montlivaltia burckhardti</i></b> Wells, n.sp. - p. 5, pl. 2: 7-14	pl. 2: 13, 14	syntype	(not available)
23344	Wells (1946): <i>Montlivaltia burckhardti</i> Wells, n.sp. - p. 5, pl. 2: 7-14	pl. 2: 11		(unassigned)
23345	Wells (1946): <i>Montlivaltia burckhardti</i> Wells, n.sp. - p. 5, pl. 2: 7-14	pl. 2: 10		<i>Montlivaltia subturbinata</i>
23346#1	Wells (1946): <i>Montlivaltia burckhardti</i> Wells, n.sp. - p. 5, pl. 2: 7-14	pl. 2: 14		<i>Trochophyllia communis</i>
23346#2	Wells (1946): <i>Montlivaltia burckhardti</i> Wells, n.sp. - p. 5, pl. 2: 7-14			(unassigned)
23347	Wells (1946): <i>Placocoenia</i> n.sp. p. 5			<i>Stylina bucheti</i>
23348	Wells (1946): <b><i>Axosmia mexicana</i></b> Wells, n.sp. - p. 6, pl. 1: 6-11	pl. 1: 8, 9	holotype	<i>Trochophyllia communis</i>
23349	Wells (1946): <b><i>Axosmia mexicana</i></b> Wells, n.sp. p. 6, pl. 1: 6-11		paratype	<i>Rayasmilia fromenteli</i>
23350#1	Wells (1946): <i>Axosmia mexicana</i> Wells, n.sp. - p. 6, pl. 1: 6-11			<i>Trochophyllia</i> sp.
23350#2	Wells (1946): <i>Axosmia mexicana</i> Wells, n.sp. - p. 6, pl. 1: 6-11	pl. 1: 10, 11		<i>Rayasmilia bangoinensis</i>

### 5. Discussion

During the period when the Cretaceous corals were established by Imlay and Wells, the general opinion was that material from a hitherto unknown area must be taxonomically new. Therefore, both scientists established new taxa or – in the case of Wells – referred to taxa that he established independently. Wells and Imlay did not invest much time in comparing their finds to species established in Europe, Africa, or Asia. After comparing the coral specimens to type material from other areas, it turned out that all new species are synonyms or could not be assigned to a genus because of the poor state of preservation and/or the absence of thin sections. One species – *Rayasmiliidae* indet. *coahuilensis* Wells 1946 – could remain in use, if it could be assigned to a genus. Table 1 gives an overview of the revised collection material.

The number of species is too low to be able to discuss their palaeobiogeographic relationships.

The Valanginian to Hauterivian corals have most joint species with the Barremian to Aptian faunas from the northern margin of the Rhodopes, which corresponds to Bulgaria and Serbia, and also corresponds to the Kimmeridgian of the Northern Tethys (Upper Jurassic of Southern Germany) and the Berriasian to Hauterivian of the epicontinental seas of the NW margin of the African continent (Algeria, Tunisia). The found species have a distribution from the middle Oxfordian to Cenomanian (Figure 5), showing more relationships to Hauterivian to Albian faunas. This is due the rarity of Berriasian to Valanginian faunas. The Berriasian and Valanginian was a time span of a global sea-level fall resulting in regressions and the following erosion, which limited the conservation of shallow marine Berriasian and Valanginian faunas (see also Löser *et al.* (2021) for discussion).




Figure 5 Summarised distribution of species in localities outside of the study area. The thickness of bars corresponds to the number of localities where the species was found. The green bars mark the age of the investigated corals.

## Contribution of the author

The author has done all work alone.

## Funding

The study has not receive any funding except the yearly assignment of the Universidad Nacional Autónoma de México.

## Acknowledgements

I am very grateful to Daniel J. Miller (Ann Arbor) who received me at the Museum of Paleontology of the University of Michigan, and who assisted me greatly during my visit. A grammatical check was carried out by Matthew Copley (Barcelona). Reviews by Ricardo Garberoglio (Buenos Aires) and an anonymous reviewer helped to improve the manuscript.

## Conflict of interests

It does not exist any conflict of interests.

## Handling editor

Francisco J. Vega.

## References

- Alloiteau, J., 1952, Madréporaires post-paléozoïques, in Piveteau, J. (ed.), *Traité de Paléontologie* (1): Paris, Masson, 539–684.
- Alloiteau, J., 1954, Sur cinq genres nouveaux de madréporaires post-paléozoïques: *Bulletin de la Société géologique de France*, (6), 3, 873–887.
- Angelis d'Ossat, G. de, 1905, Coralli del Cretacico inferiore della Catalogna: *Palaeontographia Italica*, 9, 169–251.
- Babaev, R.G., Krasnov, E.V., 1977, [Morphologic peculiarities of scleractinia in the connection of their habitats in Jurassic seas of the South-Eastern Caucasus and Gornogo Crimea.], in Betekhtina, O.A., Zhuravleva, I.T. (eds.), [Environment and life in the geological past.]: *Trudy (Institut geologii i geofiziki. Sibirskoe otd. AN SSSR)*, 302: 137–148.
- Baron-Szabo, R.C., 1997, Die Korallenfazies der ostalpinen Kreide (Helvetikum: Allgäuer Schrattenkalk; Nördliche Kalkalpen: Brandenberger Gosau) *Taxonomie, Palökologie: Zitteliana*, 21, 3–97.
- Baron-Szabo, R.C., Hamedani, A., Senowbari-Daryan, B., 2003, Scleractinian corals from Lower Cretaceous deposits north of Esfahan (Central Iran): *Facies*, 48, 199–216. <https://doi.org/10.1007/bf02667539>
- Beauvais, L., 1986, Monographie des madréporaires du Jurassique inférieur du Maroc: *Palaeontographica*, (A), 194, 1/3, 1–68.
- Beauvais, M., M'Rabet, A., 1977, Les Madréporaires du Berriasien supérieur du Djebel Siouf (Axe Nord-Sud, Tunisie centrale): *Notes du Service géologique*, 43, 11, 103–137.
- Bölsche, W., 1871, Die Korallen des unteren Pläner im Sächsischen Elbthale, in Geinitz, H.B. (ed.), *Das Elbthalgebirge in Sachsen* (1: Der untere Quader: *Palaeontographica*, 20, 46–57.
- Dietrich, W.O., 1926, Steinkorallen des Malms und der Unterkreide im südlichen Deutsch-Ostafrika: *Palaeontographica*, (suppl.7), 1, 43–62.
- Eliášová, H., 1976, Les coraux de l'ordre Hexanthinaria Montanaro-Gallitelli, 1975, Zoantharia de Blainville, 1830, dans les calcaires de Stramberk (Tithonien, Tchécoslovaquie): *Vestník Ústředního ústavu geologického*, 51, 5: 357-366, 3 pls..
- Felix, J., 1891, Versteinerungen aus der mexicanischen Jura und Kreideformation., in Felix, J., Lenk, H. (eds.), *Beiträge zur Geologie und Paläontologie der Republik Mexico* (3): *Palaeontographica*, 37, 140–194.
- Felix, J., 1900, Über die Gruppe der Montlivaltiaceae: *Sitzungsberichte der*

- Naturforschenden Gesellschaft zu Leipzig, (February, 6 1900), 20–24.
- Felix, J., 1925, Anthozoa eocaenica et oligocaenica: Fossilium Catalogus, (1: Animalia), 28, 1–296.
- Fritzsche, C.H., 1924, Neue Kreidefaunen aus Südamerika (Chile, Bolivien, Peru, Kolumbien) (3): Eine neokome Schwamm- und Korallenfauna aus Chile: Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beil.-Band, 50, 313–334.
- Fromentel, E., 1856, Note sur les polypiers fossiles de l'étage portlandien de la Haute-Saône: Bulletin de la Société géologique de France, (2), 13, 851–865.
- Fromentel, E., 1861, Introduction à l'étude des polypiers fossiles: Mémoires de la Société d'émulation du Doubs, (3), 5, 1–357.
- Fromentel, E., 1863, Zoophytes, terrain crétacé (4): Paléontologie française (A.d'Orbigny ed.), 8, 145–192.
- Garberoglio, R.M., Löser, H., Lazo, D.G., 2021, Lower Cretaceous corals from the Agrio Formation, Neuquén Basin, west-central Argentina: Family Columastraecidae: Cretaceous Research, 124, 104817, 1–19. doi.org/10.1016/j.cretres.2021.104817
- Garberoglio, R.M., Löser, H., Lazo, D.G., 2022, Lower Cretaceous corals from the Agrio Formation, Neuquén Basin, west-central Argentina: Families Latomeandriidae, Madreporidae, Thamnasteriidae, and Holocoenia Group: Cretaceous Research, 135, 105195, 1–21. doi.org/10.1016/j.cretres.2022.105195
- Imlay, R.W., 1939, Paleogeographic studies in northeastern Sonora: Geological Society of America bulletin, 50, 11, 1723–1744.
- Imlay, R.W., 1940, Neocomian fauna of northern Mexico: Geological Society of America bulletin, 51, 117–190.
- Kellum, L.B., 1936, Evolution of the Coahuila Peninsula, Mexico (3): Geology of the mountains west of the Laguna district: Geological Society of America bulletin, 47, 7, 1039–1090.
- Kelly, W.A., 1936, Evolution of the Coahuila Peninsula, Mexico (2): Geology of the mountains bordering the valley of Acatita and Las Delicias: Geological Society of America bulletin, 47, 7, 1009–1038.
- Koby, F., 1897, Monographie des polypiers crétacés de la Suisse (2): Abhandlungen der Schweizerischen Paläontologischen Gesellschaft, 23, 29–62.
- Kuzmicheva, E.I., 2002, [Skeletal morphology, systematics and evolution of the Scleractinia.]: Trudy Paleontologicheskogo instituta, 286, 1–211.
- Lamarck, J.B.P., 1816, Histoire naturelle des animaux sans vertèbres (2): Paris, Verdière, 568 pp.
- Lamouroux, J.V.F., 1821, Exposition méthodique des genres de l'ordre des polypiers: Paris, Agasse, 115 pp.
- Lathuilière, B., 1989, Répertoire objectif des coraux jurassiques: Nancy, Presses universitaires, 76 pp.
- Liao, Wei-hua, Xia, Jin-bao, 1985, Upper Jurassic and Lower Cretaceous Scleractinia from Bangoin district of northern Xizang (Tibet): Memoirs of the Nanjing Institute of Geology and Palaeontology, 21, 119–174.
- Liao, Wei-hua, Xia, Jin-bao, 1994, Mesozoic and Cenozoic scleractinian corals from Tibet: Palaeontologia Sinica (Zhongguo-gushengwu-zhi), 184, 1–252.
- Löser, H., 1994 (ed.), The Mesozoic corals. Bibliography 1758-1993: Coral Research Bulletin, 1, 1–99.
- Löser, H., 2000, Répertoire of Species: Catalogue of Cretaceous Corals, 1, 1–137.
- Löser, H., 2004, PaleoTax - a database program for palaeontological data: Computer & Geosciences, 30, 5, 513–521. https://doi.org/10.1016/j.cageo.2004.03.009
- Löser, H., 2008, Early Cretaceous coral faunas from East Africa (Tanzania, Kenya; Late Valanginian-Aptian) and revision of the Dietrich collection (Berlin, Germany): Palaeontographica, 285, 1/3, 23–75. https://

- doi.org/10.1127/pala/285/2008/23
- Löser, H., 2009, Morphology, taxonomy and distribution of the Early Cretaceous coral genus *Holocoenia* (Scleractinia) and its first record in the Caribbean: *Revista mexicana de ciencias geológicas*, 26, 1, 93–103.
- Löser, H., 2012, Intraspecific variation in the genus *Stelidioseris* (family Actinastreae, suborder Archeocaeniina, order Scleractinia; Jurassic-Cretaceous): *Geologica Belgica*, 15, 4, 382–387.
- Löser, H., 2013a, Late Aptian (Cretaceous) corals from Central Greece: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 267, 1, 89–116. <https://doi.org/10.1127/0077-7749/2012/0299>
- Löser, H., 2013b, Revision of the Hauterivian (Early Cretaceous) corals of the Paris Basin, France: a work in progress: *Bulletin d'information des géologues du Bassin de Paris*, 50, 1, 17–24.
- Löser, H., 2016, Systematic part: Catalogue of Cretaceous Corals, 4, 1–710.
- Löser, H., 2022, A new coral family and three new genera (Scleractinia) from the Lower Cretaceous of Puebla and Sonora, Mexico: *Revista mexicana de ciencias geológicas*, 39, 3, 220–229. <http://dx.doi.org/10.22201/cgeo.20072902e.2022.3.1698>
- Löser, H., Arias, C., Vilas, L., 2019, Upper Valanginian to Lower Hauterivian coral faunas from the Sierra Larga (Prebetic zone, SE Spain): *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 292, 3, 259–290. [10.1127/njgpa/2019/0820](https://doi.org/10.1127/njgpa/2019/0820)
- Löser, H., Mendicoa, J., Fernández Mendiola, P.A., 2020, Early Aptian corals from Peñascal (Bilbao; N Spain): *Spanish Journal of Palaeontology*, 35, 2, 209–228. <https://doi.org/10.7203/sjp.35.2.18484>
- Löser, H., Nieto, L.M., Castro, J.M., Reolid, M., 2021, A Lower Valanginian coral fauna from the South Iberian Palaeomargin (Internal Prebetic, SE Spain): *Palaeontologia Electronica*, 24, 1, 1–57. <https://doi.org/10.26879/1030>
- Löser, H., Werner, W., Darga, R., 2023, Middle Cenomanian coral fauna from the Roßsteinalmen (Northern Calcareous Alps, Bavaria, Southern Germany) – a revised and extended version: *Zitteliana*, 97, 89–147. [10.3897/zitteliana.97.113796](https://doi.org/10.3897/zitteliana.97.113796)
- Löser, H., Zell, P., 2015, Revision of the family Columastraecidae (Scleractinia; Cretaceous): *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 277, 2, 153–166. <https://doi.org/10.1127/0077-7749/2008/0250-0045>
- Marchal, D., 1991, Répertoire objectif des coraux du Trias. Rapport préliminaire. Mémoire de Diplôme supérieur: Nancy, Université Henri Poincaré Nancy 1, 80 pp.
- Matthews, S.C., 1973, Notes on open nomenclature and on synonymy lists: *Palaeontology*, 16, 4, 713–719.
- Milne Edwards, H., Haime, J., 1848, Observations sur les polypiers de la famille des astréides: *Comptes rendus hebdomadaires des séances de l'Académie des Sciences*, 27, 19, 465–469.
- Milne Edwards, H., Haime, J., 1849, Mémoire sur les polypiers appartenant à la famille des oculinides, au groupe intermédiaire des Pseudoastréides et à la famille des Fongides: *Comptes rendus hebdomadaires des séances de l'Académie des Sciences*, 29, 4, 67–73.
- Milne Edwards, H., Haime, J., 1851, A monograph of the British fossil corals (2:) Corals from the oolitic formations: *Palaeontographical Society monographs*, 5, 74–146.
- Morycowa, E., 1964, Hexacoralla des couches de Grodziszczce (Néocomien Carpathes): *Acta Palaeontologica Polonica*, 9, 1, 1–114.
- Morycowa, E., 1971, Hexacorallia et Octocorallia du Crétacé inférieur de Rarau (Carpathes orientales roumaines): *Acta Palaeontologica Polonica*, 16, 1/2, 1–149.
- Oppenheim, L.P., 1930, Die Anthozoen der Gosauschichten in den Ostalpen: Berlin, privately published, 604 pp.
- Orbigny, A., 1849, Note sur les polypiers fossiles:

- Paris, Masson, 12 pp.
- Orbigny, A., 1850, Prodrôme de Paléontologie stratigraphique universelle des animaux mollusques et rayonnés (1-2): Paris, Masson, 394 pp., 428 pp.
- Orbigny, A., 1851, Cours élémentaire de Paléontologie (3:) Polypiers ou Zoophytes: Paris, Masson, 2, 151–189.
- Prever, P.L., 1909, Anthozoa, in Parona, C.F. (ed.), La fauna coralligena del Cretaceo dei Monti d'Ocre nell'Abruzzo Aquilano: Memorie descrittive della carta geologica d'Italia, 5, 1, 51–147.
- Reig Oriol, J., 1991, Fauna coralina cretácica del nordeste de España: Barcelona, privately published, 53 pp.
- Reuss, A.E., 1864, Die fossilen Foraminiferen, Anthozoen und Bryozoen von Oberburg in Steiermark: Denkschriften der Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Klasse, 23, 1–38.
- Roemer, F.A., 1841, Die Versteinerungen des norddeutschen Kreidegebirges: Hannover, Hahn'schen Hofbuchhandlung, 145 pp.
- Roniewicz, E., 1987, Correction of homonymy of the generic name Etallonia Roniewicz, 1966 into Etallonasteria nom.n: Acta Palaeontologica Polonica, 32, 1/2, 152.
- Roniewicz, E., 2008, Kimmeridgian-Valanginian reef corals from the Moesian platform from Bulgaria: Annales Societatis Geologorum Poloniae, 78, 2, 91–134.
- Samaniego-Pesqueira, A., Löser, H., Moreno Bedmar, J.A., 2023, Middle Albian corals from the Espinazo del Diablo Formation (Cretaceous; Lampazos area, Sonora, Mexico): Bulletin of Geosciences, 98, 2, 111–159. 10.3140/bull.geosci.1872
- Schöllhorn, E., 1998, Geologie und Paläontologie des Oberapt im Becken von Organyà (Nordspanien): Coral Research Bulletin, 6, 1–139.
- Sikharulidze, G.Ya., 1972, [The new genus Paretallonia (Hexacorallia) from Lower Cretaceous sediments in western Georgia.]: Soobshcheniya AN Gruzinskoy SSR, 68, 3, 641–644.
- Sikharulidze, G.Ya., 1985, [Hexacorals from the Urgonian facies of the Dzirul Massif and its northern frame.]: Trudy Akademija Nauk Gruzinskoy SSR, Geologiceskij Institut, 59, 1–110.
- Tomes, R.F., 1885, Observations on some imperfectly known Madreporaria from the Cretaceous formation of England: Geological Magazine, (N.S., 3) 2, 12, 541–553.
- Tomes, R.F., 1893, Observations on the affinities of the genus Astrocoenia: Quarterly Journal of the Geological Society of London, 49, 569–573.
- Toula, F., 1882, Grundlinien der Geologie des Westlichen Balkan: Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Physikalische Klasse, 44, 2, 1–58.
- Toula, F., 1889, Geologische Untersuchungen im centralen Balkan: Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Physikalische Klasse, 55, 1–108.
- Vasseur, R., Lathuilière, B., 2021, Pliensbachian corals from the Western Tethys: Geodiversitas, 43, 22, 1187–1291. doi.org/10.5252/geodiversitas2021v43a22
- Wells, J.W., 1932, Corals of the Trinity Group of the Commanchean of central Texas: Journal of Paleontology, 6, 3, 225–256.
- Wells, J.W., 1933, Corals of the Cretaceous of the Atlantic and Gulf Coastal Plains and Western Interior of the United States: Bulletins of American Paleontology, 18, 67, 83–292.
- Wells, J.W., 1946, Some Jurassic and Cretaceous corals from Northern Mexico: Journal of Paleontology, 20, 1, 1–7.