

Long-term clinical and radiological outcomes after valgus extension osteotomy and tectoplasty for advanced Legg-Calve-Perthes disease

Resultados clínicos y radiológicos a largo plazo después de osteotomía y tectoplastia de extensión en valgo para la enfermedad de Legg-Calve-Perthes avanzada

Mehmet E. Baki, Muhammet Kalkışım*, and Celal Baki

Orthopedic and Traumatology Department, School of Medicine, Karadeniz Technical University, Trabzon, Turkey

Abstract

Objective: We evaluated the long-term clinical and radiographic outcomes of femoral valgus extension osteotomy combined tectoplasty in Herring group C Perthes disease patients with hinge abduction. **Method:** A total of 13 patients underwent this procedure between 2002 and 2009. In this retrospective study, all patients were classified as Herring group C. The mean age at diagnosis was 8.6 ± 2.2 years. The mean age at surgery was 9.3 ± 1.7 years. The mean age at the most recent follow-up was 23.3 years (range 19-29 years). All patients were followed at least 11.5 years after surgery. **Results:** All patients had hinge abduction deformity preoperatively. The mean Harris hip score improved from 70.23 ± 10.43 points preoperatively to 91.76 ± 7.25 points at the final follow-up. No patient had a limping gait at the final follow-up. The pre-operative visual analog scale score was 7.84 ± 0.22 , and the last follow-up was 1.03 ± 0.25 ($p < 0.001$). According to the Stulberg classification, good radiological outcomes were obtained in 7 hips, whereas fair or poor outcomes were noted in 6 hips. **Conclusions:** In severely affected hips of Perthes, patients who underwent femoral valgus extension osteotomy combined tectoplasty revealed satisfactory long-term clinical and radiological outcomes.

Keywords: Tectoplasty. Femoral valgus extension osteotomy. Hinge abduction. Herring group C.

Resumen

Objetivo: Evaluar los resultados clínicos y radiográficos a largo plazo de la tectoplastia combinada con osteotomía de extensión en valgo femoral en pacientes con enfermedad de Perthes del grupo C de Herring con abducción en bisagra. **Método:** Entre 2002 y 2009 se sometieron a este procedimiento 13 pacientes. En este estudio retrospectivo, todos los pacientes fueron clasificados como Herring C. La edad media en el momento del diagnóstico fue de 8.6 años. La edad media en el momento de la cirugía fue de 9.3 años. La edad media en el seguimiento más reciente fue de 23.3 años. **Resultados:** Todos los pacientes tenían deformidad en abducción en bisagra preoperatoriamente. La puntuación media de cadera de Harris mejoró de 70.23 puntos antes de la operación a 91.76 puntos en el seguimiento final. La puntuación en la EVA preoperatoria fue de 7.84 y en el último seguimiento fue de 1.03. Según la clasificación de Stulberg, se obtuvieron buenos resultados radiológicos en 7 caderas, mientras que se observaron resultados regulares o malos en 6 caderas. **Conclusiones:** En caderas gravemente afectadas de Perthes, los pacientes sometidos a osteotomía de extensión femoral en valgo y tectoplastia combinada revelaron resultados satisfactorios.

Palabras clave: Tectoplastia. Osteotomía de extensión en valgo femoral. Abducción en bisagra. Herring grupo C.

*Correspondence:

Muhammet Kalkışım

E-mail: muhammetkalkisim@gmail.com

0009-7411/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 23-09-2023

Date of acceptance: 20-02-2024

DOI: 10.24875/CIRU.23000483

Cir Cir. 2025;93(3):302-308

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

Perthes disease impaired vascularization in the femoral head and may cause incongruence of the hip joint during the hip remodeling period¹. Hip joint osteonecrosis leads to various deformities of the hip joint, and in later life, secondary end-stage degenerative osteoarthritis may occur². Despite numerous studies, an algorithm could not be established for this clinical entity.

Maintaining hip containment during the disease reduces the deforming forces on the femoral head. During the disease process, the shape of the femoral head and its congruency with the acetabulum affect the prognosis. The “Biological plasticity” concept allows the acetabulum to serve as a mold to the femoral head during the healing process. The treatment goal is to obtain a congruent hip joint with a good range of movement by achieving a spherical femoral head and a well-shaped acetabulum at skeletal maturity. The risk of developing coxarthrosis in the joint may be reduced or delayed^{3,4}.

In the late stage of LCP (Legg-Calve-Perthes) disease, the anterolateral deformed femoral head protrudes outside the acetabulum, and during hip abduction, impingement occurs. Hinge abduction causes incongruity of the hip joint and restricts hip movement. Various surgical procedures ranging from cheilectomy to shelf acetabuloplasty for surgical containment have been described to overcome “hinge abduction.” In the presence of severe femoral head deformity, valgus extension osteotomy may be required as a salvage procedure because it overcomes the hinging and creates a more congruent surface of the femoral head within the acetabulum^{5,6}. Tectoplasty provides an extra-articular weight-bearing surface and a more congruent acetabular roof⁶.

The basis of treatment in Perthes disease is to provide and maintain containment in the hip joint. According to the lateral column classification, Herring group C patients have a poor prognosis. Femoral head deformation and decreased acetabular depth cause containment problems and create discongruence in the hip joint. In case of discongruence in the hip joint, remodeling is impaired. In the combined treatment we performed femoral valgus extension osteotomy to overcome the hinged abduction while the anterolateral part of the femoral head impinges during abduction. The aim of tectoplasty is to create an extra-articular load-bearing surface area on the acetabular side.

The purpose of this study was to evaluate the long-term clinical and radiographic outcomes of femoral valgus extension osteotomy combined with tectoplasty in severe LCP patients. In a previous study, we evaluated the short-term clinical and radiographic outcomes of this technique in 11 Herring group C LCP patients⁷.

Method

The study had institutional review board approval (reference number 2020/170). We evaluated retrospectively the long-term clinical and radiographic outcomes of 13 LCP patients who underwent femoral valgus extension osteotomy combined with tectoplasty between 2002 and 2009. In our study, all patients were diagnosed with LCP and classified according to lateral pillar classification as Herring group C. All patients had intraoperative radiographic and clinically diagnosed hinge abduction.

The study group of 13 patients analyzed consisted of one female and 12 males; in eight cases, the right hip was affected, and in five, the left (Table 1). The mean age at diagnosis was 8.6 ± 2.2 years. The mean age at surgery was 9.3 ± 1.7 years. The mean age at the most recent follow-up was 23.3 years. All patients were followed at least 11.5 years after surgery (mean 15.5 ± 3.3 years). Clinical assessments of all patients were made in terms of pain, range of hip motion, fixed hip contractures, leg length discrepancy, and presence of limp. All patients were examined preoperatively under general anesthesia to confirm the presence of hinge abduction. With the presence of hinge abduction femoral valgus extension, osteotomy was performed in all the patients in our study. A lateral approach was performed to the proximal femur. The femoral neck was prepared for an angled blade plate or K wires. A transversal cut between the femoral shaft and lesser trochanter was performed, and an extension and valgus were built into the osteotomy. The correction angles required for adequate femoral coverage, were determined by comparison of the contralateral side. After femoral valgus extension osteotomy, fluoroscopic evaluation was performed intraoperatively. Tectoplasty procedure was added to patients who were detected to have inadequate femoral head coverage. The tectoplasty procedure aims to enlarge the weight-bearing area at the hip joint and obtain a more congruent joint. It was performed in a supine position by an anterior Smith-Peterson incision with the technique described by Saito et al.⁸. In this

Table 1. Patients demographics

Number of patients	13
Number of hips	13
Age at diagnosis (years)*	8.6 ± 2.2
Age at surgery (years)*	9.3 ± 1.7
Gender (M/F)	12/1
Involved site (R/L)	(8/5)
Length of follow-up (years)*	15.5 ± 3.3
Length of post-op (months)*	188.4 ± 34.6
Age at last follow-up (years)*	15.5 ± 3.3
Lateral pillar classification (A/B/B/C/C)	-/-/13
Patients with Hinge abduction	13

*The values are given as the mean, with the standard deviation in parentheses. M: male; F: female; R: right; L: left.

technique, care was taken not to damage the lateral acetabular growth cartilage. All surgical procedures were led by a single experienced pediatric orthopedic surgeon (C, B). Patients were mobilized non-weight bearing with crutches to the operated side until graft healing was observed. After graft healing and osteotomy union were observed radiologically, partial weight-bearing was permitted, and commencing full weight-bearing began at 3-4 months.

All the patients had pre-operative, early post-operative, and final follow-up clinical and radiographic evaluations (Fig. 1). The radiographic examination included an anteroposterior view of the pelvis in neutral and frog-leg position. Pre-operative staging and severity of disease were determined using modified lateral pillar Herring classification⁹. These subjective assessments were carried out independently by two of the authors (MK and ME-B). The radiographic parameters included Center-Edge angle (CE angle), Sharp angle, femoral head size ratio, neck shaft angle, acetabular depth width index, caput index, subluxation ratio, femoral head coverage ratio, and femoral head extrusion index. Osteoarthritic changes were evaluated using the Tönnis classification¹⁰. At the final follow-up, the radiographic outcome was assessed using the Stulberg classification to evaluate femoral head sphericity and hip remodeling¹¹. The spherical femoral head (Stulberg 1 or 2) was rated as good, the ovoid femoral head (Stulberg 3) as moderate, and the flat femoral head (Stulberg 4 or 5) was rated as poor

outcomes. The clinical assessment includes; the hip range of motion, limb-length discrepancy, the presence of a Trendelenburg sign, and a visual analog scale (VAS) (ranging from 0 to 10). All participants completed the Harris hip scoring system (maximum score, 100 points). A total Harris hip score below 80 points was considered a poor or fair result, and 80-100 was a good or excellent result¹².

Statistical analysis

The statistical analysis was performed using Statistical Package for Social Sciences (SPSS) Version 22.0 (SPSS, Chicago, IL) statistical analysis software. Normal distribution was defined by the Shapiro-Wilk test. All values were calculated as the mean and the standard deviation. The pre-operative and post-operative comparisons were performed using repeated measures analysis of variance. Data were analyzed for statistical significance using the paired t-test, and $p < 0.05$ was considered statistically significant. Spearman correlation analysis was conducted to analyze the pre-operative factors that were strongly associated with patient outcomes. Patients who were ≤ 8 years old at the time of diagnosis were compared with those who were > 8 years old. Harris hip score and the Stulberg classification were used to determine patient outcomes.

Results

All children presented with a painful limping gait preoperatively. At the time of surgery, all hips were classified according to lateral pillar classification as Herring group C. The mean Harris hip score improved from 70.23 ± 10.43 preoperatively to 91.76 ± 7.25 points at the final follow-up. The mean leg length discrepancy preoperatively was 1.46 ± 0.51 cm and after skeletal maturity at the last follow-up, was 0.53 ± 0.59 cm ($p < 0.001$). No patient had a limping gait or Trendelenburg sign at the final follow-up. The mean last follow-up hip flexion was $122.30^\circ \pm 11.65^\circ$. There were no significant complications or need for additional surgery. The pre-operative VAS score was 7.84 ± 0.22 , and the last follow-up VAS score was 1.03 ± 0.25 ($p < 0.001$).

Follow-up radiographs were analyzed for 13 patients (13 hips) (Table 2). According to the Stulberg classification, good radiological outcomes were obtained in seven hips (seven hips were class 2), whereas fair or poor outcomes were noted in six hips (four hips were




Figure 1. Plain radiographs of an 11-year-old boy following femoral valgus extension osteotomy combined tectoplasty. **A:** pre-operative antero-posterior radiograph. Herring group C Perthes. Femoral head subluxation and flattening **B:** pre-operative lateral radiograph **C:** early post-operative anterior-posterior radiograph **D:** post-operative 1-year anterior-posterior radiograph **E:** post-operative 4-years anterior-posterior radiograph **F:** last follow-up post-operative anterior-posterior radiograph. There is good and lasting coverage with satisfaction of the femoral head.

Table 2. Clinical outcomes according to age at diagnosis

Measure	Age at diagnosis		p
	≤ 8 years (n = 5)	> 8 years (n = 8)	
Stulberg classification			
1 or 2	4	2	0.159*
3, 4 or 5	1	5	
Last follow-up Harris hip score			
≥ 80 points	4	7	0.742*
< 80 points	1	1	
Tönnis grade			
0 or 1	3	3	0.471*
2 or 3	2	5	

*No significant difference between ≤ 8 and > 8 (p > 0.05).

class 3 and two hips class 4). Six hips had mild degenerative changes (Tönnis grade 1), five had moderate (Tönnis grade 2), and two had severe changes (Tönnis grade 3). The mean subluxation ratio was 1.66 preoperatively and 1.23 at the last follow-up ($p < 0.001$). The mean femoral head coverage ratio was 67.53% preoperatively and 84.61% at the last follow-up ($p = 0.009$). The mean femoral head size ratio was 1.1 preoperatively and 1.20 at the last follow-up ($p = 0.077$).

Sharp angle decreased from 46.76 to 35.77 ($p < 0.001$). Centre edge angle increased from 14.30 to 40.07 ($p < 0.001$). It was no significant change in neck-shaft angle ($p = 0.477$). The acetabular depth-to-width index improved from 272 to 308 ($p = 0.008$).

Patients who were ≤ 8 years old at the time of diagnosis were more likely to have a lower Stulberg class; however, the differences were not significant (Table 3). There was no significant relation between age at diagnosis and last follow-up Harris hip score and last follow-up Tönnis classification. There were no cases of failure of fixation, and all osteotomies went on to radiologically proven union. There were no significant deep infections.

Discussion

We evaluated the long-term clinical and radiographic outcomes of femoral extension valgus osteotomy combined tectoplasty as a means of surgical containment for Herring group C LCP patients with hinge abduction. In our study, we aimed to provide a spherical femoral head within a well-shaped acetabulum in the final process. Even in the case of collapse and deformation, the femoral head tends to be

Table 3. Comparison of pre-operative and post-operative radiological outcomes

Measure	Pre-operative		1 year post-operative		Last follow-up		p
	Mean	SD	Mean	SD	Mean	SD	
Subluxation ratio	1.66	0.33	1.47	0.26	1.23	0.12	< 0.001*
Femoral head coverage ratio (%)	67.53	11.68	100.30	6.27	84.61	11.75	= 0.009*
Femoral head size ratio	1.13	0.11	1.23	0.18	1.20	0.13	= 0.077
Sharp angle	46.76	6.85	33.38	4.92	35.77	2.95	< 0.001*
Sharp ratio**	1.13	0.11	1.22	0.18	1.19	0.12	= 0.077
Center-Edge angle	14.30	5.21	43.23	6.33	40.07	9.38	< 0.001*
Neck shaft angle	142.61	8.55	152.46	7.63	145.38	10.16	= 0.477
Caput index	0.69	0.12	0.67	0.15	0.74	0.12	= 0.166
Acetabular depth-to-width index	0.27	0.03	0.31	0.05	0.30	0.04	= 0.008*
Femoral head extrusion index	32.13	5.66	0.57	0.22	13.38	6.23	< 0.001*
Acetabular depth-to-width ratio	272.3	31.6	315.3	50.4	308.4	39.1	= 0.008*

*Statistically significant (p < 0.05).

**The ratio of the Sharp angle of the affected hip versus the contralateral normal hip. SD: standard deviation.

remodeled if it is well-contained in the cartilage¹³. All patients in our study were severely affected and had a massive femoral head involvement with lateral subluxation.

The treatment algorithm in severely affected Perthes patients is still controversial. In a multicentric prospective study, it was shown that the lateral pillar classification and age at the time of onset of the disease are strongly correlated with outcomes¹³. Some reports stated that group C hips frequently had poor outcomes unrelated to treatment^{14,15}. However, in those studies, the surgically treated group received either a femoral or acetabular osteotomy, not a combination of these procedures. It was shown in a study of severely affected patients, that a femoral varus osteotomy improved the sphericity of the femoral head compared to conservative treatment¹⁶. Combined procedures in severely affected patients provide better congruency of the hip joint¹⁷. In another study, the combined procedure group obtained significantly better outcomes both clinically and radiologically in severely affected hips compared with the single procedure group¹⁸.

Hinge abduction is the process of severe Perthes disease, causing the deformed and extruded antero-lateral femoral head impingement to the lateral lip of the acetabulum during the abduction. Operative and non-operative treatments were reported to deal with this entity. Traction and bed rest, Shelf

acetabuloplasty, and triple pelvic osteotomy treatment methods were performed. Another treatment method is femoral valgus extension osteotomy, which was first described by Catterall. It rotates the deformed extruded femoral head away from the lateral lip of the acetabulum, and the abnormal hinge movement aimed to get resolved. This relieves pain, reestablishes the abductor mechanism, and improves leg length. Unloading the lateral part of the femoral head and the lateral acetabular physis may produce favorable hip remodeling. We added the tectoplasty procedure to enlarge the weight-bearing surface without decreasing the volume of the acetabulum. Thus, the development of femoroacetabular impingement was prevented. A deformed femoral head and acetabulum cause an obliquity of the weight-bearing joint surface that changes the rotation center of the femoral head. As a consequence, the loading pressure in the reduced articular area increases. Osteotomies are designed to reduce the loading of the joint surface by increasing the area of the weight-bearing surface. Femoral valgus extension osteotomy stretches the superior zone of the joint capsule and its synovial membrane. It stimulates the remodeling capacity of the hip¹⁹. There was no other study that reported the long-term outcomes in the literature of the femoral valgus extension osteotomy combined tectoplasty procedure for hinge abduction in Herring group C perthes patients²⁰.

Stulberg classification has been used to evaluate the surgical clinical outcomes in the literature^{21,22}. In the treatment procedure, we aimed to prevent femoral head subluxation, obtain a congruent hip joint, eliminate hip irritation, restore adequate hip movement, and relieve pain. Treatment modalities performed on patients with severely affected hips vary widely in the literature^{20,23,24}. These are femoral valgus osteotomy, shelf acetabuloplasty, tectoplasty, triple pelvic osteotomy, Chiari's pelvic osteotomy, Salter pelvic osteotomy, and a combination of them. Chang et al. evaluated the outcomes of patients with 21 severely affected hips undergoing the Staheli procedure. According to Stulberg's classification, 33.3% were good (class 1 or 2), 38.1% were fair (class 3), and 28.6% were poor and bad (class 4 or 5)²⁵. In another study, the results of patients who underwent Shelf acetabuloplasty were reported as 51.8% good, 29.6% fair, and 18.6% poor⁵. Huang and Huang reported 14 patients' results who underwent triple pelvic osteotomy as good at 35.7%, moderate at 42.8%, and poor at 21.4%²⁶. We performed a femoral valgus extension osteotomy combined with tectoplasty, which had not been reported in the literature before. In our study, we evaluated the clinical outcomes as 53.8% good, 30.7% moderate, and 15.3% bad. This rate is comparable to reports using other surgical procedures in severely affected hips.

In our study, the femoral head coverage ratio improved from 67.53% preoperatively to 84.61% at the last follow-up. Chang et al. performed a Shelf acetabuloplasty procedure in severely affected hips, and in this study, the femoral head coverage ratio improved from 74% preoperatively to 98% postoperatively²⁵. Huang and Huang performed triple pelvic osteotomy, and improvement was observed from 66% to 101%²⁶. Our long-term outcomes concluded that the procedure improves the femoral head coverage ratio and may contribute to remodeling the hip joint. Daly et al. performed Shelf acetabuloplasty and observed that the femoral head coverage ratio decreased from 83% to 75%⁵. In our study, the femoral head size ratio increased from 1.13 preoperatively to 1.19 at the last follow-up. This result can be associated with the development of coxa magna in the follow-ups. In our study, the subluxation ratio decreased from 1.66 to 1.23. Chang et al. performed Shelf acetabuloplasty, and in this study, the subluxation ratio decreased from 1.6 to 1.17²⁵. It was shown that the Sharp angle in Perthes' hips is steeper than the unaffected side during the course of the disease²⁷. In our study, the Sharp

angle decreased from 46.76° preoperatively to 35.77° and the CE angle increased from 14.30° to 40.07°. Ghanem et al. performed Shelf acetabuloplasty in severely affected patients, and the Sharp angle decreased from 45° preoperatively to 35° postoperatively, and the CE angle increased from 12.5° to 44°³. In a study, it was shown that the acetabulum is wider and shallower at all stages of Perthes²⁷. In our study, our results (272 increased preoperatively-308 postoperatively) show that the combined procedure had a stimulatory effect on the acetabular depth growth with the capsular tension effect.

Age at onset is considered one of the most important predictors of prognosis in perthes patients. Obtaining a spherical femoral head and establishing its congruence within the acetabulum is one of the main goals in Perthes disease treatment²⁸. In our study, patients who were ≤ 8 years old at the time of diagnosis were more likely to have a lower Stulberg class, better outcomes, and lower Tönis class, but the differences were not statistically significant (Table 3).

If hinge abduction develops in LCP disease, hip containment is impaired. In this situation, valgus extension osteotomy is an effective method used as a salvage procedure. It is a procedure that reduces the uncovered femoral head into the hip joint and obstructs impingement during the abduction. One of our goals was to induce hip remodeling with this procedure, which we performed in cases where the acetabular growth capacity continued, and the triradiate cartilage was open.

The study has some limitations. The lack of a control group, the relatively small number of patients, and the retrospective study design are some limitations of our study. However, the follow-up was 15 years, which was long enough to evaluate the clinical and radiographical combined surgical procedure. In addition, this study has limited value for femoroacetabular impingement evaluation.

Many reports emphasized that severely affected hips in LCP had poor outcomes. In our study, we aimed to increase joint congruency and achieve satisfactory clinical results. It improved hip functional scores at skeletal maturity and relieved pain scores dramatically. We concluded that valgus extension osteotomy combined tectoplasty procedure in severely affected hips with hinge abduction is an effective treatment method and had satisfactory 15-year follow-up outcomes. Our method could be an alternative treatment option for late-stage Perthes patients.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients, and received approval from the Ethics Committee. The SAGER guidelines were followed according to the nature of the study.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Joseph B. Natural history of early onset and late-onset Legg-calvé-perthes disease. *J Pediatr Orthop.* 2011;31:S152-5.
2. Shah H. Perthes disease: evaluation and management. *Orthop Clin North Am.* 2014;45:87-97.
3. Ghanem I, Haddad E, Haidar R, Haddad-Zebouni S, Aoun N, Dagher F, et al. Lateral shelf acetabuloplasty in the treatment of Legg-Calvé-Perthes disease: improving mid-term outcome in severely deformed hips. *J Child Orthop.* 2010;4:13-20.
4. Kim HK, Herring JA. Pathophysiology, classifications, and natural history of perthes disease. *Orthop Clin North Am.* 2011;42:285-95, v.
5. Daly K, Bruce C, Catterall A. Lateral shelf acetabuloplasty in Perthes' disease. *J Bone Joint Surg Br.* 1999;81-B:380-4.
6. De Gheldere A, Eastwood DM. Valgus osteotomy for hinge abduction. *Orthop Clin North Am.* 2011;42:349-54, vi-vii.
7. Mehmet Emre Baki MD, Duygu Baki MD, Hafız Aydın MD, Servet Keri-moğlu MD, Celal Baki MD. Radiological evaluation of combined valgus extension osteotomy and tectoplasty for the treatment of Herring group C Perthes patients. *Jt Dis Relat Surg.* 2011;22-2:64-8.
8. Saito S, Takaoka K, Ono K. Tectoplasty for painful dislocation or subluxation of the hip. Long-term evaluation of a new acetabuloplasty. *J Bone Joint Surg Br.* 1986;68:55-60.
9. Herring JA, Kim HT, Browne RP. Legg-calvé-perthes disease part I: classification of radiographs with use of the modified lateral pillar and stulberg classifications. *J Bone Jt Surg Am.* 2004;86:2103-20.
10. Tönns D. Normal values of the hip joint for the evaluation of X-rays in children and adults. *Clin Orthop Relat Res.* 1976;119:39-47.
11. Stulberg SD, Cooperman DR, Wallensten R. The natural history of legg-calvé-perthes disease. *J Bone Jt Surg.* 1981;63:1095-108.
12. Marchetti P, Binazzi R, Vaccari V, Girolami M, Morici F, Impallomeni C, et al. Long-term results with cementless fitek (or fitmore) cups. *J Arthroplasty.* 2005;20:730-7.
13. Salter RB. The present status of surgical treatment for legg-perthes disease. *J Bone Joint Surg.* 1984;66:961-6.
14. Grzegorzewski A, Bowen JR, Guille JT, Glutting J. Treatment of the collapsed femoral head by containment in legg-calvé-perthes disease. *J Pediatr Orthop.* 2003;23:15-9.
15. Hefti F, Clarke NM. The management of Legg-Calvé-Perthes' disease: is there a consensus?: A study of clinical practice preferred by the members of the European paediatric orthopaedic society. *J Child Orthop.* 2007;1:19-25.
16. Terjesen T, Wiig O, Svenningsen S. Varus femoral osteotomy improves sphericity of the femoral head in older children with severe form of Legg-Calvé-perthes disease. *Clin Orthop Relat Res.* 2012;470:2394-401.
17. Kamegaya M, Morita M, Saisu T, Kakizaki J, Oikawa Y, Segawa Y. Single versus combined procedures for severely involved legg-calvé-perthes disease. *J Pediatr Orthop.* 2018;38:312-9.
18. Kamegaya M, Saisu T, Ochiai N, Hisamitsu J, Moriya H. A paired study of Perthes' disease comparing conservative and surgical treatment. *J Bone Joint Surg Br.* 2004;86:1176-81.
19. Bambelli R. Valgus-extension osteotomy. In: *Osteoarthritis of the Hip Pathogenesis and Consequent Therapy.* Berlin, Heidelberg, New York: Springer-Verlag; 1976, 1983. p. 49-59.
20. Wenger DR, Pring ME, Hosalkar HS, Caltoun CB, Lalonde FD, Bastron TP. Advanced containment methods for Legg-Calvé-Perthes disease: results of triple pelvic osteotomy. *J Pediatr Orthop.* 2010;30:749-57.
21. Wiig O, Terjesen T, Svenningsen S. Prognostic factors and outcome of treatment in Perthes' disease: a prospective study of 368 patients with five-year follow-up. *J Bone Jt Surg Br.* 2008;90:1364-71.
22. Ibrahim T, Little DG. The pathogenesis and treatment of legg-calvé-perthes disease. *JBJS Rev.* 2016;4:e4.
23. Freeman RT, Wainwright AM, Theologis TN, Benson MK. The outcome of patients with hinge abduction in severe perthes disease treated by shelf acetabuloplasty. *J Pediatr Orthop.* 2008;28:619-25.
24. Farsetti P, Benedetti-Valentini M, Potenza V, Ippolito E. Valgus extension femoral osteotomy to treat "hinge abduction" in Perthes' disease. *J Child Orthop.* 2012;6:463-9.
25. Chang JH, Kuo KN, Huang SC. Outcomes in advanced legg-calvé-perthes disease treated with the staheli procedure. *J Surg Res.* 2011;168:237-42.
26. Huang MJ, Huang SC. Surgical treatment of severe perthes disease: comparison of triple osteotomy and shelf augmentation. *J Formos Med Assoc.* 1999;98:183-9.
27. Huhnstock S, Svenningsen S, Pripp AH, Terjesen T, Wiig O. The acetabulum in Perthes' disease: a prospective study of 123 children. *J Child Orthop.* 2014;8:457-65.
28. Quain S, Catterall A. Hinge abduction of the hip. Diagnosis and treatment. *J Bone Joint Surg Br.* 1986;68:61-4.