

The recent instrumental seismicity of Syria and its implications

Mohamad Khir Abdul-Wahe* and Jamal Asfahani

Received: March 14, 2017; accepted: March 06 2018; published on line: April 02, 2018

Resumen

Esta contribución es un intento por ampliar el conocimiento actual sobre la reciente sismicidad instrumental que fue registrada durante el período 1995-2012 por la red sismológica nacional Siria, así como sus escenarios sismotectónicos. La reciente sismicidad instrumental ha demostrado que se han producido un pequeño número de eventos de baja magnitud. En consecuencia, en comparación con la sismicidad histórica, esto indica que la actividad está pasando por una relativa inactividad.

La correlación entre la sismicidad instrumental y las características sismotectónicas se realizó mediante el análisis de distribuciones espaciales de eventos sísmicos y mecanismos focales de algunos eventos más fuertes. Los resultados actuales permiten observar muchos tipos de actividad sísmica de la siguiente manera: tipo de enjambre, tipo de conglomerado y tipo ocasional, lo que podría mejorar la comprensión del comportamiento de las fallas sísmicamente activas. Los largos períodos de retorno de grandes terremotos ($M \geq 5$) y la brevedad de la sismicidad instrumental nos impiden distinguir por completo la actividad sísmica y descubrir todas las fallas activas en el país.

Palabras clave: sismicidad instrumental, Sistema de falla del Mar Muerto, Swarm, Siria.

Abstract

This contribution is an attempt to enlarge the current knowledge about the recent instrumental seismicity, recorded during the period 1995-2012 by the Syrian national seismological network, as well as the seismotectonic settings in Syria. The recent instrumental seismicity has shown that the earthquake activity has produced a little number of low magnitude events. Consequently, it indicates that this activity is actually passing through a relative quiescence in comparison with the historical seismicity.

The correlation between the instrumental seismicity and the seismotectonic features was performed by analyzing spatial distributions of seismic events and focal mechanisms of some strongest events. The current results, allow observing many types of the seismic activity as follows: Swarm-type, Cluster-type, and Occasional-type, which could improve the understanding of the behavior of the seismically active faults. The long return periods of large earthquakes ($M \geq 5$) and the shortness of instrumental seismicity, prevent us to completely characterize the seismic activity and to discover all the active faults in the country.

Key words: Instrumental seismicity, Dead Sea Fault System, Swarm, Syria.

M. K. Abdul-Wahed*
Jamal Asfahani
Dept. of Geology the Atomic Energy Commission of
Syria (AECS)
P.O. Box 6091 Damascus-Syria

*Corresponding author: cscientific@aec.org.sy

Introduction

Syria has suffered from several destructive earthquakes along the history, where the historical documents show many large earthquakes along the Dead Sea Fault System (DSFS). During the past decade, palaeoseismic investigations along the DSFS have proven successful in identifying surface ruptures associated with historical earthquakes (Gómez *et al.*, 2003; Wechsler *et al.*, 2014), and in documenting evidences for long-term earthquake behavior such as temporal clustering (Klinger *et al.*, 2015; El-Isa *et al.*, 2015). The focus in the present study corresponds to the recent instrumental seismic activity in Syria. The Syrian National Seismological Network (SNSN) has been installed in 1995 and covered the Syrian territories. Thanks to the SNSN, this work intends to be the first careful and detailed analysis of the recent instrumental seismicity of Syria, which has been recorded during the period from 1995 to 2012. The main purpose of this work is to characterize the seismicity pattern of Syria by locating earthquakes and calculating the focal mechanism of some strongest events, and then, relate the earthquake activity with the well-known tectonic structures in the region. This would improve and strengthen the knowledge of the

actual seismotectonic deformations taking place now. Thereby, the instrumental seismic activity should be documented, and it is hoped to improve the seismic hazard assessment and the development of appropriate risk-mitigation strategies.

Seismotectonic setting

Syria is located on the northern part of the Arabian plate (Figure 1). It is bounded from the west by the northern section of the DSFS. Northeast of Antioch, the DSFS intersects the Eastern Anatolian Fault System (EAFS) and the Bitlis Suture zone, both of which comprise the northern border of the Arabian plate. At a gross scale, Syria can be spatially divided into four major 'tectonic zones' and intervening structural highs include the Palmyrides Fold Belt (PFB), the Abd-el-Aziz Faults, the Euphrates Fault System (EFS) and the DSFS (Figure 2). These major structural zones have accommodated most of the tectonic deformation in the country throughout the Phanerozoic, whereas the intervening stable areas remained structurally high and relatively undeformed (Barazangi *et al.*, 1993; Brew *et al.*, 2001). Additionally, the style of structural reactivation, in Syria, is dependent on the orientation of the aforesaid tectonic regions to the prevailing stress




Figure 1. Map showing simplified tectonic setting of the Arabian plate and nearby tectonic features around Syria. Abbreviations: EAF: Eastern Anatolian Fault system, NAF: Northern Anatolian Fault system, Med: Mediterranean.

pattern. Furthermore, (Barazangi *et al.*, 1993) inferred that the tectonism within the country is concentrated in those four major tectonic zones, which include a fold and thrust belt, a plate boundary transform fault, inverted basins and an extensive aborted rift. The DSFS is an extremely important tectonic feature, which accounts for the bulk of seismic activity in the eastern Mediterranean and accommodates the relative sinistral motion between the African and Arabian plates (Reilinger and McClusky, 2011). Recent GPS studies have characterized the first order geodetic velocity field around the DSFS (Alchalbi *et al.*, 2010; Al-Tarazi *et al.*, 2011; Saleh *et al.*, 2012; Mahmoud *et al.*, 2013; Masson *et al.*, 2015). Focal mechanisms of moderate-to-large earthquakes show a sinistral motion along the DSFS (Salamon *et al.*, 1996; Hofstetter *et al.*, 2007). The state of stress is characterized by the coexistence of a normal faulting stress regime with the primarily strike-slip one (Palano *et al.*, 2013). In a review of the slip and seismicity of the DSFS, Garfunkel (2011) concludes that the slip rate is slowing from an average rate of 6–7 mm/year over the last 5 Ma to 4–5.5 mm/year in the Pleistocene together with a slight eastward shift of the Euler

pole of rotation between Sinai and Arabia. A quick review of the historical seismic activity in the eastern Mediterranean clearly demonstrates that this part of the world has been shaken since 2000 B.C. by strong earthquakes that destroyed thousands of structures, and caused severe casualties and loss of human life (Darawchek *et al.*, 2000; Khair *et al.*, 2000; Guidoboni *et al.*, 2004; Guidoboni & Comastri, 2005). The return periods of large earthquakes ($M \geq 5$) in Syria was estimated to be about 200–350 years (Ambraseys and Barazangi, 1989).

Data and Methods

In 1985 the General Establishment of Geology and Mineral Resources (GEGMR), responding to an initiative proposed by the Program for Assessment and Mitigation of Earthquake Risk in the Arab Region (PAMERAR), started the design and implementation phases for the construction of the SNSN. The network's design objective was to monitor all discernible earthquake activity along the DSFS and its related branches in Syria and nearby Lebanon such as Serghaya fault. The 9 seismic stations of the southern sub-network became operational in late 1994 and early 1995




Figure 2. Tectonic zones of Syria (modified from Brew *et al.* (2001)).

and allowed to record the micro-earthquake activity that occurred on November 22, 1995, along the Serghaya fault. The 11 stations of the northern sub-network followed the next year to complete the 20 stations of the western SNSN. The SNSN has been officially operated since January 14, 1995 (Dakkak *et al.*, 2005). The 7 eastern stations were added in April 2002, where 2 of them were three-component seismic stations. In 2003, the SNSN consists of 27 short period (1 Hz) stations of ~50 km seismograph spacing (Figure 3). More information about the SNSN data, such as acquisition, transmitting, processing, recording, can be found in (Dakkak *et al.*, 2005).

The SNSN has provided the local instrumental data for the first time in Syria. During the study period 1995-2012, the SNSN has digitally recorded about 5000 local events. Hypocentral locations for the recorded local events were determined using a program based on HYP071 algorithm (Lee, 1990), in which, the change in the arrival time due to change in hypocentral position is linearized. This program takes the station locations, crustal seismic velocity model, and the phase information in the input file. It assumes a trial origin time and hypocentral location for the earthquake, and improve the origin time and hypocentral location by iteratively minimizing the least square error of the travel time computed from the input

station, crustal seismic velocity model and phase information. Because of the linearization of the problem several iteration steps are necessary to obtain the smallest misfit of observed and predicted travel-times, which is represented by the minimum average location root mean square error (RMS).

The hypocentral coordinates and the origin time are iteratively estimated using stepwise statistical regression procedure. That is an inversion process, which need many control parameters. Some of them have been fixed as the following: minimum number of stations to attempt a solution to be 4, minimum number of phases to be 3, cut-off RMS value to be 0.99 s, maximum number of iterations to be 99.

It is very common to express the overall location quality by the RMS as a value of the error parameters. Since seismic events are located with arrival times that contain observational errors and the travel times are calculated under the wrong assumption that the exact model is known, all hypocenter determinations will have errors. The RMS of the final solution is very often used as a criterion for 'goodness of fit'. Although it can be an indication, RMS depends on the number of stations and does not in itself give any indication of errors. It is therefore not sufficient to give one number for the hypocenter error since it varies spatially in horizontal and vertical




Figure 3. Map showing geographical distribution of the Syrian National Seismological Network (▲) and the principal faults. Fault geometry is summarized from Brew *et al.* (2001). Abbreviations: fault (F.), basin (B.).

directions. The accuracy of events' location is a critical factor for reliably identifying faulting parameters. To improve the accuracy of location, the records were filtered by using Butterworth order 3 filter, in order to get the best signal-noise ratio, where all P and S phases were manually picked. As a result, the number of arrival time readings was increased. Events were relocated with the new arrival times.

The magnitudes have been calculated from the coda wave duration via the formula (Bulletin of SNSN, 1995-2009):

$$M_c = -3.0 + 2.6 * \log(T) + 0.001 * D \quad (1)$$

where T is the coda duration (in sec) and D is the epicentral distance (in km).

A quantitative idea about magnitude and frequency of events could be obtained by adopting the empirical Gutenberg and Richter relation:

$$\log N = a - bM \quad (2)$$

where N is the number of earthquakes of magnitude equal or greater than M, and a and b are constants for the region, where a is a measure of the level of seismic activity and b (b-value) is the rate at which events occur within a given magnitude range. Different methodologies for assessing the b-value of the Gutenberg and Richter relation are available in literature. The least-squares method (LSM) is often used, although not formally suitable since magnitude is not error free, cumulative event counts are not independent, and the error distribution of the number of earthquake occurrences does not follow a Gaussian distribution. The maximum likelihood method (MLM) has been widely applied to assess the b-value. In this study, the Gutenberg and Richter relation is evaluated from the available recorded events by the LSM and MLM methods using the code "bvalue.exe" from the SEISAN 10.3, software (2015).

The low seismicity in Syria and the insufficient coverage of seismic stations of SNSN (Dakkak *et al.*, 2005) limit the number of fault plane solutions that can be obtained from the available records. The greater the magnitude, the better the seismogram trace and the higher number of seismic stations registering the event, are allowing an improved location and a more reliable focal mechanism determination. However, small events often exhibit a large scatter due to local structure and stress perturbations and are poorly representative of the regional

stress field. Therefore, a probabilistic method, proposed by Zollo & Bernard (1991), was applied to calculate the focal mechanisms of some largest events. This non-linear inversion method assumes a double-couple point-source model, a well determined earthquake location, and the model parameters to be the fault angle orientations: strike, dip and slip (or rake). The posterior probability of these parameters for the given observational data set, P-wave polarities, is computed by using Bayesian approach. The method is based on the estimation, by an exhaustive search of the posterior probability, of model parameters. The maximum likelihood solutions are only represented by its lower hemisphere projection. The available data and the reliability of the solutions are carefully examined in order to get the best results.

Detailed analysis of results

The availability of recent instrumental data, allows the knowledge about the seismotectonic of Syria to be improved. For this instance, the spatial analysis is an important key, where a spatial comparison between seismicity and mapped fault locations allows to assess the possible activity along a given fault or group of faults. The hypocentral locations for the recorded events have been determined with a location uncertainty of less than 10 km (Figure 4). Thus, the epicentral map (Figure 5) can be considered reliable for the following interpretation of the present work. According to the epicentral map, the recent instrumental seismicity inside Syria is characterized by many clusters of weak events ($M < 4$). Most of the recorded events are low magnitude and can be qualified as weak, where their average magnitude is approximately 2.0 (Figure 6). The focal mechanism of some strongest events (Figure 7) could be helpful to relate the earthquake activity with the well-known tectonic structures in the region. A detailed analysis on the recent instrumental seismicity, carried out in the present paper, allows observing some seismic types as follows: Swarm-type, Cluster-type, Occasional-type. This classification could help to present in detail the properties of seismic activity and its relationship with the tectonic features of some specific areas, and to focus on the most important fault zones.

1- Swarm-type seismic activity

Earthquake swarms are generally observed in volcanic regions, and defined as prolonged series of earthquakes of small to moderate magnitude without a distinct major event and strike in a relatively short period of time. The seismicity of the Jordan Dead Sea Transform




Figure 4. Histograms of location uncertainty in horizontal and vertical directions, and the RMS.

region is characterized by a relatively high proportion of seismic sequences and swarms, some of which are related to subsurface volcanic activities (El-Isa, 2012; Salamon, 2009). About 57 % of historical seismicity along the DSFS occurred in the form of sequences and swarms that lasted for variable periods of time, ranging from a few hours to a few days, weeks, months, and a few years (El-Isa *et al.*, 2015). Relying on the instrumental seismicity of the Gulf of Aqaba region, El-Isa (2012) deduced that not less than 98 % of the local seismicity occurred in form of sequences and swarms. Swarms are often characterized by multiple or double earthquakes with very similar waveforms. The multiples or doublets have been interpreted as repeated slip on fault asperities or clusters of asperities. It offers the opportunity to re-examine the relationship between pre-existing

fault zones and current seismicity in order to improve thereafter our understanding of the intracratonic seismicity patterns (Uski *et al.*, 2006). Many swarms were observed in the recent instrumental seismicity in Syria, where the correlation between the observed swarms and the nearby faults was interpreted and analyzed through the spatial distributions of the hypocenters, as follows:

a- Serghaya fault

The NE–SW striking Serghaya fault branches out from the main transform of DSFS. It is a prominent structure located approximately along the Syrian–Lebanese border. Combined field investigations in geomorphology and paleoseismology have detected young fault scarps, mole tracks, pressure ridges and offset




Figure 5. Earthquake activity in Syria and nearby countries recorded by Syrian National Seismological Network from 1995 to 2012.




Figure 6. Frequency-magnitude distribution of the recorded events during the period of 1995-2012 in cumulative and non-cumulative plots.

streams along the fault trace, which attest to recent coseismic ruptures (Nemer *et al.*, 2008). The results of trenching (Gómez *et al.*, 2001; 2003) demonstrated that the Serghaya fault has important implications in terms of regional tectonics and earthquake hazard. Serghaya fault was generally regarded, until 1995, as inactive since the Pliocene (Walley, 1988; Girdler, 1990; Butler *et al.*, 1997). Asfahani and Abdul-wahed (2012) established an instrumental earthquake catalogue for the Serghaya fault for the period extending from 1995 till 2009, in order to characterize and evaluate its seismic hazard and behavior. They found that the earthquake activity of this fault produces little number of low magnitude events, at least during the study period. On November 22, 1995, a swarm of events was observed along the Serghaya fault (Figure 8). Mohamad *et al.* (2000) reported that this swarm was immediately triggered following the 1995 Aqaba earthquake at a distance of about 500 km. A total of 21 events occurred as swarms during the first three and half hours following the mainshock of Aqaba, where the strongest one was of magnitude $M_s=3.7$. The Serghaya swarm is the first

instrumentally swarm recorded by SNSN. The similarity of waveforms between the swarm events was examined, where a multiplet of 9 earthquakes and 3 of double earthquakes were found (Abdul-Wahed *et al.*, 2006; Abdul-Wahed, 2007; Abdul-Wahed *et al.*, 2011). The composite focal mechanism of the biggest multiplet (Fig. 7) demonstrates a good agreement between the fault plane solutions with NE Serghaya sinistral fault motion.

b- Southwest of Palmyrides fold-thrust belt

The Palmyrides fold-thrust belt, a NE-SW-trending ~400 km long and 100 km wide, is divided into southwestern and northeastern regions separated by Jhar fault (Figure 2). In the southwest of Palmyrides fold-thrust belt, 3 swarms of events were observed in 2004 in the west of Jayrud city (Figure 8). The first swarm occurred on February 8, 2004, including 6 events. The second occurred in the same site on May 16 and 17, 2004, including 22 events. In spite of the small size of the events ($M_c=2.6-3.3$), this swarm was widely felt at Ma'alula city and its surroundings. The third swarm of




Figure 7. The focal mechanisms map of some largest events: a) an earthquake of Palmyra which happened on December 24, 1996 $M_c=5.6$, b) an event from the Rouj basin on May 20, 2003 $M_c=4.0$, c) an event from the Ghab basin on January 3, 1997, $M_c=3.5$, d) an event from Latakia-Killis faults system on July 19, 1996 $M_c=3.9$, e) an event from Abd-el-Aziz uplift faults on January 11, 1996 $M_c=3.5$, f) an earthquake of Bishri fault on December 18, 1994 $M_c=5.0$, g) the composite focal mechanism of the detected multiplet in Asharneh Plain, h) the composite focal mechanism of the biggest multiplet detected in Serghaya swarm on November 22, 1995, i) an earthquake of Roum fault on March 26, 1997, $M_c=4.9$, j) the composite focal mechanism of the detected doublet in Zobeideh fault. M_c : coda magnitude.




Figure 8. Detailed epicenters map of some important seismic activities observed in Damascus region.

8 events occurred on June 7, 2004, in which, an event attend the magnitude of 3.9. Another swarm of 8 events was observed on April 21, 2011, in Jayrud region. Those swarms can be related to southwest of Palmyrides faults as an extension of Damascus NE fault.

c- Baer-Bassit region faults

The Baer-Bassit region is located in a critical region near the boundary of the African and the Eurasian plates and the interaction between DSFS and EAFS. A Latakia-Killis faults system (Figure 2) was mapped as a thrust fault bordering the Baer-Bassit massif to the northeast, parallel to EAFS, where unusually steep dips were noted. This system was seen as connecting Latakia city area to the town of Killis in southern Turkey and the Kurd Dagh region, ~150 km to the northeast. The Baer-Bassit region is seismically considered as the most active in Syria (Figure 5) with high density of faults extending in the sea toward the southwest according the Cyprus arc. A swarm of 4 events started on June 3, 1996, in the sea against the Baer-Bassit region, where the biggest one has a magnitude of 4.5 and was felt in the whole coastal region (Sbeinati and Darawchah, 1997). Three swarms have been observed around Kassab city. The first of 5 events occurred on June 19, 1996. The second swarm of 6 events occurred on January 3, 2003, in which, an event had magnitude of 4.1. The third swarm of 3 events

occurred on February 1, 2007. These swarms have been probably generated on Latakia-Killis faults system.

d- Kurd-Dagh region faults

The Kurd-Dagh region is located in the northwest of Aleppo plateau (Figure 2). This region is seismically active with high density of faults extending toward the northeast according to Latakia-Killis faults system. A swarm of 5 events started on March 23, 2010, in the southwest of Rajo city, and a swarm of 3 events occurred on September 21, 2011, in the south of Afrin city. Another swarm of 4 events occurred on November 20, 2011, in the south of Azaz city. Although the events of those swarms are of low magnitude, however, they indicate the activity of Kurd-Dagh region.

e- Roum fault

Fresh fault scarps and pressure ridges visible along the Roum fault trace attest to recent coseismic ruptures (Nemer and Meghraoui, 2006). On this fault, branching out from the DSFS in south Lebanon at the southern deflection point of the Yammouneh bend, a swarm of events started on March 26, 1997, by two events, with magnitudes 4.9 and 4.5, followed by 6 micro-events the next day. The first two events were felt in western Damascus (about 50 km far away), where the felt effects

were rattling of windows till some windows were broken in Serghaya (Sbeinati and Darawchah, 1998). The composite focal mechanism of these two events (Abdul-Wahed and Al-Tahan, 2010) shows a good agreement between the fault plane solutions, and the NW Roum fault indicating a left-lateral motion on an NW striking fault (Figure 7).

2- Cluster-type seismic activity

The seismicity inside Syria (Figure 5) is characterized by many clusters of weak events ($M < 4$) such as on southwest Aleppo plateau faults. The correlation between different seismic clusters and the seismotectonic features is analyzed through the spatial distributions of seismic events and the focal mechanisms. The cluster-type seismic activity allows to distinguish between the next faults:

a- Latakia-Killis faults system

Many clusters of events can be associated to this faults system and some of them extend in the sea toward the southwest according to the Cyprus arc. The focal mechanisms of many events demonstrate a sinistral strike slip common mechanism trending NE in agreement with the known fault system (Figure 7).

b- Northern segments of DSFS

The epicentral map (Figures 5 and 9) has shown that the instrumental seismicity is very low along the northern segments of DSFS: Masyaf, Apamea and Ghab. A cluster of 4 events was located in the middle of the Ghab basin. One

event of the cluster had a normal mechanism (Abdul-Wahed, 2012) in agreement with the NW normal fault associated with the local trans-tensional features observed by Brew *et al.* (2001b). The Rouj basin, ~24-km-long and up to ~5-km-wide, is bounded by left-lateral faults (Fig. 3). A cluster of 42 events was observed in this basin (Figure 9). The focal mechanism, calculated for the biggest events, has shown a sinistral mechanism trending NE in agreement with the main existent faults (Figure 7). A cluster of 17 events was observed at the northeast extension of St. Simeon fault around Shran city. At the northern end of the Ghab Basin, the El-Wastani mountain is bounded by Armanaz fault and Salqin fault in the east and west respectively. A cluster of 41 events was observed at the El-Wastani mountain. This cluster could be related to Salqin-Armanaz faults. Asharneh plain is located in the east of the south part of the Ghab basin (25 Km northwest of Hama city) (Figure 9). A cluster of about 20 events was located in the Asharneh plain. Three of them were aligned in the NW direction and form a multiplet. The composite focal mechanism of these 3 events shows a normal fault mechanism (Figure 7) which could also be associated with local trans-tensional features. A cluster of 142 events was located between Hama and Rastan cities and aligned in the NW direction. It could be related to unmapped Hama fault.

c- Abd-el-Aziz uplift faults

The Abd-el-Aziz uplift faults, in the northeast of Syria, were roughly east-west striking and were active almost exclusively in the latest Cretaceous. These faults extend from the west




Figure 9. Detailed epicenters map of the seismic activity recorded in central part of Syria.

through Abd-el-Aziz mountain (~100 km length, max. elevation 920 m) and eastwards well into Iraq (Figure 2). The most recognized earthquake along the Abd-el-Aziz uplift faults occurred in 1975 with magnitude of $M \approx 5.2$ (Ahmad, 2013). A cluster of 43 events was observed in the middle part of the Abd-el-Aziz uplift faults (Figure 5), where an event reached magnitude 4.7. Another cluster of 31 events was observed in the eastern part. Four events, aligned in the EW direction along the Abd-el-Aziz uplift, show nearly identical source mechanism solutions and have a sinistral mechanism trending EW (Figure 7). They might indicate shearing process in this direction.

d- Palmyrides Faults

The Palmyrides fold-thrust belt separates the Rutbah uplift to the south from the Aleppo plateau to the north (Figure 2). Many sets of weak events were located in the northern region of Palmyrides. They could be related to the faults: Ithrya, Okeirbat, Bishri, Jhar, Khnefice, Palmyra and others. The scattered seismicity, observed in the current study (Figure 5) and the previous ones (Brew *et al.*, 2001) suggests that the Palmyrides region is still tectonically active. The Bishri fault system is probably a dextral strike-slip fault similar to the Jhar Fault. A magnitude $M_s=4.9$ earthquake, occurred on November, 20, 1994, was located near this fault, at a depth of 15 km, northeast of the mapped extent of the fault (Litak *et al.*, 1997). The focal mechanism for a 1970 earthquake ($M_b=4.8$) was determined to be dextral but the initial solution for the 1994 event suggests sinistral motion (Dziewonski *et al.*, 1995). Moreover, field evidence and kinematic models (Chaimov *et al.*, 1993) are consistent with dextral motion along the Bishri fault. The strongest event that occurred on December 18, 1994 ($M_c=5.0$), was located at the northeast of Bishri mountain. It shows dextral motion on the NE striking nodal plane (Figure 7). It could be related to the Bishri fault. A cluster of 20 events was observed during the study period in the northeast of the mapped extent of the fault (Figure 5). In the southwestern regions of Palmyrides, a cluster of 10 events is located at the northeast extension of Damascus fault near Khan-Arus site (Figure 8), where this fault interacts with a secondary north-west fault. This cluster might indicate the activity of the secondary fault rather than Damascus fault. Another cluster of 7 events are aligned in the NE direction and located along the Zobeideh fault (Figure 7), where a doublet has been detected. This doublet has a sinistral mechanism trending NE, in agreement with the Zobeideh fault (Abdul-Wahed *et al.*, 2011).

A cluster of 773 micro-events has been observed in Jayrud region. In this cluster, 660 micro-events have a magnitude of less than 2.0, they could be therefore induced by quarrying exploitation. The early works in Jayrud's quarries started in 2001 (Al-Tahan, 2014) but the induced seismicity didn't appear until 2003 (Figure 10), where the exploitation has been extended to many quarries. Similarly, a cluster of 304 micro-events has been observed around Hyssia city and could be induced by quarry exploitations.

e- Northern extension of Serghaya fault

Serghaya fault can be traced approximately 125 km through the Anti-Lebanon mountains to the eastern edge of the Bekaa valley (Figures 8 and 10). Field evidence for Quaternary strike-slip faulting along the Serghaya fault is observed as far north as the village of Aarsal, near the edge of the Bekaa valley (Gómez *et al.*, 2003). The trace of the Serghaya fault is obscure towards to the north of this point. Its northern termination suggests a possible linkage with the Jhar fault as an oblique ramp or tear fault that bounds the Palmyrides fold belt (Gómez *et al.*, 2006). Walley (1998) suggested that the Serghaya fault continues also in NNE into Syria, past Al Qusayr and SE of Homs, before splaying into the Jhar and Bishri faults within the Palmyrides fold belt. However, geological mapping in the vicinity of Al Qusayr has revealed no evidence of any such fault, and none has been noted during recent fieldwork in this area (Bridgland *et al.*, 2003). In this study, a cluster of 60 events, aligned to the NE until the village of Alqaa near the Syrian-Lebanese border, was observed around the village of Aarsal (Figure 8). This cluster could be associated to the northern part of Serghaya fault. The seismic activity in northern Lebanon, shown on the epicenters map, is concentrated on the northern part of Serghaya fault, which supports the finding of Walley (1998). Other cluster of 5 events was observed around Al Qusayr and makes probable the suggestion of linkage of the Serghaya fault with the Jhar and Bishri faults.

f- Aleppo plateau

The Aleppo plateau forms a central more stable region of low relief (Figure 2). The eastern and western margins of the plateau are poorly defined, especially near the DSFS, where several N-S to NE-SW strike-slip and normal faults splay out from the master fault. The Aleppo plateau is regarded as a platform with minor amounts of deformation compared to the Palmyrides (Zanchi *et al.*, 2002). It is cut by some NE-SW faults which are probable zones




Figure 10. Detailed epicenters map of the seismic activity observed in Damascus region during the period 1995-2012.

of strike-slip (Al-Saad *et al.*, 1992). The recently known earthquake in this plateau occurred in 1987 with magnitude of 4.8 (Ahmad, 2013). The seismic activity in the western margin along the DSFS, discussed above, could be considered as relatively high (Figure 11). However, it is very low in the eastern margin. In northern Aleppo, a cluster of 102 events is aligned in the NE direction and could be associated with NE Ikhrin fault; another cluster of 26 events could be related to Manbeg fault. In the west of Aleppo plateau, Az-zawiye mountain overlooks

the Ghab basin. It is bounded by NE-striking sinistral strike-slip fault. In this mountain, a cluster of 100 events was located.

g- Golan plateau faults

The Golan, situated at the SW of Syria, is an elevated basalt-covered plateau. To the north, it is bounded by the Hermon mountain anticline. About 180 events are located in Golan plateau and could be associated to DSFS. Some of them formed multiplets with events aligned in




Figure 11. Detailed epicenters map of northern Aleppo.

NS direction. Two multiplets have a sinistral mechanism trending NS in agreement with DSFS (Abdul-Wahed *et al.*, 2011). Another multiplet has a normal mechanism trending NE and could be related to local trans-tensional faults that extend along the margins of DSFS.

h- Coastal range

The Syrian coastal range, occupies most of the Syrian onshore area at the west of the DSFS and Ghab basin. The stratigraphic relationships indicate that the uplift of the coastal range is a part of the extensive Syrian arc deformation (compressional folding along the eastern Mediterranean coast), documented in the Levant (Walley, 2001; Salamon *et al.*, 2003, 2007; Yolsal *et al.*, 2007; Carton *et al.* 2009; Eyal, 2011). The coastal range has clearly been affected by the propagation of the DSFS and the formation of the Ghab basin, resulting in the steep eastern limb (Brew *et al.*, 2001), that created the extremely steep scarp on the eastern face of the coastal range alongside the Ghab basin. A cluster of about 46 events is located in northern of Tartous city. Another cluster of 81 events is located in south and south east of this city.

3- Occasional-type seismic activity

Some significant events can be classified as individual such as the earthquake of Palmyra 1996 and the events of Rutbah uplift. Those events seem to be apparently scattered during the study period, but they might be in reality linked with other events that occurred before or after this period. The study period of about 17 years, from 1995 to 2012, is very short in comparison with the estimated long return period for whole Syria of 200-350 years (Ambraseys and Barazangi, 1989). It is therefore not possible to conclude that some faults, such as the northern segments of DSFS, are inactive, depending on the recent instrumental seismicity qualified as short term observations. The high magnitude of some events of occasional-type, such as the earthquake of Palmyra, 1996, could be interpreted by accumulation the seismic energy during a long period of quiescence. One can conclude that the long period of quiescence along the northern segments of DSFS might accumulate the seismic energy and could at any time produce a big earthquake especially for the reasons mentioned above (geomorphic evidence for Pliocene, recent tectonic activity on the fault together with historical seismicity and GPS measurements). The scattered seismicity in the Palmyrides fold belt suggests that this region is still tectonically active.

Discussion

During the period of studyd 1995-2012, about 5000 local events have been recorded by the SNSN (Bulletins of SNSN, 1995-2012 2009 on the reference list!!!!). In total, nearly 3800 earthquakes were located with a location uncertainty of less than 10 km. Figure 4 shows that about 50% of the entire events are reliable locations with an average location uncertainty less than 5 km in horizontal and vertical directions. The average RMS for the whole dataset is about 0.5 sec. Most of the recorded events are located within the network and have relative small location uncertainty values. Therefore, locations of these events can be considered reliable for the following interpretation of the present work. The epicentral map (Figure 5) presents the geographical distribution of the seismic activity. The study of this local instrumental data has illuminated the general appearance of the recent seismicity in Syria and spotlighted some important observations:

- The little amount of events, where the annual average of recorded events is about 292, which could be considered very low as compared with other parts of DSFS, such as Yammounah or Aqaba Gulf, indicates that Syria is actually passing through a relative quiescence period. For example, 2414 events were recorded in Aqaba Gulf in 1995 (Jordan Seismological Observatory, 1995) and 948 events in 1997 (Jordan Seismological Observatory, 1997). Furthermore, the instrumental seismicity of the eastern Mediterranean region shows that the southern part of DSFS is seismically the most active (Hofstetter *et al.*, 2007; Imprescia, 2010; Imprescia *et al.*, 2012). The most active area since the early 1980s was the Gulf of Aqaba, where thousands of small cluster earthquakes occurred in different regions of the Gulf of Aqaba during several months to few years (Baer *et al.*, 1999; Klinger *et al.*, 1999).
- The epicentral map has shown that the main instrumental seismicity with many moderate earthquakes ($M_c > 4$) is apparently concentrated on the Syrian borders: along the EAFS extending on the northwest borders and along the central segments of DSFS in Lebanon such as Yammouneh and Roum. The EAFS, including Latakia-Killis faults system, could be characterized by the highest instrumental seismicity in Syria.
- The seismic activity inside Syria has produced events of low magnitude (Fig.

6), which consequently indicates that it is actually passing within a relative quiescence period as compared with other parts of DSFS (Meghraoui *et al.*, 2003). Most of the recorded events are qualified as weak, where their average magnitude is approximately 2.0. Fig. 6 shows the frequency of the recorded events according to their magnitude, which varies between 1.0 and 5.6. Only 49 events have exceeded the magnitude of 3.0 during the period 1995-2003 (Abdul-Wahed and Al-Tahan, 2010). In the present study, only 470 events, which make 9% of the total 5000 events, have exceeded the magnitude of 3.0 during the period 1995-2012. The fitting Gutenberg and Richter relation to the observed frequency-magnitude distribution has shown a magnitude completeness down to $M_{\text{completeness}} = 2.5$ (Figure 6). This indicates that not all micro-earthquakes ($M_c < 2.5$) have been detected by the SNSN. Some events could be too small to be recorded on a sufficiently large number of stations to trigger the recording system in the SNSN, to initiate the location procedure, and thus the report of the event. The network operators decided that some micro-earthquakes below a certain threshold ($M_c < 2.5$) are not of interest. The constants of the empirical Gutenberg and Richter relationship, a - and b - values, were evaluated using the LSM method ($\log N = 5.43 - 0.91M$) and MLM method ($\log N = 5.2 - 0.73M$).

- According to the epicentral map, the recent instrumental seismicity inside Syria is characterized by many clusters of weak events ($M < 4$) such as in southwest of Aleppo plateau. Some clusters were observed as swarm indicating the behavior of some faulting zones such as Serghaya fault, southwest Palmyrides faults and Bassit region faults (Figure 2).

- The most important event was the earthquake of Palmyra which happened on December 24, 1996, with a magnitude of about 5.6 (Tan and Taymaz, 2003; Alchalbi, 2004). This event is located in Palmyrides fold belt that separates two relatively stable zones, Rutbah uplift, in the southeastern of Syria from Aleppo plateau in the northwestern of Syria. The earthquake of Palmyra has a sinistral mechanism trending NE (Figure 7), and can be related to Palmyra fault (Abdul-Wahed, 2012).

Conclusion

The instrumental seismicity of Syria during the period 1995-2012, has been characterized

and documented in the present paper. In total, nearly 3800 micro-earthquakes were located with a location uncertainty of less than 10 km. Therefore, locations of these events can be considered reliable for the following interpretation. The correlation between the instrumental seismicity and the seismotectonic features was performed through the analysis of spatial distributions of seismic events and some focal mechanisms, where most epicenters appear to correlate with the regional strike-slip faults. The epicentral map has shown that the main instrumental seismicity with many moderate earthquakes ($M > 4$) is apparently concentrated on the Syrian northwest borders along the EAFS, including Latakia-Killis faults system, where the highest instrumental seismicity in Syria has been observed.

This research allowed observing many types of seismic activity as follows: Swarm-type, Cluster-type, and Occasional-type. Many observed swarms have been correlated through the spatial distributions of the hypocenters with some active faults, such as Serghaya fault and the southwest of Palmyrides. The seismicity inside Syria was characterized by many clusters of weak events ($M < 4$) such as the clusters on Latakia-Killis faults, Rouj basin faults, and northern Aleppo plateau faults. However, some significant events could be classified as individual, such as the earthquake of Palmyra, 1996, regarding the shortness of the study period. The geographical distribution of epicenters of a cluster could be related to an active fault. This correlation could be confirmed by the focal mechanism of some strongest events. As a consequence, many faults can be considered active in Syria. The observed swarms could give an insight on the seismic behavior of some active faults such as Serghaya fault, where many events striking in a relatively short period of time are followed by a long period of quiescence. The occasional-type of seismic activity could be interpreted as a very long behavior of some active faults.

The instrumental seismicity of Syria could be classified in general as low magnitude events. Most of the recorded events are qualified as weak, where their average magnitude is approximately 2.0. In the present study, only 470 events, which make 9% of the total 5000 events, have exceeded their magnitude of 3.0 during the period 1995-2012. The fitting Gutenberg and Richter relation to the observed frequency-magnitude distribution has shown a magnitude completeness down to 2.5. This indicates that not all micro-earthquakes ($M_c < 2.5$) have been detected by the SNSN. Regarding the low magnitude level of the recorded events, we can deduce that the seismic activity in Syria is

actually passing by a relative quiescence. Such result may therefore be taken to support the finding of previous studies indicating that the seismicity in Syria tends to fade out with time, in line with the decrease of major seismic intensity which has occurred during the last millennium. This view, however, is inconsistent with both the historical records of large, devastating earthquakes, and the field observations that attest to the seismic activity and seismogenic potential of the DSFS. The discrepancy between measured rate of small-magnitude earthquakes ($M < 4$) from instrumental records and large earthquake rates from paleoseismic records, as also observed on other faults, makes the fault's behavior unpredictable. This discrepancy can be explained if we consider that the instrumental record lacks both moderate-to-large events and their aftershocks. Since the beginning of instrumental seismicity, the DSFS experienced only one large earthquake ($M=7.2$) that occurred in the Gulf of Aqaba on November 22, 1995. The characterization of the recent seismic activity in Syria is not an easy task, due to the long return periods of large earthquakes, estimated in previous studies to be about 200-350 years, and to the shortness of instrumental seismicity, about 17 years. It is therefore difficult to discover all the active faults in the country. The long period of quiescence along some faults, such as the Ghab fault, accumulates probably the seismic energy and could produce a big earthquake at any time. The nonperiodic behavior of the DSFS over the millennial timescale makes it more difficult to meaningfully predict the probability for a large earthquake soon. Therefore, it is imperative to prepare for a large earthquake on the DSFS, which will occur sooner or later. The actual results, documented in this research, highlight the need and necessity of more detailed researches for a longest possible period of time.

Acknowledgements

The authors wish to express their thanks to Prof. Ibrahim Othman Director General of the Atomic Energy Commission of Syria for his constant support. They also wish to thank the General Establishment of Geology and Mineral Resources for providing some important information and data. The anonymous reviewers are cordially thanked for their constructive remarks and suggestions that considerably improve the quality of this paper.

References

Abdul-Wahed M. K., 2012, The focal mechanism of recent largest seismic events in Syria. The 7th Gulf Seismic forum, Saudi Geological

Survey; 22-25 January 2012, Jeddah, Saudi Arabia.

Abdul-Wahed M. K., Asfahani J., Al-Tahan I., 2011, A combined methodology of multiplet and composite focal mechanism techniques for the identification of the seismological active zones in Syria. *Acta Geophysica*, 59, 967-992, DOI:10.2478/s11600-011-0024-2.

Abdul-Wahed M. K. & Al-Tahan I., 2010, Preliminary outlining of the seismological active zones in Syria. *Annals of geophysics*, 53, 1-9.

Abdul-Wahed M. K., 2007, The multiplet analysis and some of its applications. The 4th Gulf Seismic forum, Kuwait Institute for Scientific Research; Kuwait, 24-27 March, 2007.

Abdul-Wahed M. K., Al Heib M., Senfaute G., 2006, Mining-induced seismicity: Seismic measurement using multiplet approach and numerical modeling. *International Journal of Coal Geology*, 66, 137-147.

Ahmad R. A., 2013, Seismic Hazard Assessment of Syria. *Journal of Seismology and Earthquake Engineering*, 15, No. 1.

Alchalbi A., 2004, Workshop on "Earthquake Hazard Assessment in Syria and Lebanon", September 7-9, 2004, Damascus – Syria.

Alchalbi A., Daoud M., Gómez F., McClusky S., Reilinger R., Romeyeh M. A., Alsouod A., Yassminh R., Ballani B., Darawcheh R., 2010, Crustal deformation in northwestern Arabia from GPS measurements in Syria: slow slip rate along the northern Dead Sea Fault. *Geophys. J. Int.*, 180, 125-135.

Al-saad D., Sawaf T., Gebran A., Barazangi M., Best J. & Chaimov T., 1992, Crustal structure of central Syria: The intracontinental Palmyride mountain belt. *Tectonophysics*, 207, 345-358.

Al-Tahan I., 2014, Personal communication. General Establishment of Geology and Mineral Resources.

Al-Tarazi E., Abu-Rajab J., Gómez F., Cochran W., Jaafar R., Ferry M., 2011, GPS measurements of near-field deformation along the southern Dead Sea Fault System. *Geochem. Geophys. Geosyst.* 12, Q12021, <http://dx.doi.org/10.1029/2011GC003736>.

Ambraseys N. N. & Barazangi M., 1989, The 1759 Earthquake in the Bekaa Valley: implications for earthquake hazard assessment in the

- Eastern Mediterranean region. *J. geophys. Res.*, 94, 4007–4013.
- Ambraseys N. N. and C. P. Melville, 1995, Historical evidence of faulting in Eastern Anatolia and Northern Syria. *Annali Di Geofisica*, 38, 337.
- Asfahani J. and Abdul-wahed M. K., 2013, Evaluation of Earthquake Activity Along the Serghaya Fault, Syria, from Instrumental Seismic Data, *Acta Geophysica*, 61, 37–59, DOI: 10.2478/s11600-012-0059-z.
- Baer G., Sandwell D., Williams S., Bock Y., Shamir G., 1999, Coseismic deformation associated with the November 1995, Mw=7.1 Nuweiba earthquake, Gulf of Elat (Aqaba), detected by synthetic aperture radar interferometry. *J. Geophys. Res.*, 104, 25221– 25232.
- Barazangi M., Seber, D., Chaimov, T., Best, J. and Sawaf, T., 1993, Tectonic evolution of the northern Arabian plate in western Syria. In: E. Boschi, E. Mantovani and A. Morelli (Eds.), Recent Evolution and Seismicity of the Mediterranean Region. Kluwer Academic Publishers, p.117–140.
- Brew G., Barazangi M., Al-Maleh A. K., Sawaf T., 2001, Tectonic and geologic evolution of Syria, *GeoArabia*, 6, 573–616.
- Brew G., J. Lupa, M. Barazangi, T. Sawaf, A. Al-Imam and T. Zaza, 2001b, Structure and tectonic development of the Ghab Basin and in Dead Sea Fault System Syria. *Journal of the Geological Society London*, 158, 665–674.
- Bridgland D. R., Philip G., Westaway R., White M., 2003, A long Quaternary terrace sequence in the valley of the River Orontes, near Homs, Syria. *Current Science*, 84, 1080–1089.
- Bulletin of the Syrian National Seismological Network (SNSN), 1995–2009, National Earthquake Center, Ministry of Petroleum and mineral resources, Syrian Arab Republic.
- Butler R. W. H., Spencer S. & Griffiths H. M., 1997, Trans current fault activity on the Dead Sea Transform in Lebanon and its implications for plate tectonics and seismic hazard. *J. geol. Soc. Lond.*, 154, 757–760.
- Carton H., S. C. Singh, P. Tapponnier, A. Elias, A. Briaies, A. Sursock, R. Jomaa, G. C. P. King, M. Dae`ron, E. Jacques, and L. Barrier, 2009, Seismic evidence for Neogene and active shortening offshore of Lebanon (Shalimar cruise). *J. Geophys. Res.*, 114, B07407, doi:10.1029/2007JB005391.
- Chaimov T.A., M. Barazangi, D. Al-saad, T. Sawaf and M. Khaddour, 1993, Seismic fabric and 3-D structure of the southwestern intracontinental Palmyride fold belt, Syria. *Am. Assoc. Petro Geol. Bull.*, 77, 2032–2047.
- Dakkak R., Daoud M., Mreish M., Hade G., 2005, The Syrian National Seismological Network (SNSN): Monitoring a major continental transform fault. *Seismological Research Letters*, 76, 437–445.
- Darawcheh R., Sbeinati M. R., Margottini C. & Paolini S., 2000, The 9 July 551 AD Beirut earthquake, eastern Mediterranean region. *J. Earthquake Eng.*, 4, 403–414.
- Dziewonski A. M. , G. Ekström, M. P. Salganik, 1995, Centroid-moment tensor solutions for October–December 1994. *Physics of the Earth and Planetary Interiors*, 91, 187–201.
- El-Isa Z. H., 2012, Seismicity and seismotectonics of the Gulf of Aqaba region. *Arab J Geosci.*, doi:10.1007/s 12517-012-0604-8.
- El-Isa Z. H., S. McKnight & D. Eaton, 2015, Historical seismicity of the Jordan Dead Sea Transform region and seismotectonic implications. *Arab J Geosci.*, DOI 10.1007/s12517-014-1483-y.
- Eyal Y., 2011, The Syrian Arc Fold system: Age and rate of folding. *Geophysical Research*, 13, EGU2011-7401, 2011, EGU General Assembly 2011.
- Garfunkel Z., 2011, The long- and short-term lateral slip and seismicity along the Dead Sea Transform: an interim evaluation. *Isr J Earth Sci*, 58, 217–235.
- Girdler R.W., 1990, The Dead Sea transform fault system. *Tectonophysics*, 180, 1–13.
- Gómez F., Meghraoui M., Darkal A. N., Sbeinati R., Darawcheh R., Tabet C., Khawlie M., Charabe M., Khair K. & Barazangi M., 2001, Coseismic displacements along the Serghaya fault : an active branch of the Dead Sea Fault System in Syria and Lebanon. *J. Geol. Soc. Lond.*, 158, 405–408.
- Gómez F., Meghraoui M., Darkal A.N., Hijazi F., Mouty M., Suleiman Y., Sbeinati R., Darawcheh R., Al-ghazzi R., Barazangi M., 2003, Holocene faulting and earthquake

- recurrence along the Serghaya branch of the Dead Sea fault system in Syria and Lebanon. *Geophys. J. Int.*, 153, 658–674.
- Gómez F., Khawlie M., Tabet C., Darkal A. N., Khair K., Baranzangi M., 2006, Late Cenozoic uplift along the northern Dead Sea transform in Lebanon and Syria. *Earth Planet. Sci. Lett.*, 241, 913–931.
- Guidoboni E., Bernardini F., Comastri A., 2004, The 1138–1139 and 1156–1159 destructive seismic crises in Syria, south-eastern Turkey and northern Lebanon. *Journal of Seismology*, 8, 105–127.
- Guidoboni E. & Comastri A., 2005, Catalogue of earthquakes and tsunamis in the Mediterranean area from the 11th to the 15th century. *Istituto Nazionale di Geofisica e Vulcanologia*, 1037 pp.
- Hofstetter R., Klinger Y., Amrat A. Q., Rivera L. and Dorbath L., 2007, Stress tensor and focal mechanisms along the Dead Sea fault and related structural elements based on seismological data. *Tectonophysics*, 429, 165–181, doi:10.1016/j.tecto.2006.03.010.
- Imprescia P., 2010, Meccanismi focali di terremoti come strumento per la definizione della sismotettonica dell'area mediterranea. *Ph.D. thesis*, University of Catania.
- Imprescia P., Palano Mimmo, Gresta Stefano, 2012, Mapping crustal strain and stress fields across the Dead Sea Fault System by GPS observations and focal plane solutions. Conference paper: GNGTS November 2012, Volume: Extended Abstract.
- Jordan Seismological Observatory, 1995, Earthquakes in Jordan and adjacent areas, Nat. Res. Auth., *Jordan Seis. Obs., Bul.*, No. 27.
- Jordan Seismological Observatory, 1997, Earthquakes in Jordan and adjacent areas, Nat. Res. Auth., *Jordan Seis. Obs., Bul.*, No. 29.
- Khair K., Karakaisis G. F. and Papadimitriou E. E., 2000, Seismic zonation of the Dead Sea Transform Fault area. *Annali di Geofisica*, 43, 61–79.
- Klinger Y., Rivera L., Haessler H., Maurin J.C., 1999, Active faulting in the Gulf of Aqaba: new knowledge from the Mw7.3 earthquake of 22 November 1995. *Bull. Seismol. Soc. Am.*, 89, 1025–1036.
- Klinger Y., M. Le Beon and M. Al-Qaryouti, 2015, 5000 yr of paleoseismicity along the southern Dead Sea fault. *Geophys. J. Int.*, 202, 313–327, doi: 10.1093/gji/ggv134.
- Lee W. H. K., 1990, Hypo71 Program. vol. 1. IASPI Software Library.
- Litak R. K., M. Barazangi, W. Beauchamp, D. Seber, G. Brew, T. Sawaf and W. Al-Youssef, 1997, Mesozoic-Cenozoic Evolution of the Intraplate Euphrates Fault System, Syria: Implications for Regional Tectonics. *Journal of the Geological Society*, 154, 653–666.
- Mahmoud Y., F. Masson, M. Meghraoui, Z. Cakir, A. Alchalbi, H. Yavasoglu, O. Yönlü, M. Daoud, S. Ergintav, S. Inan, 2013, Kinematic study at the junction of the East Anatolian fault and the Dead Sea fault from GPS measurements. *Journal of Geodynamics*, 67, 30–39.
- Masson F, Y. Hamiel, A. Agnon, Y. Klinger, A. Deprez, 2015, Variable behavior of the Dead Sea Fault along the southern Arava segment from GPS measurements. *C. R. Geoscience*, <http://dx.doi.org/10.1016/j.crte.2014.11.001>.
- Meghraoui M., Gómez F., Sbeinati R., van derWoerd J., Mouty M., Darkal A. N., Radwan Y., Layyous I., Al Najjar H., Darawcheh R., Hijaz F., Al-Ghazzi R., Barazangi M., 2003, Evidence for 830 years of seismic quiescence from palaeoseismology, archeoseismology and historical seismicity along the Dead Sea fault in Syria. *Earth Planet. Sci. Lett.*, 210, 35–52.
- Mohamad R., Darkal A. N., Seber D., Sandvol E., Gómez F., Barazangi M., 2000, Remote earthquake triggering along the Dead Sea fault in Syria, following the 1995 Gulf of Aqaba earthquake ($M_s = 7.3$). *Seismol. Res. Lett.*, 71, 47–52.
- Nemer T. & Meghraoui M., 2006, Evidence of coseismic ruptures along the Roum fault (Lebanon): a possible source for the AD 1837 earthquake. *Journal of Structural Geology*, 28, 1483–1495.
- Nemer T, Meghraoui M, Khair K., 2008, The Rachaya-Serghaya fault system (Lebanon): evidence of coseismic ruptures, and the AD 1759 earthquake sequence. *J. Geophys. Res.*, 113:B05312
- Palano M., Imprescia P., and Gresta S., 2013, Current stress and strain-rate fields across the Dead Sea Fault System: Constraints from seismological data and GPS observations. *EPSL*, 369, 305–316.

- Preliminary Seismological Bulletin, 1995, SNSN: Syrian National Seismological Network, General Establishment of Geology and Mineral Resources, Syria.
- Reilinger R., and S. McClusky, 2011, Nubia Arabia Eurasia plate motions and the dynamics of Mediterranean and Middle East tectonics. *Geophys. J. Int.*, 186, 971–979.
- Salamon A., Hofstetter A., Garfunkel Z., Ron H., 1996, Seismicity of the eastern Mediterranean region: perspective from the Sinai subplate. *Tectonophysics*, 263, 293–305.
- Salamon A., Hofstetter A., Garfunkel Z., Ron H., 2003, Seismotectonics of the Sinai subplate–the eastern Mediterranean region. *Geophys. J. Int.*, 155, 149–173.
- Salamon A., T. Rockwell, S. N. Ward, E. Guidoboni, and A. Comastri, 2007, Tsunami hazard evaluation of the eastern mediterranean: historical analysis and selected modeling. *Bul. Seis. Soc. Am.*, 97, 1–9, doi: 10.1785/0120060147.
- Salamon A., 2009, Patterns of seismic sequences in the Levant interpretation of historical seismicity. *J. Seismology*, 14, 339–367.
- Saleh M., Hamiel Y., Ziv A., Bock Y., Fang P., Wdowinski A., 2012, Crustal deformation along the Dead Sea Transform and the Carmel Fault inferred from 12 years of GPS measurements. *J. Geophys. Res.*, 117, B08410, <http://dx.doi.org/10.1029/2012JB009241>.
- Sbeinati M. R. & Darawcheh R, 1997, Earthquakes in and around Syria during 1996, Bulletin No. 6, AECS-GRSS 233
- Sbeinati M. R. & Darawcheh R, 1998, Earthquakes in and around Syria during 1997, Bulletin No. 7, AECS-GRSS 279.
- SEISAN: THE EARTHQUAKE ANALYSIS SOFTWARE, Version 10.3, Jens Havskov and Lars Ottemöller, Department of Earth Science, University of Bergen, Allégaten 41, 5007 Bergen, Norway, 2015.
- Tan O. and Taymaz T., 2003, Source Parameters of November 20, 1994 and December 24, 1996 Palmyra (Syria) Earthquakes, and Analogy to the Dead Sea Transform Fault Zone (DSTFZ). International Workshop on the North Anatolian, East Anatolian and Dead Sea Fault Systems: Recent Progress in Tectonics and Paleoseismology and Field Training Course in Paleoseismology, P.118, Middle East Technical University (METU), 31 August – 12 September 2003, Ankara, Turkey.
- Uski M., Tiira, Korja, Elo S., 2006, The 2003 earthquake swarm in Anjalankoski, south-eastern Finland. *Tectonophysics*, 422, 55–69.
- Walley C. D., 1988, A braided strike-slip model for the northern continuation of the Dead Sea fault and its implications for Levantine tectonics. *Tectonophysics*, 145, 63–72.
- Walley C. D., 1998, Some outstanding issues in the geology of Lebanon and their importance in the tectonic evolution of the Levantine region. *Tectonophysics*, 298, 37–62.
- Walley C. D., 2001, The Lebanon passive margin and the evolution of the Levantine Neotethys. In, W. Cavazza, A.H.F. Robertson and P. Ziegler (Eds.), PeriTethyan rift/wrench basins and margins. Memoires du Muséum National d'Histoire Naturelle, Paris, PeriTethys Memoir 6.
- Wechsler N., T. K. Rockwell, Y. Klinger, P. Štěpančíková, M. Kanari, S. Marco, and A. Agnon, 2014, A Paleoseismic Record of Earthquakes for the Dead Sea Transform Fault between the First and Seventh Centuries C. E.: Nonperiodic Behavior of a Plate Boundary Fault. *Bul. Seis. Soc. Am.*, 104, No. 3, doi: 10.1785/0120130304.
- Yolsal S., Taymaz T., Yalçiner A. C., 2007, Understanding Tsunamis, Potential Source Regions and Tsunami Prone Mechanisms in the Eastern Mediterranean. The Geological Society of London, Special Publications Book, Vol: 291, ISBN: 978-1-86239-239-7, pp. 201–230.
- Zanchi a., G. B. Crosta , A. N. Darkal, 2002, Paleostress analyses in NW Syria: constraints on the Cenozoic evolution of the northwestern margin of the Arabian plate. *Tectonophysics*, 357, 255–278.
- Zollo A. & Bernard P., 1991, Fault mechanisms from near-source data : joint inversion of S polarization and P polarities. *Geophys. J. Int.*, 104, 441–451.