



What we know about the breeding biology of the endemic Gray-barred Wren (*Campylorhynchus megalopterus*)

Lo que sabemos sobre la biología reproductiva del endémico Matraca barrada (*Campylorhynchus megalopterus*)

Vladislav Marcuk^{1*}

¹ Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany.

* Autor de correspondencia: vladislav.marcuk@googlemail.com

Abstract

The Gray-barred Wren (*Campylorhynchus megalopterus*) remains among the most poorly known representatives of the family Troglodytidae endemic to Mexico. Little information exists on the species' ecology and breeding biology, with only two known egg records in museum collections. Here I synthesized data from the literature and obtained 14 photographic records from citizen science databases to enhance our understanding of breeding behavior, egg morphology, and nesting habits of the Gray-barred Wren. I also critically assessed two egg collections in museum records. Only two, decades-old studies provided data on Gray-barred Wren breeding parameters, which I compared with information on 7 endemic wren species. According to the review, the Gray-barred Wren has an extended 10-month breeding season from November to August. The globular nest, cream-white egg interspersed with magenta to brownish spots, and evidence of cooperative breeding, were similar to closely-related species. However, the Gray-barred Wren had a smaller clutch size (1 – 3 eggs) and larger eggs (26.87 × 19.43 mm) than other endemic wren species. Notably, my critical reassessment of an egg set in the Western Foundation of Vertebrate Zoology collection that is attributed to the Gray-barred Wren, determined discrepancies in egg morphology, collection location, and associated nest description, suggesting a misidentification and incorrect species allocation. These findings highlight the importance of citizen science and museum records in uncovering essential biological data for poorly-known taxa.

Keywords: Citizen science, eBird, egg collections, Gray-barred Wren, nesting, Troglodytidae.

Resumen

La Matraca Barrada (*Campylorhynchus megalopterus*) sigue siendo uno de los miembros menos conocidos de la familia Troglodytidae endémico de México. Existe poca información sobre la ecología y la biología reproductiva de la especie, y solo se conocen dos registros de huevos en colecciones de museos. Aquí, sintetice datos de la literatura, y obtuve 14 registros fotográficos de bases de datos de ciencia ciudadana para mejorar nuestra comprensión sobre el comportamiento reproductivo, la morfología de los huevos y los

INFORMACIÓN SOBRE EL ARTÍCULO

Recibido:

26 octubre 2024

Aceptado:

10 diciembre 2025

Editor Asociado:

Adolfo Gerardo Navarro Sigüenza

Contribución de cada uno de los autores:

El autor realizó completamente el estudio y escritura del manuscrito.

Cómo citar este documento:

Marcuk V. 2025. What we know about the breeding biology of the endemic Gray-barred Wren (*Campylorhynchus megalopterus*). Huitzil Revista Mexicana de Ornitología 26(2):e692. <https://doi.org/10.28947/hrmo.2025.26.2.822>



Esta obra está bajo una licencia de Creative Commons Reconocimiento No Comercial-Sin Obra Derivada 4.0 Internacional.

hábitos de anidación de *C. megalopterus*. También evalué críticamente dos colecciones de huevos en registros de museos. Sólo dos estudios realizados hace décadas proporcionaron datos sobre los parámetros reproductivos de *C. megalopterus*, que comparé con información de 7 especies endémicas. De acuerdo con la revisión, *C. megalopterus* tiene una temporada reproductiva extendida de 10 meses, desde noviembre a agosto. El nido globular, el huevo blanco crema intercalada con manchas magenta a marrones, y la evidencia de reproducción cooperativa, eran similares a especies estrechamente relacionadas. Sin embargo, *C. megalopterus* tuvo un tamaño de puesta más pequeña (1–3 huevos) y huevos más grandes (26.87×19.43 mm) que otras especies endémicas de Troglodytidae. Notablemente, mi reevaluación crítica de un conjunto de huevos en la colección del Western Foundation of Vertebrate Zoology, atribuida a *C. megalopterus*, determinó discrepancias en la morfología de los huevos, el sitio de colecta, y la descripción del nido, sugiriendo una identificación errónea y una asignación incorrecta de la especie. Estos hallazgos destacan la importancia de la ciencia ciudadana y los registros de museos para descubrir datos biológicos esenciales de especies poco conocidas.

Palabras clave: anidación, ciencia ciudadana, colecciones de huevos, eBird, Matraca Barrada, Troglodytidae.

Introduction

Mexico is home to 37 different species of wren, 12 of which are endemic to the country, including nine terrestrial and three insular species (Clements et al. 2025). The Gray-barred Wren (*Campylorhynchus megalopterus*) is predominantly found in the highlands of the Mexican central plateau, and two subspecies are currently recognised (Márquez-Valdelamar 1998, Brewer 2001, Kroodsma and Brewer 2020). The nominate form (*C. m. megalopterus*) is found in the Sierra de Colima (Jalisco), extending from northern Michoacán to western Puebla. The smaller race *C. m. nelsoni* is confined to eastern Puebla, the Sierra Madre in Veracruz, and northern Oaxaca (Márquez-Valdelamar 1998, Brewer 2001, Kroodsma and Brewer 2020). This distinctive wren inhabits a broad spectrum of montane forest types at altitudes between 1800–3230 m asl, including epiphytic forests, tall fir forests, and humid Pine-Oak forests. It occasionally penetrates degraded forest areas, indicating that the species tolerates disturbance to

some extent (Márquez-Valdelamar 1998, Brewer 2001, Kroodsma and Brewer 2020).

The ecology of the species remains poorly studied. The Gray-barred Wren is often seen in groups of 6–12 individuals (Brewer 2001), in the medium vertical strata, probing for different invertebrates and rarely descending to the understorey (Márquez-Valdelamar 1998; Brewer 2001). Analysis of stomach contents revealed that the diet consists mainly of insects (larvae, Formicidae, Orthoptera, Lepidoptera, Hemiptera, Diptera and Coleoptera), arachnids and plant parts (seeds and small fruits; Márquez-Valdelamar 1998).

Information about the species' breeding biology is scarce (Márquez-Valdelamar 1998, Kroodsma and Brewer, 2020), with only a few nests documented in the literature (Schaldach 1963, Rowley 1984, Márquez-Valdelamar 1998, Brewer 2001). The main objective of this study was to summarise information on breeding biology of the species by reviewing existing literature, museum records, and citizen science databases. Additionally, the alleged breeding record of the species, based on a single egg set collected by Fred B. Nyc., Jr. and preserved in the Western Foundation of Vertebrate Zoology (WVZ), was critically reviewed by comparing the egg set with an actual egg of the species preserved in the Colección Nacional de Aves (CNAV), Instituto de Biología, Universidad Nacional Autónoma de México. Finally, the results are compared with those of other congeneric species native to Mexico.

Methods

Literature review

For the literature review, I consulted 14 published records to retrieve breeding data of closely-related species (12 publications), and compared this to data for the Gray-barred Wren (4 publications). I used key words like nest/egg descriptions, breeding biology, wrens Mexico, filtering my results by peer-reviewed publications (journals), grey literature (thesis) and books. For the Gray-barred Wren, the following information was retrieved from literature data: nest and eggs (Márquez-Valdelamar 1998); breeding records, seasonality (Schaldach 1963, Rowley 1984, Brewer 2001).

For the comparative literature review, I considered breeding records published in the literature for seven congeneric species with distribution confined to Mexico. Clarion Wren *Troglodytes tanneri*: breeding season, nest, eggs

and clutch size (Sosa-López et al. 2012), eggs and clutch size (Brattstrom and Howell 1956). Boucard's Wren *Campylorhynchus jocosus*: breeding season, nest and eggs (Márquez-Valdelamar 1998, Brewer 2001). Spotted Wren *C. gularis*: breeding seasonality, nest, eggs, breeding behavior and clutch size (Calderón-Torres 2011, Ceja-Madrigal and Salgado-Ortiz 2013, Ceja-Madrigal 2014). Yucatan Wren *C. yucatanicus*: breeding seasonality, nest, clutch size, incubation and nestling period (Vargas-Soriano et al. 2010). Sumichrast's Wren *Hylorchilus sumichrasti*: breeding seasonality, nest description, breeding behavior, clutch size (Pérez-Villafan 1997), eggs, clutch size (Bangs and Peters 1927). Happy Wren *Pheugopedius felix*: breeding seasonality, nest morphology, clutch size (Grant 1966, Brewer 2001), breeding seasonality, eggs, nest (Márquez-Valdelamar 1998). Sinaloa Wren *Thryophilus sinaloa*: breeding seasonality, nest morphology, clutch size (Grant 1966), breeding seasonality, eggs, nest (Márquez-Valdelamar 1998), egg size (Schönwetter 1979). No information was found for the Socorro Wren (*Troglodytes sissonii*)

and Nava's Wren (*Hylorchilus navai*). Detailed studies are also lacking for the Veracruz Wren *Campylorhynchus rufinucha* (Bradley et al. 2024) and the Cozumel Wren *Troglodytes beani* (Juárez and Johnson 2024).

Citizen science databases

I consulted citizen science records on the eBird database (14 photographic records in total), and only considered records as valid which allowed clear species identification, i.e. photos showing Gray-barred Wren adults. I obtained 13 records of photographs of Gray-barred Wren adults either at the nest site or close to it (Figure 1). On nine occasions, adults were observed collecting nest material, and on seven occasions, adults were pictured in close proximity to a nest. On two occasions, adults were observed feeding chicks. While the behavioural notes provided alongside the photographic records could not be confirmed directly, none of the observations appeared unusual, and are therefore presented here as reported by the original authors.




Figura 2. Gray-barred Wren (*Campylorhynchus megalopterus*) distribution map (after Birdlife Distribution Map Data 2022.02) with the associated breeding records based on citizen science data and the egg set at the Instituto de Biología, Universidad Nacional Autónoma de México (CNAV catalogue number 004917, label number IBUNAM H000141). Numbers correspond to ID records in Table 1. Illustration by Vladislav Marcuk.

Museum records

I also consulted the only known breeding records existing in museum collections. These consisted of a single egg preserved in the Colección Nacional de Aves (CNAV) of the Instituto de Biología, Universidad Nacional Autónoma de México (Hernán Vázquez Miranda pers. comm.), and one alleged clutch preserved in the Western Foundation of Vertebrate Zoology (WVZ) egg collection (René Corado pers. comm.). However, it is possible that other records in regional or undisclosed collections were overlooked here.

Results

Literature review

Breeding season

According to the literature review, the breeding season of the Gray-barred Wren seems to last from November to August. Nest construction was reported between November and December, and nests containing eggs and chicks were discovered in August (Schaldach 1963, Márquez-Valdelamar 1998). Adults at the nest were observed between December and May, and feeding of the chicks was documented between April and August (Márquez-Valdelamar 1998). Females in breeding condition were collected in April (Brewer 2001), and several nests were located on 8 June and 27 June (Brewer 2001). Juveniles were collected in June, and a nest containing a single chick and egg was found in mid-August in the State of Mexico (Schaldach 1963). The literature also reports an instance in which an adult female was observed feeding three fully grown juveniles on 26 April (Rowley 1984).

Nest

The nest is described as globular or dome-shaped, with a lateral entrance composed of grass, pine needles, twigs, roots and mosses (Márquez-Valdelamar 1998). Literature records indicate that the nest is placed 1.5-18.2 m above the ground in conifers, oaks or mandrillo (*Arbustus xalapensis*) trees (Márquez-Valdelamar 1998). One nest is recorded by Márquez-Valdelamar (1998) as having the following dimensions: total length 71 mm; external diameter 82 × 72 mm; internal diameter 76 × 61 mm; depth 48.5 mm. Another nest was reported to have a diameter of 36 cm and to be open at the top (Edwards and Martin 1955).

Eggs and clutch size

The average egg size reported by Márquez-Valdelamar (1998) is 25.2 × 19.9 mm ($n = NA$), with no further information given. Little is known of clutch size. One nest found in August contained a single egg and chick (Schaldach 1963), while Rowley (1984) observed a female feeding three fully sized juveniles, indicating that the clutch size at least consists of 2-3 eggs.

Parental behavior

The species was reported as a cooperative breeder by Márquez-Valdelamar (1998). Observations of ten individuals in Quiroga (Michoacán) indicated that at least two individuals assisted with nest construction, while two others rearranged the nest (Edwards and Martin 1955). Further evidence of a cooperative breeding system was obtained from Rowley (1984), who observed three individuals participating in constructing of a nest, which was used as a roosting site. Otherwise, nothing else is known about the breeding biology, and the incubation and nestling periods remain unknown.

Citizen science database

In addition to the available literature, citizen science data expanded the breeding season from 9 November to 16 September (Table 1). Several photographic records supplemented the literature by providing information on breeding season based on months when photographs were taken (Table 1). Photographs also documented various stages of nesting, including nest construction, the presence of adults at the nest, and feeding of nestlings (Table 1).

Nest construction and nest records

Birds collecting nest material were photographed between 9 November and 20 May (Table 1), with most individuals observed collecting material between November and March. However, as the individuals were not continuously monitored, or at least the details are not documented in the supplementary information, it remains unclear whether a nest was present when the photographs were taken.

Eight photographic records documented birds near a nest (Table 1). Four records documented two birds at or near a nest, frequently reported as nest building (Table 1), although the photographs do not show material in their beaks. On three occasions, one bird was documented at a nest, and on 28 February

Table 1. Summary of museum and citizen science (eBird database) breeding data for the Gray-barred Wren (*Campylorhynchus megalopterus*) with the associated breeding behavior, date, locality, catalogue number or link and collector/photographer. ID = locality number in Figure 1 map; Ind = Individual; Ad = Adult; MID = Misidentified.

ID	Description of the behavior	Date	Locality	Coordinates	Links/Catalogue number	Collector/Photographer
1	single egg from a nest	18 Aug 1988	Temascaltepec, Valle de Bravo	19.155, -100.02666	Catalogue number CNAV 004917	Moisés Sanchez
2	1 Ind collecting nest material	9 Nov 2016	Volcán Nevado de Colima, San Gabriel, Jalisco	19.6137224, -103.5677934	ML284437521	Tim Ludwick
3	2 Ind at nest, reported bringing material to the 30 cm wide nest for 1 hour	9 June 2018	San Jose de la Montaña, Huitzilac, Morelos	19.0129494, -99.2303127	ML103872041	Rafael Rodríguez Brito
4	1 Ind collecting moss, report 5 Ind doing this in tree	20 May 2019	El Temascal, Charo, Michoacán	19.6527321, -100.9545171	ML159960951	Paul Lewis
5	2 Ind near nest	22 Apr 2019	Camino La Cumbre, Ixtlán, Oaxaca	17.1771118, 96.6071078	ML201644531	Hervé Jacob
6	3 Ind: 2 Ind adding material to a nest and third begging for food, but was not fed	18 Aug 2019	San Jose de la Montaña, Huitzilac, Morelos,	19.012949, -99.2303127	ML173085591	Rafael Rodríguez Brito
7	1 Ad at 'communal' nest	16 Sep 2019	Camino La Cumbre, Ixtlán, Oaxaca	17.1771118, -96.6071078	ML177688261	Yve Morrell
8	1 Ad collecting nest material	13 Dec 2021	Pino Real, Charo, Michoacán	19.6485895, -101.0142338	ML398525781	Paul Lewis
9	1 Ad with feather-like material in beak at nest entrance	28 Feb 2022	Corral de Piedra, Ixtlán, Oaxaca	17.1851291, -96.6430978	ML420936141	Carlos Gonzalez
10	1 Ad at nest, reported as nest building	6 Mar 2024	Camino La Cumbre, Ixtlán, Oaxaca	17.1863, -96.61795	ML615732300	Manuel Grosselet
11	1 Ad collecting nest material	10 Mar 2024	Corral de Piedra, Ixtlán, Oaxaca	17.1851291, -96.6430978	ML615901796	Fátima Santana

ID	Description of the behavior	Date	Locality	Coordinates	Links/Catalogue number	Collector/ Photographer
12	2 Ad at nest, carrying food and feeding chicks	24 Mar 2024	Parque Hermenegildo Galeana Tenancingo, Estado de México	19.0215387, -99.6167114	ML616656440	C. Salgado-Miranda and E. Soriano-Vargas
13	1 Ad collecting nest material	10 Nov 2024	Camino La Cumbre-Este Ixtlán, Oaxaca	17.163157, -96.5948953	ML626948020 ; ML626678913	Serge Horellou; Manuel Grosselet
14	2 Ad at nest, reported nest building	17 Mar 2025	Camino Viejo a Coajomulco, Huitzilac	19.033135, -99.230224	ML632278345	Nancy Dunn
MID	Clutch of four eggs	12 May 1956	Abovaro road, Veracruz		WFVZ Catalogue number EN-145380	Fred B. Nyc Jr

2022 one bird was photographed at the nest entrance with feathery material in its beak (Table 1). At least one pair was recorded bringing nest material to a nest, estimated as 30 cm wide, and chasing away a Bronzed Cowbird (*Molothrus aeneus*) on 9 June (Table 1). At another nest, the presence of at least three individuals in close proximity was documented on 18 August, of which two were actively adding nest material and one was begging for food (Table 1). A recent record stated that a pair was observed

collecting nest material and feeding chicks on 24 March 2024 (Table 1). This suggests that eggs and chicks may be produced early in the season, or that multiple broods may be produced, as is known to occur in other *Campylorhynchus* taxa (Brewer 2001).



Figure 2. The only known egg of the Gray-barred Wren (*Campylorhynchus megalopterus*) collected by Moisés Sanchez in Temascaltepec on the 18 August 1983, preserved in the Colección Nacional de Aves of the Instituto de Biología, Universidad Nacional Autónoma de México (CNAV catalogue number 004917, IBUNAM label number H000141). Photo by Violeta Andrade.



Figure 3. Alleged clutch of the Gray-barred Wren (*Campylorhynchus megalopterus*) in the Western Foundation of Vertebrate Zoology (WFVZ catalogue number EN-145380), collected by Fred B. Nyc Jr on 12 May 1956, one-mile SE of the Aborado road, in Veracruz, Mexico. Photo by René Corado.

Table 2. Comparison of breeding parameters among endemic Troglodytidae taxa in Mexico, for which the nest and eggs are described. No information was found for the Socorro Wren (*Troglodytes sissonii*) and Nava's Wren (*Hylorchilus navai*). Suffix letters indicate literature sources for each species. a: This study, b: Márquez-Valdelamar 1998, c: Bangs and Peters 1927, d: Pérez-Villafan 1997, e: Brewer 2001, f: Grant 1966, g: Sosa-López et al. 2012, h: Ceja-Madrigal and Salgado-Ortiz 2013, i: Schönwetter 1979, j: Vargas-Soriano et al. 2010, k: Brattstrom and Howell 1956, l: Calderón-Torres 2011, m: Ceja-Madrigal 2014.

Species	Breeding season	Nest form	Clutch size	Egg color	Egg measurements (L x W mm)	Incubation (days)	Fledging (days)	Breeding system
Gray-barred Wren								
Campylorhynchus megalopterus ^{a,b}	Dec-Aug	globular	1 – 3	White to cream, white ground color with magenta to brownish spots	26.87 × 19.43 (n = 1) or 25.2 × 19.9 (n = NA)	NA	NA	Cooperative
Happy Wren <i>Pheugopedius felix</i> ^{b,f,i}	May-Aug	retort shaped (teardrop or flask shaped)	3 – 5	Bluish white	21.1 × 14 (n = NA) or 20.5 × 14 (18.6 – 21.8 x 13.9 – 14.2; 2.88 g estimate, n = 4)	NA	NA	Not cooperative
Sumichrast's Wren <i>Hylorchilus sumichrasti</i> ^{b,c,d}	Mar-Jul	cup-shaped in nest-cavity	3	White	23 – 23.4 × 17 (3.59 g estimated; n = 3) or 25 x 15	NA	NA	Not cooperative
Spotted Wren Campylorhynchus <i>gularis</i> ^{b,h}	Mar-Sep	globular	2 – 4 or 3-5?	Buff white, with red-brown spots, concentrated stronger towards the larger egg pole	20.8 × 16.2 (3.3 g; n = 4) or 23.0 ± 0.41 × 16.5 ± 0.30 (n = 5)	12 (n = 2)	NA	Instance of cooperative
Clarion Wren <i>Thryophilus tanneri</i> ^{b,g,k}	Oct-May	nest cavity	3 – 4	Pink to whitish ground colour with small brownish to pink spots concentrated at the larger end of the egg	17.7 × 14 or 19.8 × 14	NA	NA	Instance of cooperative
Boucard's Wren Campylorhynchus <i>jocosus</i> ^b	Apr-Jul	globular	3 – 4	White or pale pink densely covered with grey-brown or reddish-brown speckles, concentrated stronger on the larger egg pole	22.86 × 16.63 (n = 18)	NA	NA	Cooperative
Sinaloa Wren <i>Thryophilus sinaloa</i> ^{b,f,i}	May-Sep	retort shaped (teardrop or flask shaped)	4 – 5	Immaculate white	19 × 13.5; 20 × 14 (1.92 g estimated, n = 2)	NA	NA	Not cooperative

Species	Breeding season	Nest form	Clutch size	Egg color	Egg measurements (L x W mm)	Incubation (days)	Fledging (days)	Breeding system
Yucatan Wren <i>Campylorhynchus yucatanicus</i> ^j	May-Jul	ovoid globular	3 ± 1.5 (1 – 5; n = 99)	Buff white, less or more densely covered with brown speckles, stronger concentrated on the larger egg pole	NA	16.0 ± 1.0 (15 – 17; n = 47)	16.5 ± 1.9 (14 – 19; n = 51)	Cooperative

Museum records

One egg collected by Moisés Sanchez (CNAV catalogue number 004917) from an active nest in Temascaltepec on 18 August 1983 (Figure 2) measured 26.87 × 19.43 mm (0.02 mm precision digital caliper). The egg was elongated and ovate, and white with pinkish, magenta or light brown spots that were unevenly dispersed over its surface but concentrated more heavily towards the larger end (Figure 2).

A second presumed egg set (Figure 3), currently preserved at the Western Foundation of Vertebrate Zoology (WVFZ catalogue number EN-145380), was collected on 12 May 1956 by Fred B. Nyc Jr, at a location one mile south-east of the Abovaro road leading towards the city of Veracruz. According to notes on the specimen label, Fred B. Nyc Jr found several nests situated in cacti and small bushes, around 4–5 feet above the ground. He described the nests as being dome-shaped with a lateral entrance and composed predominantly of grass and weed stems. They were lined with fine grass, small feathers, and plant down. The clutch size consisted of four eggs, averaging 21.20 ± 0.62 × 15.42 ± 0.07 mm (range: 20.23–21.87 × 15.35–15.53 mm; n = 4; René Corado pers. comm.). The eggs were elongated and ovate with an immaculate white base colour, and have small to large (c. 3–4 mm) clusters or blotches of dark brown, grey or black colouration that are distributed rather irregularly along the egg surface and are more concentrated towards the large egg pole (Figure 3). The collection location, egg morphology, and nest description are inconsistent with reports for the species, therefore this egg set seems to have been wrongly allocated to the Gray-barred Wren.

Comparison with congenics

I obtained formal and qualitative descriptions of the nest and eggs for six of the nine congeneric endemic wren species (Table 2), with no literature records found for the Socorro Wren and Nava’s Wren, and little information for the Veracruz Wren. Information on nest and eggs of the Gray-barred Wren was obtained from only one Master’s thesis (Márquez-Valdelamar 1998), and the present study (Table 2). Four congeneric Mexican species in the genus *Campylorhynchus*, the Gray-barred Wren, Spotted Wren, Boucard’s Wren, and Yucatan Wren, showed striking similarities in nest and egg morphology and breeding system (Table 2). All four

species were recorded to build globular nests, have similar egg colors, and breed in a cooperative system (Table 2). Contrasting nest shapes were reported for congeneric species outside the genera: retort shaped (flask or teardrop) in the Happy and Sinaloa wrens; and cavity nests for the Clarion and Sumichrast's wrens (Table 2). Egg morphology seemed to show similarity in base color, but with more variability in the presence and absence of spot patterns (Table 2). However, the Gray-barred Wren differed from other species by having a smaller clutch size (1–3 eggs), and laying larger eggs (Table 2). Other congeneric species tended to lay 3 to 4 or 5 eggs, although the Spotted Wren may lay 2–4 or 3–5 eggs (Table 2). Nevertheless, records on clutch size of the Gray-barred Wren are far too limited to allow adequate comparability.

Discussion

Gray-barred Wren nesting and egg collections

Through a revision of published literature, citizen science database, and museum egg collections, I provide new information on the seasonality, nesting habits, egg morphology, and clutch size of a species whose breeding biology remains poorly understood. This short communication also highlights the importance of carefully evaluating the authenticity of egg and nest sets currently preserved in museum collections, paying particular attention to morphological traits, collection locality, and supplementary information on the specimen label. Investigative work can help confirm or refute species identification and prevent errors in modern literature.

As a result of this evaluation, I conclude there is a high likelihood that the egg set preserved and collected by Fred B. Nyc., Jr (WFVZ catalogue EN-145380) was wrongly allocated to the Gray-barred Wren. This is supported by inconsistencies in the collection location, egg morphology, and nest description. These contradict previous literature records for the species (Márquez-Valdelamar 1998), particularly with regard to nest site, and the vertical placement of the nest. The egg morphology further supports the inconsistencies in species identification, as the egg size and morphology of the WFVZ collection does not resemble the single egg preserved in the CNAV (catalogue number 004917, label number IBUNAM H000141), which is considerably smaller and differs in secondary coloration.

Furthermore, following the chronology of other egg sets that were collected by Fred B. Nyc., Jr on 14 May 1956 in Villahermosa, Tabasco, Mexico (Barred Parakeet *Bolborhynchus lineola lineola*, WFVZ catalogue number EN-145379, 18.02166666666667, -92.92666666666667), and on 22 May 1956 in Gomez Farias, Tamaulipas, Mexico (Plain Chachalaca *Ortalis vetula mcallii*, WFVZ catalogue number EN-145393, 29.36, -107.72833333333333), there is no doubt that the eggs were in fact collected in Veracruz. The records indicate that Fred B. Nyc., Jr was following the coastal roads and associated sites, not descending to any proximate areas within the native range of the Gray-barred Wren, supporting the proposal that the eggs were wrongly assigned to the Gray-barred Wren. Considering the magnitude of divergence in egg, nest morphology and distance to the actual distribution trajectory of the Gray-barred Wren, it is very likely that the eggs in question belong to the Veracruz Wren, which shows strong resemblance in egg morphology and distribution (Bradley et al. 2024). Very little is known about the breeding biology of the Veracruz Wren (Bradley et al. 2024), therefore the WFVZ egg set may provide the first quantitative data on egg and clutch size for the species.

Comparison with congenics

My literature review highlights that the breeding biology of most endemic species remains poorly studied, except for the closely-related Yucatan Wren, where breeding biology was studied in more detail (Vargas-Soriano et al. 2010, Armiger 2014). For the internationally near-threatened Sumichrast's Wren (Pérez-Villafan 1997) and threatened Clarion Wren (Sosa-López et al. 2012) a more comprehensive description of the nest, eggs and breeding behavior were published only recently. For the Happy Wren and Sinaloa Wren the vast majority of information originated from a single study published decades ago (Grant 1966) and a thesis study (Márquez-Valdelamar 1998).

My comparison of breeding parameters among congeneric species indicated strong similarity among members of the genera *Campylorhynchus*, where traits are likely to show less variability due to taxonomic affinities (Barker 2007). Cooperative breeding was recorded in all *Campylorhynchus* taxa, which aligns with the cooperative breeding recorded in many congeneric species in the Neotropics (Rabenold 1990). In general, nest

structure showed intra-generic consistency among species of the same genus *Campylorhynchus*. Although the composition of nests varies, they tend to contain predominantly grass, plant fibres, twigs, leaves or feathers (Márquez-Valdelamar 1998). Evidently, nest and egg morphology can differ to some extent due to differences in environmental and biotic factors.

Breeding seasons showed high variability among species and therefore comparison is difficult. However, the Gray-barred Wren seemed to have an extended breeding season spanning over ten months. A possible cause for such a prolonged breeding season could be due to double brooding, either after successfully fledging a first brood or replacing a clutch after failure, however this remains highly speculative. There is no evidence of double broods in the Yucatan Wren, whereas multiple broods have been recorded in other congeneric species (Brewer 2001). Some wren species also construct nests as dormitories that are used for roosting (Brewer 2001, Bradley et al. 2024), therefore further research is required to determine whether double broods are present in the Gray-barred Wren.

Notable differences were observed in the egg size and clutch size, although considering the small sample size this needs to be interpreted with caution. Further field data collection is required to understand the true range of clutch size. The egg size, however, is substantially larger in comparison to the Spotted Wren and Boucard's Wren. The Gray-barred Wren has overall a larger body size (length: 16.6-21.8 cm, mass: 32.92 g; $n = 55$) in contrast to the Spotted Wren (length: 17.9 cm, mass: 27.08 g; $n = 27$) and Boucard's Wren (length: 18.5 cm, mass: 28.82 g, $n = 9$; Márquez-Valdelamar 1998). These differences in body size could explain the variation in egg size. In fact, if clutch size of the Gray-barred Wren is smaller, it is possible that larger eggs are produced in an overall trade-off.

Facilitating research into the biology of Mexico's endemic wren species remains highly desirable, particularly for species of conservation interest and those that are poorly known. While undocumented museum records, such as the one discussed in detail here, can contribute significantly to our understanding of the breeding biology of lesser-known species, field studies are still essential for revealing different aspects of their biology. Nevertheless, this study highlights the importance of using citizen science and museum specimens

to provide fundamental information about poorly known species.

Acknowledgements

I would like to thank René Corado (WFVZ) and Patricia Escalante, Hernán Vázquez Miranda, and Violeta Andrade (CNAV) from the Instituto de Biología, Universidad Nacional Autónoma de México for providing information about the respective egg sets, including pictures of the egg sets and allowing me to use them for the publication. In addition, I would like to thank Alexandre Courtiol, the reviewers and editors that significantly improved the quality of the manuscript.

Literature cited

- Armiger JW. 2014. Breeding biology of a tropical cooperative passerine: The Yucatan Wren (*Campylorhynchus yucatanicus*). M.Sc. Thesis. Villanova University, Pennsylvania, US.
- Bangs O, Peters JL. 1927. Birds from the rain forest region of Vera Cruz. Bulletin of the Museum of Comparative Zoology 67: 471-487.
- Barker FK. 2007. Avifaunal interchange across the Panamanian isthmus: insights from *Campylorhynchus* wrens. Biological Journal of the Linnean Society 90:687-702. <https://doi.org/10.1111/j.1095-8312.2007.00758.x>
- Brewer D. 2001. Wrens, dippers, and thrashers. Yale University Press, New Haven.
- Brattstrom BH, Howell TR. 1956. The birds of the Revilla Gigedo islands, México. Condor 58: 107-120. <https://doi.org/10.2307/1364977>
- Bradley DW, Mennill DJ, van Dort J 2024. Veracruz Wren (*Campylorhynchus rufinucha*), version 1.0. In Keeney BF, Schulenberg TS (eds). Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.runwre2.01>
- Calderón-Torres L. 2011. Éxito de anidación de una comunidad de aves residentes en el Área Natural Protegida "Cerro de Punhuato", Morelia, Michoacán, México. B.Sc. thesis. Facultad de Biología. Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Mexico.
- Ceja-Madrigal A. 2014. Contribución al conocimiento de la biología reproductiva

- y estructura social de la matraca encinera (*Campylorhynchus gularis*) en el Área Natural Protegida “Cerro Punhuato”, Morelia, Michoacán, México. B.Sc. thesis. Facultad de Biología. Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Mexico.
- Ceja-Madrigal A, Salgado-Ortiz J. 2013. Descripción del nido y huevos de la matraca serrana (*Campylorhynchus gularis*). Huitzil 14:113-116. <https://doi.org/10.28947/hrmo.2013.14.2.202>
- Clements JF, Rasmussen PC, Schulenberg TS, Iliff MJ, Gerbracht JA, Lepage D, Spencer A, Billerman SM, Sullivan BL, Smith M, Wood CL. 2025. The eBird/Clements checklist of Birds of the World. version 2025. Cornell Lab of Ornithology, Ithaca, NY, USA. <https://www.birds.cornell.edu/clementschecklist/download/> Consulted 31 December 2025.
- Edwards EP, Martin PS. 1955. Further notes on birds of the Lake Patzcuaro region, Mexico. The Auk 72:174-178. <https://doi.org/10.2307/4081422>
- Grant PR. 1966. The coexistence of two wren species of the genus *Thryothorus*. Wilson Bulletin 78:266-278.
- Juárez R, Johnson LS. 2024. Cozumel Wren (*Troglodytes beani*). version 1.0. In Keeney BK, Billerman SM (eds). Birds of the World. Cornell Lab of Ornithology. Ithaca, NY, USA. <https://doi.org/10.2173/bow.houwre3.01>
- Kroodsma DE, Brewer D 2020. Gray-barred Wren (*Campylorhynchus megalopterus*), version 1.0. In del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (eds) Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.grbwre1.01>
- Márquez-Valdelamar LM. 1998. Monografía de las especies mexicanas de la familia Troglodytidae (Aves). M.Sc. Thesis. Facultad de Ciencias, Universidad Nacional Autónoma de México. México, DF
- Pérez-Villafan AM. 1997. Contribución al conocimiento de la historia de vida de *Hylorchilus sumichrasti* (Aves: Troglodytidae) en el norte del estado de Oaxaca. B.Sc. Thesis, Universidad Nacional Autónoma de México, Campus Iztacala, México, DF.
- Rowley JS. 1984. Breeding records of land birds in Oaxaca, Mexico. Proceedings of the Western Foundation of Vertebrate Zoology 2:76–221.
- Rabenold KN. 1990. *Campylorhynchus* wrens: the ecology of delayed dispersal and cooperation in the Venezuelan savanna. Pp 159-196. In Stacey PB, Koenig WD (eds). Cooperative breeding in birds. Cambridge University Press, Cambridge, United Kingdom.
- Schaldach WJ. 1963. The avifauna of Colima and adjacent Jalisco, Mexico. Proceedings of the Western Foundation of Vertebrate Zoology 1:1-100.
- Schönwetter M. 1979. Handbuch der Oologie. Band II (Passeriformes 1). Akademie Verlag, Berlin, Germany.
- Sosa-López JR, Martínez-Gómez JE, Hernández-Yáñez H, Mennill DJ. 2012. Description of the nest, eggs, and parental behavior of the Clarion Wren (*Troglodytes tanneri*), a vulnerable island-endemic songbird. Ornitología Neotropical 23:291-298.
- Vargas-Soriano J, Ortiz JS, Segura GE. 2010. Breeding phenology and nesting success of the Yucatan Wren in the Yucatan Peninsula, Mexico. The Wilson Journal of Ornithology 122:439-446. <https://doi.org/10.1676/08-164.1>