



**Diversidad y análisis germinativo de especies arbóreas y
arbustivas de interés ecológico en un área incendiada**
**Diversity and germination analysis of tree and shrub species of
ecological interest in a burned area**

Homero Gárate-Escamilla¹, Aldo Tovar-Cárdenas¹, Enrique Jurado-Ybarra¹,
Mauricio Cotera-Correa¹, Eduardo Alanís-Rodríguez^{1*}, Maritza Gutiérrez-
Gutiérrez¹

Fecha de recepción/Reception date: 8 de marzo de 2023.

Fecha de aceptación/Acceptance date: 22 de julio de 2023.

¹Universidad Autónoma de Nuevo León, Facultad de Ciencias Forestales. México.

*Autor de correspondencia: eduardoforestal@gmail.com

*Corresponding author; e-mail: eduardoforestal@gmail.com

Abstract

The vegetation of an ecosystem can be degraded by forest fires; which are caused by human or natural actions and generally reduce the diversity and composition of species, as well as their natural regeneration. This work was carried out in a temperate forest degraded by forest fires in the *El Tejocote* Natural Protected Area, *Santiago* municipality, *Nuevo León* State. The objectives were to calculate the Importance Value Index, analyze species diversity and determine the percentage of germination. Sampling was carried out in 10 random plots of 1 000 m². The Importance Value Index (*IVI*), diversity indexes (*Margalef*, *Shannon-Weiner* and *Simpson*) and the germination percentages were determined for each species. *Pinus teocote* had the highest *IVI* (45.5 %), it was also the most dominant, covering 69 % of the sampled area. *Quercus laeta* had 25 % *IVI* and was the most abundant of all species (39 %). *Agave scabra* and *Arbutus xalapensis* had *IVIs* of 7.9 % and 6.9 %, respectively; both were classified as shrubs. A low diversity was obtained in all the sites sampled; the average values were: *Margalef* (0.74), *Shannon-Weiner* (0.46), and *Simpson* (0.77). The highest germination percentages were presented by *Pinus pseudostrobus* (97.40 %) and *Agave scabra* (84 %). The difference in the germination capacity of taxa could change the diversity and species composition of the *El Tejocote* community after its degradation by fire.

Key words: Natural Protected Area, temperate forest, degradation, germination, forest fire, Importance Value Index.

Resumen

La vegetación de un ecosistema puede ser degradada por los incendios forestales; estos se originan por acciones humanas o de manera natural y por lo general reducen la diversidad y composición de especies, así como su regeneración natural. El presente trabajo se desarrolló en un bosque templado degradado por incendios forestales en el Área Natural Protegida *El Tejocote*, municipio *Santiago*, *Nuevo León*. Los objetivos fueron calcular el Índice de Valor de Importancia, analizar la diversidad de especies y determinar el porcentaje de germinación. El muestreo se realizó en 10 parcelas al azar de 1 000 m². El Índice de Valor de Importancia (*IVI*), índices de diversidad (*Margalef*, *Shannon-Weiner* y *Simpson*) y los porcentajes de germinación se determinaron para cada especie. *Pinus teocote* tuvo el mayor *IVI* (45.5 %), también fue la más dominante al cubrir 69 % de la superficie muestreada. *Quercus laeta* tuvo 25 % de *IVI* y fue la más abundante de todas las especies (39 %). *Agave scabra* y *Arbutus xalapensis* tuvieron *IVI* de 7.9 % y 6.9 %, respectivamente; ambas se clasificaron como arbustivas. Se obtuvo una baja diversidad en todos los sitios muestreados, y los valores

promedio fueron: *Margalef* (0.74), *Shannon-Weiner* (0.46) y *Simpson* (0.77). Los porcentajes de germinación más altos los presentaron *Pinus pseudostrobus* (97.40 %) y *Agave scabra* (84 %). La diferencia en la capacidad germinativa de los taxones podría cambiar la diversidad y composición de las especies de la comunidad El Tejocote después de su degradación por incendio.

Palabras clave: Área Natural Protegida, bosque templado, degradación, germinación, incendio, Índice de Valor de Importancia.

Temperate forests provide a variety of ecosystem services, and possess high biodiversity (Prieto-Amparán *et al.*, 2019). Forest degradation negatively affects their functional and structural characteristics (Vásquez-Grandón *et al.*, 2018), it refers to processes that result in long-term and drastic changes in the ecosystems (Thompson *et al.*, 2013) and is generally associated with a reduction in biomass, changes in the structure and composition of species, as well as in their natural regeneration (Chazdon *et al.*, 2008).

Forest fires are considered one of the main natural or anthropogenic factors causing the degradation of temperate forests (Pausas and Keeley, 2009). In Mexico, the fire regime is characterized by its moderate intensity, occurring at intervals of two to ten years (Santini *et al.*, 2019). Some temperate forest (TF) species can benefit from low to moderate fires, because clearing the land stimulates seed release and promotes natural regeneration; however, when fires are severe, the seed viability declines (Rodríguez-Trejo and Fulé, 2003).

Multiple areas of the *Cumbres de Monterrey* National Park (CMNP) have been degraded due to forest fires that have occurred over time (Cantú *et al.*, 2013). The level of species diversity and germination capacity after a fire of the *El Tejocote Laguna de Sánchez* community, located in the central zone of the CMNP, is unknown. For this reason, the following objectives were proposed: (1) To describe the ecological importance and diversity of tree and shrub species in a burned area, and (2) To determine the percentage of germination of tree and shrub species surviving after a fire.

The burned site has a surface area of 10 ha, and the geographical coordinates of the center of the area are 25°19'54" N and 100°15'09" W, at an average altitude of 1 905 masl. The average rainfall there is 640 mm, and the average temperature is 13 °C (Cantú *et al.*, 2013).

A ten percent of the study area was sampled one year after the fire occurred. Ten circular sampling units (SU) of 1 000 m² were established, randomly located at a minimum distance of 20 m between sites (Silva-García *et al.*, 2021). Within each SU, all shrub and tree species were recorded in a census and identified; total height, crown diameter and basimetric diameter at 10 cm were measured for each species. The crown diameters were measured with a model TP20ME Truper® longimeter, the basimetric area with a model 283d Forestry Suppliers® diametric tape, and the total height, with a model B084G8TRG9 Nikon® laser rangefinder.

In order to determine the relative ecological value of the species, the relative frequency (*RF*; Equation 1), relative abundance (*RA*; Equation 2), relative dominance (*RD*; Equation 3) and Importance Value Index were calculated (*IVI*; Equation 4). The *IVI* value was divided by three to standardize an interval of 0 to 100, as the maximum value is 300 when the *RF*, *RA*, and *RD* are added together (Manzanilla *et al.*, 2020). The equations for their calculation were as follows:

$$RF = \left(\frac{n}{N} \right) \times 100 \quad (1)$$

Where:

RF = Relative frequency of the species

n = Number of the SU where the species is present

N = Total number of SUs

$$RA = \left(\frac{n}{N} \right) \times 100 \quad (2)$$

Where:

RA = Relative abundance per species

n = Number of individuals of the species

N = Overall number of all the species

$$RD = \left(\frac{i}{N} \right) \times 100 \quad (3)$$

Where:

RD = Relative dominance of the species

i = Crown area of the species

N = Overall crown area of all the species

$$IVI = \frac{\sum_{i=1}^n (RA_i, RD_i, RF_i)}{3} \quad (4)$$

Where:

IVI = Importance Value Index

RA_i = Relative abundance per species

RD_i = Relative dominance of the species

RF_i = Relative frequency of the species

The indexes to measure species diversity were: (1) Margalef (D_{Mg} ; Equation 5), whose values above 5 indicate high diversity, while values below 2 suggest low diversity (Margalef, 1972), (2) Simpson's complement (λ ; Equation 6), the range of whose values fluctuates between 0 and 1, high diversity is associated with values close to 1, and a value close to 0 indicates a site with low diversity (Soler *et al.*, 2012; Salmerón *et al.*, 2017), and (3) Shannon-Weiner (H' ; Equation 7), whose values range from 0.5 to 5, values above 3 indicate high diversity, while values below 2 denote the opposite (Somarriba, 1999). The equations used to calculate the diversity indexes were as follows (Moreno, 2001):

$$D_{MG} = \frac{S-1}{n \ln N} \quad (5)$$

Where:

D_{Mg} = Margalef diversity Index

S = Number of species present

$n \ln$ = Natural logarithm

N = Total number of individual

$$\lambda = \frac{1}{\sum p_i^2}$$

$$\text{Simpson's complement} = 1 - \sum p_i^2 \quad (6)$$

Where:

λ = Simpson Index

p_i = Abundance according to species i

$$H' = - \sum_{i=1}^s p_i \times \ln(p_i)$$

$$p_i = n_i / N \quad (7)$$

Where:

H' = Shannon-Wiener Index

P_i = Proportional abundance of the species

S = Total number of the species

\ln = Natural logarithm

n_i = Number of individuals of species i

N = Total number of individuals

An analysis of the percentage of germinated seeds was carried out; the experiment consisted of the following: (1) Collecting seeds from living individuals after the fire, (2) Identifying seeds with no apparent damage, (3) Performing buoyancy tests to detect empty seeds, (4) Applying a pre-germinative process depending on the type of seed: (i) *Pinus* L. and *Quercus* L. were immersed in water for 24 hours, (ii) *Juniperus* L. seeds were placed in 100 % sulfuric acid for 15 minutes, while (iii) *Agave* L., *Arbutus* L. and *Rhus* L. underwent no treatment (Antonio *et al.*, 2020), and (5) Sowing 144 seeds of each of the seven identified species in a 288-cavity model ch288 GLL® semi-rigid plastic container.

A completely randomized design distributed in four replications (36 seeds per replication) was used; the response variable was germination, and seven treatments were considered for each of the tree and shrub species. For the germination data, a Shapiro-Wilk normality test was performed (González-Estrada y Cosmes, 2019), and no normality was found ($p \leq 0.00097$). Therefore, a Kruskal-Wallis test was performed to determine the presence of differences between treatments (McKight y Najab, 2010); the comparison between the means of the different species was carried out by means of a Mann-Whitney U test (McKight y Najab, 2010).

The taxa present in the study area were *Pinus teocote* Schltdl. & Cham. (twisted leaf or Aztec pine), *Pinus pseudostrobus* Lindl. (smooth-bark Mexican pine), *Juniperus flaccida* Schltdl. (Mexican juniper), *Quercus laeta* Liebm. (white oak), *Rhus virens* Lindh. ex A. Gray (evergreen sumac), *Arbutus xalapensis* Kunth (Texas madrone), and *Agave scabra* Ortega (rough agave or maguey).

The white oak was the most abundant species (Table 1), which is related to the behavior of other species of the same genus (*Quercus*) that survived after the severe fires in *Chipinque* Ecological Park (Alanís-Rodríguez *et al.*, 2011). The most dominant tree species was the Aztec pine (Table 2); this agrees with what has been reported in other temperate forests in the state of *Nuevo León* with fire damage, where *Pinus* is the most dominant tree over all other species (Alanís-Rodríguez *et al.*, 2011). The most frequent tree species were *Pinus teocote* and *Quercus laeta* (Table 1); this behavior is normal in this type of ecosystems because in other temperate forests of the *Sierra Madre Occidental* with different environmental conditions and floristic compositions, the frequency percentages for pines and oaks are similar to those obtained in the present research (Hernández-Salas *et al.*, 2013). Both species registered a higher *IVI* (Table 2); this relationship differs from the one reported in *Chipinque* Ecological Park after a fire, where *Quercus* had a higher *IVI* compared to the value of *Pinus* (Alanís-Rodríguez *et al.*, 2011). However, this coincides with what was reported by Méndez *et al.* (2014) in a fire-affected TF

in the *Sierra Madre de Guerrero*, where the *IVI* of *Pinus radiata* D. Don was high, while that of *Quercus glaucescens* Bonpl. was low.

Table 1. Ecological indicators of tree and shrub species.

Species	Abundance		Dominance		Frequency		IVI
	N ha ⁻¹	RA	m ² ha ⁻¹	RD	N/Site	RF	
<i>Pinus teocote</i> Schltld. & Cham.	280	32.18	5816.08	69.91	9	34.62	45.57
<i>Quercus laeta</i> Liebm.	340	39.08	956.37	11.49	7	26.92	25.83
<i>Agave scabra</i> Ortega	100	11.49	78.45	0.94	3	11.54	7.99
<i>Pinus pseudostrabus</i> Lindl.	50	5.75	875.71	10.53	2	7.69	7.99
<i>Arbutus xalapensis</i> Kunth	40	4.6	390.44	4.69	3	11.54	6.94
<i>Rhus virens</i> Lindh. ex A. Gray	40	4.6	32.77	0.39	1	3.85	2.95
<i>Juniperus flaccida</i> Schltld.	20	2.3	170.09	2.04	1	3.85	2.73
Total	870	100	8 319.92	100	26	100	100

RA = Relative abundance; N ha⁻¹ = Number of individuals per hectare; RD = Relative dominance; m² ha⁻¹ = Crown cover in square meters per hectare; RF = Relative frequency; N/Site = Number of sites where the species is present; IVI = Importance Value Index.

Table 2. Species diversity indexes.

Sampling unit	Margalef Index	Simpson's Complement	Shannon-Weiner Index
1	0.40	0.49	0.68
2	0.42	0.30	0.47
3	0.91	0.37	0.68
4	0.48	0.50	0.69
5	0.62	0.32	0.50
6	0.00	0.00	0.00
7	0.62	0.48	0.67

8	1.74	0.78	1.56
9	0.80	0.65	1.08
10	1.44	0.72	1.32
Average	0.74	0.46	0.77

One year after the fire, species diversity indices indicate low values in *El Tejocote* (Table 2), consistently with the findings of Méndez *et al.* (2014) and Méndez *et al.* (2018), who also cite low values for the Margalef and Shannon-Weiner indices in a burned temperate forest of the *Sierra de Guerrero*.

Significant differences were obtained between the germination of shrub and tree species ($H=17.42$, $H_c=17.62$, $p=0.0015$). Pine species had contrasting germination percentages (Table 3); similar values have been documented for *Pinus pseudostrobus* in *Michoacán* State (Sáenz *et al.*, 2011) and for *P. teocote* in Veracruz State (Ramírez, 2000). The low germination of *Quercus laeta* (Table 3), can be associated with the fact that *Quercus* seeds are recalcitrant and lose their viability quickly due to fast desiccation (Wawrzyniak *et al.*, 2022).

Table 3. Germination percentage of tree and shrub species.

Species	Common name	Germination percentage
<i>Pinus pseudostrobus</i> Lindl.	Smooth-bark Mexican pine	94.4±10.09a
<i>Agave scabra</i> Ortega	<i>Maguey</i>	84.00±7.96a
<i>Pinus teocote</i> Schltl. & Cham.	Aztec pine	31.9±3.49b
<i>Rhus virens</i> Lindh. ex A. Gray	Evergreen sumac	10.40±1.32c
<i>Quercus laeta</i> Liebm.	White oak	7.60±0.74c
<i>Arbutus xalapensis</i> Kunth	Texas madrone	0±0
<i>Juniperus flaccida</i> Schltl.	Mexican juniper	0±0

The letters indicate differences between treatments.

Agave scabra reached a high germination percentage (Table 3), which has been recorded in several *Agave* species (average germination of 91 %) under different environmental conditions (Campos *et al.*, 2020; Castillo *et al.*, 2022). The low germination of *Rhus virens* and the null germination of *Juniperus flaccida* and *Arbutus xalapensis* suggest a change in the scarification used: For *J. flaccida* it is proposed to increase the exposure time to sulfuric acid to 5 hours (Martínez-Pérez *et al.*, 2006), as well as cold stratification (Yucedag *et al.* 2021). A similar treatment is recommended for *R. virens* and *Arbutus xalapensis*, but with sulfuric acid exposure times of 30 minutes (Olmez *et al.*, 2007).

Acknowledgments

The authors are grateful to the *Consejo Nacional de Humanidades, Ciencia y Tecnología* for granting Aldo Tovar-Cárdenas a master's degree scholarship, and to the inhabitants of the community of *El Tejocote, Santiago* municipality, *Nuevo León*, for allowing this research.

Conflict of interest

The authors declare that they have no conflicts of interest.

Contribution by author

Homero Gárate-Escamilla: development of the manuscript and statistical analysis; Aldo Tovar-Cárdenas: interpretation of results; Enrique Jurado-Ybarra and Mauricio Cotera-Correa: review of the manuscript; Eduardo Alanís-Rodríguez and Maritza Gutiérrez-Gutiérrez: data analysis.

References

- Alanís-Rodríguez, E., J. Jiménez-Pérez, A. Valdecantos-Dema, M. Pando-Moreno, O. Aguirre-Calderón y E. J. Treviño-Garza. 2011. Caracterización de regeneración leñosa post-incendio de un ecosistema templado del Parque Ecológico Chipinque, México. *Revista Chapingo Serie Ciencias Forestales y del Ambiente* 17(1):31-39. https://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S2007-40182011000100004. (20 de mayo de 2023).
- Antonio B., A., C. V. Velarde N. y O. L. Rumayor R. 2020. Manual para el análisis físico y biológico de semillas forestales. Quintanilla Ediciones. Saltillo, Coah., México. 131 p. <https://sma.gob.mx/wp-content/uploads/2022/08/Manual-semillas-.pdf>. (22 de mayo de 2023).
- Campos, H., C. Trejo, C. B. Peña-Valdivia, R. García-Nava, F. V. Conde-Martínez and R. Cruz-Ortega. 2020. Water availability effects on germination, membrane stability and initial root growth of *Agave lechuguilla* and *A. salmiana*. *Flora* 268:151606. Doi: 10.1016/j.flora.2020.151606.
- Cantú A., C., S. Ortiz H. y M. Rovalo M. 2013. Consideraciones Finales. In: Cantú A., C., M. Rovalo M., J. Marmolejo M., S. Ortiz H. y F. Serriñá G. (Edits.). *Historia natural del Parque Nacional Cumbres de Monterrey, México*. Universidad Autónoma de Nuevo León y Comisión Nacional de Áreas Naturales Protegidas. Linares, NL, México. pp. 410-414.

- Castillo R., F., D. Castillo Q., J. E. Sáenz C., A. Rueda S. and J. T. Sáenz R. 2022. Effects of *Trichoderma* and *Bacillus* pre-treatments on the germination of *Agave victoriae-reginae* T. Moore seeds. *Revista Mexicana de Ciencias Forestales* 13(69):56-72. Doi: 10.29298/rmcf.v13i69.844.
- Chazdon, R. L. 2008. Beyond deforestation: restoring forests and ecosystem services on degraded lands. *Science* 320(5882):1458-1460. Doi: 10.1126/science.1155365.
- González-Estrada, E. and W. Cosmes. 2019. Shapiro–Wilk test for skew normal distributions based on data transformations. *Journal of Statistical Computation and Simulation* 89(17):3258-3272. Doi: 10.1080/00949655.2019.1658763.
- Hernández-Salas, J., Ó. A. Aguirre-Calderón, E. Alanís-Rodríguez, J. Jiménez-Pérez, ... y L. A. Domínguez-Pereda. 2013. Efecto del manejo forestal en la diversidad y composición arbórea de un bosque templado del noroeste de México. *Revista Chapingo Serie Ciencias Forestales y del Ambiente* 19(2):189-199. Doi: 10.5154/r.rchscfa.2012.08.052.
- Manzanilla Q., G. E., J. M. Mata B., E. J. Treviño G., Ó. A. Aguirre C., E. Alanís R. y J. I. Yerena Y. 2020. Diversidad, estructura y composición florística de bosques templados del sur de Nuevo León. *Revista Mexicana de Ciencias Forestales* 11(61):94-123. Doi: 10.29298/rmcf.v11i61.703.
- Margalef, R. 1972. Homage to Evelyn Hutchinson, or why there is an upper limit to diversity. *Connecticut Academy of Arts and Sciences* 44:211-235. <https://digital.csic.es/handle/10261/166281>. (26 de junio de 2022).
- Martínez-Pérez, G., A. Orozco-Segovia and C. Martorell. 2006. Effectiveness of some pre-germination treatments eight woody species of the Oaxacan Mixteca Alta with relevant features for restoration. *Boletín de la Sociedad Botánica de México* (79):9-20. Doi: 10.17129/botsci.1729.

- McKight, P. E. and J. Najab. 2010. Kruskal-Wallis test. In: Weiner, I. B. and W. E. Craighead (Edits). *The Corsini Encyclopedia of Psychology*. John Wiley & Sons, Inc. New York, NY, United States of America. pp. 1413-1422.
- Méndez O., C., C. A. Mora D., E. Alanís R., J. Jiménez P., ... y M. Á. Pequeño L. 2018. Fitodiversidad y estructura de un bosque de pino-encino en la Sierra Madre del Sur, México. *Revista Mexicana de Ciencias Forestales* 9(50):35-53. Doi: 10.29298/rmcf.v9i50.236.
- Méndez O., C., E. Alanís R., J. Jiménez P., Ó. A. Aguirre C. y E. J. Treviño G. 2014. Análisis de la regeneración postincendio en un bosque de pino-encino de la Sierra de Guerrero, México. *CIENCIA UANL* 17(69):63-70. <https://cienciauanl.uanl.mx/wp-content/uploads/2014/10/Art.-de-la-Sierra-Guerrero-a-colorts.pdf>. (3 de abril de 2023).
- Moreno, C. E. 2001. *Métodos para medir la biodiversidad*. Sociedad Entomológica Aragonesa (SEA). Zaragoza, Z, España. 84 p. https://www.researchgate.net/publication/304346666_Metodos_para_medir_la_biodiversidad. (12 de diciembre de 2022).
- Olmez, Z., A. Gokturk and F. Temel. 2007. Effects of some pretreatments on seed germination of nine different drought-tolerant shrubs. *Seed Science and Technology* 35(1):75-87. Doi: 10.15258/sst.2007.35.1.07.
- Pausas, J. G. and J. E. Keeley. 2009. A burning story: The role of fire in the history of life. *BioScience* 59(7):593-601. Doi: 10.1525/bio.2009.59.7.10.
- Prieto-Amparán, J. A., F. Villarreal-Guerrero, M. Martínez-Salvador, C. Manjarrez-Domínguez, G. Vázquez-Quintero and A. Pinedo-Alvarez. 2019. Spatial near future modeling of land use and land cover changes in the temperate forests of Mexico. *PeerJ* 7:e6617. Doi: 10.7717/peerj.6617.
- Ramírez G., E. O. 2000. Variación de semillas y plántulas de tres procedencias de *Pinus teocote* Schl. & Cham. Tesis de Maestría en Ecología Forestal. Instituto de Genética Forestal, Universidad Veracruzana. Xalapa, Ver., México. 78 p.

<https://www.uv.mx/iif/files/2014/10/Tesis-Elba-Pinus-teocote-MC.pdf>. (22 de marzo de 2022).

Rodríguez-Trejo, D. A. and P. Z. Fulé. 2003. Fire ecology of Mexican pines and a fire management proposal. *International Journal of Wildland Fire* 12(1):23-37. Doi: 10.1071/WF02040.

Sáenz R., J. T., H. J. Muñoz F. y A. Rueda S. 2011. Especies promisorias de clima templado para plantaciones forestales comerciales en Michoacán. INIFAP-Campo Experimental Uruapan. Uruapan, Mich., México. 64 p. <https://es.scribd.com/document/401504925/Especies-promisorias-de-clima-templado-para-plantaciones-forestales-comerciales-en-Michoacan-J-Trinidad-Saenz-Reyes-pdf>. (10 de enero de 2023).

Salmerón L., A., G. Geada L. y M. C. Fagilde E. 2017. Propuesta de un índice de diversidad funcional: Aplicación a un bosque semideciduo micrófilo de Cuba Oriental. *Bosque* 38(3):457-466. Doi: 10.4067/S0717-92002017000300003.

Santini, N. S., M. F. Adame, R. H. Nolan, Y. Miquelajauregui, ... and D. Eamus. 2019. Storage of organic carbon in the soils of Mexican temperate forests. *Forest Ecology and Management* 446:115-125. Doi: 10.1016/j.foreco.2019.05.029.

Silva-García, J. E., O. A. Aguirre-Calderón, E. Alanís-Rodríguez, E. Jurado-Ybarra, J. Jiménez-Pérez y B. Vargas-Larreta. 2021. Estructura y diversidad de especies arbóreas en un Bosque templado del Noroeste de México. *Polibotánica* (52):88-102. Doi: 10.18387/polibotanica.52.7.

Soler, P. E., J. L. Berroterán, J. L. Gil y R. A. Acosta. 2012. Índice valor de importancia, diversidad y similaridad florística de especies leñosas en tres ecosistemas de los llanos centrales de Venezuela. *Agronomía Tropical* 62(1-4):25-37. http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S0002-192X2012000100003. (16 de febrero de 2023).

Somarriba, E. 1999. Diversidad Shannon. *Agroforestería en las Américas* 6(23):72-74. <http://bco.catie.ac.cr:8087/portal-revistas/index.php/AGRO/article/view/707>. (28 de marzo de 2022).

Thompson, I. D., M. R. Guariguata, K. Okabe, C. Bahamondez, ... and C. Sabogal. 2013. An operational framework for defining and monitoring forest degradation. *Ecology and Society* 18(2):20. <https://www.jstor.org/stable/26269330>. (28 de marzo de 2023).

Vásquez-Grandón, A., P. J. Donoso and V. Gerding. 2018. Forest degradation: when is a forest degraded? *Forests* 9(11):726. Doi: 10.3390/f9110726.

Wawrzyniak, M. K., E. M. Kalemba, T. P. Wyka and P. Chmielarz. 2022. Changes in reserve materials deposited in cotyledons of *Pedunculate* Oak (*Quercus robur* L.) Seeds during 18 Months of Storage. *Forests* 13(12):2142. Doi: 10.3390/f13122142.

Yucedag, C., M. Cetin, H. B. Ozel, A. E. S. A. Aisha, O. B. M. Alrabiti and A. M. O. Al Jama. 2021. The impacts of altitude and seed pretreatments on seedling emergence of Syrian juniper (*Juniperus drupacea* (Labill.) Ant. et Kotschy). *Ecological Processes* 10(1):1-6. Doi: 10.1186/s13717-020-00276-z.



Todos los textos publicados por la **Revista Mexicana de Ciencias Forestales** –sin excepción– se distribuyen amparados bajo la licencia *Creative Commons 4.0 Atribución-No Comercial (CC BY-NC 4.0 Internacional)*, que permite a terceros utilizar lo publicado siempre que mencionen la autoría del trabajo y a la primera publicación en esta revista.