

ASSESSMENT OF DIESEL EXHAUST POLLUTANTS EFFECTS IN *Tillandsia capillaris* AND *Ramalina celastri* BY LABORATORY TRIALS

Evaluación de los efectos de contaminantes emitidos por motores diésel en *Tillandsia capillaris* y *Ramalina celastri* mediante ensayos de laboratorio

Ana Carolina MATEOS*, Iván TAVERA BUSSO,
Hebe Alejandra CARRERAS and Claudia María GONZÁLEZ

Universidad Nacional de Córdoba, Área de Contaminación y Bioindicadores (IMBIV-CONICET), Av. Vélez Sarsfield 1611, Córdoba X5016CGA, Argentina.

*Author for correspondence: ac.mateos@unc.edu.ar

Received: December 2020; accepted: March 2022

Key words: vehicular air pollutants, physiochemical damage, PM bound metals, biomonitor, controlled experimental conditions.

ABSTRACT

Traffic-related air pollution is one of the most relevant environmental problems in urban areas. Several cryptogams (i.e., lichens and mosses) and vascular species have been employed to monitor urban air pollution since they allow the assessment of air quality in a large number of sampling sites simultaneously at low cost. In large urban cities, vehicle emissions are frequently the major source of air pollution along with residential energy (for cooking and heating), industry, power generation, and waste incineration. Biomonitor in these urban environments are exposed to a mixture of pollutants making it difficult to identify which pollutant causes the greatest damage to organisms. However, studies that analyze the effect of pollutants emitted by vehicle exhaust are scarce and in the particular case of the most used biomonitor species in Argentina, no analysis of how they are affected by vehicle emissions has been carried out so far. So, the aim of this work was to analyze changes in physiochemical parameters (pigment content, pro-oxidant products, and sulfur accumulation) in *Ramalina celastri*, and heavy metal accumulation in *Tillandsia capillaris*, exposed to diesel exhausts under laboratory conditions. A strong damage in the photosynthetic apparatus of *R. celastri* was observed as well as metal concentration in *T. capillaris* after 20 min of exposure and 48 h of permanence in the exposure chambers. The results indicate that not only the particles and metals cause damage to these two well-known biomonitor, but the interaction of these pollutants with other components of the atmosphere that form different secondary pollutants, together with a longer exposure time, could cause the highest level of damage in them.

Palabras clave: contaminantes vehiculares del aire, daño físico y químico, metales unidos a partículas, biomonitor, condiciones experimentales controladas.

RESUMEN

La contaminación del aire relacionada con el tráfico es uno de los problemas ambientales más relevantes en las zonas urbanas. Se han empleado varias criptógamas (líquenes y musgos) y especies vasculares para monitorear la contaminación del aire urbano, ya que permiten evaluar la calidad de éste en una gran cantidad de sitios de muestreo simultáneamente y a bajo costo. En las grandes ciudades urbanas, las emisiones vehiculares son generalmente la principal fuente de contaminación del aire junto con la energía residencial (para cocinar y calentar), la industria, las centrales de energía y la incineración de residuos. Los biomonitoros en estos entornos urbanos están expuestos a una mezcla de contaminantes, lo que dificulta identificar qué contaminante causa el mayor daño a los organismos. Sin embargo, los estudios que analizan el efecto de los contaminantes emitidos por los gases de escape de los vehículos son escasos y, en el caso particular de las especies de biomonitoros más utilizadas en Argentina, hasta el momento no se ha realizado ningún análisis del efecto de dichas emisiones. Por lo tanto, el objetivo de este trabajo fue analizar en dos biomonitoros ampliamente utilizados, los cambios en los parámetros físicos y químicos (contenido de pigmentos, productos de peroxidación y acumulación de azufre) en *Ramalina celastri* y la acumulación de metales pesados en *Tillandsia capillaris*, expuestos a gases de escape de diésel en condiciones controladas de laboratorio. Se observó un marcado daño en el aparato fotosintético de *R. celastri*, así como concentración de metales en *T. capillaris* después de 20 min de exposición y 48 h de permanencia en las cámaras de exposición. Los resultados indican que no sólo las partículas y los metales causan daño a los biomonitoros, sino que la interacción de estos contaminantes con otros componentes de la atmósfera que forman diferentes contaminantes secundarios, junto con un mayor tiempo de exposición, podría causar el mayor nivel de daño en los biomonitoros.

INTRODUCTION

In urban environments, motor vehicles are one of the main emission sources that contribute to air pollution at local, regional, and global scale along with residential energy (for cooking and heating), industry, power generation, and waste incineration (Jain et al. 2016, Álvarez-Vázquez et al. 2017, Goyal et al. 2021). Over the last years, many urban areas from developing countries, such as Córdoba city (Argentina), showed a strong population growth that derived in an intense traffic flow. This greater number of vehicles generated an increase in the levels of urban air pollutants, which is heightened during the frequent traffic congestions (Puliafito et al. 2011, Morales et al. 2012, Mateos et al. 2018a).

One of the main pollutants emitted by vehicles are suspended particles with aerodynamic diameters less than 10 and 2.5 μm (PM_{10} and $\text{PM}_{2.5}$) (Maricq et al. 1999, Puliafito et al. 2011), although a major contribution of particles can be attributed to emission from diesel-powered vehicles (Maher et al. 2008, Giordano et al. 2010). PM are produced by the engine due to incomplete fuel combustion, lubricant volatilization, and wear and tear on auto parts. Also, brake lining, tire wear, and road dust contribute to

vehicle-related PM emissions (Vouktsis et al. 2009). It has been proven that in urban areas the major source of ultrafine particles are low-mass aerosols from diesel combustion exhausts (Verma et al. 2014), and over 80 % of PM_{10} present in large cities originates from freight and passenger transport that are mainly diesel power vehicles.

PM emissions can be classified as exhaust and non-exhaust (Lawrence et al. 2016). The first category considers particles produced due to incomplete fuel combustion and lubricant volatilization (Vouktsis et al. 2009), while non-exhaust emissions are particles generated by re-suspension of road dust but also by corrosion of vehicle components or during mechanical processes, such as braking, using the clutch, or tire wear (Lawrence et al. 2013, Ravindra et al. 2015). According to a national transport report (Puliafito and Cartesana 2010), in the 1960s 56 % of freight and passenger transportation in Argentina used diesel as fuel and the other 44 % gasoline, while in 2008, 74 % of the transportation was diesel-powered and only 26 % used gasoline. Thus, diesel consumption has been increasing over the years, being always the most consumed fuel in the country. Although a partial control of PM emissions is performed with after-treatment devices, such as catalytic converters

and particle filters, several studies have shown an increase in particle numbers due to enhanced nucleation downstream (Giordano et al. 2010). The disadvantage of these devices is that they release into the air large amounts of platinum group elements (Pt, Pd, Rh) due to thermal and chemical mechanisms that occur while the vehicle is running (Moldovan et al. 2002). Moreover, the replacement of the catalytic converter is not a usual practice in the country due to the lack of regulations (Sbarato and Rubio 2017). Organic matter and elemental carbon account for most of the exhaust particles, being the polycyclic aromatic hydrocarbons (oxy-, and nitro-PAHs) the dominant organic compounds identified (Valavanidis et al. 2006, Phuleria et al. 2007, Najmeddin and Keshavarzi 2018) as the most widespread mutagenic and carcinogenic particulate environmental pollutants. The United States Environmental Protection Agency (US-EPA) has listed 16 PAHs as priority pollutants due to their photomutagenicity (Yan et al. 2004). For example, when the exposure concentration to B[a]P (benzopyrene), one of the 16 PAHs, exceed 1 ng/m³, the DNA would be damaged (Han et al. 2021).

Also, vehicular emissions contribute to atmospheric levels of several toxic metals (Monaci et al. 2000, Singh et al. 2002, Lough et al. 2005). An example of this is the emission of Zn, Cu and Fe (which are used as a coating due to their heat conduction properties) during mechanical abrasion of vehicular brakes (Zechmeister et al. 2006, Raparthi and Phuleria 2021). Zn is also released in the combustion of motor oil and tire wear (Huang et al. 1994, Akbar et al. 2006, Manno et al. 2006). Ni is also related to vehicular air pollution due to the corrosion of bearings, shafts and crankshafts, and it also is emitted from vehicular exhaust (Kumar et al. 2021). Even though the use of leaded gasoline has been banned since the 1990s, vehicular traffic continues to be a source of emission of this metal (Mishra et al. 2004, Kabata-Pendias and Mukherjee 2007) due to the removal of residues accumulated for years in the ground and to resuspension in what is called vehicular dust (Pandey et al. 2014). Another metal associated with vehicular emissions is Mn, since it is used as an additive to increase the octane levels of gasoline and in the manufacture of brake tapes (Keskin et al. 2007, Swietlik et al. 2015). There is also a group of heavy metals related to non-tailpipe emissions, like Co (Wang et al. 2021), which are present in auto parts. Another group of toxic species emitted are gaseous compounds, such as carbon monoxide (CO), nitrogen oxides (NOx), volatile organic compounds (VOCs),

and sulfur dioxide (SO₂), mainly from diesel exhaust (Dogruparmak et al. 2014), as well as some greenhouse gases such as carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) (Hu et al. 2009, Nel and Cooper 2009, Zhang et al. 2016).

In Argentina, as in many other developing countries, despite the harmful effects of the aforementioned pollutants emitted by diesel exhaust, air pollutants are scarcely monitored and only isolated data from researchers is available. The lack of instrumental monitoring has led to the growing use of biomonitoring since they allow the assessment of air quality in a large number of sampling sites simultaneously, with relatively very low cost (Gombert et al. 2006, Augusto et al. 2010, Wannaz et al. 2013, Abril et al. 2014a, Mateos et al. 2018b). In addition, biomonitoring like mosses and lichens can evidence the presence of substances that are difficult to monitor in real time with instruments, such as PM-bound metals (Monaci et al. 2000). Despite its great utility, it is very difficult to find species that can be used globally, so it is interesting to test biomonitoring that are widely distributed in the study region. In particular, *Tillandsia capillaris* has been frequently employed in many air quality studies in Argentina (Pignata et al. 2002, Wannaz and Pignata 2006, Wannaz et al. 2006, Bermúdez et al. 2009, Rodríguez et al. 2011, Abril et al. 2014b, Mateos et al. 2018b), and *Ramalina celastri* has demonstrated to be an excellent biomonitor in urban environments (Garty et al. 1993, 2007, González and Pignata 1994, 1999, Levin and Pignata 1995, González et al. 1998, 2003, Carreras and Pignata 2002, Augusto et al. 2007, Pignata et al. 2007, Carreras et al. 2009, Álvarez et al. 2012, Mateos and González 2016).

Previous studies have demonstrated that vehicular emissions are the main source of pollutants in Córdoba (Stein and Toselli 1996, Olcese and Toselli 2002, Mateos et al. 2018a). On the other hand, other studies with *T. capillaris* (Mateos et al. 2018b) and *R. celastri* in the city (Mateos and González 2016) have demonstrated severe damage to this biomonitor after exposition to the urban atmosphere. However, they were exposed to the mixture of urban pollutants, therefore it is not possible to attribute the observed damage to a specific pollutant. In the present study we focused on the effect of diesel exhaust emissions on physicochemical parameters and heavy metal accumulation in two frequently used biomonitoring. After a bibliographic review, we decided to analyze the variations in physicochemical parameters in *R. celastri* and the accumulation of heavy metals in *T. capillaris*, since

in several studies lichen is indicated as an efficient physiological biomonitor, and the vascular epiphyte as a good biomonitor for heavy metal accumulation. Our hypothesis is that the pollutants emitted by diesel exhaust induce adverse changes in the values of the physicochemical parameters in *R. celastri* and cause an increase in the levels of heavy metals accumulated in *T. capillaris*.

MATERIALS AND METHODS

Biomonitor

Tillandsia capillaris Ruiz and Pav.

Bromeliaceae is a monocotyledon family with a wide distribution in South America that includes the *Tillandsioideae* subfamily, with numerous slow growth epiphytic species that live in trees or inert substrates. They are completely independent from soil to get nutrients, being the adhesion to substrate the only function of their roots (Papini et al. 2010). Due to the epiphytic nature of the *Tillandsia* genus, they are suitable for atmospheric quality monitoring studies (de Souza et al. 2007).

Several samples of *T. capillaris* (Fig. 1) were collected from tree branches using plastic gloves to prevent any risk of sample contamination in a non-polluted, natural reserve area (La Quebrada) located 38 km NW from Córdoba city (Abril et al. 2014a). Part of the collected material was separated to be analyzed as baseline or control group (unexposed) and the rest was used as test organisms in the different trials with diesel exhausts.



Fig. 1. *Tillandsia capillaris* Ruiz and Pav.

Ramalina celastri (Spreng.) Krog and Swinsc

Ramalina celastri (Fig. 2) is an abundant lichen in Argentina growing mainly on tree branches and shrubs, although it is common to find posts covered by *Ramalina thalli* (Nash 2008, Estrabou et al. 2011). It is a very sensitive lichen species, one of the first to disappear when air pollution increases (González and Pignata 1994).



Fig. 2. *Ramalina celastri* (Spreng.) Krog and Swinsc.

R. celastri was collected in Despeñaderos, within a native relict forest located 45 km SW from Córdoba city. The collection of lichens was carried out using plastic gloves in order to avoid any contamination (González and Pignata 1994). Part of the collected material was separated as baseline or control group (unexposed) and the rest was used as test organisms in the different trials with diesel exhausts.

Exposure chambers and experimental setup

To investigate the effects of diesel exhaust emissions, we exposed the biomonitor in closed glass chambers connected to a diesel truck engine (Fig. 3a). Each chamber consisted of a closed glass jar (5 L) with an air intake for the input of diesel emissions and an air output for the exhaust of gases, which generates a constant current flow inside the chambers (Fig. 3b). Thus, 10 net bags (per trial) containing samples either of *R. celastri* (8-10 g/bag) or *T. capillaris* (200 g/bag) were put inside the chambers and exposed to diesel emissions during different time periods. A water-cooling system (Fig. 3c) was employed to reduce the temperature of the emissions and avoid a high temperature stress on the biomonitor. The exhaust temperature after cooling ranged between 22-26 °C. The diesel engine was an MWM-International



Fig. 3. Exposure chambers and experimental equipment. (a) Diesel engine; (b) glass chambers with biomonitor bags; (c) cooling system.

Maxion S4, with the following characteristics: displacement: 3990 cm³; compression ratio: 18.5:1; power: 88 hp in 2800 rpm; cuplemax: kg × m in 1600 rpm: 27.5; consumption every 100 cm³: 96 s; specific weight of diesel fuel: 0.85 g/cm³; volumetric performance: 0.7; number of rpm: 1200 rpm.

First, a trial test was performed to adjust the exposure times until differences were observed with the baseline or control group (unexposed). Then, three trials were conducted twice varying the exposure time and the permanence of biomonitor inside the chamber after exposition: (i) 10 min exposure (ii) 20 min exposure, and (iii) 20 min exposure + 48 h inside the exposure chambers, closed hermetically. All the physiological and chemical determinations in the biomonitor after the trials, were performed also in samples of the control groups that were not exposed to any of the trials, in order to obtain baseline conditions.

Physiological variables

Ramalina celastri

Pigments content

One hundred milligrams of lichen material were homogenized in 10 mL of ethanol at 96 % V/V with an Ultra Turrax homogenizer, T18 (1KA Works, USA). By centrifugation, the supernatant was separated and HCl 0.06 M was added to the clear chlorophyll extract (1 mL HCl and 5 mL chlorophyll extract) in order to produce phaeophytin formation. The absorption of chlorophyll a (Chl a), chlorophyll b (Chl b), carotenoids (Carot), phaeophytin a (Ph a), phaeophytin b (Ph b) and phaeophytins alone (after addition of HCl) was measured with a DU 7000 spectrophotometer (Beckman, USA). Concentrations of chlorophylls, carotenoids and phaeophytins were calculated considering the dry weight (DW). The results were expressed as mg g⁻¹ DW (Wintermans and de Mots 1965). The Ph a/Chl a ratio was also calculated (Carreras et al. 1998, González et al. 1998).

Peroxidation products

Lichen samples were freeze-dried and 100 mg were homogenized in 2.5 mL of distilled H₂O. An equal volume of 0.5 % 2-thiobarbituric acid in a 20 % trichloroacetic acid solution was added, and samples were incubated at 95 °C for 30 min. The reaction was cut by placing the tubes in an ice bath and then samples were centrifuged for 30 min at 410 rad/s. After that, the supernatant was removed and the absorption at 532 nm was registered; the value for non-specific absorption at 600 nm was subtracted. Malondialdehyde (MDA) concentration was calculated from the extinction coefficient of 155 mM/cm (Kosugi et al. 1989) with results being expressed in μmol/g DW. Hydroperoxy conjugated dienes (HPCD) were extracted by homogenization of 50 mg of the lichen in 96 % v/v ethanol at 1:50 FW/v ratio with an Ultra Turrax homogenizer. The supernatant was separated and the absorption was measured at 234 nm; the concentration was calculated with the formula $\Sigma = 2.65 \times 10^4$ M/cm (Boveris et al. 1980) and the results expressed as μmol/g DW.

Bioaccumulation variables

Sulfur content in R. celastri

A Mg(NO₃)₂ saturated solution was used where 5 mL were added to 0.5 g of freeze-dried lichen and dried in a stove. Then, the sample was heated in an oven for 30 min at 500 °C and ashes were suspended in 10 mL of a 6 M HCl solution. Then, they were filtered and the resulting solution boiled for 3 min. The solution was brought up to 50 mL with distilled H₂O and the amount of SO₄²⁻ in the solution was determined by the turbidimetric method using barium chloride (González and Pignata 1994). Results were expressed in mg of total sulfur [(S)/g DW].

Heavy metal content in T. capillaris

Approximately 2.5 g of *T. capillaris* dry weight leaf were put at 450 °C for 4 h in a muffle furnace,

TABLE I. MEAN TEMPERATURE (°C), ATMOSPHERIC PRESSURE (hPa), FUEL CONSUMPTION (cm³) AND VOLUME OF GASES EMITTED (m³) DURING EACH TRIAL.

Trial	Temperature (°C)	Atmospheric pressure (hPa)	Fuel consumption (cm ³)	Gases volume (m ³)
10 min	19.1	955.0	625	16.75
20 min	20.0	954.8	1250	33.50
20 min + 48 h	19.8	951.1	1237	33.20

then digested with 5 mL concentrated HNO₃ (65 % Merck, Germany) and kept 24 h in the dark (Wannaz and Pignata 2006). Samples were filtered twice using a 2 µm filter paper (Munktell, Germany) and brought to a final volume of 25 mL with ultrapure water (MilliQ). The accumulation of Pb, Cu, Co, Fe, Mn, Zn and Ni were determined by atomic absorption spectrophotometry (AAS, Perkin-Elmer AA3110). The same procedure was done in baseline samples, certified reference material (CTA-OTL-1, Institute of Nuclear Chemistry and Technology) and laboratory blanks to acknowledge the baseline heavy metal content of the species and to evaluate the digestion procedure, respectively. Only residual standard deviation values less than 10 % were accepted with samples outside this range being re-analyzed. Results were expressed in µg/g DW.

Pollution index

The pollution index (PI) for *R. celastri* was determined using the equation 1 (González et al. 1996):

$$\text{PI} = \frac{[(\text{Ph a} / \text{Chl a}) + (\text{S}_E / \text{S}_F)]}{[(\text{MDA}_E / \text{MDA}_F) + (\text{HPCD}_E / \text{HPCD}_F)]} \quad (1)$$

Where subindex E indicates the exposed samples and subindex F refers to the freshly picked material (baseline). This index has been widely checked and used in biomonitoring studies with the same species (González and Pignata 1994, González et al. 1996, 1998, 2003, Olcese and Toselli 2002, Rodríguez et al. 2007, Mateos and González 2016).

Data analysis

Statistical analyses were carried out with mean values corresponding to three subsamples from each bag. Assumptions for normality were tested by the modified Shapiro-Wilks normality test and the homogeneity of variance by the Levene's test. The mean and standard deviation for each physicochemical parameter determined in *R. celastri* and heavy metals measured in *T. capillaris* in each trial, were calculated. An analysis of the variance (ANOVA) was performed to compare the results of the four different

trials (unexposed samples, exposition during 10 min, exposition during 20 min and exposition during 20 min + 48 h). When the ANOVA null hypothesis was rejected ($p < 0.05$) post-hoc comparisons were performed using the LSD (least significant difference) Fisher posteriori test. In order to analyze relationships between the parameters determined in the biomonitor and the different trials, a multivariate analysis (principal component analysis [PCA]) was carried out. All the analyses were performed employing InfoStat (Di Rienzo et al. 2011).

RESULTS AND DISCUSSION

Trials parameters

Table I shows the temperature, atmospheric pressure, fuel consumption and gas volumes during the different expositions. Emitted gases concentrations were estimated according to engine characteristics, combustion period and amount of fuel consumed (**Table II**). These results are in good agreement with the data showed in Mbuligwe and Kassenga (1997) for a 3.5 t truck/bus with a diesel engine similar to the one employed in this study.

TABLE II. CONCENTRATION RANGES OF DIESEL EXHAUST GAS COMPONENTS IN IDLING CONDITIONS.

Components	NOx (ppm)	NO ₂ (ppm)	CO (ppm)
This study	120-150	2-10	520-640
Mbuligwe and Kassenga 1997	< 1.5-300	< 0.5 > 60	440-600

Physicochemical parameters

We first run a 10 min trial exposure, but no significant differences were observed with unexposed samples in any of the parameters measured in *R. celastri* (**Table III**). When we exposed the biomonitor

TABLE III. MEAN VALUES \pm STANDARD DEVIATION OF PHYSIOCHEMICAL PARAMETERS MEASURED IN *Ramalina celastri* AND ANALYSES OF VARIANCE (ANOVA) BETWEEN TRIALS.

	Unexposed	10 min	20 min	20 min + 48 h	ANOVA
Chl a	1.129 \pm 0.067 A	1.126 \pm 0.180 A	1.211 \pm 0.142 A	0.834 \pm 0.101 B	***
Chl b	0.350 \pm 0.017 A	0.382 \pm 0.071 A	0.379 \pm 0.044 A	0.276 \pm 0.033 B	**
Ph a	1.405 \pm 0.090	1.200 \pm 0.183	1.198 \pm 0.249	1.563 \pm 0.003	ns
Ph b	0.360 \pm 0.019	0.367 \pm 0.058	0.374 \pm 0.036	0.330 \pm 0.026	ns
Carot	0.304 \pm 0.012 A	0.303 \pm 0.046 A	0.315 \pm 0.039 A	0.206 \pm 0.027 B	***
Ph a/Chl a	1.062 \pm 0.004 B	1.067 \pm 0.014 B	1.065 \pm 0.013 B	1.205 \pm 0.060 A	***
Chl b/Chl a	0.339 \pm 0.008 AB	0.338 \pm 0.012 AB	0.325 \pm 0.016 B	0.402 \pm 0.067 A	*
HPCD	5.347 \pm 0.060 B	11.831 \pm 0.769 AB	12.495 \pm 1.817 A	15.318 \pm 0.255 A	***
MDA	0.109 \pm 0.015 C	0.112 \pm 0.017 C	0.136 \pm 0.012 B	0.156 \pm 0.013 A	**
Sulfur	0.808 \pm 0.034 C	0.895 \pm 0.097 BC	1.003 \pm 0.111 B	1.281 \pm 0.030 A	**
PI	3.787 \pm 0.052 B	3.854 \pm 0.047 B	3.727 \pm 0.156 B	4.209 \pm 0.307 A	*

Chl a: chlorophyll a; Chl b: chlorophyll b; Ph a: phaeophytin a; Ph b: phaeophytin b; Carot: carotenoids; HPCD: hydroperoxy conjugated dienes; MDA: malondialdehyde; PI: pollution index; ns: not significant.

Values in each horizontal line followed by the same letter do not differ significantly.

*Significant at the 0.05 probability level; **significant at the 0.01 probability level; ***significant at the 0.001 probability level.

to a 20 min period, no significant differences were observed in pigment contents. However, an increase in MDA levels, PI and sulfur content was observed, indicating the presence of compounds that produced some physiological damage in the biomonitor. When lichens were exposed for 20 min and left 48 h in the closed chamber, all physiological parameters were significantly altered. A notable decrease in Chl a, Chl b and carotenoids concentration was observed, which is consistent with the findings of Langmann et al. (2014), who observed in another lichen species that diesel exhaust affects the photosynthetic apparatus, decreasing pigments content and altering its ability to perform photosynthesis. Chl a is the most important photosynthetic pigment, while Chl b and carotenoids function as photoprotective pigments (Han et al. 2017). Besides, carotenoids are capable of eliminating some free radicals from the chloroplast, therefore a decrease in these pigments increases the vulnerability of the chloroplast to the oxidative damage of free radicals, causing a severe damage to membranes and associated molecules. Moreover, an alteration of antioxidant systems in plants exposed to atmospheric pollutants have been already observed (González and Pignata 1994, Levin and Pignata 1995, Carreras et al. 1998, Munzi et al. 2012). The decrease in carotenoids concentration could explained the increase in biomarkers related to lipid peroxidation of cell membranes, MDA and HPCD, in the 20 min + 48 h trial.

One of the main gases emitted by diesel exhaust is SO₂ (Liu et al. 2016), in addition to sulfates formed by oxidation of sulfur in fuels and lubricant (Tan et al.

2017); therefore, the highest sulfur values measured in the 20 min + 48 h trial could be attributed to diesel emissions. Like all the parameters analyzed above, the PI also showed the highest values in the 20 min + 48 h trial. This prolonged permanence of biomonitor in the exposed chamber could be compared with the exposition in heavy polluted urban environments.

In order to identify the parameters that best explain the variability of the data, a multivariate analysis (PCA) was performed using the physiochemical determinations as variables and the different trials as classification criteria (excluding the 10 min that showed no effect). Eigenvalues corresponding to the first two components are shown in **table IV**. The eigenvalues associated with the axis and the different trials are indicators of the extent to which the lichen response explains laboratory trial variables. The first two axes (PC1 and PC2) explained most of the total variance, where the first axis (PC1) was mostly determined by individual pigments, the ratios and the sulfur accumulation, and the second (PC2) was driven by the PI and HPCD (**Table IV**), reflecting that these two biomarkers are less useful to identify diesel exhaust effects on lichens. The biplot (**Fig. 4**) shows that all damage indicators (Chl b/Chl a, Ph a/Chl a, sulfur, MDA, PI and in a lesser extent HPCD) were associated with the 20 + 48 h trial, in agreement with the results informed in **table III**. It is interesting to note that the PI was associated to a lesser extent to the 20 min exposure, confirming this indicator is sensitive enough to reflect physiological damage caused by vehicular emissions, as seen in previous studies with the same species (González et al. 1996,

TABLE IV. EIGENVECTORS CORRESPONDING TO THE FIRST TWO PRINCIPAL COMPONENTS OBTAINED IN THE PRINCIPAL COMPONENT ANALYSIS FOR THE PHYSICOCHEMICAL PARAMETERS DETERMINED IN *Ramalina celastri* EXPOSED TO DIESEL EXHAUST.

Parameters	Component	
	1	2
Chl a	0.31	0.10
Chl b	0.31	-0.03
Ph a	0.31	-0.08
Ph b	0.29	0.30
Carot	0.31	0.09
HPCD	-0.12	0.69
MDA	-0.31	0.15
Ph a/Chl a	-0.30	-0.25
Chl b/Chl a	-0.28	-0.32
Sulfur	-0.31	-0.09
PI	-0.26	0.42

Chl a: chlorophyll a; Chl b: chlorophyll b; Ph a: phaeophytin a; Ph b: phaeophytin b; Carot: carotenoids; HPCD: hydroperoxy conjugated dienes; MDA: malondialdehyde; PI: pollution index.

1998, 2003, Rodríguez et al. 2007, Mateos and González 2016). These results indicate that 20 min of high exposure to diesel exhaust are not enough to produce a considerably damage in the lichen.

Heavy metals accumulation

The mean concentrations ($\mu\text{g/g DW}$) of Pb, Cu, Ni, Co, Fe, Mn and Zn accumulated in samples of

T. capillaris as well as the ANOVA results between trials (unexposed; 10 min; 20 min; 20 min + 48 h) are shown in **table V**. A significant increase in the concentration of all elements was observed in the 20 min + 48 h trial. These results could be explained by the fact that besides primary particles that are directly emitted from a source, PM can also be formed in the air due to some physical and chemical processes (secondary particles) where condensation or homogenous nucleation of gas species onto existing particles are among the physical processes that can result in secondary particle formation (Esmaeilirad and Hosseini 2018). Moreover, particles of diesel exhaust have large and irregular surfaces that facilitate the adsorption of different organic and inorganic materials from the environment or from the engine exhaust itself (Wernke 2014). Also, the values found in this study are in good agreement with those shown in Mateos et al. (2018b) for the same specie exposed for six months in the downtown area of Córdoba city.

The levels of Ni and Fe were also high in the 20 min trial, suggesting that even short exposure periods can induce changes in elemental concentrations. These results are in good agreement with studies that indicate that besides gases, particle-bound heavy metals are also abundant in diesel emissions (Wernke 2014, Wang et al. 2018). Even though the Co emission from diesel fuel is not frequently reported (Fuga et al. 2008), in the present study we observed an increase even in samples exposed during 10 min to diesel emissions. This difference could be attributed to the fact that the in the present study we did not use a catalytic converter neither a




Fig. 4. Biplot based on the two principal components (PC1 and PC2) of the principal component analysis for physicochemical parameters determined for *Ramalina celastri* in the trials' categories (unexposed; 20 min; 20 min + 48 h).

TABLE V. MEAN VALUES (μg/g DW) ± STANDARD DEVIATION OF HEAVY METAL ACCUMULATED IN *Tillandsia capillaris* AND ANALYSES OF VARIANCE (ANOVA) BETWEEN TRIALS.

	Unexposed			10 min			20 min			20 min + 48 h			ANOVA				
Pb	10.73	±	0.81	C	12.15	±	0.90	BC	14.20	±	1.49	B	18.31	±	0.66	A	***
Cu	5.37	±	0.86	B	5.09	±	0.66	B	5.19	±	0.99	B	7.35	±	1.42	A	***
Ni	2.11	±	0.01	B	3.02	±	0.70	B	5.60	±	1.32	AB	7.31	±	3.30	A	*
Co	1.16	±	0.12	B	2.15	±	0.93	AB	2.36	±	1.18	A	2.90	±	0.66	A	*
Fe	974.68	±	102.10	B	1598.23	±	365.12	B	3773.33	±	378.25	A	3962.54	±	321.65	A	**
Mn	49.44	±	2.82	B	94.48	±	15.850	AB	96.97	±	47.26	AB	137.60	±	18.30	A	*
Zn	12.28	±	1.07	C	18.27	±	1.460	BC	23.39	±	1.79	B	29.82	±	2.07	A	***

Values in each horizontal line followed by the same letter do not differ significantly.

*Significant at the 0.05 probability level; **significant at the 0.01 probability level; ***significant at the 0.001 probability level.

particle filter, as is the situation of public transport in Córdoba city.

It is noteworthy that the concentrations of Pb, Mn and Zn were significantly higher in exposed samples, even during short time periods in agreement with the fact that these metals have been found in high levels in particles collected from areas with intense traffic (Mateos et al. 2018b). Metals in diesel emissions can derive from different sources such as fuel alone or from the use of additives; however, most of them derive from lubricating oil and by-products of engine wear that enter the combustion chamber (Giordano et al. 2010). Elements that can be attributed to traffic have been found at remarkably high levels in the exhaust of diesel vehicles and to a lesser extent in gasoline powered vehicles (Monaci et al. 2000). Particularly, the presence of Pb, Cu, Ni, Fe, Mn and Zn in diesel emissions has been confirmed before (Monaci et al. 2000, Geller et al. 2006, Fuga et al. 2008, Hu et al. 2009); therefore, the effect of some of these metals together with some powerful oxidant gases, such as SO₂, could be the responsible for the noticeable damage observed in the biomonitor *R. celastri*, as reflected earlier by the PI. To globally observe the relationship of heavy metals accumulated in the biomonitor with the different trials, a PCA was performed. **Table VI** shows the eigenvectors values, where PC1 is mostly explained by the concentrations of Pb, Ni, Co, Mn, and Zn, all of which have the greatest association with diesel emissions (Giordano et al. 2010). PC2 was associated with Cu and Fe concentrations in the biomonitor, which is consistent with the findings of Bermúdez et al. (2012), who suggested that Fe and Cu concentrations are indicators of anthropogenic emissions related to soil resuspension and metallurgical and metal-mechanical activities, respectively. **Figure 5** presents the biplot obtained with the PCA analysis showing the strong association of heavy metals with

TABLE VI. EIGENVECTORS CORRESPONDING TO THE FIRST TWO COMPONENTS OBTAINED IN THE PRINCIPAL COMPONENT ANALYSIS FOR THE HEAVY METAL ACCUMULATION DETERMINED IN *Tillandsia capillaris* EXPOSED TO DIESEL EXHAUST.

Heavy metal	Component	
	1	2
Pb	0.41	-0.16
Cu	0.31	-0.58
Ni	0.42	0.05
Co	0.42	0.07
Fe	0.29	0.63
Mn	0.38	0.37
Zn	0.39	-0.33

the 20 + 48 h trial, confirming what was observed in the ANOVA test.

CONCLUSIONS

The laboratory trials performed in the present study demonstrate that pollutants coming from diesel engines might cause physiological damage in biomonitor, even with exposures as short as 20 min. In addition, an increase in the damaging effect and larger metal accumulation was observed when the exposure time of biomonitor was augmented. Thus, larger concentrations of secondary pollutants and more damaging effects on biomonitor are expected with time inside the chamber, since secondary pollutants are much more reactive and damaging than primary ones.

On the other hand, a clear effect of pollutants from diesel engines was observed on the integrity




Fig. 5. Biplot based on the two principal components (PC1 and PC2) of the principal component analysis for heavy metals accumulation determined for *Tillandsia capillaris* in the trials' categories (unexposed; 20 min; 20 min + 48 h).

of the photosynthetic apparatus. This is consistent with previous studies where it was observed that diesel emissions were significantly related to the chlorophyll degradation index. In addition, diesel exhaust pollutants cause a marked decrease in carotenoid content, which increases the susceptibility of biomonitoring to oxidative damage caused by free radicals.

Another noticeable result were the high levels of Pb accumulated by *T. capillaris* after a 20 min exposition to vehicle emissions. Despite the fact that this metal is no longer used in fuels since the 1990s, it is possible that vehicles are still sources of Pb from lead plates that line the fuel tanks, lead in vulcanized gasoline tubes or in the lining of pistons, valve seats and spark plugs. The results of the present study evidence the need to regulate Pb emissions coming from vehicles, considering that this metal has known toxicological and carcinogenic effects. On the other hand, the results obtained here are a first approximation to establish that pollutants emitted by diesel exhaust have a notable effect on biomonitoring. This confirms the assumption that traffic is one of the main and most worrying sources of air pollutants in urban areas. More studies evaluating other air pollutants present in urban areas are necessary to complement the preliminary results of this work.

Overall, the results presented in here are particularly relevant for cities with a large proportion of old vehicles, whose atmospheres could be similar in composition to the one simulated in the chambers employed in this study.

ACKNOWLEDGMENTS

The authors would like to acknowledge the financial support granted by the Agencia Nacional de Promoción Científica y Tecnológica (FONCyT), the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), and the Secretaría de Ciencia y Técnica de la Universidad Nacional de Córdoba (SECyT-UNC). We especially thank Ing. Mario Spinoza and all the engineers from the FCEFyN-UNC Engine Testing Area, whose collaboration was fundamental for the realization of this study.

REFERENCES

- Abril G.A., Wannaz E.D., Mateos A.C., Invernizzi R., Plá R. and Pignata M. (2014a). Characterization of atmospheric emission sources of heavy metals and trace elements through a local-scale monitoring network using *T. capillaris*. *Ecological Indicators* 40, 153-161. <https://doi.org/10.1016/j.ecolind.2014.01.008>
- Abril G.A., Wannaz E.D., Mateos A.C. and Pignata M.L. (2014b). Biomonitoring of airborne particulate matter emitted from a cement plant and comparison with dispersion modelling results. *Atmospheric Environment* 82, 154-63. <https://doi.org/10.1016/j.atmosenv.2013.10.020>
- Akbar K.F., Hale W.H.G., Headley A.D. and Athar M. (2006). Heavy metal contamination of roadside soils. *Journal of Soil and Water Research* 1 (4), 158-163. <https://doi.org/10.17221/6517-SWR>

Álvarez R., del Hoyo A., García-Breijo F., Reig-Armiñana J., del Campo E.M., Guéra A. and Casano L.M. (2012). Different strategies to achieve Pb-tolerance by the two *Trebouxia algae* coexisting in the lichen *Ramalina farinacea*. *Journal of Plant Physiology* 169 (18), 1797-1806. <https://doi.org/10.1016/j.jplph.2012.07.005>

Álvarez-Vázquez L., García-Chan N., Martínez A. and Vázquez-Méndez M. (2017). Numerical simulation of air pollution due to traffic flow in urban networks. *Journal of Computational and Applied Mathematics* 326, 44-61. <https://doi.org/10.1016/j.cam.2017.05.017>

Augusto S., Mágua C., Matos J., Pereira M.J. and Branquinho C. (2010). Lichens as an integrating tool for monitoring PAH atmospheric deposition: A comparison with soil, air and pine needles. *Environmental Pollution* 158 (2), 483-489. <https://doi.org/10.1016/j.envpol.2009.08.016>

Augusto S., Catarino F. and Branquinho C. (2007). Interpreting the dioxin and furan profiles in the lichen *Ramalina canariensis* Steiner for monitoring air pollution. *Science of the Total Environment* 377 (1), 114-123. <https://doi.org/10.1016/j.scitotenv.2007.01.089>

Bermúdez G.M., Rodríguez J.H. and Pignata M.L. (2009). Comparison of the air pollution biomonitoring ability of three *Tillandsia* species and the lichen *Ramalina celastri* in Argentina. *Environmental Research* 109 (1), 6-14. <https://doi.org/10.1016/j.envres.2008.08.014>

Bermúdez G.M., Jasan R., Plá R. and Pignata M.L. (2012). Heavy metals and trace elements in atmospheric fallout: their relationship with topsoil and wheat element composition. *Journal of hazardous materials* 213, 447-456. <https://doi.org/10.1016/j.jhazmat.2012.02.023>

Boveris A., Cadenas E. and Chance B. (1980). Low-level chemi-luminescence of the lipoxygenase reaction. *Photobiochemistry and Photobiophysics* 1 (3), 175-82.

Carreras H., Gudino G. and Pignata M. (1998). Comparative biomonitoring of atmospheric quality in five zones of Córdoba city (Argentina) employing the transplanted lichen *Usnea* sp. *Environmental Pollution* 103 (2-3), 317-25. [https://doi.org/10.1016/S0269-7491\(98\)00116-X](https://doi.org/10.1016/S0269-7491(98)00116-X)

Carreras H.A. and Pignata M.L. (2002). Biomonitoring of heavy metals and air quality in Córdoba City, Argentina, using transplanted lichens. *Environmental Pollution* 117 (1), 77-87. [https://doi.org/10.1016/S0269-7491\(01\)00164-6](https://doi.org/10.1016/S0269-7491(01)00164-6)

Carreras H.A., Wannaz E.D. and Pignata M.L. (2009). Assessment of human health risk related to metals by the use of biomonitoring in the province of Córdoba, Argentina. *Environmental Pollution* 157 (1), 117-122. <https://doi.org/10.1016/j.envpol.2008.07.018>

De Souza Pereira M., Heitmann D., Reifenhäuser W., Meire R.O., Santos L.S. and Torres J.P.M. (2007). Persistent organic pollutants in atmospheric deposi-
tion and biomonitoring with *Tillandsia usneoides* (L.) in an industrialized area in Rio de Janeiro state, southeast Brazil—Part II: PCB and PAH. *Chemosphere* 67 (9), 1736-1745. <https://doi.org/10.1016/j.chemosphere.2006.05.141>

Di Rienzo J.A., Casanoves F., Balzarini M.G., González L., Tablada M. and Robledo C.W. (2011). InfoStat v. 2011.

Dogruparmak S.C., Keskin G.A., Yaman S. and Alkan A. (2014). Using principal component analysis and fuzzy c-means clustering for the assessment of air quality monitoring. *Atmospheric Pollution Research* 5 (4), 656-663. <https://doi.org/10.5094/APR.2014.075>

Esmaeilirad S. and Hosseini V. (2018). Modeling the formation of traditional and non-traditional secondary organic aerosols from in-use, on-road gasoline and diesel vehicles exhaust. *Journal of Aerosol Science* 124, 68-82. <https://doi.org/10.1016/j.jaerosci.2018.07.003>

Estrabou C., Filippini E., Soria J.P., Schelotto G. and Rodríguez J.M. (2011). Air quality monitoring system using lichens as bioindicators in Central Argentina. *Environmental Monitoring and Assessment* 182 (1-4), 375-383. <https://doi.org/10.1007/s10661-011-1882-4>

Fuga A., Saiki M., Marcelli M.P. and Saldiva P.H. (2008). Atmospheric pollutants monitoring by analysis of epiphytic lichens. *Environmental Pollution* 151 (2), 334-340. <https://doi.org/10.1016/j.envpol.2007.06.041>

Garty J., Karary Y., Harel J. and Lurie S. (1993). Temporal and spatial fluctuations of ethylene production and concentrations of sulfur, sodium, chlorine and iron on/in the thallus cortex in the lichen *Ramalina duriaeae* (De Not.) Bagl. *Environmental and Experimental Botany* 33 (4), 553-563. [https://doi.org/10.1016/0098-8472\(93\)90030-J](https://doi.org/10.1016/0098-8472(93)90030-J)

Garty J., Tamir O., Levin T. and Lehr H. (2007). The impact of UV-B and sulphur- or copper-containing solutions in acidic conditions on chlorophyll fluorescence in selected *Ramalina* species. *Environmental Pollution* 145 (1), 266-273. <https://doi.org/10.1016/j.envpol.2006.03.022>

Geller M.D., Ntziachristos L., Mamakos A., Samaras Z., Schmitz D.A. and Froines J.R. (2006). Physicochemical and redox characteristics of particulate matter (PM) emitted from gasoline and diesel passenger cars. *Atmospheric Environment* 40 (36), 6988-7004. <https://doi.org/10.1016/j.atmosenv.2006.06.018>

Goyal P., Gulia S. and Goyal S.K. (2021). Identification of air pollution hotspots in urban areas - An innovative approach using monitored concentrations data. *Science of The Total Environment* 798, 149143. <https://doi.org/10.1016/j.scitotenv.2021.149143>

Giordano S., Adamo P., Spagnuolo V. and Vaglieco, B.M. (2010). Instrumental and bio-monitoring of heavy metal and nanoparticle emissions from diesel engine

exhaust in controlled environment. *Journal of Environmental Sciences* 22 (9), 1357-1363. [https://doi.org/10.1016/S1001-0742\(09\)60262-X](https://doi.org/10.1016/S1001-0742(09)60262-X)

Gombert S., Asta J. and Seaward M.R.D. (2006). Lichens and tobacco plants as complementary biomonitoring of air pollution in the Grenoble area (Isère, southeast France). *Ecological Indicators* 6, 429-443. <https://doi.org/10.1016/j.ecolind.2005.06.001>

González C.M. and Pignata M.L. (1994). The influence of air pollution on soluble proteins, chlorophyll degradation, MDA, sulphur and heavy metals in a transplanted lichen. *Chemistry and Ecology* 9 (2), 105-113. <https://doi.org/10.1080/02757549408038568>

González C.M., Casanovas S.S. and Pignata M.L. (1996). Biomonitoring of air pollutants from traffic and industries employing *Ramalina ecklonii* (Spreng.) Mey. and Flot. in Cordoba, Argentina. *Environmental Pollution* 91 (3), 269-277. [https://doi.org/10.1016/0269-7491\(95\)00076-3](https://doi.org/10.1016/0269-7491(95)00076-3)

González C., Orellana L., Casanovas S. and Pignata M. (1998). Environmental conditions and chemical response of a transplanted lichen to an urban area. *Journal of Environmental Management* 53 (1), 73-81. <https://doi.org/10.1006/jema.1998.0194>

González C. and Pignata M. (1999). Effect of pollutants emitted by different urban-industrial sources on the chemical response of the transplanted *Ramalina ecklonii* (Spreng.) Mey. and Flot. *Toxicological and Environmental Chemistry* 69 (1-2), 61-73. <https://doi.org/10.1080/02772249909358688>

González C., Pignata M. and Orellana L. (2003). Applications of redundancy analysis for the detection of chemical response patterns to air pollution in lichen. *Science of the Total Environment* 312 (1), 245-253. [https://doi.org/10.1016/S0048-9697\(03\)00253-5](https://doi.org/10.1016/S0048-9697(03)00253-5)

Han K., Ran Z., Wang X., Wu Q., Zhan N., Yi Z. and Jin T. (2021). Traffic-related organic and inorganic air pollution and risk of development of childhood asthma: A meta-analysis. *Environmental Research* 194, 110493. <https://doi.org/10.1016/j.envres.2020.110493>

Han P.P., Shen S.G., Guo R.J., Yao S.Y., Sun Y., Tan Z.L. and Jia S.R. (2017). The regulation of photosynthetic pigments in terrestrial *Nostoc flagelliforme* in response to different light colors. *Algal Research* 25, 128-135. <https://doi.org/10.1016/j.algal.2017.04.009>

Hu S., Herner J.D., Shafer M., Robertson W., Schauer J.J. and Dwyer H. (2009). Metals emitted from heavy-duty diesel vehicles equipped with advanced PM and NOX emission controls. *Atmospheric Environment* 43 (18), 2950-2959. <https://doi.org/10.1016/j.atmosenv.2009.02.052>

Huang X., Ilhan O. and Aras A. (1994). Emissions of trace elements from motor vehicles: potential marker elements and source composition profile. *Atmospheric Environment* 28, 1380-1392. [https://doi.org/10.1016/1352-2310\(94\)90201-1](https://doi.org/10.1016/1352-2310(94)90201-1)

Jain S., Aggarwal P., Sharma P. and Kumar P. (2016). Vehicular exhaust emissions under current and alternative future policy measures for megacity Delhi, India. *Journal of Transport and Health* 3 (3), 404-412. <https://doi.org/10.1016/j.jth.2016.06.005>

Kabata-Pendias A. and Mukherjee A.B. (2007). Trace elements from soil to human. Springer-Verlag, Berlin, 550 pp. https://doi.org/10.1007/978-3-540-32714-1_7

Keskin A., Gürü M. and Altiparmak D. (2007). Biodiesel production from tall oil with synthesized Mn and Ni based additives: effects of the additives on fuel consumption and emissions. *Fuel* 86 (7-8), 1139-1143. <https://doi.org/10.1016/j.fuel.2006.10.021>

Kosugi H., Kojima T. and Kikugawa K. (1989). Thio-barbituric acid-reactive substances from peroxidized lipids. *Lipids* 24 (10), 873-881. <https://doi.org/10.1007/BF02535762>

Kumar A., Kumar P., Singh H. and Kumar N. (2021). Adaptation and mitigation potential of roadside trees with bio-extraction of heavy metals under vehicular emissions and their impact on physiological traits during seasonal regimes. *Urban Forestry and Urban Greening* 58, 126900. <https://doi.org/10.1016/j.ufug.2020.126900>

Langmann U., Madl P., Türk R., Hofmann W. and Brunauer G. (2014). Sensitivity of lichens to diesel exhaust under laboratory conditions. *Journal of Environmental Protection* 5 (13), 1331-1341. <https://doi.org/10.4236/jep.2014.513127>

Lawrence S., Sokhi R., Ravindra K., Mao H., Prain H.D. and Bull I.D. (2013). Source apportionment of traffic emissions of particulate matter using tunnel measurements. *Journal of Atmospheric Environment* 77, 548-557. <https://doi.org/10.1016/j.atmosenv.2013.03.040>

Lawrence S., Sokhi R. and Ravindra K. (2016). Quantification of vehicle fleet PM₁₀ particulate matter emission factors from exhaust and non-exhaust sources using tunnel measurement techniques. *Environmental Pollution* 210, 419-28. <https://doi.org/10.1016/j.enpol.2016.01.011>

Levin A.G. and Pignata M.L. (1995). *Ramalina ecklonii* as a bioindicator of atmospheric pollution in Argentina. *Canadian Journal of Botany* 73 (8), 1196-1202. <https://doi.org/10.1139/b95-129>

Liu X., Osaka Y., Huang H., Kodama A., He Z., Yang X. and Chen Y. (2016). Development of high-performance SO₂ trap materials in the low-temperature region for diesel exhaust emission control. *Separation and Purification Technology* 162, 127-133. <https://doi.org/10.1016/j.seppur.2016.02.010>

Lough G.C., Schauer J.J., Park J., Shafer M.M., DeMinter J.T. and Weinstein J.P. (2005). Emissions of metals associated with motor vehicle roadways. *Environmental Science and Technology* 39 (3), 826-836. <https://doi.org/10.1021/es048715f>

Maher B.A., Moore C. and Matzka J. (2008). Spatial variation in vehicle-derived metal pollution identified by magnetic and elemental analysis of roadside tree leaves. *Atmospheric Environment* 42 (2), 364-373. <https://doi.org/10.1016/j.atmosenv.2007.09.013>

Manno E., Varrica D. and Dongarra G. (2006). Metal distribution in road dust samples collected in an urban area close to a petrochemical plant at Gela, Sicily. *Atmospheric Environment* 40, 5929-5942. <https://doi.org/10.1016/j.atmosenv.2006.05.020>

Maricq M.M., Podsiadlik D.H. and Chase R.E. (1999). Examination of the size-resolved and transient nature of motor vehicle particle emissions. *Environmental Science and Technology* 33, 1618-1626. <https://doi.org/10.1021/es9808806>

Mateos A.C. and González C. (2016). Physiological response and sulfur accumulation in the biomonitor *Ramalina celastri* in relation to the concentrations of SO₂ and NO₂ in urban environments. *Microchemical Journal* 125, 116-23. <https://doi.org/10.1016/j.microc.2015.11.025>

Mateos A.C., Amarillo A.C., Busso I.T. and González C.M. (2018a). Evaluación espacial y temporal de la contaminación por SO₂, NO₂, O₃ y CO en la ciudad de Córdoba. *Revista de la Facultad de Ciencias Exactas, Físicas y Naturales* 5 (2), 47.

Mateos A.C., Amarillo A.C., Carreras H.A. and González C. (2018b). Land use and air quality in urban environments: Human health risk assessment due to inhalation of airborne particles. *Environmental Research* 161, 370-380. <https://doi.org/10.1016/j.envres.2017.11.035>

Mbuligwe S.E. and Kassenga G.R. (1997). Automobile air pollution in Dar es Salaam city, Tanzania. *Science of the Total Environment* 199 (3), 227-235. [https://doi.org/10.1016/S0048-9697\(97\)05461-2](https://doi.org/10.1016/S0048-9697(97)05461-2)

Mishra V.K., Kim K.H., Kang C.H. and Choi K.C. (2004). Winter time and distribution of airborne lead in Korea. *Atmospheric Environment* 38, 2653-2664. <https://doi.org/10.1016/j.atmosenv.2004.02.025>

Moldovan M., Palacios M.A., Gómez M.M., Morrison G., Rauch S., McLeod C. and Santamaría J. (2002). Environmental risk of particulate and soluble platinum group elements released from gasoline and diesel engine catalytic converters. *Science of the Total Environment* 296 (1-3), 199-208. [https://doi.org/10.1016/S0048-9697\(02\)00087-6](https://doi.org/10.1016/S0048-9697(02)00087-6)

Monaci F., Moni F., Lanciotti E., Grechi D. and Bargagli R. (2000). Biomonitoring of airborne metals in urban environments: New tracers of vehicle emission, in place of lead. *Environmental Pollution* 107 (3), 321-327. [https://doi.org/10.1016/S0269-7491\(99\)00175-X](https://doi.org/10.1016/S0269-7491(99)00175-X)

Morales Pinzón T., Martínez Carmona J.A. and Varela Montoya S.M. (2012). Valoración económica del efecto sobre la salud de la contaminación atmosférica por fuentes móviles en Pereira. *Scientia et Technica* 1 (52), 225-234.

Munzi S., Paoli L., Fiorini E. and Loppi S. (2012). Physiological response of the epiphytic lichen *Evernia prunastri* (L.) Ach. to ecologically relevant nitrogen concentrations. *Environmental Pollution* 171, 25-29. <https://doi.org/10.1016/j.envpol.2012.07.001>

Najmeddin A. and Keshavarzi B. (2018). Health risk assessment and source apportionment of polycyclic aromatic hydrocarbons associated with PM₁₀ and road deposited dust in Ahvaz metropolis of Iran. *Environmental Geochemistry and Health* 41, 1267-1290. <https://doi.org/10.1007/s10653-018-0209-6>

Nash III TH. (2008). Lichen sensitivity to air pollution. In: *Lichen biology* (Nash T.H. III, Ed.). Cambridge University Press, Cambridge, UK, 299-314. <https://doi.org/10.1017/CBO9780511790478.016>

Nel W.P. and Cooper C.J. (2009). Implications of fossil fuel constraints on economic growth and global warming. *Energy Policy* 37 (1), 166-180. <https://doi.org/10.1016/j.enpol.2008.08.013>

Olcese L.E. and Toselli B.M. (2002). Some aspects of air pollution in Córdoba, Argentina. *Atmospheric Environment* 36 (2), 299-306. [https://doi.org/10.1016/S1352-2310\(01\)00336-3](https://doi.org/10.1016/S1352-2310(01)00336-3)

Pandey B., Agrawal M. and Singh S. (2014). Assessment of air pollution around coal mining area: emphasizing on spatial distributions, seasonal variations and heavy metals, using cluster and principal component analysis. *Atmospheric Pollution Research* 5 (1), 79-86. <https://doi.org/10.5094/APR.2014.010>

Papini A., Tani G., Di Falco P. and Brighigna L. (2010). The ultrastructure of the development of *Tillandsia* (Bromeliaceae) trichome. *Flora-Morphology, Distribution, Functional Ecology of Plants* 205 (2), 94-100. <https://doi.org/10.1016/j.flora.2009.02.001>

Phuleria H.C., Sheesley R.J., Schauer J.J., Fine P.M. and Sioutas C. (2007). Roadside measurements of size-segregated particulate organic compounds near gasoline and diesel-dominated freeways in Los Angeles, CA. *Atmospheric Environment* 41 (22), 4653-4671. <https://doi.org/10.1016/j.atmosenv.2007.03.031>

Pignata M., Gudiño G., Wannaz E., Plá R., González C. and Carreras H. (2002). Atmospheric quality and distribution of heavy metals in Argentina employing *Tillandsia capillaris* as a biomonitor. *Environmental Pollution* 120 (1), 59-68. [https://doi.org/10.1016/S0269-7491\(02\)00128-8](https://doi.org/10.1016/S0269-7491(02)00128-8)

Pignata M., Plá R., Jasan R., Martínez M., Rodríguez J. and Wannaz E. (2007). Distribution of atmospheric trace elements and assessment of air quality in Argentina employing the lichen, *Ramalina celastri*, as a passive biomonitor: detection of air pollution emission sources. *International Journal of Environment and Health* 1 (1), 29-46. <https://doi.org/10.1504/ijenvh.2007.012223>

Puliafito S.E. and Castesana P. (2010). Emisiones de carbono del sector transporte en Argentina. *Avances en Energías Renovables y Medio Ambiente* 14, 1-8.

Puliafito S.E., Allende D., Fernández R., Castro F. and Cremades P. (2011). New approaches for urban and regional air pollution modelling and management. *Advanced Air Pollution* 429-454. <https://doi.org/10.5772/16673>

Raparthi N. and Phuleria H.C. (2021). Real-world vehicular emissions in the Indian megacity: Carbonaceous, metal and morphological characterization, and the emission factors. *Urban Climate* 39, 100955. <https://doi.org/10.1016/j.uclim.2021.100955>

Ravindra K., Sidhu M., Mor S., John S. and Pyne S. (2015). Air pollution in India: Bridging the gap between science and policy. *Journal of Hazardous, Toxic and Radioactive Waste* 20 (4), A4015003. [https://doi.org/10.1061/\(ASCE\)HZ.2153-5515.0000303](https://doi.org/10.1061/(ASCE)HZ.2153-5515.0000303)

Rodríguez J., Carreras H., Pignata M. and González C. (2007). Nickel exposure enhances the susceptibility of lichens *Usnea amblyoclada* and *Ramalina celastri* to urban atmospheric pollutants. *Archives of Environmental Contamination and Toxicology* 53 (4), 533-540. <https://doi.org/10.1007/s00244-006-0034-2>

Rodríguez J., Weller S., Wannaz E., Klumpp A. and Pignata M. (2011). Air quality biomonitoring in agricultural areas nearby to urban and industrial emission sources in Córdoba province, Argentina, employing the bioindicator *Tillandsia capillaris*. *Ecological Indicators* 11 (6), 1673-1680. <https://doi.org/10.1016/j.ecolind.2011.04.015>

Sbarato R.D. and Rubio M. (2017). Estimación de COVs emitidos por fuentes fijas y móviles en el aire de la ciudad de Córdoba, Argentina. *Revista de Salud Pública* 21 (3), 75-81. <https://doi.org/10.31052/1853.1180.v21.n3.18568>

Singh M., Jaques P.A. and Sioutas C. (2002). Size distribution and diurnal characteristics of particle-bound metals in source and receptor sites of the Los Angeles basin. *Atmospheric Environment* 36 (10), 1675-1689. [https://doi.org/10.1016/S1352-2310\(02\)00166-8](https://doi.org/10.1016/S1352-2310(02)00166-8)

Stein A. and Toselli B. (1996). Street level air pollution in Córdoba City, Argentina. *Atmospheric Environment* 30 (20), 3491-3495. [https://doi.org/10.1016/1352-2310\(96\)00097-0](https://doi.org/10.1016/1352-2310(96)00097-0)

Swietlik R., Trojanowska M., Strzelecka M. and Bocho- Janiszewska A. (2015). Fractionation and mobility of Cu, Fe, Mn, Pb and Zn in the road dust retained on noise barriers along expressway—A potential tool for determining the effects of driving conditions on speciation of emitted particulate metals. *Environmental Pollution* 196, 404-413. <https://doi.org/10.1016/j.envpol.2014.10.018>

Tan P., Li Y. and Shen H. (2017). Effect of lubricant sulfur on the morphology and elemental composition of diesel exhaust particles. *Journal of Environmental Sciences* 55, 354-362. <https://doi.org/10.1016/j.jes.2017.01.014>

Valavanidis A., Fiotakis K., Vlahogianni Th., Bakeas E.B., Triantafyllaki S., Paraskevopoulou V. and Dassenakis M. (2006). Characterization of atmospheric particulates, particle-bound transition metals and polycyclic aromatic hydrocarbons of urban air in the centre of Athens (Greece). *Chemosphere* 65, 760-768. <https://doi.org/10.1016/j.chemosphere.2006.03.052>

Verma M.K., Chauhan L.K.S., Sultana S. and Kumar S. (2014). The traffic linked urban ambient air superfine and ultrafine PM₁ mass concentration, contents of pro-oxidant chemicals, and their seasonal drifts in Lucknow, India. *Atmospheric Pollution Research* 5 (4), 677-685. <https://doi.org/10.5094/APR.2014.077>

Vouitsis E., Ntziachristos L., Pistikopoulos P., Samaras Z., Chrysikou L. and Samara C. (2009). An investigation on the physical, chemical and ecotoxicological characteristics of particulate matter emitted from light-duty vehicles. *Environmental Pollution* 157, 2320-2327. <https://doi.org/10.1016/j.envpol.2009.03.028>

Wang J.M., Jeong C.H., Hilker N., Healy R.M., Sofowote U., Debosz J. and Evans G.J. (2021). Quantifying metal emissions from vehicular traffic using real world emission factors. *Environmental Pollution* 268, 115805. <https://doi.org/10.1016/j.envpol.2020.115805>

Wang X., Wang Y., Bai Y., Wang P. and Zhao Y. (2018). An overview of physical and chemical features of diesel exhaust particles. *Journal of the Energy Institute* 92 (6), 1864-1888. <https://doi.org/10.1016/j.joei.2018.11.006>

Wannaz E.D. and Pignata M.L. (2006). Calibration of four species of *Tillandsia* as air pollution biomonitor. *Journal of Atmospheric Chemistry* 53 (3), 185-209. <https://doi.org/10.1007/s10874-005-9006-6>

Wannaz E.D., Carreras H.A., Pérez C.A. and Pignata M.L. (2006). Assessment of heavy metal accumulation in two species of *Tillandsia* in relation to atmospheric emission sources in Argentina. *Science of the Total Environment* 361 (1), 267-278. <https://doi.org/10.1016/j.scitotenv.2005.11.005>

Wannaz E., Abril G., Rodríguez J. and Pignata M. (2013). Assessment of polycyclic aromatic hydrocarbons in industrial and urban areas using passive air samplers

and leaves of *Tillandsia capillaris*. Journal of Environmental Chemical Engineering 1 (4), 1028-1035. <https://doi.org/10.1016/j.jece.2013.08.012>

Wernke M.J. (2014). Diesel exhaust. In: Encyclopedia of toxicology, 3rd ed. (Wexler P., Ed.). Elsevier, USA, 111-114.

Wintermans J. and de Mots A. (1965). Spectrophotometric characteristics of chlorophylls a and b and their phenophytins in ethanol. Biochimica et Biophysica Acta (BBA)-Biophysics including Photosynthesis 109 (2), 448-453. [https://doi.org/10.1016/0926-6585\(65\)90170-6](https://doi.org/10.1016/0926-6585(65)90170-6)

Yan J., Wang L., Fu P.P. and Yu H. (2004). Photomutagenicity of 16 polycyclic aromatic hydrocarbons from the US EPA priority pollutant list. Mutation Research/Genetic Toxicology and Environmental Mutagenesis 557 (1), 99-108. <https://doi.org/10.1016/j.mrgentox.2003.10.004>

Zechmeister H.G., Hagendorfer H., Hohenwallner D., Hanus-Illnar A. and Riss A. (2006). Analyses of platinum group elements in mosses as indicators of road traffic emissions in Austria. Atmospheric Environment 40 (40), 7720-7732. <https://doi.org/10.1016/j.atmosenv.2006.08.018>

Zhang Q., Wu L., Yang Z., Zou C., Liu X., Zhang K. and Mao H. (2016). Characteristics of gaseous and particulate pollutants exhaust from logistics transportation vehicle on real-world conditions. Transportation Research Part D: Transport and Environment 43, 40-48. <https://doi.org/10.1016/j.trd.2015.09.005>