

NEW PREDATION RECORDS ON *THOROPA MILIARIS* (ANURA: CYCLORAMPHIDAE), WITH A REVIEW OF KNOWN PREDATORSNUEVOS REGISTROS DE DEPREDACIÓN DE *THOROPA MILIARIS* (ANURA: CYCLORAMPHIDAE), CON UNA REVISIÓN DE LOS DEPREDADORES CONOCIDOSHENRIQUE J. OLIVEIRA^{1*}, CARLOS H. O. NOGUEIRA², PAULO S. AMORIM¹, ANDRÉ C. GUARALDO^{1,3} & HENRIQUE C. COSTA¹¹Universidade Federal de Juiz de Fora, Programa de Pós-Graduação em Biodiversidade e Conservação da Natureza. 36036-900, Juiz de Fora, MG, Brazil.²Universidade Federal de Mato Grosso do Sul, Programa de Pós-Graduação em Biologia Animal. 79070-900, Campo Grande, MS, Brazil.³Universidade Federal do Paraná, Departamento de Zoologia. 81531-980, Curitiba, PR, Brazil.*Correspondence: henrique.bio22@gmail.com

Received: 2022-09-29. Accepted: 2022-11-09. Published: 2022-12-29.

Editor: Katyuscia Araujo-Vieira, Brasil.

Resumen.— La rana de las piedras *Thoropa miliaris* (Cycloramphidae) es una especie saxícola endémica del Bosque Atlántico. La información sobre los posibles depredadores de esta especie es escasa en la literatura, un hecho posiblemente relacionado con la dificultad de registrar eventos de interacción entre predadores y presas. En este trabajo registramos diversos eventos predatorios oportunistas sobre *T. miliaris* por la serpiente *Chironius laevis* (Colubridae), por arañas del género *Ctenus* (Ctenidae) y por el ave *Furnarius rufus* (Furnariidae). Además, recopilamos los registros de depredación de las especies publicados hasta ahora.

Palabras claves.— déficit eltoniano, dieta, ecología trófica, Bosque Atlántico.

Abstract.— The rock river frog *Thoropa miliaris* (Cycloramphidae) is a saxicolous species endemic to the Atlantic Forest. There is scarce literature on potential predators of this species, which is likely related to the difficulty of recording its predatory interactions. In this paper, we report opportunistic records of predation on *T. miliaris* by the snake *Chironius laevis* (Colubridae), spiders of the genus *Ctenus* (Ctenidae), and the bird *Furnarius rufus* (Furnariidae). In addition, we compiled the reports of predation on *T. miliaris* from current published data.

Key words.— Atlantic Forest, diet, Eltonian shortfall, trophic ecology.

Data on predation are important to reduce the Eltonian shortfall— the lack of knowledge on biotic interactions (Hortal et al., 2015) —, but in nature, such data are recorded only occasionally (Menegucci et al., 2020). Anurans are an important part of the food web, as the prey of many vertebrate and invertebrate species (Toledo et al., 2007; Hocking & Babbitt, 2014). Brazil is the country with the greatest anuran diversity (Segalla et al., 2021; Frost, 2022), but the knowledge of biotic interactions of Brazilian anurans is deficient, even for common species like *Thoropa miliaris* (Spix, 1824), an endemic species to rocky environments in the Brazilian Atlantic Forest (Giarretta & Facure, 2004; Feio et al., 2006). In this work, we report opportunistic records of predation on *T. miliaris* by a snake,

spiders, and a bird. We also compile the literature data on predators recorded for this species.

On 18 August 2014, at 20:04, CHON found a dead male snake *Chironius laevis* (Wied, 1824) (Colubridae) of 2.10 m snout-vent length at the Tombo D'água waterfall (Fig. 1A) (district of Mocotó do Imbé, municipality of Campo dos Goytacazes, state of Rio de Janeiro, Brazil; 21.81367° S, 41.75552° W, WGS84, 282 m a.s.l.). Dissection of the snake's abdomen revealed a partially digested *T. miliaris* (Fig. 1B). Predator and prey were not collected due to the degree of decomposition, but we were able to identify both species based on their overall morphology and color pattern (Dixon et al., 1993; Feio et al., 2006).



Figura 1. A) Individuo de *Chironius laeviscolis* (Colubridae) encontrado muerto en el municipio de Campos dos Goytacazes, estado de Rio de Janeiro, Brasil. B) Individuo de *Thoropa miliaris* (Cycloramphidae) encontrado en el estómago de *C. laeviscolis*, Campos dos Goytacazes, Rio de Janeiro, Brasil. Fotos de CHON. C) Individuo de *T. miliaris* siendo presa de una araña del género *Ctenus* sp. (Ctenidae) en Campos dos Goytacazes, Rio de Janeiro, Brasil. D) Individuo de *T. miliaris* siendo presa de una araña del género *Ctenus* cf. *medius* (Ctenidae) en Campos dos Goytacazes, Rio de Janeiro, Brasil. Fotos de CHON. E) y F) Individuo de *Furnarius rufus* (Furnariidae) predando un individuo de *T. miliaris* en el municipio de Juiz de Fora, estado de Minas Gerais, Brasil. Fotos de PSA.

Figure 1. A) Specimen of *Chironius laeviscolis* (Colubridae) found dead in Campos dos Goytacazes, Rio de Janeiro, Brazil. B) *Thoropa miliaris* (Cycloramphidae) removed from the stomach of *C. laeviscolis*. Photos by CHON. C) *Thoropa miliaris* being preyed upon by a *Ctenus* sp. (Ctenidae) spider in Campos dos Goytacazes, Rio de Janeiro, Brazil. D) *Thoropa miliaris* being preyed on by a *Ctenus* cf. *medius* (Ctenidae) in Campos dos Goytacazes, Rio de Janeiro, Brazil. Photos by CHON. E) and F) *Furnarius rufus* (Furnariidae) preying on *T. miliaris* in Juiz de Fora, Minas Gerais, Brazil. Photos by PSA.

At this same locality, CHON recorded two additional predation events on *T. miliaris* by spiders of the family Ctenidae: *Ctenus* sp. and *Ctenus* cf. *medius* Keyserling, 1891, identified by the spider taxonomist Leonardo Carvalho, Universidade Federal do Piauí, Brazil. The first record was on 19 February 2020, at 18:35 (21.82724° S, 41.74726° W; WGS84; 174 m a.s.l.; Fig. 1C), and the second on 16 June 2021, at 19:46 (21.81449° S, 41.75748° W; Fig. 1D). The *T. miliaris* individuals were collected and deposited in the Coleção Herpetológica do Norte Fluminense (CHNF 0897 and CHNF 0975, respectively).

On 21 February 2020, at 12:10, PSPA recorded a *T. miliaris* being preyed by a juvenile Rufous hornero, *Furnarius rufus* (Gmelin,

1788) (Furnariidae) in an urban environment near the hospital of the Universidade Federal de Juiz de Fora, municipality of Juiz de Fora, state of Minas Gerais, Brazil (21.78553° S, 43.36787° W, WGS84, 878 m a.s.l.; Fig. 1E). The *T. miliaris* was initially observed beside a paved road with little traffic, when the bird appeared and repeatedly pecked the frog on its dorsum, venter, and head. The frog attempted to escape several times, but the bird held it with its beak. The bird immobilized the frog and swallowed it alive (Fig. 1F). The bird was subsequently observed for approximately 40 minutes from a distance of 30 cm, but no regurgitation was recorded (see record video at <https://doi.org/10.5281/zenodo.6954386>). Prey identity was confirmed by

Tabla 1. Lista de depredadores de *Thoropa miliaris* (Anura: Cycloramphidae) basada en los datos propios de los autores y en una revisión de la literatura.

Table 1. List of predators of *Thoropa miliaris* (Anura: Cycloramphidae) based on the authors' data and literature review.

Predator identification	Life stage of prey	Environment	Source
Arachnida			
<i>Cteniza</i> sp. Latreille, 1829	Post-metamorphic	Forest	Pertel et al., 2010
<i>Ctenus</i> cf. <i>medius</i> Keyserling, 1891	Post-metamorphic	Forest	This study
<i>Ctenus</i> sp.	Post-metamorphic	Forest	This study
<i>Trechalea</i> sp.	Tadpole	Forest	Diniz et al., 2020
Anura			
<i>Cycloramphus boraceiensis</i> (Heyer, 1983)	Tadpole	Forest	Hartmann et al., 2003
<i>Thoropa miliaris</i> (Spix, 1824) (Tadpole)	Egg	Forest	Giaretta & Facure, 2004
Aves			
<i>Furnarius rufus</i> (Gmelin, 1788)	Post-metamorphic	Urban	This study
Insecta			
<i>Tropisternus</i> sp. (Larvae)	Tadpole	Forest	Siqueira et al., 2006
Squamata			
<i>Bothrops jararaca</i> (Wied, 1824)	Post-metamorphic	Forest	Sazima, 1992
<i>Chironius bicarinatus</i> (Wied, 1820)	Post-metamorphic	Forest	Roberto & Souza, 2020
<i>Chironius flavolineatus</i> (Jan, 1863)	Post-metamorphic	Not informed	Pinto et al., 2008; Roberto & Souza, 2020
<i>Chironius fuscus</i> (Linnaeus, 1758)	Post-metamorphic	Forest	Marques & Sazima, 2004; Hartmann, 2005; Roberto & Souza, 2020
<i>Chironius laevis</i> (Wied, 1824)	Post-metamorphic	Forest	This study
<i>Erythrolamprus miliaris</i> (Linnaeus, 1758)	Post-metamorphic	Forest	Albolea, 1998; Marques & Sazima, 2004; Mônico et al., 2016; Burg, 2020
<i>Erythrolamprus poecilogyrus</i> (Wied, 1825)	Post-metamorphic	Not informed	Pinto & Fernandes, 2004
<i>Tropidurus torquatus</i> (Wied, 1825)	Unknown - Museum specimen	Urban	Mônico & Mônico, 2022

Diego J. Santana (Universidade Federal do Mato Grosso do Sul, Brazil).

We reviewed literature records of predators of *T. miliaris* in Google Scholar on 28 April 2022, searching for the expression “predation OR predator OR prey AND ‘*Thoropa miliaris*’” in English, Spanish (depredación OR presa OR depredador), and Portuguese (predação OR presa OR predador), and found 267 results. Of these, only eight showed *T. miliaris* as the prey. We also searched issues of Herpetological Review from 1967-2022, finding four additional records. In addition, we searched the authors' pdf libraries. All reports are shown in Table 1.

Snakes are among the most important predators of post-metamorphic anurans (Toledo et al., 2007). Most reports of predation on *T. miliaris* were by *Chironius* spp., a genus of frog-eating species (Marques et al., 2016). Conversely, lizards are the least frequent vertebrate predators of frogs (Toledo et al., 2007). We found a single record of *T. miliaris* being preyed by a lizard (Mônico & Mônico, 2022): *Tropidurus torquatus*, a saxicolous species with a generalist diet (Gomides et al., 2013; Guimarães & Sbrek-Araujo, 2018). We also found two records of frogs preying on *T. miliaris*: i) tadpoles being preyed upon by *Cycloramphus boraceiensis* (Hartmann et al., 2003), and ii) egg cannibalism by conspecific tadpoles (Giaretta & Facure, 2004). Tadpoles were also preyed upon by larvae of the beetle *Tropisternus* sp., the only record of an insect preying on *T. miliaris*, despite Coleoptera being one of the most recorded invertebrate groups of anuran predators (Toledo, 2005). Arachnida is another frequently recorded invertebrate group of anuran predators (Toledo, 2005), although we found only two predation events, by a *Cteniza* sp. and *Trechalea* sp. (Pertel et al., 2010; Diniz et al., 2020).

Our records improve the information about the predation of post-metamorphic specimens of *T. miliaris* by the spider genus *Ctenus*, whose species often actively forage for anurans in leaf litter (Salvestrini & Gasnier, 2001; Menin et al., 2005). We also report the first avian predator of *T. miliaris*, the Rufous hornero, *Furnarius rufus*. Birds are common predators of post-metamorphic anurans (Toledo et al., 2007), but the absence of records for *T. miliaris* highlights the knowledge gaps on the biotic interactions of this relatively common, conspicuous, and abundant species.

Acknowledgments. - The authors are grateful to Diego J. Santana (Universidade Federal de Mato Grosso do Sul) and Leonardo S. Carvalho (Universidade Federal do Piauí) for confirming species identification. To Juan Carrillo for the review of the Spanish text and Ross MacCulloch for the review of the English text. To the

editor Katyuscia Araujo-Vieira and two anonymous reviewers for suggestions in a first version of this article. This study is partially funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) – Finance Code 001 (Master's scholarship to HJO and CHON, and PhD scholarship to PSA).

CITED LITERATURE

- Albolea, A.B.P. 1998. Padrões de atividade em serpentes não peçonhentas de interesse médico: *Helicops modestus* (Colubridae: Xenodontinae) e *Liophis miliaris* (Colubridae: Xenodontinae) e sua relação com a epidemiologia. Master Dissertation. Universidade Guarulhos, São Paulo. Brasil.
- Burg, M.P.V.D. 2020. How to source and collate natural history information: a case study of reported prey items of *Erythrolamprus miliaris* (Linnaeus, 1758). Herpetology Notes 13:739-746.
- Diniz, H.S., C.L. Assis, J.J.M. Guedes & R.N. Feio. 2020. *Thoropa miliaris* (Military River Frog). Predation. Herpetological Review 51:571-572.
- Dixon, J.R., J.A. Wiest & J.M. Cei. 1993. Revision of the Neotropical snake genus *Chironius* Fitzinger (Serpentes, Colubridae). Museo Regionale di Scienze Naturali - Monografie 13:1-279.
- Feio, R.N., M.F. Napoli & U. Caramaschi. 2006. Considerações taxonômicas sobre *Thoropa miliaris* (Spix, 1824), com revalidação e redescritção de *Thoropa taophora* (Miranda-Ribeiro, 1923) (Amphibia, Anura, Leptodactylidae). Arquivos do Museu Nacional 64:41-60.
- Frost, D.R. 2022. Amphibian Species of the World: an Online Reference. Version 6.1. <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA. [Accessed on 01 August 2022]
- Giaretta, A.A. & K.G. Facure. 2004. Reproductive ecology and behavior of *Thoropa miliaris* (Spix, 1824) (Anura, Leptodactylidae, Telmatobiinae). Biota Neotropica 4:1-10.
- Gomides, S.C., L.B. Ribeiro, V.M. Peters & B.M. Sousa. 2013. Feeding and reproduction ecology of the lizard *Tropidurus torquatus* (Squamata: Tropiduridae) in a rock outcrop area in southeastern Brazil. Revista Chilena de Historia Natural 86:137-151.
- Guimarães, L.J. & A.C. Sbrek-Araujo. 2018. Attempted predation of *Ameivula nativo* by *Tropidurus torquatus* (Reptilia: Squamata,

- Sauria) with ingestion of tail fragments in southeastern Brazil. *Herpetology Notes* 11:709-713.
- Hartmann, P.A. 2005. História natural e ecologia de duas taxocenoses de serpentes na Mata Atlântica. Doctoral Thesis. Instituto de Biociências da Universidade Estadual Paulista Júlio de Mesquita Filho. Brasil.
- Hartmann, M.T., P.A. Hartmann, C.P.A. Prado & P.C.A. Garcia. 2003. *Cycloramphus boraceiensis* (flattened waterfall frog). Defensive behavior. *Herpetological Review* 34:50.
- Hocking, D.J. & K.J. Babbitt. 2014. Amphibian contributions to ecosystem services. *Herpetological Conservation and Biology* 9:1-17.
- Hortal, J., F. de Bello, J.A.F. Diniz-Filho, T.M. Lewinsohn, J.M. Lobo & R. J. Ladle. 2015. Seven shortfalls that beset large-scale knowledge of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* 46:523-549.
- Marques, O.A.V. & I. Sazima. 2004. História natural dos répteis da estação ecológica Juréia-Itatins. Pp. 257-277 In O.A. V. Marques & W. Duleba (Eds.), *Estação Ecológica Juréia-Itatins. Ambiente Físico, Flora e Fauna. Holos. Ribeirão Preto, Brasil.*
- Marques, R., K. Mebert, É. Fonseca, D. Rödder, M. Solé & M.S. Tinoco. 2016. Composition and natural history notes of the coastal snake assemblage from Northern Bahia, Brazil. *ZooKeys* 611:93-142.
- Menegucci, R.C., E. Muscat, R.M. Tanaka & D. Stuginski. 2020. Predation on *Hylodes phyllodes* (Anura: Hylodidae) by the harvestman *Heteromitobates discolor* (Arachnida: Opiliones). *Phyllomedusa: Journal of Herpetology* 19:121-124.
- Menin, M., D.J. Rodrigues & C.S. Azevedo. 2005. Predation on amphibians by spiders (Arachnida, Araneae) in the Neotropical region. *Phyllomedusa: Journal of Herpetology* 4:39-47.
- Mônico, A.T., E.H. Krause & R.B. Ferreira. 2016. *Erythrolamprus miliaris* (Cobra D'água; Military Ground Snake). Prey. *Herpetological Review* 47:311.
- Mônico, A.T. & A.T. Mônico. 2022. *Tropidurus torquatus* (Calango; Amazon Lava Lizard). Diet. *Herpetological Review* 53:332.
- Pertel, W., R.L. Teixeira & R.B. Ferreira. 2010. Comparison of diet and use of bromeliads between a bromelicolous and a bromeligenous anuran at an inselberg in the southeastern of Brazil. *Caldasia* 32:149-159.
- Pinto, R.R. & R. Fernandes. 2004. Reproductive biology and diet of *Liophis poecilogyrus poecilogyrus* (Serpentes, Colubridae) from southeastern Brazil. *Phyllomedusa: Journal of Herpetology* 3:9-14.
- Pinto, R.R., R. Fernandes & O.A.V. Marques. 2008. Morphology and diet of two sympatric colubrid snakes, *Chironius flavolineatus* and *Chironius quadricarinatus* (Serpentes: Colubridae). *Amphibia-Reptilia* 29:149-160.
- Roberto, I.J. & A.R. Souza. 2020. Review of prey items recorded for snakes of the genus *Chironius* (Squamata, Colubridae), including the first record of *Osteocephalus* as prey. *Herpetology Notes* 13:1-5.
- Salvestrini, F.M.D. & T.R. Gasnier. 2001. Differences in the activity of juveniles, females and males of two hunting spiders of the genus *Ctenus* (Araneae, Ctenidae): active males or inactive females? *The Journal of Arachnology* 29:276-278.
- Sazima, I. 1992. Natural history of the jararaca pitviper, *Bothrops jararaca*, in southeastern Brazil. Pp. 199-216 In J.A. Campbell & E. D. Brodie (Eds.), *Biology of the Pitvipers*. Selva Publishing.
- Segalla, M.V., B. Berneck, C. Canedo, U. Caramaschi, C.A.G. Cruz, P.C.A. Garcia, T. Grant, C.F.B. Haddad, A.C.C. Lourenço, S. Mângia, T. Mott, L.B. Nascimento, L.F. Toledo, F.P. Werneck & J. A. Langone. 2021. List of Brazilian amphibians. *Herpetologia Brasileira* 10:121-216.
- Siqueira, C.C., C.V. Ariani, M. Van-Sluys & C.F.D. Rocha. 2006. *Thoropa miliaris* (Rock River Frog). Tadpole predation. *Herpetological Review* 37:451-452.
- Toledo, L.F. 2005. Predation of juvenile and adult anurans by invertebrates: current knowledge and perspectives. *Herpetological Review* 36:395-399.
- Toledo, L.F., R.S. Ribeiro & C.F. Haddad. 2007. Anurans as prey: an exploratory analysis and size relationships between predators and their prey. *Journal of Zoology* 271:170-177.

