

## *uvby* – $\beta$ PHOTOELECTRIC PHOTOMETRY OF THE OPEN CLUSTER NGC 2343<sup>1</sup>

J. H. Peña<sup>2</sup> and A. Martínez<sup>3</sup>

Received 2013 July 25; accepted 2014 January 10

### RESUMEN

A partir de fotometría fotoeléctrica *uvby* –  $\beta$  del cúmulo abierto NGC 2343 (20 estrellas) realizamos la determinación de distancias de cada una y, por ende, la pertenencia de las estrellas al cúmulo. Asimismo se determinaron el enrojecimiento y la edad.

### ABSTRACT

From *uvby* –  $\beta$  photometry of the open cluster NGC 2343 (20 stars) we were able to determine membership of the stars to the cluster and fix its age and reddening.

*Key Words:* open clusters and associations: individual (NGC 2343) — techniques: photometric

### 1. MOTIVATION

As a continuation of a study of open clusters in *uvby* –  $\beta$ , we now present a membership study of the open cluster NGC 2343.

Despite the fact that there are a large number of articles devoted to open clusters in general and to NGC 2343 in particular, there are very few previous papers that present photometric data. The last one was that of Clariá (1972).

For NGC 2343, WEBDA<sup>4</sup> reports a distance of 1056 pc (DM of 10.48) with reddening  $E(B - V)$  of 0.118, a log age of 7.104 and a determined metallicity  $[\text{Fe}/\text{H}]$  of  $-0.30$ . It has been amply studied by *UBV* photometry, with 64 measured stars. Spectroscopy for the stars in its direction involves a sample of merely three stars. In the WEBDA compilation these measurements are reported as done by Mofat (1974) who states it is a moderately young open cluster in Canis Major. The previous *uvby* –  $\beta$  photometry was carried out for only two stars reported by Gray & Olsen (1991). McSwain & Gies (2005) determined a reddening  $E(b - y)$ , a DM  $V_0 - M_V$  and a log age of 0.11, 10.12, and 8, respectively.

### 2. OBSERVATIONS

These were all done at the Observatorio Astronómico Nacional, SPM, Mexico. The 1.5 m telescope to which a spectrophotometer was attached was utilized. The observing season was from December 2–13, 2009, but the cluster was observed along with standard stars to transform the cluster measurements into the standard *uvby* –  $\beta$  photometric system on only one night, December 10th. The identification utilized the ID chart provided by Clariá (1972); we followed his numerical identification system. Despite the large extension shown by Clariá (1972) and WEBDA, the region we observed was centered at  $(X, Y) = (0, 0)$  in the WEBDA chart and extended only in 200 in WEBDA units, in the region in which the cluster shows a larger concentration of stars. Most of the brightest stars in this zone were observed in this study.

#### 2.1. Data acquisition

During all the observational nights the following procedure was utilized: each star was measured, with at least five ten-second integrations of the star and one ten-second integration of the sky for the *uvby* filters and the narrow and wide filters that define  $H\beta$ . Individual uncertainties were determined by calculating the standard deviations of the fluxes in each filter for each star. The percent error in each measurement is, of course, a function of both the spectral

<sup>1</sup>Based on observations collected at the San Pedro Mártir Observatory, Mexico.

<sup>2</sup>Instituto de Astronomía, Universidad Nacional Autónoma de México, México.

<sup>3</sup>Universidad de Panamá, Panamá.

<sup>4</sup><http://webda.physics.muni.cz/>.

type and the brightness of each star, but they were observed long enough to a secure sufficient number of photons ( $N$ ) to get a signal to noise ratio of accuracy of  $N/\sqrt{(N)}$  of at least 0.01 mag. A series of standard stars was also observed on the night the cluster was observed, so as to transform the data into the standard system. Generally the standard stars were counted to much higher values. These standard stars will serve to evaluate the goodness of the night when comparing the calculated values to those reported in the literature. Atmospheric extinction coefficients were not calculated but taken from the mean values determined by Schuster & Parrao (2001) and Schuster, Parrao, & Guichard (2002). In this study they consider the individual nightly determinations of the atmospheric extinction above San Pedro Mártir from 82 nights of 13-color observations from 1980 to 1983 and from 287 nights of *uvby* observations from 1984 to 1999.

### 2.2. Data reduction

The reduction was done with the numerical package NABAPHOT (Arellano-Ferro & Parrao 1988) which reduces the data into a standard system. The chosen standard system was that defined by the standard values of Grönbech & Olsen (1976, 1977) and Olsen (1983) and the transformation equations are those defined by Crawford & Barnes (1970) and by Crawford & Mander (1966) which are shown below.

$$\begin{aligned} V &= A + B(b-y)_{\text{st}} + y_{\text{inst}} , \\ (b-y)_{\text{st}} &= C + D(b-y)_{\text{inst}} , \\ m_{1\text{st}} &= E + Fm_{1\text{ inst}} + G(b-y)_{\text{inst}} , \\ c_{1\text{st}} &= H + Im_{1\text{ inst}} + J(b-y)_{\text{inst}} , \\ \beta_{1\text{ st}} &= K + L\beta_{1\text{ inst}} . \end{aligned}$$

In these equations the coefficients  $D$ ,  $F$ ,  $I$  and  $L$  are the slope coefficients for  $(b-y)$ ,  $m_1$ ,  $c_1$  and  $\beta$ , respectively. The coefficients  $B$ ,  $G$  and  $J$  are the color terms of  $V$ ,  $m_1$ , and  $c_1$ . The standard stars were taken from Grönbech & Olsen (1976, 1977) and Olsen (1983), but some of the standard bright stars were also taken from the Astronomical Almanac for the year 2006 (2005).

### 2.3. Obtained data

The averaged transformation coefficients are listed in the corresponding equations. The standard deviations are the following:  $V$ : 0.0122;  $(b-y)$ : 0.007;  $m_1$ : 0.014;  $c_1$ : 0.028 and  $\beta$ : 0.0193. Emphasis is made on the large range of the standard stars in the magnitude and color values:  $V$ : (5.2, 8.8);

$(b-y)$ : (-0.00, 0.79);  $m_1$ : (0.123, 0.659);  $c_1$ : (0.153, 1.052) and  $\beta$ : (2.53, 2.89). A set of 16 standard stars were measured.

$$\begin{aligned} V &= 19.0809 + 1.00(b-y)_{\text{st}} + y_{\text{inst}} , \\ (b-y)_{\text{st}} &= 1.5329 + 0.9856(b-y)_{\text{inst}} , \\ m_{1\text{st}} &= -1.2977 + 1.0125m_{1\text{ inst}} + 0.0283(b-y)_{\text{inst}} , \\ c_{1\text{st}} &= -0.1918 + 1.0526m_{1\text{ inst}} + 0.1499(b-y)_{\text{inst}} , \\ \beta_{1\text{ st}} &= 2.4351 + -1.0763\beta_{1\text{ inst}} . \end{aligned}$$

The errors of this night were evaluated by means of the standard stars observed. These uncertainties were calculated through the differences between the standard values reported in the literature and the transformed values in magnitude and colors, for  $(V, b-y, m_1, c_1 \text{ and } H\beta)$ . These differences were evaluated and the standard deviations provided a measure of the quality of the data and the night. Table 1 lists these differences. For the whole sample the standard deviations of the differences were calculated and are listed at the bottom of the table. As can be seen, the quality of the night was good. The standard deviations for  $V$ ,  $(b-y)$ ,  $m_1$ ,  $c_1$  and  $\beta$  are 0.011, 0.005, 0.011, 0.016 and 0.019, respectively.

Uncertainties for each star were evaluated in the following way: for each filter, mean values and standard deviations were calculated. With these values the percentual errors are given as standard deviation/average value for each filter. Finally, the uncertainties for the color indexes were evaluated through propagation of errors from the definitions for each index as  $\delta m_1 = \delta v + 2\delta b + \delta y$  and  $\delta c_1 = \delta u + 2\delta b + \delta v$ . All the stars were measured five times with integration times of ten seconds for the star, and three times for the sky. The only exceptions were stars 23 and 29 which, due to their faintness, were measured ten times. Table 2 lists the photometric values of the observed stars for this cluster. In this table, Column 1 reports the ID of the stars as listed by WEBDA (Paunzen & Mermilliod 2007), Columns 2 to 5 the Strömgren values  $V$ ,  $(b-y)$ ,  $m_1$  and  $c_1$ , respectively; Column 6, the  $\beta$  values. The calculated uncertainties are listed in Columns 7 to Column 10. Column 11 lists the spectral types determined from the Strömgren photometry and the last column, the reported spectroscopy for the one star in common with the three measured stars compiled by WEBDA from The Henry Draper catalogue (Cannon & Pickering 1918). Hence, the spectral types determined and reported here are new and valuable data.

TABLE 1  
DIFFERENCES BETWEEN LITERATURE VALUES AND CALCULATED  
VALUES FOR THE OBSERVED STANDARD STARS

| ID       | $V$   | $(b - y)$ | $m_1$ | $c_1$ | $\delta V$ | $\delta(b - y)$ | $\delta m_1$ | $\delta c_1$ |
|----------|-------|-----------|-------|-------|------------|-----------------|--------------|--------------|
| BS1321   | 6.960 | 0.422     | 0.244 | 0.294 | 0.020      | 0.000           | 0.018        | -0.016       |
| HD36003  | 7.624 | 0.629     | 0.659 | 0.188 | -0.005     | -0.014          | 0.026        | -0.010       |
| HD64702  | 8.500 | 0.790     | 0.638 | 0.373 | -0.011     | 0.008           | -0.029       | 0.023        |
| HD42807  | 6.440 | 0.415     | 0.228 | 0.292 | -0.004     | 0.003           | 0.007        | -0.004       |
| HD57006  | 5.915 | 0.340     | 0.168 | 0.472 | 0.004      | 0.005           | -0.004       | 0.003        |
| HD60803  | 5.916 | 0.377     | 0.186 | 0.387 | 0.017      | 0.005           | -0.002       | 0.018        |
| HD77354  | 7.943 | 0.443     | 0.202 | 0.360 | 0.017      | 0.000           | 0.017        | 0.029        |
| HD85217  | 6.250 | 0.305     | 0.162 | 0.415 | 0.017      | 0.003           | -0.001       | 0.039        |
| BS3555   | 5.450 | 0.084     | 0.205 | 0.972 | -0.003     | -0.004          | -0.010       | -0.021       |
| HD51530  | 6.203 | 0.345     | 0.137 | 0.401 | 0.001      | 0.007           | -0.005       | -0.007       |
| HD57517  | 6.568 | 0.342     | 0.174 | 0.390 | -0.011     | -0.011          | -0.005       | 0.005        |
| HD59984  | 5.933 | 0.356     | 0.123 | 0.336 | -0.018     | -0.007          | -0.014       | 0.003        |
| BS3297   | 5.600 | 0.311     | 0.138 | 0.400 | -0.004     | 0.000           | -0.012       | 0.035        |
| BS3538   | 6.010 | 0.410     | 0.239 | 0.325 | -0.015     | -0.006          | 0.002        | -0.021       |
| BS4378   | 6.660 | 0.024     | 0.190 | 1.052 | -0.011     | 0.003           | 0.006        | 0.000        |
| $\sigma$ |       |           |       |       | 0.011      | 0.005           | 0.011        | 0.016        |

TABLE 2  
 $uvby - \beta$  PHOTOELECTRIC PHOTOMETRY OF THE OPEN CLUSTER NGC 2343

| ID       | $V$    | $(b - y)$ | $m_1$  | $c_1$ | $\beta$ | $\delta V$ | $\delta(b - y)$ | $\delta m_1$ | $\delta(c_1)$ | SpTyp | SpTyp |
|----------|--------|-----------|--------|-------|---------|------------|-----------------|--------------|---------------|-------|-------|
| N2343-03 | 9.511  | 0.045     | 0.091  | 0.795 | 2.718   | 0.005      | 0.008           | 0.013        | 0.015         | B7V   | A0    |
| N2343-04 | 10.084 | 0.098     | 0.069  | 0.421 | 2.667   | 0.003      | 0.005           | 0.009        | 0.012         | BB5V  |       |
| N2343-05 | 10.120 | 0.151     | 0.085  | 0.872 | 2.772   | 0.009      | 0.014           | 0.019        | 0.022         | BB8V  |       |
| N2343-06 | 10.350 | 0.123     | 0.082  | 0.563 | 2.678   | 0.002      | 0.005           | 0.012        | 0.018         | B6V   |       |
| N2343-10 | 10.789 | 0.143     | 0.090  | 0.658 | 2.707   | 0.004      | 0.008           | 0.016        | 0.022         | B5V   |       |
| N2343-11 | 8.488  | 0.796     | 0.350  | 0.278 | 2.595   | 0.029      | 0.071           | 0.197        | 0.268         | M0V   |       |
| N2343-15 | 11.353 | 0.695     | 0.240  | 0.399 | 2.560   | 0.008      | 0.012           | 0.034        | 0.059         | K2V   |       |
| N2343-17 | 11.493 | 0.147     | 0.176  | 1.041 | 2.827   | 0.012      | 0.019           | 0.034        | 0.033         | A4V   |       |
| N2343-18 | 11.579 | 0.186     | 0.143  | 1.054 | 2.857   | 0.014      | 0.019           | 0.030        | 0.036         | B4V   |       |
| N2343-19 | 11.648 | 0.719     | -0.056 | 0.149 | 2.628   | 0.012      | 0.017           | 0.031        | 0.049         | B     |       |
| N2343-20 | 7.263  | -0.071    | 0.095  | 0.215 | 2.767   | 0.014      | 0.021           | 0.032        | 0.034         | B2V   |       |
| N2343-23 | 11.914 | 0.466     | 0.146  | 0.421 | 2.601   | 0.010      | 0.016           | 0.024        | 0.049         | G0V   |       |
| N2343-24 | 12.022 | 0.214     | 0.090  | 1.020 | 2.815   | 0.013      | 0.024           | 0.043        | 0.044         | B9V   |       |
| N2343-26 | 12.154 | 0.211     | 0.158  | 0.702 | 2.807   | 0.131      | 0.269           | 0.516        | 0.530         | A9V   |       |
| N2343-27 | 12.443 | 0.318     | 0.120  | 0.853 | 2.770   | 0.010      | 0.017           | 0.031        | 0.042         | A7V   |       |
| N2343-28 | 12.459 | 0.246     | 0.115  | 0.958 | 2.853   | 0.026      | 0.038           | 0.065        | 0.094         | B5V   |       |
| N2343-29 | 12.626 | 0.452     | 0.114  | 0.438 | 2.616   | 0.033      | 0.052           | 0.085        | 0.092         | F8V   |       |
| N2343-30 | 12.625 | 0.227     | 0.202  | 1.005 | 2.900   | 0.023      | 0.044           | 0.070        | 0.116         | A5Vp  |       |
| N2343-32 | 12.834 | 0.296     | 0.158  | 0.978 | 2.933   | 0.058      | 0.067           | 0.099        | 0.101         | A5Vp  |       |
| N2343-34 | 12.826 | 0.196     | 0.214  | 0.924 | 2.783   | 0.034      | 0.047           | 0.067        | 0.066         | A5Vp  |       |




Fig. 1. Magnitude difference vs. magnitude for NGC 2343.

### 3. COMPARISON WITH OTHER PHOTOMETRIES

To test the goodness of our observations a comparison with other previous measurements was done. Since practically no  $uvby - \beta$  data have been obtained for these clusters, a comparison of our values is made with the available  $UBV$  photometry.

There are two references with observations of this cluster: (Clariá 1972; McSwain & Gies 2005) that merely compile previously observed data. This cluster was compared with those  $UBV$  values reported by WEBDA from Clariá (1972). The set of intersection between the two sets is constituted by 20 stars and there are two (11 and 20) which show a remarkable difference. The rest of the sample is in good agreement. The average of the  $V$  differences without considering these two stars is  $-0.042$  mag. The linear fit among the two sets gave  $V_{pp} = -0.114 + 1.006V_{Clariá}$  with a correlation coefficient of 0.996 and a standard deviation of 0.096, a very good agreement. The difference between the magnitudes obtained by Clariá (1972) and the present paper is shown schematically in Figure 1 in which the zero point shift of  $-0.114$  is clearly seen. Only two stars, 17 and 26, have large differences. The two stars with large differences were not included in the plot. The color indexes were related by the relation  $(b - y) = -0.073 + 1.518(B - V)$  with a correlation coefficient of 0.975 and a standard deviation of 0.069. In the fit the same two stars (11 and 20) were discarded making it evident that there was an unexplained misidentification or a variable nature of the stars. The latter is quite possible for NGC2343-11 which has a spectral type of M0V, and could have

remarkable variations. With respect to the spectral classification of the stars, as was mentioned before, it was carried out by locating the unreddened indexes in the  $[m_1] - [c_1]$  plane, comparing with the stars determined by Golay (1974), and corroborated from the results of a study of Peña & Sareyan (2006) for the open cluster Alpha Per which has stars with well-determined spectral types. From the comparison, seven stars are of spectral types earlier than A0 and three of late spectral types; the rest lie between A3V and F7V. Two, 30 and 34, lie in the region which corresponds to the Ap stars. Star 32 is located in the boundary between the Ap and normal stars. In Table 2 the photometrically determined spectral class has been indicated. That spectral type compiled by WEBDA is also presented.

### 4. METHODOLOGY

In order to determine the physical characteristics of the member stars in each cluster, the following procedure was undertaken:

The evaluation of the reddening was done by first establishing, as stated above, to which spectral class the stars belong: early types (B and early A stars) or late types (late A and F stars); the late class stars (later than G) were not considered in the analysis because no reddening determination calibration can be done for MS stars.

The reddening determination was obtained from the spectral types through the Strömgren photometry. The application of the calibrations for each spectral type either Shobbrook (1984) for O and early A type:

$$(b - y)_0 = -0.1146 - 0.0805c_1 + 0.0616c_1^2 + 0.2719c_1^3 + 0.7801c_1^4 + 0.4679c_1^5,$$

or Nissen (1988) for late A and F stars respectively, allows us to determine their reddening and hence, the desired physical properties of the stars:

$$(b - y)_0 = K + 1.11\Delta\beta + 2.7\Delta\beta^2 - 0.05\delta c_0 - (0.1 + 3.6\Delta\beta)\delta m_0,$$

for A stars (with  $2.72 < \beta < 2.88$ ):

$$(b - y)_0 = 2.946 - 1.00\beta - 0.1\delta c_0,$$

with

$$\begin{aligned} \Delta\beta &= 2.72 - \beta, \\ m_0 &= m_{1,Hyades(\beta)} - m_0, \\ c_0 &= c_0 - c_{1,std(\beta)}. \end{aligned}$$

TABLE 3

REDDENING AND UNREDDENED PARAMETERS OF THE OPEN CLUSTER NGC 2343

| ID       | $E(b-y)$ | $(b-y)_0$ | $m_0$ | $c_0$ | $\beta$ | $V_0$ | $M_V$ | DM   | DST  | Mbr |
|----------|----------|-----------|-------|-------|---------|-------|-------|------|------|-----|
| N2343-11 | 0.279    | 0.517     | 0.434 | 0.222 | 2.595   | 7.29  | 4.83  | 2.4  | 31   | NM  |
| N2343-20 | 0.023    | -0.094    | 0.102 | 0.211 | 2.767   | 7.17  | -0.45 | 7.6  | 334  | NM  |
| N2343-26 | 0.063    | 0.148     | 0.177 | 0.689 | 2.807   | 11.88 | 3.57  | 8.3  | 460  | NM  |
| N2343-23 | 0.103    | 0.363     | 0.177 | 0.400 | 2.601   | 11.47 | 2.75  | 8.7  | 553  | NM  |
| N2343-29 | 0.119    | 0.333     | 0.150 | 0.414 | 2.616   | 12.12 | 2.78  | 9.3  | 737  | M:  |
| N2343-05 | 0.186    | -0.035    | 0.141 | 0.837 | 2.772   | 9.32  | -0.12 | 9.4  | 772  | M:  |
| N2343-18 | 0.194    | -0.008    | 0.201 | 1.017 | 2.857   | 10.74 | 0.82  | 9.9  | 967  | M   |
| N2343-27 | 0.155    | 0.163     | 0.167 | 0.822 | 2.770   | 11.78 | 1.65  | 10.1 | 1057 | M   |
| N2343-03 | 0.084    | -0.039    | 0.116 | 0.779 | 2.718   | 9.15  | -1.01 | 10.1 | 1078 | M   |
| N2343-28 | 0.275    | -0.029    | 0.197 | 0.906 | 2.853   | 11.28 | 0.85  | 10.4 | 1217 | M   |
| N2343-17 | 0.049    | 0.098     | 0.191 | 1.031 | 2.827   | 11.28 | 0.78  | 10.5 | 1257 | M   |
| N2343-24 | 0.232    | -0.018    | 0.160 | 0.976 | 2.815   | 11.02 | 0.34  | 10.6 | 1370 | M   |
| N2343-10 | 0.191    | -0.048    | 0.147 | 0.622 | 2.707   | 9.97  | -1.0  | 10.9 | 1562 | M   |
| N2343-04 | 0.170    | -0.072    | 0.120 | 0.389 | 2.667   | 9.35  | -1.62 | 10.9 | 1568 | M   |
| N2343-19 | 0.834    | -0.115    | 0.194 | -0.01 | 2.628   | 8.06  | -3.01 | 11.0 | 1635 | M   |
| N2343-06 | 0.179    | -0.056    | 0.136 | 0.529 | 2.678   | 9.58  | -1.51 | 11.1 | 1655 | M   |
| N2343-34 | 0.051    | 0.145     | 0.229 | 0.914 | 2.783   | 12.61 | 1.21  | 11.4 | 1904 | M   |

The absolute magnitude is

$$M_V = M_{V,ZAMS(\beta)} - f\delta c_0.$$

In these calibrations reddening, absolute magnitude and distance were determined through the calibrations described by Nissen (1988). In this procedure, the intrinsic color index  $(b-y)_0$  was first calculated by the expression for F stars (with  $2.59 < \beta < 2.72$ ).

## 5. RESULTS

The application of the above mentioned numerical packages gave the results listed in Table 3 for NGC 2343, in which the ID, reddening, unreddened indexes, absolute magnitude, distance modulus, DM and distance, in parsecs, are listed in consecutive columns. The distance modulus distribution has been plotted in a histogram, Figure 2. A Gaussian fit was calculated and the mean value and the standard deviation were obtained from the Gaussian fit. We consider those stars within one sigma from the mean to be members. The results from this calibration gave a mean distance modulus of  $10.4 \pm 0.9$  mag. From that we assigned the memberships listed in the last columns of the table. There are four stars at a small distance, two that are marginal members, denoted by M: and eleven stars that were established to be cluster members.




Fig. 2. Histogram of the distance modulus ( $X$  axis, in magnitudes) found for the stars in the direction of NGC 2343.

Once the member stars have been defined, a mean reddening determination can be obtained. The numerical value is of  $E(b-y)$  equal to  $0.157 \pm 0.07$ .

TABLE 4  
COMPILED CHARACTERISTICS FOR NGC 2343

|                            | WEBDA     | Newly Determined |
|----------------------------|-----------|------------------|
| Right Ascension            | 07 08 06  |                  |
| Declination                | −10 37 00 |                  |
| Galactic longitude         | 224.268   |                  |
| Galactic latitude          | −1.173    |                  |
| Distance [pc]              | 1056      | 1291±362         |
| Reddening [mag] $E(b - y)$ |           | 0.157 ± 0.070    |
| Reddening [mag] $E(B - V)$ | 0.118     | 0.212 ± 0.094    |
| Distance modulus [mag]     | 10.48     | 10.47 ± 0.65     |
| Log age                    | 7.104     | 7.09             |
| Metallicity                | −0.30     | −0.38            |




Fig. 3. Location of the unreddened points of NGC 2343 in the LGK86 grids.

Equally, the mean distance is  $1291 \pm 362$  pc. Metallicity was obtained for only two F type stars, 23 and 29, but the first one was found to be a non-member. Hence, the metallicity of NGC 2343 can be determined as  $[\text{Fe}/\text{H}] = -0.38$ .

## 6. AGE DETERMINATION

The age was fixed for the cluster once we determined the brightest and hence the hottest, main sequence stars. The effective temperature of these hottest stars was determined by plotting the location of all member stars on the theoretical grids of Lester, Gray, & Kurucz (1986, hereafter LGK86) for the corresponding chemical composition. There is a reported metallicity for NGC 2343 of  $-0.30$  in WEBDA. This value is consistent with the results

obtained from Nissen (1988), who determined values of  $[\text{Fe}/\text{H}]$  of  $-0.37$  and  $-0.21$  for stars 23 and 29, respectively, although the first one is non-member and the other, a marginal member. A solar chemical composition was assumed for the models. We have utilized the  $(b - y)_0$  vs.  $c_0$  diagram which allows the determination of the temperatures with an accuracy of a few hundreds of degrees. Once the membership had been established, the temperature for the hottest stars was determined to be around 25000 K, see Figure 3.

The age is determined after calculating the effective temperature through the calibrations of Meynet, Mermilliod, & Maeder (1993) for open clusters; from the relation  $\log(\text{age}) = -3.499 \log T_e + 22.476$ , valid in the effective temperature ( $T_e$ ) range  $[4.25, 4.56]$ , which applies for NGC 2343. The application of the above mentioned calibration yields the following age:  $\log(\text{age})$  of 7.1 ( $1.2 \times 10^7$  yr) for NGC 2343 for which Meynet et al. (1993) assign an uncertainty of less than  $3 \times 10^3$ . Table 4 shows the newly derived characteristic values of NGC 2343 compared with those previously compiled.

## 7. CONCLUSIONS

New  $uvby - \beta$  photoelectric photometry has been acquired and is presented for the brightest stars in the direction of the open cluster NGC 2343. For the observed stars in the field spectral types were assigned. Using the calibrations to calculate reddening and distance for these stars, the distance to the cluster has been fixed. Membership in the cluster has been determined by constructing a histogram for the distances of the stars of this cluster. A Gaussian fit was calculated and the distance was assigned given

the mean value and the standard deviation. Those stars are denoted by the letter M. Those which were close but beyond the limits are represented by M:. Unreddened indexes in the LGK86 grids allowed us to determine the effective temperature of the hottest stars and hence, the cluster age.

Since no previous  $uvby - \beta$  exists, previous knowledge of the NGC 2343 cluster rested on  $UBV$  photometry. Both the color-color and the color-magnitude diagrams do not show a clear main sequence which make the distance, age and reddening determinations ambiguous. We only measured a few stars, the brightest in the direction of the cluster, but with this small sample we determined those that could belong to the cluster. We found that it is at a closer distance than previously assumed and with a slightly larger reddening, but of the same age. Since the sample is complete up to magnitude 10, we have a good estimation of the properties although, of course, more data would, unambiguously, establish the true nature of this cluster.

We would like to thank the staff of the OAN for their assistance in securing the observations. This work was partially supported by PAPIIT IN104612. This analysis was presented as part of the requirements for obtaining a Bachelor degree by A. Martínez at University of Panama. Typing and proofreading were done by J. Orta, and J. Miller, respectively. C. Guzmán, F. Ruiz and A. Díaz assisted us in the computing. This research has made use of the Simbad databases operated at CDS, Strasbourg, France and NASA ADS Astronomy Query Form.

## REFERENCES

- Arellano-Ferro, A., & Parrao, L. 1988, Reporte Técnico 57 (Mexico: Instituto de Astronomía, Univ. Nacional Autónoma de México)
- Cannon, A. J., & Pickering, E. C. 1918, *Ann. Harvard Coll. Obs.*, 91
- Clariá, J. J. 1972, *AJ*, 77, 868
- Crawford, D. L., & Barnes, J. V. 1970, *AJ*, 75, 978
- Crawford, D. L., & Mander, J. 1966, *AJ*, 71, 114
- Golay, M. 1974, *Introduction to Astronomical Photometry* (Dordrecht: Reidel)
- Gray, R. O., & Olsen, E. H. 1991, *A&AS*, 87, 541
- Grønbech, B., & Olsen, E. H. 1976, *A&AS*, 25, 213
- \_\_\_\_\_. 1977, *A&AS*, 27, 443
- Lester, J. B., Gray, R. O., & Kurucz, R. I. 1986, *ApJS*, 61, 509 (LGK86)
- McSwain, M. V., & Gies, D. R. 2005, in *ASP Conf. Ser. 337, The Nature and Evolution of Disks Around Hot Stars*, ed. R. Ignace & K. G. Gayley (San Francisco: ASP), 270
- Meynet, G., Mermilliod, J.-C., & Maeder, A. 1993, *A&AS*, 98, 477
- Moffat, A. F. J. 1974, *A&AS*, 16, 33
- Nissen, P. 1988, *A&A*, 199, 146
- Olsen, E. H. 1983, *A&AS*, 54, 55
- Paunzen, E., & Mermilliod, J. C. 2007, *WEBDA: A Site Devoted to Stellar Clusters in the Galaxy and the Magellanic Clouds* (<http://webda.physics.muni.cz>)
- Peña, J. H., & Sareyan, J. P. 2006, *RevMexAA*, 42, 179
- Schuster, W. J., & Parrao, L. 2001, *RevMexAA*, 37, 187
- Schuster, W. J., Parrao, L., & Guichard, J. 2002, *J. Astronomical Data*, 8, 2
- Shobbrook, R. R. 1984, *MNRAS*, 211, 659
- U.S. Government Printing Office 1995, *The Astronomical Almanac for the Year 1996, Data for Astronomy, Space Sciences, Geodesy, Surveying, Navigation and other applications* (Washington, D. C.: USGPO)

A. Martínez: Universidad de Panamá, Panama (adams.martinez@gmail.com).

J. H. Peña: Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 70-264, C.P. 04510, México, D.F., Mexico (jhpena@astro.unam.mx).