



## Research note

# Helminth fauna of *Lithobates brownorum* (Anura: Ranidae) at three localities in the state of Yucatán, Mexico

## Helmintofauna de *Lithobates brownorum* (Anura: Ranidae) en tres localidades del estado de Yucatán, México

Carlos A. Yáñez-Arenas and Sergio Guillén-Hernández\*

Departamento de Biología Marina, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Yucatán, Carretera Mérida-Xmatkuil, Km 15.5. Apartado postal 4-116, Itzimná, Mérida, Yucatán, México.

\*Correspondencia: ghernand@uady.mx

**Abstract.** Between July 2004 and June 2005, 84 specimens of *Lithobates brownorum* (Sanders, 1973) were collected and examined for helminths. Hosts came from 3 localities in the state of Yucatán, Mexico: Celestún (n= 35), Lagunas de Yalahau (n= 33), and Ría Lagartos (n= 16). Twelve helminth taxa were found: 7 nematodes, 4 digeneans, and 1 acanthocephalan. With the exception of *Haematoloechus floedae* and *Megalodiscus temperatus*, all helminth taxa found have a Neotropical distribution. Our results differ from those obtained in previous studies dealing with amphibians in Mexico because we found higher richness and abundance of nematodes than digenean species. The relatively low mean intensity and mean abundance values reached by digenean species in this study may be related to the generalist host diet and with the vagility of frogs (from the aquatic to the terrestrial environment or vice versa).

Key words: parasites, anurans, infection levels.

**Resumen.** Entre julio del 2004 y junio del 2005, se recolectaron y examinaron 84 ejemplares de *Lithobates brownorum* (Sanders, 1973) en busca de helmintos en 3 localidades del estado de Yucatán: Celestún (n= 35), Lagunas de Yalahau (n= 33) y Ría Lagartos (n= 16). Los ejemplares estuvieron parasitados por 12 taxa de helmintos, 7 de nematodos, 4 digéneos y 1 acantocefalo. Con excepción de *Haematoloechus floedae* y *Megalodiscus temperatus*, todos los taxa presentan distribución neotropical. Nuestros resultados difieren de los obtenidos por otros autores que han reportado más especies de digéneos que de nematodos en este grupo de hospederos. Los bajos valores de intensidad y abundancia media de los digéneos encontrados en este estudio pueden estar relacionados con la amplia dieta del hospedero y la vagilidad que éste presenta entre el medio acuático y el terrestre.

Palabras clave: parásitos, anuros, niveles de infección.

In southern Mexico, *Lithobates brownorum* (Sanders, 1973) is a widely distributed frog that is commonly found in and around water bodies throughout the Yucatán Peninsula (Zaldívar-Riverón et al., 2004). However, information concerning the helminth parasites associated to this frog species in the Yucatán Peninsula (Moravec, et al. 2002; León-Règagnon et al., 2005), or elsewhere in Mexico (Paredes-León et al. 2008) is scarce. The present work was aimed at characterizing the helminth fauna present in *L. brownorum* specimens collected at 3 different localities in Yucatán.

From July 2004 to June 2005, *L. brownorum* specimens were collected from the following localities: 1), Celestún, located on the northwest coast of the Yucatán Peninsula (20°46', 21°06' N; 90°11', 90°25' W); 2), Yalahau, an inland lagoon found in the central portion of the state (20°40'37.3'', 20°34'59.7''N, 89°10'49.6'', 89°15'00.5''W); and 3), Ría Lagartos (21°32', 21°34' N; 87°35', 88°15'W) found on the northeast coast of the Yucatán Peninsula.

Frogs were captured either by hand or with seine nets, and specimens were kept alive until parasitological examination was conducted. Specimens were sacrificed with an overdose of sodium pentobarbital. All organs

were examined under a stereo-microscope, and helminths were removed, counted, and identified. Standard techniques were employed to prepare parasite specimens for microscopic examination (see Lamothe-Argumedo, 1997). Parasite voucher specimens were deposited in the Colección Nacional de Helminths (CNHE) of the Instituto de Biología, Universidad Nacional Autónoma de México (UNAM), Mexico City. Ecological parameters such as prevalence, mean intensity, and mean abundance were calculated following Bush et al. (1997).

A total of 84 *L. brownorum* specimens were collected and examined (Celestún, n= 35; Yalahau, n= 33; and Ría Lagartos, n= 16). Four hundred and forty helminths (296 nematodes, 139 digeneans, and 5 acanthocephalans) were found, representing 12 taxa (3 of them in larval stage). Seven taxa were nematodes, 4 digeneans, and 1 acanthocephalan. Nine of these taxa are recorded for the first time in Yucatán (Table 1), and 9 represent new records for *L. brownorum* (*Langeronia macrocirra* Caballero y Bravo-Hollis, 1949, *Megalodiscus temperatus* (Stafford, 1923), *Aplectana incerta* Caballero, 1949, *Foleyellides striatus* (Ochoterena and Caballero, 1932), *Rhabdias füelleborni* Travassos, 1926, *Subulascaris falcaustriformis* Teixeira de Freitas and Dobbin, 1957, *Contraeaecum* sp., Physalopteridae gen.

sp., and *Oncicola* sp.).

The greatest number of helminth taxa was found in specimens from Celestún, while the lowest numbers of parasites were found in specimens from Ría Lagartos. Overall, more nematode species were found than any other helminth taxonomic group. The greatest prevalence and mean abundance values were observed for the nematodes *S. falcaustriformis* and *A. incerta* in specimens from Celestún and Yalahau, respectively. On the other hand, the digenean *L. macrocirra* showed the highest prevalence and mean abundance values for specimens in Ría Lagartos. Finally, the digeneans *G. brownorumae* and again *L. macrocirra* showed the highest mean intensity values for specimens from Ría Lagartos; *Glypthelmins brownorumae* also was the species with the highest mean intensity value for specimens from Celestún, and *L. macrocirra* had the greatest value for this parameter for specimens from Yalahau (Table 2).

Previous to this work, only 2 helminth species had been recorded for *L. brownorum* in Yucatán (Moravec et al., 2002; León-Règagnon et al., 2005, Paredes-León et al., 2008); our study increased this number to 13. Except for *H. floedae* and *M. temperatus*, all helminth species found have Neotropical distribution. The presence of *H. floedae*

**Table 1.** Helminth parasites found in *L. brownorum* specimens (n= 84) collected at 3 localities in Yucatán, Mexico. RBC= Reserva de la Biosfera de Celestún, PELY= Parque Estatal Lagunas de Yalahau, RBRL= Reserva de la Biosfera de Ría Lagartos

Helminth	RBC	PELY	RBRL	Distribution	Hábitat
DIGENEA					
<i>Glypthelmins brownorumae</i> Razo-Mendivil, et al. 2004†	X		X	Neotropical	Intestine
<i>Haematoloechus floedae</i> Harwood, 1932		X		Nearctic	Lungs
<i>Langeronia macrocirra</i> Caballero y Bravo-Hollis, 1949* †	X	X	X	Neotropical	Intestine
<i>Megalodiscus temperatus</i> (Stafford, 1923)*†	X			Nearctic	Intestine
ACANTHOCEPHALA					
<i>Oncicola</i> sp.*	X		X	Neotropical	Mesentery
NEMATODA					
<i>Aplectana incerta</i> Caballero, 1949 *†	X	X	X	Neotropical	Intestine
<i>Contraeaecum</i> sp. * †	X	X	X	?	Body cavity
<i>Foleyellides striatus</i> (Ochoterena and Caballero, 1932)* †	X	X	X	Neotropical	Body cavity
Caballero, 1935* †					
<i>Oswaldocruzia subauricularis</i> (Rudolphi 1819) *		X		Neotropical	Intestine
Physalopteridae gen. sp. * †	X	X			Body cavity
<i>Rhabdias füelleborni</i> Travassos, 1926 *†	X	X		Neotropical	Lungs
<i>Subulascaris falcaustriformis</i> Teixeira de Freitas y Dobbin, 1957 * †	X	X	X	Neotropical	Intestine

\* New host record

† New record in the state of Yucatán.

**Table 2.** Helminth infection levels from *L. brownorum* specimens collected at 3 localities in Yucatán, Mexico. RBC= Reserva de la Biosfera de Celestún, PELY= Parque Estatal Lagunas de Yalahau, RBRL= Reserva de la Biosfera de Ría Lagartos. %= Prevalence, MI= Mean Intensity ± SD, MA=Mean Abundance ± SD

Locality	RBC (n=35)			PELY (n=33)		
	%	MA (± SE)	MI (± SD)	%	MA (± SD)	MI (± SD)
<b>DIGENEA</b>						
<i>Megalodiscus temperatus</i>	2.0	0.02 (± 0.16)	1 (± 0)	0.0	0.0	0.0
<i>Haematoloechus floedae</i>	0.0	0.0	0.0	3.0	0.03 (± 0.17)	1.0 (± 0.0)
<i>Langeronia macrocirra</i>	14.0	0.54 (± 1.54)	3.80 (± 2.16)	15.0	1.42 (± 6.13)	9.40 (± 14.39)
<i>Glypthelmins brownorumae</i>	20.0	1.17 (± 3.61)	5.85 (± 6.49)	0.0	0.0	0.0
<b>ACANTHOCEPHALA</b>						
<i>Oncicola</i> sp.	8.0	0.11 (± 0.40)	1.33 (± 0.57)	0.0	0.0	0.0
<b>NEMATODA</b>						
<i>Rhabdias fülleborni</i>	20.0	0.80 (± 2.38)	4.14 (± 4.017)	9.0	0.24 (± 0.86)	2.66 (± 1.52)
<i>Aplectana incerta</i>	5.0	0.14 (± 0.69)	2.50 (± 2.12)	39.0	1.54 (± 2.39)	3.92 (± 2.28)
<i>Foleyellides striatus</i>	14.0	0.42 (± 1.24)	3.0 (± 1.87)	6.0	0.09 (± 0.38)	1.50 (± 0.70)
<i>Subulascaris falcaustriformis</i>	25.0	1.48 (± 3.95)	5.77 (± 6.22)	18.0	0.96 (± 3.03)	5.33 (± 5.57)
<i>Contraecum</i> sp.	2.0	0.02 (± 0.16)	1.0 (± 0)	6.0	0.06 (± 0.24)	1.0 (± 0)
<i>Oswaldocruzia subauricularis</i>	0.0	0.0	0.0	18.0	0.57 (± 1.41)	3.16 (± 1.72)
Physalopteridae gen. sp.	8.0	0.20 (± 0.75)	2.33 (± 1.52)	15.0	1.27 (± 5.41)	8.40 (± 12.64)

  

Locality	RBRL (n=16)		
	%	MA (± SD)	MI (± SD)
<b>DIGENEA</b>			
<i>Megalodiscus temperatus</i>	0.0	0.0	0.0
<i>Haematoloechus floedae</i>	0.0	0.0	0.0
<i>Langeronia macrocirra</i>	25.0	1.50 (± 2.96)	6.0 (± 2.82)
<i>Glypthelmins brownorumae</i>	6.0	0.37 (± 1.5)	6.0 (± 0)
<b>ACANTHOCEPHALA</b>			
<i>Oncicola</i> sp.	6.0	0.06 (± 0.25)	1.0 (± 0.0)
<b>NEMATODA</b>			
<i>Rhabdias fülleborni</i>	0.0	0.0	0.0
<i>Aplectana incerta</i>	18.0	0.75 (± 2.04)	4.0 (± 3.46)
<i>Foleyellides striatus</i>	12.0	0.43 (± 1.31)	3.50 (± 2.12)
<i>Subulascaris falcaustriformis</i>	6.0	0.12 (± 0.5)	2.0 (± 0)
<i>Contraecum</i> sp.	12.0	0.56 (± 1.99)	4.50 (± 4.94)
<i>Oswaldocruzia subauricularis</i>	0.0	0.0	0.0
Physalopteridae gen. sp.	0.0	0.0	0.0

in native amphibian species in Yucatán may be due to the introduction of *Lithobates catesbeiana* (Shaw, 1802) in the region for culture purposes (León-Règagnon, et al., 2005). On the other hand, *M. temperatus* has a wide distribution and has been recorded from the southern United States of America to Costa Rica, where it is a common parasite of several genera of anurans (Yamaguti, 1971; Bursey and Golberg, 2005). The widespread distribution of final hosts of *M. temperatus* suggests the presence of a species complex which needs further research.

Numerous studies have shown that in general more digeneans species are found in amphibians with either aquatic or semiaquatic habits, while nematode species are more common in terrestrial amphibians (Aho, 1990; Guillén-Hernández et al., 2000; Pérez-Ponce de León et al., 2000; Paredes-Calderón, et al., 2004; Espínola-Novelo and Guillén-Hernández, 2008). Overall, results from this work contradict these findings; nematode species richness and mean abundance in *L. brownorum* were greater compared to that of digeneans in 2 of the localities (Table 2). This difference in the number of nematode and digenean species and their relative abundances in *L. brownorum* may be due to the specific biotic and abiotic conditions present in those localities. A low richness of suitable invertebrate species acting as intermediate hosts (e.g., snails and insects) throughout the life cycle of digeneans at those localities may explain our results. Nematode species collected in this study are monoxenous and most of them infect their host via skin penetration. Amphibians are generally considered generalist feeders and their diets usually reflect the availability of suitable prey (Duellman and Trueb, 1994). In particular, *L. brownorum* has been shown to feed on a wide variety of terrestrial and aquatic arthropods, although it does not show a special preference for any given item (Ramírez-Bautista and Lemos-Espinal, 2004). Digeneans usually require an aquatic intermediate host species in addition to a snail species in their life cycle. The presence of *M. temperatus* and *G. brownorum* in this amphibian species suggests the possibility that it feeds on its own skin after molting since larval stages (metacercariae) of these parasites occur in amphibian skin (Smith, 1980); another possibility is that adults prey on tadpoles.

The relatively low mean intensity and mean abundance values found for digeneans in this study may be related to this amphibian's generalist diet, as well as its mobility between aquatic and terrestrial habitats. The high prevalence values observed for nematodes relative to digeneans suggests that *L. brownorum* spends a significant amount of time out of the water, resulting in high rates of nematode recruitment via skin penetration.

Results for *L. brownorum*, the mostly aquatic host that we examined in this study and those reported for the

terrestrial hosts *C. marinus* and *C. valliceps* in Yucatán (Espínola-Novelo and Guillén-Hernández, 2008) strongly differ from results reported in a study which examined specimens of both terrestrial species and 2 aquatic (*L. berlandieri* and *L. vaillanti*) species closely related to *L. brownorum* from Los Tuxtlas, Veracruz (Guillén-Hernández, 1992; Guillén-Hernández, et al. 2000). These findings suggest that digeneans species are an important component which structures the parasite community of amphibians in Veracruz, while in Yucatán nematodes appear to take their place as we found them to be more species-rich and abundant. Further studies which record helminth species for other species of amphibians in the Yucatan Peninsula are necessary to provide insight on this issue.

We thank Jorge Cerón, Alfredo Barrera, Karla Rodríguez, Juan Espínola, and Mara Bravo for field assistance. The authors also thank Rosario Mata, David Osorio, and Elizabeth Martínez (Laboratorio de Helminología, UNAM) for their assistance in the identification of some helminth species. Financial support for this study was granted by CONACYT (FOMIX-YUC-2003-CO3-036) as part of the project titled "Valoración de la diversidad de helmintos en anfibios de tres áreas protegidas del estado de Yucatán"

#### Literature cited

- Aho, J. M. 1990. Helminth communities of Amphibians and Reptiles: comparative approaches to understanding patterns and processes. In Parasites communities: patterns and processes, G. Esch, A. Bush and J. Aho (eds.). Chapman and Tall, London. p. 157-196.
- Brooks, D., V. León-Règagnon, D. McLennan and D. Zelmer. 2006. Ecological fitting as a determinant of the community structure of platyhelminth parasites of anurans. *Ecology* 87: S76-S85.
- Bursey, C. R. and S. R. Golberg. 2005. New species of *Oswaldocruzia* (Nematoda: Molineoidea), new species of *Rhabdias* (Nematoda: Rhabdiasidae), and other helminthes of *Rana* cf. *forreri* (Anura: Ranidae) from Costa Rica. *Journal of Parasitology* 91:600-6005.
- Bush, A. O., K. D. Lafferty, J. M. Lotz and A. W. Shostak. 1997. Parasitology Meets Ecology on its own terms: Margolis et al. Revisited. *Journal of Parasitology* 83:575-583.
- Duellman, W. E. and L. Trueb. 1986. Biology of amphibians. The Johns Hopkins University Press. Baltimore and London. 670 p.
- Espínola-Novelo, J. F. and S. Guillén-Hernández. 2008. Helminth parasites in *C. marinus* and *C. valliceps* (Anura: Bufonidae) from Lagunas Yalahau, Yucatán, Mexico. *Journal of Parasitology* 94:672-674.
- Guillén-Hernández, S. 1992. Comunidades de helmintos de algunos anuros de "Los Tuxtlas", Veracruz. Thesis, Facultad

- de Ciencias, UNAM. 90 p.
- Guillén-Hernández, S., G. Salgado-Maldonado and R. Lamothe-Argumedo. 2000. Digeneans (Platyhelminthes: Trematoda) of seven sympatric species of anurans from Los Tuxtlas, Veracruz, México. *Studies of Neotropical Fauna and Environment* 35:10-13.
- Lamothe-Argumedo, R. 1997. Manual de técnicas para preparar y estudiar los parásitos de animales silvestres. AGT, México, D.F. 43 p.
- León-Règagnon, V., S. Guillén-Hernández and M. A. Arizmendi-Espinosa. 2005. Intraespecific variation of *Haematoloechus floedae* Harwood, 1932 (Digenea: Plagiorchiidae), from *Rana* spp. in North and Central America. *Journal of Parasitology* 91:915-921.
- Moravec, F., E. Mendoza-Franco, C. Vivas-Rodríguez, J. Vargas-Vázquez and D. González-Solís. 2002. Observations on seasonal changes in the occurrence and maturation of five helminth species in the Pimelodid catfish, *Rhamdia guatemalensis*, in the cenote (=sinkhole) Ixin-há, Yucatán, Mexico. *Acta Societate Zoologicae Bohemia* 66:121-140.
- Paredes-Calderón, E. L., V. León-Regagnon and L. García-Prieto. 2004. Helminth infracommunities of *Rana vaillanti* Brocchi (Anura: Ranidae) in Los Tuxtlas, Veracruz, Mexico. *Journal of Parasitology* 90:692-696.
- Paredes-León, R., L. García-Prieto, C. Guzmán-Cornejo, V. León-Regagnon and T. M. Pérez. 2008. Metazoan parasites of Mexican amphibians and reptiles. *Zootaxa* 1904:1-166.
- Pérez-Ponce de León, G., V. León-Règagnon, L. García-Prieto, U. Razo Mendivil and A. Sánchez-Álvarez. 2000. Digenean fauna of amphibians from Central Mexico: Nearctic and Neotropical Influences. *Comparative Parasitology* 67:92-106.
- Ramírez-Bautista, A. and J. A. Lemos-Espinal. 2004. Diets of two syntopic populations of frogs, *Rana vaillanti* and *Rana brownorum*, from a tropical rain forest in southern Veracruz, México. *Southwestern Naturalist* 49:316-320.
- Razo-Mendivil, U., V. León-Règagnon and G. Pérez-Ponce de León. 2006. Monophyly and systematic position of *Glyphelmis* (Digenea), based on partial 18S rDNA sequences and morphological evidence. *Organisms, Diversity & Evolution* 6:308-320.
- Smith, J. D. and M. M. Smith. 1980. Frogs as host-parasites system I. The MacMillan Press LTD, London. 101 p.
- Yamaguti, S. 1971. Synopsis of Digenetic Trematodes of Vertebrates. Vol. I and III. Keigaku Publishing Co. Tokyo, Japan. 1974 p.
- Zaldívar-Riverón, A., V. León-Règagnon and A. Nieto-Montes de Oca. 2003. Phylogenetic relationships of Mexican coastal frogs (*Rana berlandieri* group) based on mtDNA sequences. *Molecular Phylogenetics and Evolution* 26:38-49.