



## Research note

# New molecular data on subcutaneous *Philornis* (Diptera: Muscidae) from southern South America suggests the existence of a species complex

## *Datos moleculares nuevos sobre Philornis (Diptera: Muscidae) subcutáneas del sur de Sudamérica sugieren la existencia de un complejo de especies*

Martín A. Quiroga<sup>a,b</sup>, Lucas D. Monje<sup>a,\*</sup>, Juan P. Arrabal<sup>c</sup>, Pablo M. Beldomenico<sup>a,d</sup>

<sup>a</sup> Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral (ICIVET-Litoral), Universidad Nacional del Litoral – Consejo Nacional de Investigaciones Científicas y Técnicas (UNL-CONICET), R.P. Kreder 2805, Esperanza, Santa Fe (3080), Argentina

<sup>b</sup> Facultad de Ciencia y Tecnología – Universidad Autónoma de Entre Ríos (UAdER), Oro Verde (3100), Argentina

<sup>c</sup> Instituto Nacional de Medicina Tropical, Ministerio de Salud de la Nación, Neuquén y Jujuy s/n, Puerto Iguazú (3370), Argentina

<sup>d</sup> Facultad de Ciencias Veterinarias, Universidad Nacional del Litoral, R.P. Kreder 2805, Esperanza, Santa Fe (3080), Argentina

Received 4 February 2016; accepted 31 May 2016

Available online 16 November 2016

### Abstract

The taxonomic identification of species in the genus *Philornis* Meinert (Diptera: Muscidae) has been based primarily upon analysis of morphological characters. However, relying solely on morphology has led to frequent species misidentification. This has been empirically demonstrated when specimens identified morphologically as *Philornis seguyi* have been reclassified as *Philornis torquans* based on DNA sequences of the second ribosomal internal transcribed spacer (ITS2), showing that the morphologic distinction between these 2 species is extremely challenging. Herein, new molecular data on specimens assigned to *P. torquans* and *P. seguyi* from the type locality of *P. seguyi* (Misiones province, Argentina) were analyzed. The ITS2 sequences obtained differed from those previously reported for *Philornis*, increasing to 3 the number of documented genotypes of flies morphologically indistinguishable from *P. torquans*. These results strongly suggest that subcutaneous *Philornis* spp. from southern South America could represent a species complex (*P. torquans* complex).

© 2016 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** *Philornis seguyi*; *Philornis torquans*; ITS2; Species complex; Argentina

### Resumen

La identificación taxonómica de especies del género *Philornis* Meinert (Diptera: Muscidae) ha estado basada principalmente en el análisis de los caracteres morfológicos. Sin embargo, el uso exclusivo de información morfológica ha llevado con frecuencia a errores en la identificación de las especies. Esto fue demostrado de forma empírica cuando especímenes morfológicamente identificados como *P. seguyi* fueron reclasificados como *P. torquans* utilizando secuencias del segundo espaciador interno ribosomal transcrito (ITS2), lo que demuestra que la distinción morfológica entre estas 2 especies es extremadamente complicada. En el presente trabajo se analizaron nuevos datos moleculares de especímenes identificados como *P. torquans* y *P. seguyi* provenientes de la región reportada como localidad tipo para *P. seguyi* (provincia de Misiones, Argentina). Las secuencias de ITS2 obtenidas fueron diferentes a aquellas previamente reportadas para otras especies de *Philornis* y se incrementa a 3 el número de genotipos documentados de moscas morfológicamente indistinguibles asignadas a *P. torquans*. Estos resultados indican que los organismos de *Philornis* spp. subcutáneos del sur de Sudamérica podrían representar un complejo de especies (complejo *P. torquans*).

© 2016 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Palabras clave:** *Philornis seguyi*; *Philornis torquans*; ITS2; Complejo de especies; Argentina

\* Corresponding author.

E-mail address: [lmonje@fcv.unl.edu.ar](mailto:lmonje@fcv.unl.edu.ar) (L.D. Monje).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

The genus *Philornis* Meinert (Diptera: Muscidae) comprises about 50 known species mainly distributed in the Neotropical region. Adults are free living, whereas larvae are strictly associated with bird nestlings, establishing different types of associations (coprophagous, semihaematophagous or subcutaneous; Dudaniec & Kleindorfer, 2006). The study of ecology and systematics of this group is highly relevant, since *Philornis* flies are parasites of some endangered species of conservation concern (e.g. Darwin finches parasitized by *Philornis downsi* Dodge & Aitken; Koop, Huber, Laverty, & Clayton, 2011), have the potential to be favored by global environmental change (Antoniazzi et al., 2010), and also are excellent models for the study of parasite ecology (Manzoli et al., 2013). To date, one semi-haematophagous (*P. downsi*) and 3 subcutaneous *Philornis* species (*Philornis blanchardi* García, *Philornis torquans* Nielsen and *Philornis seguyi* García) were reported for Argentina (Couri, Antoniazzi, Beldomenico, & Quiroga, 2009; Silvestri, Antoniazzi, Couri, Monje, & Beldomenico, 2011). The taxonomic identification of *Philornis* spp. is currently based mainly upon analysis of morphological characters of adult specimens (descriptions of larvae and pupae are available for a reduced number of species). According to the last published taxonomic review for *Philornis* spp. from Argentina (Couri et al., 2009), *P. seguyi* can be distinguished from *P. torquans* by the width of frons (females) and number of setae on the anterodorsal surface of the hind tibia (males). Differentiation between females from these 2 species could be considered clear (no overlap in values of distinctive characters), though this is not the case for males (the number of setae on the anterodorsal surface of the hind tibia is 3–4 in *P. seguyi* and 4–6 in *P. torquans*). As a consequence of this limitation, such identification approaches have been inaccurate, leading to frequent species misidentification (Monje et al., 2013).

A critical issue that obscures our knowledge on the systematics of *Philornis* spp. from southern South America is that holotypes of several species described in Argentina are lost (i.e., *P. blanchardi*, *P. seguyi*, Patitucci, Mulieri, Domínguez, & Mariluis, 2015), which complicates a proper taxonomic characterization of *Philornis* species in South America. A paradigmatic example of this problem is the case of *P. seguyi*. This species was described by García (1952) based on a single female originally housed at the collection of the “Administración Nacional de Laboratorios e Institutos de Salud Dr. Carlos G. Malbrán (ANLIS-Ent)” (Buenos Aires, Argentina), though is now considered lost (Patitucci et al., 2015). Moreover, the type locality of *P. seguyi* was only reported by García (1952) as “Misiones, Argentina” (Fig. 1A) without any specific geographic location. More recently, Couri, Rabuffetti, and Reboreda (2005) reported *Philornis* specimens in Magdalena (Buenos Aires province, Argentina), which females matched the description of the single female of *P. seguyi*, together with males that were assumed to belong to that species. In this work, the authors redescribed the species including both sexes, and reported it for several areas of Argentina.

Recently, Monje et al. (2013) analyzed specimens of *Philornis* from several localities from Argentina and Uruguay, and used the second internal transcribed spacer region (ITS2) of

the ribosomal DNA as a molecular marker to differentiate adult specimens. The samples analyzed in this study were assigned to *P. seguyi* and *P. torquans* following Couri et al. (2009) work. ITS2 sequences were identical for all specimens from Uruguay and the central region of Argentina, though specimens from the southernmost locality (Magdalena city, samples provided by M.S. Couri) that were morphologically identified as *P. seguyi* had a genetic divergence of 5.9% compared to the former ones. Of the localities studied by Monje et al. (2013), Magdalena was one of the most distinctive, both in terms of distance and habitat, from the type locality of *P. seguyi* (Misiones province, Fig. 1A), suggesting that the specimens from Magdalena could not represent *P. seguyi*. Because the morphology of this particular group of flies may sometimes lead to species misidentification, here we refer to the *Philornis* genotypes identified by Monje et al. (2013) as the ‘*Philornis* sp. genotype Central Argentina’ and the ‘*Philornis* sp. genotype Magdalena’.

Herein, new material from the Misiones province mentioned by García (1952) as type locality of *P. seguyi* were analyzed to increase the number of ITS2 sequences analyzed by Monje et al. (2013), and to contribute to the reconstruction of *Philornis* systematics in southern South America.

*Philornis* larvae were collected from broods of *Cacicus haemorrhous* Linnaeus (Passeriformes: Icteridae) in the rain-forest habitat surrounding Puerto Iguazú (25°42' S, 54°27' W). Twelve third-instar larvae were collected from 3 broods and incubated at room temperature. Emerging adults were morphologically identified by a *Philornis* specialist (MAQ) following a thorough analysis of wings (ciliated/unciliated, r-m and m-m veins pigmentation), eye separation, cheek hair coloration, number of setae in fore/hind tibia, among other characters included in the taxonomic review and description of the species involved (Couri, 1999; Couri et al., 2009).

Genomic DNA from individual flies was extracted using the AccuPrep® Genomic DNA Extraction Kit (Bioneer, USA) following the manufacturer’s protocol. An ITS2 fragment was amplified using the following primers as previously described (Monje et al., 2013): ITS2-LEcEn-F and ITS2-LEcEn-R. PCR products were separated by electrophoresis in a 1.5% agarose gel stained with GelRed™ (Biotium, USA) and examined by UV transillumination. All PCR products were column purified and sequenced directly in both directions using amplifying primers. Sequences were aligned with those previously reported for *Philornis* and closely related fly species using the MUSCLE sequence alignment tool. The sequence alignment and the analyses mentioned below were carried out with the program Mega version 5.0 (Tamura et al., 2011). A phylogenetic analysis was performed with the Maximum-likelihood (ML) method. The best fitting substitution models were determined with the Akaike Information Criterion using the ML model test. Support for the topology was tested by bootstrapping over 1,000 replications, excluding gaps and missing data.

Out of the 9 adult specimens from Misiones that emerged from pupae in the laboratory, 6 were morphologically identified as *P. seguyi*, whereas the remaining 3 were identified as *P. torquans*. Simultaneous infestation of 2 morphologically distinct species of *Philornis* was previously reported by Nores




Figure 1. (A) Map of part of southern South America showing geographical distribution of *Philornis* spp. used in this study. ‘*Philornis* sp. genotype Misiones’ (star), ‘*Philornis* sp. genotype Central Argentina’ (triangle), ‘*Philornis* sp. genotype Magdalena’ (circle) and *Philornis downsi* (cross). Striped area: Misiones province. (B) Phylogenetic analysis of ITS2 sequences from adult specimens of *Philornis* species present in Argentina and closely related fly species. Bootstrap consensus tree was inferred by Maximum-likelihood method. The ITS2 sequence for *Drosophila melanogaster* (EU306667) was used as an outgroup.

(1995), who described co-infestation of *Philornis pici* and *P. seguyi* in broods of *Anumbius anumbi* Orbigny, Lafresnaye (Passeriformes: Furnariidae) and *Pseudoseisura lophotes* Reichenbach (Passeriformes: Furnariidae). We excluded from the analysis specimens with intermediate features, and only those that showed all the distinctive morphological characters for each species were processed for sequencing. Three specimens identified as *P. seguyi* and 1 identified as *P. torquans* were therefore included in the analysis.

The ITS2 sequences from the above 4 specimens were identical regardless of their morphological identification, whereas their divergence from other known *Philornis* ITS2 sequences, ‘*Philornis* sp. genotype Central Argentina’ (KC485555), ‘*Philornis* sp. genotype Magdalena’ (KC485557) and *P. downsi* (HQ834532) was of 11.7%, 9.2% and 27.9%, respectively. In the light of these results, hereafter all specimens from Misiones are referred to as ‘*Philornis* sp. genotype Misiones’.

The phylogenetic tree inferred from maximum likelihood analysis of the ITS2 sequences for *Philornis* spp. and closely related fly species showed the ‘*Philornis* sp. genotype Misiones’ nested in a clade together with ‘*Philornis* sp. genotype Central Argentina’ and ‘*Philornis* sp. genotype Magdalena’, all of which are considerably similar morphologically. The morphologically distinct *P. downsi* was, on the other hand, placed in a separate clade (Fig. 1B). The newly generated sequences were deposited on GenBank (GenBank accession numbers KJ187043–KJ187046). The sequence alignment and the ML tree can be retrieved from the Treebase website ([www.treebase.org](http://www.treebase.org); ID number 19285).

Monje et al. (2013) identified a single genotype (here named ‘*Philornis* sp. genotype Central Argentina’) in samples from 5 out of 6 localities studied (Fig. 1A), whereas a second lineage named ‘*Philornis* sp. genotype Magdalena’ was found in specimens from the southernmost locality (Magdalena city, Fig. 1A). On the basis of morphological characteristics, specimens from Magdalena were identified as *P. seguyi* (Couri et al.,

2005; Monje et al., 2013), though it is unlikely that they actually belong to this species given that Misiones is spatially and ecologically distant from Magdalena. Herein the ITS2 sequences of *Philornis* specimens identified as *P. torquans* and *P. seguyi* from Misiones province (*P. seguyi* type locality) presented differences from sequences of Argentine subcutaneous *Philornis* previously reported (Monje et al., 2013), increasing to 3 the number of documented genotypes of these morphologically undistinguishable flies.

The genetic differences observed between the ITS2 sequences of these 3 genotypes of *Philornis* are higher than the interspecific variations observed between many fly species (Song, Wang, & Liang, 2008; Thanwisai, Kuvangkadilok, & Baimai, 2006). It is unknown whether ‘*Philornis* sp. genotype Misiones’ is indeed *P. seguyi*. Unfortunately, since the holotype of *P. seguyi* is lost, further morphological examination for confirmation of this species’ validity is infeasible. Our results highlight the need to revise and reconstruct the systematics of the genus *Philornis* complementing both molecular and morphological techniques.

Cryptic *Philornis* species are being ignored while scientific literature continues to report wrongly identified species. In this regard, the simultaneous infestation by 2 species of *Philornis* reported by Nores (1995) might also be due to misidentification. Taking into consideration that currently accepted taxonomic reviews (Couri, 1999; Couri et al., 2009) could not identify the species for each of the 3 genotypes described so far, ‘*Philornis* sp. genotype Central Argentina’, ‘*Philornis* sp. genotype Magdalena’ and ‘*Philornis* sp. genotype Misiones’, and that an analysis of their ITS2 sequences showed a close genetic relationship, we believe that these species could actually represent a complex of species under the taxon *P. torquans*.

We propose that, until more extensive studies of *Philornis* species are carried out including molecular data coupled with detailed morphological analyses and cross-mating experiments, the subcutaneous *Philornis* resembling *P. seguyi* or *P. torquans*

in southern South America should be treated as the ‘*Philornis torquans* complex’, as previously proposed for other taxa (e.g. Bertrand et al., 2014; Nava et al., 2014).

We would like to thank Dr. David Anderson for proofreading this work. All procedures conducted in this study comply with the current National and Provincial laws, and were approved by the Bioethics Committee of Universidad Nacional del Litoral. This work was funded by the Argentine Council for Research and Technology (Grant No. PIP 1122010010026).

## References

- Antoniazzi, L. R., Manzoli, D. E., Rohrmann, D., Saravia, M. J., Silvestri, L., & Beldomenico, P. M. (2010). Climate variability affects the impact of parasitic flies on Argentinean forest birds. *Journal of Zoology*, *283*, 126–134.
- Bertrand, C., Janzen, D. H., Hallwachs, W., Burns, J. M., Gibson, J. F., Shokralla, S., et al. (2014). Mitochondrial and nuclear phylogenetic analysis with Sanger and next-generation sequencing shows that, in Área de Conservación Guanacaste, northwestern Costa Rica, the skipper butterfly named *Urbanus belli* (family Hesperiiidae) comprises three morphologically cryptic species. *BMC Evolutionary Biology*, *14*, 153.
- Couri, M. S. (1999). Myiasis caused by obligatory parasites, Ia. *Philornis* Meinert (Muscidae). In J. Guimaraes, & N. Papavero (Eds.), *Myiasis in man and animals in the Neotropical region* (pp. 51–70). São Paulo: Editora Pleiade.
- Couri, M. S., Antoniazzi, L. R., Beldomenico, P. M., & Quiroga, M. (2009). Argentine *Philornis* Meinert species (Diptera: Muscidae) with synonymic notes. *Zootaxa*, *2261*, 52–62.
- Couri, M. S., Rabuffetti, F. L., & Reboreda, J. C. (2005). New data on *Philornis seguyi* García (1952) (Diptera: Muscidae). *Brazilian Journal of Biology*, *65*, 631–637.
- Dudaniec, R. Y., & Kleindorfer, S. (2006). Effects of the parasitic flies of the genus *Philornis* (Diptera: Muscidae) on birds. *Emu*, *106*, 13–20.
- García, M. (1952). Las especies argentinas del género *Philornis* Mein., con descripción de especies nuevas. *Revista de la Sociedad Entomológica Argentina*, *15*, 277–293.
- Koop, J., Huber, S., Laverly, S., & Clayton, D. (2011). Experimental demonstration of the fitness consequences of an introduced parasite of Darwin's finches. *PLoS ONE*, *6*, e19706.
- Manzoli, D., Antoniazzi, L. R., Saravia, M. J., Silvestri, L., Rohrmann, D., & Beldomenico, P. M. (2013). Multi-level determinants of parasitic fly infection in forest passerines. *PLoS ONE*, *8*, e67104.
- Monje, L. D., Quiroga, M., Manzoli, D., Couri, M., Silvestri, L., Venzal, J., et al. (2013). Sequence analysis of the internal transcribed spacer 2 (ITS2) from *Philornis seguyi* (García, 1952) and *Philornis torquans* (Nielsen, 1913) (Diptera: Muscidae). *Systematic Parasitology*, *86*, 43–51.
- Nava, S., Beati, L., Labruna, M. B., Cáceres, A., Mangold, A., & Guglielmo, A. A. (2014). Reassessment of the taxonomic status of *Amblyomma cajennense* (Fabricius, 1787) with the description of three new species, *Amblyomma tonelliae* n. sp., *Amblyomma interandinum* n. sp. and *Amblyomma patinoi* n. sp., and resurrection of *Amblyomma mixtum* Koch, 1844 and *Amblyomma sculptum* Berlese, 1888 (Ixodida: Ixodidae). *Ticks and Tick Borne Diseases*, *5*, 252–276.
- Nores, A. I. (1995). Botfly ectoparasitism of the brown cacholote and the firewood-gatherer. *The Wilson Bulletin*, *107*, 734–738.
- Patitucci, L. D., Mulieri, P. R., Domínguez, M. C., & Mariluis, J. C. (2015). The type specimens of Calypttratae (Diptera) housed in non-traditional institutions in Argentina. *Zootaxa*, *3905*, 557–572.
- Silvestri, L., Antoniazzi, L. R., Couri, M. S., Monje, L. D., & Beldomenico, P. M. (2011). First record of the avian ectoparasite *Philornis downsi* Dodge and Aitken, 1968 (Diptera: Muscidae) in Argentina. *Systematic Parasitology*, *80*, 137–140.
- Song, Z., Wang, X., & Liang, G. (2008). Species identification of some common necrophagous flies in Guangdong province, southern China based on the rDNA internal transcribed spacer 2 (ITS2). *Forensic Science International*, *175*, 17–22.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, *28*, 2731–2739.
- Thanwisai, A., Kuvangkadilok, C., & Baimai, V. (2006). Molecular phylogeny of black flies (Diptera: Simuliidae) from Thailand, using ITS2 rDNA. *Genetica*, *128*, 177–204.