

Life history

## Natural history of the threatened coral snake *Micrurus altirostris* (Serpentes: Elapidae) in Argentina

### *Historia natural de la serpiente de coral amenazada Micrurus altirostris (Serpentes: Elapidae) en Argentina*

María Eugenia Rodríguez <sup>a, \*</sup>, Vanesa Arzamendia <sup>a, b</sup>, Gisela Paola Bellini <sup>a, b</sup>,  
Alejandro Raúl Giraudo <sup>a, b</sup>

<sup>a</sup> Instituto Nacional de Limnología (Consejo Nacional de Investigaciones Científicas y Técnicas - Universidad Nacional del Litoral), Ciudad Universitaria, Santa Fe 3000, Argentina

<sup>b</sup> Facultad de Humanidades y Ciencias, Universidad Nacional del Litoral, Ciudad Universitaria, Santa Fe 3000, Argentina

\*Corresponding author: mauge525@yahoo.com.ar (M.E. Rodríguez)

Received: 6 February 2018; accepted: 4 June 2018

#### Abstract

The genus *Micrurus* comprises nearly 80 endemic species in the American continent. Knowledge on its natural history is based on populations from Brazil and Uruguay, making it necessary to intensify ecological research in Argentina. We present data on morphology, diet, and reproduction for a threatened Argentine population of *M. altirostris* from the Atlantic Forest biodiversity hotspot. Ecological data were obtained from collected specimens and scientific collections. Females attained sexual maturity at a longer body size than males, but the latter reached a larger size, which is related to male-male combat, a behavior reported in this study. We report a female with 3 oviductal eggs, and 2 ovipositions of 4 and 6 eggs in the late spring-early summer period. Males seemed to have a seasonal reproductive cycle. The diet of *M. altirostris* was based on elongated reptilians and was mostly similar to the diet of populations from Brazil and Uruguay. We reported 4 new prey types for *M. altirostris* and confirm the consumption of reptile eggs for the genus *Micrurus*. Our results contribute to a better understanding of the natural history of *M. altirostris*, providing valuable information for designing strategic conservation plans.

**Keywords:** Argentina; Diet; *Micrurus altirostris*; Morphology; Natural history; Reproduction

#### Resumen

El género *Micrurus* comprende cerca de 80 especies endémicas del continente Americano. El conocimiento sobre su historia natural está basado en poblaciones de Brasil y Uruguay, siendo necesario intensificar los estudios ecológicos en Argentina. Presentamos datos de morfología, dieta y reproducción de *M. altirostris* en Argentina, poblaciones amenazadas del hotspot bosque Atlántico. Los datos ecológicos fueron obtenidos del análisis de especímenes colectados y de colecciones científicas. Las hembras maduran con un tamaño corporal mayor al de los machos, pero éstos alcanzan

un mayor tamaño, relacionado con el combate entre machos, comportamiento que fue observado en este estudio. Registramos una hembra con 3 huevos en el oviducto y 2 oviposiciones de 4 a 6 huevos a fines de la primavera y principio del verano. Los machos parecen tener un ciclo reproductivo estacional. Al igual que en las poblaciones de Brasil y Uruguay, la dieta se basa en reptiles alargados. Reportamos 4 nuevas presas para *M. altirostris* y confirmamos el consumo de huevos en el género *Micrurus*. Nuestros resultados contribuyen a una mejor comprensión de la historia natural de *M. altirostris*, información valiosa para aplicar en estrategias de conservación.

*Palabras clave:* Argentina; Dieta; *Micrurus altirostris*; Morfología; Historia natural; Reproducción

## Introduction

Coral snakes belonging to the genus *Micrurus* (Wagler, 1824) are endemic and highly venomous elapids found in the American continent (Roze, 1996). The genus comprises nearly 80 species. However, some of those species' taxonomy is poorly resolved due to both lack of information and morphological similarity among them (Giraudo et al., 2015; Silva & Sites, 1999; Uetz et al., 2017). Coral snakes constitute a relevant issue for public health due to their high toxicity; and thus, it is important to know about their biology and natural history. Even though they have potent neurotoxic venoms, the cases of envenomations caused by *Micrurus* are relatively rare due to the reclusive nature of these snakes and the fact that they generally inhabit sparsely populated areas (Corrêa-Netto et al., 2011). Cambell and Lamar (2004) stated that coral snakes are greatly diminished in many areas altered by humans. Usually, when forest areas are cleared, their habitats become drier, and thus moisture-associated species (such as *M. altirostris*) and their prey are either reduced in number or extirpated. The natural history of *Micrurus* is quite unknown because most current works in the literature are focused on a few species (Campbell & Lamar, 2004; Roze, 1996). There are hardly any studies on the ecology of *Micrurus* (Almeida-Santos et al., 2006; Ávila et al., 2010; Marques, 2002; Marques et al., 2006, 2013), whereas most of them have addressed systematic aspects and information about its venom, morphology, and taxonomy (Camargo et al., 2011; Morales-Mondoñedo, 2010; Silva & Aird, 2001; Silva & Sites, 1999, 2001).

*Micrurus altirostris* is a medium-sized coral snake with populations distributed across southeastern Brazil, Paraguay, Uruguay, and Northeastern Argentina (Silva & Sites, 1999). In the latter country, it shows a restricted distribution associated with humid habitats of Atlantic Forest and extends to the south by the riparian humid forests of the Uruguay River. Additionally, *M. altirostris* has an isolated population in the riparian humid forest of the Middle Paraná River, which is considered the southernmost expansion of the impoverished Atlantic Forest (Arzamendia & Giraudo, 2009; Giraudo, 2001;

Giraudo et al., 2012; Scrocchi, 1990; Vuoto, 2000). The Atlantic Forest is a biodiversity hotspot considered as one of those with the highest conservation priority in the world due to both its elevated rate of habitat loss and the fact that only 7-8% of the original surface remains (Galindo-Leal & Camara, 2003; Myers et al., 2000).

The natural history of *M. altirostris* in Argentina is scarce because of its fossorial habits, low frequency of capture, the danger of its venom, and the difficulty in manipulating specimens (Oliveira et al., 2005). There exist some data about reproduction and activity patterns of *M. altirostris* based on studies related to Brazilian populations (Marques et al., 2006, 2013). Additionally, there are some data about the ecological traits of *M. altirostris*, but this information is mostly anecdotal and incomplete since it is based on small samples from Brazilian and Uruguayan populations (Carreira-Vidal, 2002; Silva, 1995; Silva & Aird, 2001). In Argentina, *M. altirostris* was categorized as a vulnerable species owing to its relatively restricted distribution, which is strongly associated to the remnants of the humid Atlantic Forest. More than 60% of these habitats have been lost due to the high and still increasing deforestation rate (Giraudo et al., 2003). The poor state of conservation of *M. altirostris* is further affected by high human persecution and road kills (Giraudo et al., 2012). For all these reasons, Giraudo et al. (2012) suggested intensifying ecological research in order to broaden knowledge about the biology of this coral snake species in Argentina. In this study, we present new and basic information about the natural history of *M. altirostris* in Argentina, including morphology, sexual dimorphism, diet, and reproductive biology so that it can be considered when designing conservation strategies for this species.

## Materials and methods

The study area is located in the Northeast of Argentina and comprises Misiones, Corrientes, and Entre Ríos provinces. This area is characterized by subtropical-temperate climate, with a seasonality mainly marked by temperature. Average annual temperatures range from 22 °C in the north to 17 °C in the south. Annual precipitation

averages from 1,000 to 1,800 mm (Arzamendia et al., 2015). The study area has great environmental diversity due to the influence of the Paraná and Uruguay rivers and the confluence of Paranaense or Atlantic Forest, Chacoan, Espinal, and Pampean phytogeographic provinces (Arzamendia & Giraud, 2009).

Data were obtained by analyzing a total of 74 specimens of *M. altirostris*. We sampled the study area from January 1991 to April 2016, mainly by active search and road sampling in different habitats (Bellini et al., 2014). Road sampling allowed us to cover large areas and detect both roadkills and alive snakes. We also conducted time-constrained searches and opportunistic encounter surveys (Scott, 1994; Valdujo et al., 2002). The sampling was carried out at day and night hours. For each specimen, we recorded the exact location (using a Garmin eTrex Legend Global Positioning System device). Snakes were measured with a flexible ruler to the nearest millimeter and subsequently released at the capture site. We preserved both recently road-killed and intentionally killed snakes that were in good conditions for collecting dietary, reproductive, and morphological data. All collected specimens were housed in the collection of Instituto Nacional de Limnología (INALI, Santa Fe, Argentina). The biological material was supplemented with original data from specimens deposited in the scientific collections of Museo de Ciencias Naturales y Antropológicas Profesor Antonio Serrano (MER), Laboratorio de Genética Evolutiva, Instituto de Biología Subtropical CONICET-Universidad Nacional de Misiones (LBG), and Museo de la Plata (MLP).

We recorded the following variables: sex; snout-vent length (SVL), tail length (TL), head length (HL from the tip of the snout to the angle of the jaw), and trunk length (TRL), using a flexible ruler to the nearest millimeter; body mass, using a Pesola scale to the nearest 0.1 g; number of ventral (VEN) and subcaudal (SC) scales, using Gans' (1964) technique; length, width, and thickness of the right testis; diameter of deferent duct close to cloaca; observation of deferent duct (convoluted or non-convoluted); and length, width, and thickness of ovarian follicles (in primary or secondary vitellogenesis) or oviductal eggs. All measurements were made to the nearest 0.1 mm using digital calipers (CONNOR).

Males were considered mature if they had convoluted deferent ducts (Leite et al., 2009; Pizzatto, Jordão et al., 2008; Shine, 1978). Females were considered mature if they had follicles in secondary vitellogenesis, oviductal eggs, or folded oviductus (Pizzatto, Cantor et al., 2008). In males, increased size and mass of the testes probably reflect spermiogenesis (Pizzatto, Cantor et al., 2008), so the volume of the testis was calculated using the ellipsoid

formula (Pleguezuelos & Ferliche, 1999). Male reproductive cycle was determined by comparing variations in the volume of testes and diameter of deferent duct from one season to another, using analyses of covariance (Ancova), since both variables covary with SVL (Almeida-Santos et al., 2006; Pizzatto, Cantor et al., 2008; Pizzatto, Jordão et al., 2008). Due to the questionable accuracy of the testicles volume measure for a real evaluation of male reproductive cycle (Almeida-Santos et al., 2014), sections of testicles and deferent duct were performed. Since most specimens were run over, and the condition of their organs was not adequate, 3 specimens were dissected to obtain histological samples of the proximal region of testes and proximal, medial, and distal region of deferent ducts. Only right-side organs were extracted, thus preserving contralateral testes and deferent ducts. Samples were embedded in paraffin and sectioned at 5  $\mu$ m thickness with a rotating microtome (Thermo Scientific Rotary Microtome HM 325); then, samples were stained with hematoxylin and eosin (Loebens et al., 2018). Slides were observed using an optical microscope (Leica DM 2500).

The presence of sexual dimorphism was evaluated according to the number of ventral and subcaudal scales and SVL and between sexes by means of a nonparametric Mann-Whitney U test. As TL and body mass covariate with SVL and HL covariates with TRL, an analysis of covariance (Ancova) was used. The index of sexual size dimorphism (SSD) was calculated as (mean SVL of the larger sex) / (mean SVL of the smaller sex) - 1 (Shine, 1994). This index is expressed by positive values if females are the larger sex and in negative values if males are the larger sex. Neither juvenile nor neonatal specimens were included in the analysis of sexual dimorphism of measurements (SVL, TL, body mass), except for lepidosis variables because they do not vary with sexual maturity. All raw data were tested for normality and homoscedasticity with Shapiro-Wilks and Levene's tests (Zar, 1999). Digestive tracts of each specimen were dissected to analyze their contents and identify prey items. Prey items were identified to the lowest possible taxonomic group by comparing them with material from the reference collection of the Instituto Nacional de Limnología and guides in the literature. Number and direction of ingestion were recorded from the obtained prey. All statistical analyses were performed by using Infostat software version 2008 (Di Rienzo et al., 2008) with  $\alpha < 0.05$ .

## Results

Of the 74 individuals analyzed, 57 were males (77.1%, 54 adults and 3 juveniles) and 17 were females (22.9%, 8 adults and 9 juveniles). Females attained sexual maturity

(SVL = 570 mm) at a larger body size than males (SVL = 320). Sexual dimorphism was only recorded for the number of ventral scales ( $U = 776.5$ ;  $p = 0.0057$ ), females proving to have a greater amount of scales (Table 1). The sexual size dimorphism index for adults was  $SSD = 0.0064$ , also showing that males and females presented no differences in body size.

Considering macroscopic data, males are reproductively active throughout the whole year, since we found convoluted deferent ducts throughout all seasons; and significant differences were not found in testicular volume between seasons ( $F = 1.28$ ,  $df = 3$ ,  $p = 0.3193$ ). Neither were differences observed in deferent duct diameter between seasons ( $F = 0.59$ ,  $df = 3$ ,  $p = 0.6288$ ). At the microscopic level, 2 different developmental stages were characterized by cellular changes in testes of *M. altirostris*: 1) recrudescence in January, with spermatogonia and spermatocytes in testicles and spermatozoa in deferent duct, and 2) spermiogenesis in February and December, with spermatozoa in testicles and deferent duct. Male-male combat was registered only once on October 7, 1994 in a secondary forest in the province of Misiones.

In addition to the low number of females being studied, only 2 specimens with vitellogenic follicles were found: 1 collected in September (14.25 mm long) and another one in August (13.7 mm long) and another specimen with 3 oviductal eggs (38.2 - 40.75 mm long and 1.4 - 11.6 mm wide) in January. Additionally, 2 events of oviposition

were recorded: one of 6 eggs in November, and another of 4 eggs within 90-minute intervals in December.

From the total collected specimens (74), 27 (36%) had stomach content. In the field, 2 species of snakes were recorded as prey items of *M. altirostris* (Table 2). Three different major prey items were identified: Scolecophidia, the most frequent prey category (41.4%), Alethinophidia (27.6%), and Amphisbaenia (20.7%). Also, 2 specimens with a reptile egg in their digestive tract were recorded (Table 2). In every case, head-first ingestion was identified.

## Discussion

Males of *M. altirostris* were more frequently found than females, representing 77.1% of our sample. These results agree with those obtained by Silva and Sites (1999), Di Bernardo et al. (2007), and Marques et al. (2013), who found 60 - 81% males and 21 - 34% females. The higher activity of males could be due to the courtship ritual (male-male combat). Females would have more secretive habits, which will potentially reduce their vulnerability to predators and increase the chances of survival and, therefore, the amount of mating, which would ensure the reproductive success of the species (Marques et al., 2006).

*Micrurus altirostris* females attain sexual maturity at longer body sizes than males. This pattern has been reported for *Pseudoboina* (Pizzatto, 2005), *Xenodontini* (Pizzatto, Jordão et al., 2008), *Tachymenini* (Bellini et al.,

Table 1

Meristic and morphometric variation in mature and juvenile specimens of *Micrurus altirostris*. Sample size (n); ventral scales (VEN); subcaudal scales (SC); snout-vent length (SVL); tail length (TL); head length (HL).

	Females			Males			Statistic
	n	$\bar{X} \pm SD$	Range	n	$\bar{X} \pm SD$	Range	
VEN	16	214.81 ± 4.67	208-223	55	210.35 ± 5.77	194-224	$p = 0.0057$ ; $U = 776.5$
SC	16	19.81 ± 2.93	14-25	55	20.25 ± 2.70	15-25	$p = 0.4827$ ; $U = 532.5$
Matures							
SVL	8	626.25 ± 34.15	570-664	53	622.66 ± 200.07	320-1150	$p = 0.6926$ ; $U = 0.16$
TL	8	40 ± 7.27	32-54	53	39.17 ± 11.53	13-63	$p = 0.8859$ ; $df = 1$ ; $F = 0.02$
HL	7	18.63 ± 1.73	16.6-20.9	49	17.23 ± 4.29	9.6-26.6	$p = 0.1476$ ; $df = 1$ ; $F = 2.16$
Body mass	8	65.66 ± 25.48	29.5-114.8	49	65.89 ± 50.4	8-230	$p = 0.9712$ ; $df = 1$ ; $F = 0.01$
Juvenile							
SVL	8	340.88 ± 63.05	249-433	3	250.33 ± 17.67	230-262	-
TL	8	24.75 ± 6.94	16-33	3	16.67 ± 3.79	14-21	-
HL	8	11.58 ± 1.48	9.3-13.56	3	9.58 ± 0.95	8.5-10.25	-
Body mass	8	9.71 ± 4.98	3.5-17	3	2.9 ± 1.9	1.3-5	-

Table 2

Diet of *Micrurus altirostris* in the studied population from Argentina and compared with that of populations from Brazil (Silva Jr., 1995; Silva Jr. & Aird, 2001) and Uruguay (Carreira-Vidal, 2002). Number of prey items (N), Frequency (F). \* Prey items prey recorded by direct observation.

Population	Argentina		Brazil		Uruguay	
	N	F (%)	N	F (%)	N	F (%)
<b>Amphisbaenia</b>						
Family Amphisbaenidae						
<i>Amphisbaena mertensii</i>	1	3.5%				
<i>Amphisbaena darwini</i>					2	28.6%
<i>Amphisbaena sp.</i>			1	25%		
Unidentified Amphisbaenidae	5	17.2%				
Subtotal Amphisbaenia	6	20.7%	1	25%	2	28.6%
<b>Serpentes</b>						
<b>Scolecophidia</b>						
Family Anomalepididae						
<i>Liotyphlops beui</i>	1	3.5%				
Family Typhlopidae						
<i>Amerotyphlops brongersmianus</i>	1	3.5%				
Family Leptotyphlopidae						
<i>Epictia munoai</i>					2	28.6%
Unidentified Scolecophidia	10	34.4%				
Subtotal Scolecophidia	12	41.4%			2	28.6%
<b>Alethinophidia</b>						
Family Colubridae						
<i>Atractus reticulatus</i>	2*	6.8%				
<i>Atractus sp.</i>	1	3.5%	1	25%		
<i>Tomodon ocellatus</i>	1	3.5%				
<i>Erytrolamprus sp.</i>			1	25%		
<i>Sibynomorphus sp.</i>			1	25%		
<i>Phalotris bilineatus</i>					1	14.3%
Unidentified Colubridae	3	10.3%				
Family Viperidae						
<i>Bothrops jararaca</i>	1*	3.5%				
Subtotal Alethinophidia	8	27.6%	3	75%	1	14.3%
<b>Sauria</b>						
Family Anguidae						
<i>Ophiodes striatus</i>					1	14.3%
Family Gymnophthalmidae						
<i>Cercosaura schreibersii</i>					1	14.3%
Subtotal Sauria					2	28.6%
Reptile egg	2	6.8%				
Unidentified Vertebrate	1	3.7%				
Total	29	100%	4	100%	7	100%

2013, 2014), Colubrinae (Leite et al., 2009), Crotalinae (Valdujo et al., 2002), and Hydrodynastini (Giraudo et al., 2014). The evidence indicates that sexual maturity in females may be delayed in relation to males due to the high energy cost involved in reproduction. Reproduction in females is not worthwhile until they gain higher probabilities of reproductive success (larger litter size). In addition, early maturity of males increases their chance of mating (Madsen & Shine, 1994; Shine, 1994, 2003).

The male-male combat observed in October is a new record in *M. altirostris* behavior and was also registered in Brazilian populations but in the months of April and May (Almeida-Santos et al., 1998; Marques et al., 2006; Marques et al., 2013). For other species, in which male-male combat was also registered, males were the largest sex (Shine, 1978, 1994). Despite this trend, *M. altirostris* exhibits sexual dimorphism in the number of ventral scales, females being those with a greater number. Marques et al. (2013) analyzed only morphometric characters (SVL and TL) and males were observed to have longer bodies and tails, which does not match our data. This difference between results could be due to the disparity between sample sizes (our sample included more males than females).

Males probably have a seasonal reproductive cycle, since 2 stages of development of *M. altirostris* testes were detected: recrudescence in summer and spermatogenesis in spring and summer. Males of *M. altirostris* do not present seasonal variation of deferent duct diameter, unlike what was found for *M. corallinus* (Almeida-Santos et al., 2006). For the latter species, a correlation was found between increased sizes of the distal portion of the deferent duct and sperm storage, which was confirmed by using histological techniques (Almeida-Santos et al., 2006).

Our records of the female reproductive cycle agree with those reported by Marques et al. (2013), who recorded females with vitellogenic follicles between June and January. However, our data on gravid females and oviposition (December to January) extends the period reported by Marques et al. (2013) (November). Since mating occurs at the end of summer and early autumn, and ovulation occurs during spring, Marques et al. (2013) suggested that *M. altirostris* females could be able to store sperm during winter. For that reason, further histological analyses are necessary to assess possible sperm storage in females and complete spermatogenesis cycle in males (Almeida-Santos et al., 2014).

The diet of *M. altirostris* in Argentina is based on elongated reptiles. The same was reported of populations in Brazil and Uruguay (Carreira-Vidal, 2002; Silva, 1995; Silva & Aird, 2001). We reported 4 new genera in the diet of *M. altirostris*: *Amerotyphlops*, *Liotyphlops*, *Tomodon*, and *Bothrops*. Roze (1996) reported an egg in the stomach

of *M. nigrocinctus zunilensis*. He asserted that no further content was found in the stomach and thus the egg-eating behavior arises as an open question. Our records of reptile eggs in 2 specimens with empty stomachs allow us to confirm for the first time the consumption of reptile eggs in the genus *Micrurus*, as suggested by Roze (1996). Lizards were not registered in the diet of this coral snake in Argentina, but this prey item was reported in populations from Uruguay (Carreira-Vidal, 2002). Probably the differences in diet between our data and populations from Brazil and Uruguay are influenced by the small sample analyzed in those studies and by differences in prey availability.

For *M. altirostris*, direction of prey ingestion is head-first, as was mentioned previously for snakes in general (Bellini et al., 2013; Greene, 1976; Pinto & Lema, 2002; Prieto et al., 2012; Rodríguez-Robles, 2002). This form of ingestion reduces time and deglutition effort and minimizes the possibility of prey escape (Albarelli & Santos-Costa, 2010; Andrade & Silvano, 1996; Pinto & Lema, 2002). Additionally, in the case of ophiophagous snakes, this behavior allows them to avoid the attack of potentially dangerous prey and reduce resistance of ventral scales (Greene, 1976; Pinto & Lema, 2002).

The assessment of the conservation status of neotropical snakes is hampered by the lack of basic natural history data (Giraudo et al., 2012; Leão et al., 2014). In the present study, new data were registered about reproduction (male-male combat, female and male reproductive cycle) and diet of *M. altirostris*. This work provides the first natural history data for a threatened population of coral snake, including relevant information for designing strategic conservation plans.

## Acknowledgements

We would like to express our great appreciation to museum curators Gisela Bahler (Museo Antonio Serrano), Diego Baldo (Laboratorio de Genética Evolutiva), Jorge Williams, and Leandro Alcalde (Museo de la Plata) for allowing us to analyze specimens of the museum's collection. Special thanks are offered to Diana Alberto for her technical assistance with histological sections, and Selma Maria Almeida-Santos and Rafaela Zani Coeti for analyzing the photos of the slides and for their comments. Also, suggestions made by the reviewers and editor of this manuscript have been helpful for us to improve our work. We would also like to thank Liliana Tauber for her assistance in statistics. Our special thanks are extended to our labmates: Carla Bessa, Belén Costanzo, Romina Pavé, Daiana Ferraro, Maximiliano Cristaldi, and Andrés Sarquis for their collaboration throughout this research work.

## References

- Albarelli, L. P. P., & Santos-Costa, M. C. (2010). Feeding ecology of *Liophis reginae semilineatus* (Serpentes: Colubridae: Xenodontinae) in eastern Amazon, Brazil. *Zoologia*, 27, 87–91.
- Almeida-Santos, S. M., Aguiar, L. F. S., & Balestrin, R. L. (1998). *Micrurus frontalis* (Coral snake). Male Combat. *Herpetological Review*, 29, 242.
- Almeida-Santos, S. M., Braz, H. B., Santos, L. C., Sueiro, L. R., Barros, V. A., Rojas, C. A., & Kasperoviczus, K. N. (2014). Biología reproductiva de serpentes: recomendações para a coleta e análise de dados. *Herpetologia Brasileira*, 3, 14–24.
- Almeida-Santos, S. M., Pizzatto, L., & Marques, O. (2006). Intra-sex synchrony and inter-sex coordination in the reproductive timing of the atlantic coral snake *Micrurus corallinus* (Elapidae) in Brazil. *Herpetological Journal*, 16, 371–376.
- Andrade, R. O., & Silvano, R. A. M. (1996). Comportamento alimentar e dieta da falsa coral *Oxyrhopus guibei* (Serpentes, Colubridae). *Revista Brasileira de Zoologia*, 13, 143–150.
- Arzamendia, V., & Giraudo, A. R. (2009). Influence of large South American rivers of the Plata Basin on distributional patterns of tropical snakes: a panbiogeographical analysis. *Journal of Biogeography*, 36, 1739–1749.
- Arzamendia, V., Giraudo, A. R., & Bellini, G. (2015). Relaciones biogeográficas de los grandes ríos de la cuenca del Plata basadas en ensambles de serpientes. *Revista Mexicana de Biodiversidad*, 86, 674–684.
- Ávila, R. W., Kawashita-Ribeiro, R. A., Ferreira, V. L., & Strüssmann, C. (2010). Natural history of the coral snake *Micrurus pyrrhocryptus* Cope 1862 (Elapidae) from semideciduous forests of western Brazil. *South American Journal of Herpetology*, 5, 97–101.
- Bellini, G., Arzamendia, V., & Giraudo, A. R. (2013). Ecology of *Thamnodynastes hypoconia* in subtropical-temperate South America. *Herpetologica*, 69, 67–79.
- Bellini, G., Giraudo, A. R., & Arzamendia, V. (2014). Comparative ecology of three species of *Thamnodynastes* (Serpentes, Dipsadidae) in subtropical-temperate South America. *Herpetological Journal*, 24, 87–96.
- Camargo, T. M., de Roodt, A. R., da Cruz-Höfling, M. A., & Rodrigues-Simioni, L. (2011). The neuromuscular activity of *Micrurus pyrrhocryptus* venom and its neutralization by commercial and specific coral snake antivenoms. *Journal of Venom Research*, 2, 24–31.
- Campbell, J., & Lamar, W. (2004). *The venomous reptiles of the Western Hemisphere*. Ithaca: Cornell University Press.
- Carreira-Vidal, S. (2002). *Alimentación de los ofidios de Uruguay*. Barcelona: Asociación Herpetológica Española, Monografías de Herpetología.
- Corrêa-Netto, C., Junqueira-de Azevedo, I. L. M., Silva, D. A., Ho, P. L., Leitão-de Araújo, M., Alves, M. L. M. et al. (2011). Snake venomomics and venom gland transcriptomic analysis of Brazilian coral snakes, *Micrurus altirostris* and *M. corallinus*. *Journal of Proteomics*, 74, 1795–1809.
- Di Bernardo, M., Borges-Martins, M., & Silva, Jr., N. J. (2007). A new species of coralsnake (*Micrurus*: Elapidae) from southern Brazil. *Zootaxa*, 1447, 1–26.
- Di Rienzo, J. A., Casanoves, F., Balzarini, M. G., González, L., Tablada, M., & Robledo, C. W. (2008). *InfoStat, versión 2008*. Córdoba, Argentina: Grupo InfoStat, FCA/ Universidad Nacional de Córdoba, Argentina.
- Galindo-Leal, C., & Camara, I. G. (2003). Atlantic forest hotspot status: an overview. In Galindo-Leal, C., & Camara, I. G. (Eds.), *The Atlantic forest of South America: biodiversity status, threats, and outlook* (pp. 3–11). Washington, DC.: Center for Applied Biodiversity Science, Island Press.
- Gans, C. (1964). A redescription and geographic variation in *Liophis miliaris* Linné, the common water snake of southeastern South America. *American Museum Novitates*, 2178, 1–58.
- Giraudo, A. R. (2001). *Diversidad de serpientes de la selva Paranaense y del Chaco Húmedo. Taxonomía, biogeografía y conservación*. Buenos Aires: Literature of Latin America.
- Giraudo, A. R., Arzamendia, V., Bellini, G. P., Bessa, C. A., Calamante, C. C., Cardozo, G. et al. (2012). Categorización del estado de conservación de las serpientes de la República Argentina. *Cuadernos de Herpetología*, 26, 1–24.
- Giraudo, A. R., Arzamendia, V., Bellini, G. P., Bessa, C. A., & Costanzo, M. B. (2014). Ecología de una gran serpiente sudamericana, *Hydrodynastes gigas* (Serpentes: Dipsadidae). *Revista Mexicana de Biodiversidad*, 85, 1206–1216.
- Giraudo, A. R., Nenda, S., Arzamendia, V., Bellini, G., & Franzoy, A. (2015). Nuevos datos sobre la distribución, morfología y conservación de *Micrurus silviae* (Serpentes: Elapidae), una serpiente coral amenazada poco conocida. *Revista Mexicana de Biodiversidad*, 86, 1041–1047.
- Giraudo, A. R., Povedano, H., Belgrano, M. J., Krauczuk, E., Pardyñas, U., Miquelarena, A. et al. (2003). Biodiversity status of the interior Atlantic forest of Argentina. In Galindo-Leal, C., & I. G. Cámara (Eds.), *Atlantic forest of the South America. Biodiversity status, threats, and outlook* (pp. 160–180). Washington D.C.: Center for Applied Biodiversity Science, Island Press.
- Greene, H. W. (1976). Scale overlap, a directional sign stimulus for prey ingestion by ophiophagous snakes. *Zeitschrift für tierpsychologie*, 41, 113–120.
- Leão, S. M., Pelegrin, N., Nogueira, C. C., & Brandão, R. A. (2014). Natural history of *Bothrops itapetingae* Boulenger, 1907 (Serpentes: Viperidae: Crotalinae), an endemic species of the Brazilian Cerrado. *Journal of Herpetology*, 48, 324–331.
- Leite, P. T., Nunes, S. F., Kaefer, I. L., & Cechin, S. Z. (2009). Reproductive biology of the swamp racer *Mastigodryas bifossatus* (Serpentes: Colubridae) in subtropical Brazil. *Zoologia*, 26, 12–18.
- Loebens, L., Rojas, C. A., Almeida-Santos, S. M., & Cechin, S. Z. (2018). Reproductive biology of *Philodryas patagoniensis* (Snakes: Dipsadidae) in south Brazil: Female reproductive cycle. *Acta Zoologica*, 99, 105–114.

- Madsen, T., & Shine, R. (1994). Costs of reproduction influence the evolution of sexual size dimorphism in snakes. *Evolution*, 48, 1389–1397.
- Marques, O. (2002). Natural history of the coral snake *Micrurus decoratus* (Elapidae) from the Atlantic Forest in southeast Brazil, with comments on possible mimicry. *Amphibia-Reptilia*, 23, 228–232.
- Marques, O., Almeida-Santos, S. M., & Rodrigues, M. G. (2006). Activity patterns in coral snakes, genus *Micrurus* (Elapidae), in South and Southeastern Brazil. *South American Journal of Herpetology*, 1, 114–120.
- Marques, O., Pizzatto, L., & Almeida-Santos, S. M. (2013). Reproductive strategies of new world coral snakes, genus *Micrurus*. *Herpetologica*, 69, 58–66.
- Morales-Mondoñedo, V. R. (2010). Análisis filogenético del melanismo en las especies peruanas de *Micrurus* Wagler 1824 (Reptilia, Serpentes, Elapidae). *Biotempo*, 10, 26–31.
- Myers, N., Mittermeier, R., Mittermeier, C. G., da Fonseca, G., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.
- Oliveira, L., Almeida-Santos, S. M., Costa, A. C. O. R., Scartozzoni, R. R., Germano, V. J., & da Graça-Salomã, M. (2005). Manutenção de serpentes em cativeiro no Instituto Butantan: I A longevidade do gênero *Micrurus*. *Publicações Avulsas Do Instituto Pau Brasil*, 17, 55–62.
- Pinto, C. C., & Lema, T. (2002). Comportamento alimentar e dieta de serpentes, gêneros *Boiruna* e *Clelia* (Serpentes, Colubridae). *Iheringia. Série Zoologia*, 92, 9–19.
- Pizzatto, L. (2005). Body size, reproductive biology and abundance of the rare pseudoboine snakes genera *Clelia* and *Boiruna* (Serpentes, Colubridae) in Brazil. *Phyllomedusa*, 4, 111–122.
- Pizzatto, L., Cantor, M., de Oliveira, J. L., Marques, O. A. V., Capovilla, V., & Martins, M. (2008). Reproductive ecology of dipsadine snakes, with emphasis on South American species. *Herpetologica*, 64, 168–179.
- Pizzatto, L., Jordão, R. S., & Marques, O. A. V. (2008). Overview of reproductive strategies in Xenodontini (Serpentes: Colubridae: Xenodontinae) with new data for *Xenodon newwiedii* and *Waglerophis merremii*. *Journal of Herpetology*, 42, 153–162.
- Pleguezuelos, J. M., & Feriche, M. (1999). Reproductive ecology of the horseshoe snake (*Coluber hippocrepis*) in the Iberian peninsula. *Journal of Herpetology*, 33, 202–207.
- Prieto, Y., Giraudo, A. R., & López, M. S. (2012). Diet and sexual dimorphism of *Liophis poecilogyus* (Serpentes, Dipsadidae) from the wetland regions of Northeast Argentina. *Journal of Herpetology*, 46, 402–406.
- Rodríguez-Robles, J. A. (2002). Feeding ecology of North American gopher snakes (*Pituophis catenifer*, Colubridae). *Biological Journal of the Linnean Society*, 77, 165–183.
- Roze, J. A. (1996). *Coral snakes of the Americas: biology, identification, and venoms*. New York: Krieger Publishing Company.
- Scott, Jr., N. J. (1994). Complete species inventories. In R.W. Heyer, M. A. Donnelly, R.W. McDiarmid, L.A.C. Hayk, & Foster, M. S. (Eds.), *Measuring and monitoring biological diversity standard methods for amphibians* (pp. 78–79). Washington D.C.: Smithsonian Institution.
- Scrocchi, G. (1990). El género *Micrurus* (Serpentes: Elapidae) en la República Argentina. *Bollettino del Museo Regionale di Scienze Naturali Torino*, 8, 343–368.
- Shine, R. (1978). Sexual size dimorphism and male combat in snakes. *Oecologia*, 33, 269–277.
- Shine, R. (1994). Sexual size dimorphism in snakes revisited. *Copeia*, 1994, 326–346.
- Shine, R. (2003). Reproductive strategies in snakes. *Proceedings. Biological sciences, The Royal Society*, 270, 995–1004.
- Silva, Jr., N. J. (1995). *Molecular systematics and evolution of venoms in the South American 'triad' coral snakes (Elapidae: Micrurus) (Ph.D. Dissertation)*. Brigham Young University, Provo, UT, USA.
- Silva, Jr., N. J., & Aird, S. D. (2001). Prey specificity, comparative lethality and compositional differences of coral snake venoms. *Comparative Biochemistry and Physiology Part C*, 128, 425–456.
- Silva, Jr., N. J., & Sites, J. W. (1999). Revision of the *Micrurus frontalis* complex (Serpentes: Elapidae). *Herpetological Monographs*, 13, 142–194.
- Silva, Jr., N. J., & Sites, J. W. (2001). Phylogeny of South American triad coral snakes (Elapidae: *Micrurus*) based on molecular characters. *Herpetologica*, 57, 1–22.
- Uetz, P., Freed, P., & Hošek, J. (Eds.). (2017). *The reptile database*. Accessed: 25 October 2017. Available at <http://www.reptile-database.org/>
- Valdujo, P. H., Nogueira, C., & Martins, M. (2002). Ecology of *Bothrops newwiedi pauloensis* (Serpentes: Viperidae: Crotalinae) in the Brazilian Cerrado. *Journal of Herpetology*, 36, 169–176.
- Vuoto, J. A. (2000). Análisis de ejemplares del oeste (w) de la Provincia de Entre Ríos de *Micrurus altirostris* (Cope, 1859) (Serpentes: Elapidae) en el Museo “Profesor Antonio Serrano” de Paraná, Entre Ríos, Argentina. *Museo de Ciencias Naturales y Antropológicas “Prof. Antonio Serrano”, Memorias del Museo y Serie Nueva*, 7, 1–13.
- Zar, J. H. (1999). *Biostatistical analysis. 4th Edition*. Upper Saddle River: Prentice Hall.