

Validation of the EuroSCORE model in patients undergoing heart surgery in Regional Hospital of High Speciality of Bajío

Validación del modelo EuroSCORE en pacientes sometidos a cirugía cardíaca en el Hospital Regional de Alta Especialidad del Bajío

Raúl Teniente-Valente,* Francisco E González-Bravo,**
Miguel Ángel Chagolla-Santillán,* Iliana Acevedo-Bañuelos,*
Enrique Alfredo Bernal-Ruíz,* Luis Rey López-Luis,*
Oscar Samuel Medina-Torres,* Ricardo Elías Romo-Escamilla,*
José Esteban Landeros-Aranda**

Key words:

Cardiac surgery,
risk stratification,
EuroSCORE,
validation.

Palabras clave:

Cirugía cardíaca,
estratificación de
riesgos, EuroSCORE,
validación.

*Cardiology and cardiovascular surgery service. Regional Hospital of High Speciality of Bajío.
** Health Sciences Division. University of Guanajuato.

Received:
08/04/2018
Accepted:
10/09/2018

ABSTRACT

Introduction: EuroSCORE is a probabilistic model with good performance in the prognosis of mortality in heart surgery in many latitudes. It is recommendable to validate it in hospitals where it is employed. **Objective:** To validate the EuroSCORE model in adult patients at the Hospital Regional de Alta Especialidad del Bajío (HRAEB) of León, Guanajuato, Mexico. **Material and methods:** We conducted an observational transversal, and retrospective study, accomplishing this through the review of the clinical files of patients submitted to heart surgery with and without extracorporeal circulation pump from 01/01/2008 to 12/31/2013 at the HRAEB. This included mortality up to hospital discharge, utilizing the on-line calculator of the EuroSCORE program to estimate risk of death. In order to validate the EuroSCORE model, we assessed discrimination and calibration through the Area Under the receiver operating characteristics (ROC) curve and χ^2 test with Hosmer-Lemeshow (H-L) goodness-of-fit, respectively. **Results:** A total of 342 patients, aged 50.02 ± 16.66 years, 181 males (52.9%) and 161 women (47.1%). The area under the ROC curve of the additive model was 0.763, and of the Hosmer-Lemeshow test was 5.30, with $p = 0.62$. The area under the ROC curve of the logistic model was 0.761 and of the Hosmer-Lemeshow test, 8.78, with $p = 0.36$. **Conclusion:** The EuroSCORE model is a reliable score for estimating the probabilities of death in adult patients submitted to heart surgery with or without the pump at the HRAEB.

RESUMEN

Introducción: EuroSCORE es un modelo probabilístico con buen desempeño en el pronóstico de mortalidad en cirugía cardíaca en muchas latitudes. Es recomendable validarlo en los hospitales donde se emplea. **Objetivo:** Validar el modelo EuroSCORE en pacientes adultos en el Hospital Regional de Alta Especialidad del Bajío (HRAEB) de León, Guanajuato, México. **Material y métodos:** Se realizó un estudio observacional transversal y retrospectivo, que se completó con la revisión de los expedientes clínicos de pacientes sometidos a cirugía cardíaca con y sin bomba de circulación extracorpórea desde el 01/01/2008 hasta el 31/12/2013 en el HRAEB. Esto incluyó la mortalidad hasta el alta hospitalaria, utilizando la calculadora en línea del programa EuroSCORE para estimar el riesgo de muerte. Para validar el modelo EuroSCORE, evaluamos la discriminación y la calibración a través de la curva de características de operación del receptor (ROC) y la prueba de χ^2 con la efectividad de ajuste de Hosmer-Lemeshow (H-L), respectivamente. **Resultados:** Un total de 342 pacientes, de 50.02 ± 16.66 años, 181 hombres (52.9%) y 161 mujeres (47.1%). El área bajo la curva ROC del modelo aditivo fue de 0.763, y la de la prueba Hosmer-Lemeshow fue de 5.30, con $p = 0.62$. El área bajo la curva ROC del modelo logístico fue 0.761 y de la prueba Hosmer-Lemeshow, 8.78, con $p = 0.36$. **Conclusión:** El modelo EuroSCORE es una herramienta confiable para estimar las probabilidades de muerte en pacientes adultos sometidos a cirugía cardíaca con o sin bomba en el HRAEB.

INTRODUCTION

The practice of Medicine at present demands quality care. Multiple indicators are utilized to measure the latter; however, the clinical results are, in the end, the most relevant.¹ For some investigators, mortality is the first and most important indication of quality in the clinical practice.¹ Crude or non-adjusted mortality has its limitations in that it does not consider, among other aspects, the clinical conditions of the sick person nor his/her comorbidities.^{1,2} In heart surgery, statistical programs have been developed from a group of risk factors; these factors are capable of predicting mortality.² For comparing the mortality rate among diverse institutions or for temporary comparison at a same institution, it is recommended to employ some of the mathematical models developed that evaluate mortality adjusted to the clinical conditions and risk profile of the population cared for.^{1,2} During the last decade, the Parsonnet score, that of the U.S. Society of Thoracic Surgeons (STS), and the EuroSCORE³ score have been those most frequently utilized. The latter was obtained from the study of 19,030 adult patients submitted to heart surgery (ischemic, valvular, and congenital) with extracorporeal circulation pump in 128 hospitals in eight European countries (Germany, France, the U.K., Italy, Spain, Finland, Sweden, and Switzerland). In this study, based on its objectivity, reliability, and prevalence, the authors analyzed 97 variables or risk factors (68 preoperative and 29 operative) were analyzed, among which 18 remained with prognostic value.²⁻⁴ The EuroSCORE model has two variants: additive and logistic. The former awards a numerical value to each variable that the patient presents, and the sum of these values provides the probability of death.⁴ It entertains the advantage of being a simple and uncomplicated instrument that can be applied at the patient's bedside. The logistic version is more sophisticated. The formula for obtaining the latter is the following: estimated mortality = $e^{(\beta_0 + \sum \beta_i X_i)} / 1 + e^{(\beta_0 + \sum \beta_i X_i)}$, in which β_0 is the constant of the logistic regression model (-4.789594) and β_i is the coefficient of variable X_i of the logistic-regression information.⁵ The definition, the odds ratio (OR), the value of each

variable of the additive model, and coefficient β of each variable of the logistic model appear in *Table 1*. To utilize a risk score or a predictive model developed to predict mortality in patients submitted to heart surgery, which has shown good performance in the group of patients where it was generated, thus proving its internal validity, it is recommended that it be evaluated at the hospitals where its use is intended to confirm or discard external validity, therefore using it or not.^{1,2,4} The objective of the present study was to know the risk profile of the adult population submitted to cardiac surgery and to validate its additive as well as in its logistic version in the population of adult patients submitted to cardiac surgery with and without extracorporeal circulation pump at the *Hospital Regional de Alta Especialidad del Bajío* (HRAEB) in the city of León, Guanajuato, Mexico.

MATERIAL AND METHODS

We conducted an observational, transversal, and retrospective study by means of the review of the clinical files of patients aged of ≥ 16 years submitted to cardiac surgery with and without extracorporeal circulation pump that were effected from January 1, 2008 to December 31, 2013 at the HRAEB. We did not include incomplete files, or those with doubtful information on the variables-of-interest, nor the files of patients who died due to causes not related to the cardiac surgery index. The following variables were obtained: mortality not adjusted up to the patient's hospital discharge, defined as death occurring during the hospitalization index; type of surgery, defined as the procedure or procedures carried out during the index surgery, whether a) valvular surgery, b) aortic-coronary bypass surgery, c) corrective surgery for some congenital malformation(s), and d) surgery of another different type, including valvular + coronary bypass surgery, aortic surgery, closing of post-infarct interventricular communication, traumatic cardiac lesions, pericardium resection). The required variables were collected and the mortality-risk score was calculated of the EuroSCORE employing the EuroSCORE-program on-line calculator in its additive as well as its logistic version (www.euroscore.org/calsp.html).

Statistical treatment

The qualitative variables are presented in percentages of frequency and are compared with the χ^2 test, the Cochran Q test, and the

Friedman ranges. The numerical variables are presented as averages and standard deviations (SD). Comparison of the averages between the two groups was performed with the student *t* test, and comparison among three or more

Table 1: Risk variables of the additive and logistic EuroSCORE model and their definition.

Variable	Definition	OR	Points	β
Patient factors				
Age	Every 5 years from the age of 60 years	1.1	1	0.066635
Gender	Female	1.4	1	0.330405
Serum creatinine	> 200 $\mu\text{mol/L}$	1.9	2	0.652165
Extracardiac arteriopathy	Claudication of the lower limbs	1.9	2	0.655892
	Carotid stenosis $\geq 50\%$, prior or planned vascular surgery on the abdominal aorta, carotids o peripheral arteries			
Chronic lung disease	Requires prolonged treatment with bronchodilators or steroids	1.6	1	0.493134
Neurological dysfunction	Neurological damage that severely affects walking or daily activity	2.3	2	0.841626
Previous cardiac surgery	Cardiac surgery that required opening of the pericardium	2.6	3	1.002625
Active endocarditis	The patient is under antibiotic treatment at the time of surgery	2.5	3	1.101265
Critical preoperative state	One of the following conditions: ventricular tachycardia/ventricular fibrillation or sudden death recovered, preoperative cardiac massage, mechanical ventilation prior to anesthesia, preoperative inotropic, acute Preoperatory kidney failure (oliguria ≤ 10 mL/hour)	2.2	3	0.905813
Cardiac factors				
Unstable angina	Resting angina requiring intravenous nitrates until arrival at the operating room	1.5	2	0.567708
Recent myocardial infarction	Previous myocardial infarction in the last 90 days	1.6	2	0.546022
Expulsion fraction of the VI	Expulsion fraction less than 30%	2.5	3	1.09443
Expulsion fraction of the VI	30-50%	1.5	1	0.419643
Pulmonary artery systolic	> 60%	2.0	2	0.767692
Factors of surgery				
Emergency surgery	Required before next work day	2.8	2	0.712795
Rupture of Septum IV		3.8	4	1.462009
Surgery different from coronary revascularization	Cardiac surgery other than coronary revascularization or in addition to it	1.6	2	0.542036
Thoracic aortic surgery	Surgery in ascending aorta arch or descending aorta	3.2	3	1.159787

averages was carried out with ANalysis Of VAriance (ANOVA) and the Bonferroni *post-hoc* test. A significance level of $p \leq 0.05$ was accepted. To validate the EuroSCORE model, we evaluated discrimination and calibration. Discrimination consists of the capacity of the model to identify the patients who will survive those who will die. This was evaluated by means of the area under the receiver operating characteristics (ROC) Curve. Values of ≤ 0.5 speaks to the model that does not discriminate better than chance, and values of 1 indicate perfect discrimination. Values greater than 0.75 identify systems with a good capacity of discrimination of the model.³ Calibration consists of the comparison of the expected episodes against those observed along the entire risk range. We carried out calibration with the χ^2 test of goodness-of-fit of

the Hosmer and Lemeshow (H-L) adjustment, which calculates a statistical C, with which the difference can be evaluated between the values of mortality predicted by the model and the mortality values observed in the distinct groups of the population studied. The lower the value of this statistic, the better the calibration of the model (the predicted and the observed mortality becomes closer). A p value greater than 0.05 suggests that the model fits well, and consequently, it will predict well the probability of the patients dying.

RESULTS

From the list of cardiac-surgery and cardiovascular-surgery procedures that were carried out from January 1, 2008 to December 31, 2013, which were collected by the HRAEB Department of Statistics, we identified 350 surgical procedures. Among the latter, we eliminated eight as follows: two for not having complete data-of-interest in the file; one for having been triplicated and with different information, and four due to death unrelated to the cardiac-surgical-procedure index. A total of 342 procedures remained that complied with the selection criteria and brought together and that are those that comprise the material of the present report. The general characteristics of the study group appear in *Table 2*. The frequency of each of the variables of the population studied and its comparison with that obtained in the EuroSCORE study are depicted in *Table 3*. As we can observe, there are notable differences in the prevalence of the majority of the risk factors. The population of the EuroSCORE study is 12 years older than that of our study and presents higher rates of extracardiac arteriopathy and of the left ventricle expulsion fraction (LVEF) of $< 30\%$. On the other hand, our population has the following: a greater percentage of women; of chronic pulmonary disease; neurological dysfunction; prior cardiac surgery; creatinine values above 200 $\mu\text{mol/L}$ (2.26 mg/dL); active endocarditis; critical preoperative status; unstable angina; pulmonary arterial hypertension; emergent surgeries, and surgeries different from those of coronary bypass and thoracic aorta surgeries. In

Table 2: Clinical and demographic characteristics of the group studied.

N	342
Age	
Mean \pm SD	50.02 \pm 16.6
% \leq 50 years	40.6
51-60 years	29.8
61-70 years	21.6
\geq 70 years	7.9
Gender male/female	181/161
% Diabetes	26.6
Hypertension	45.3
Hypercholesterolemia	21.6
Smoking	30.1
% Type of surgery	
Valvular	34.2
Coronary bypass	29.5
Congenital	14.6
Another type	21.6
LVEF (mean \pm SD)	54.4 \pm 11.6 (321/342)
Systolic pulmonary pressure (mean \pm SD)	42.6 \pm 17 (298/342)
Additive EuroSCORE (mean \pm SD)	5.3 \pm 3.2
Logistic EuroSCORE (mean \pm SD)	7.6 \pm 10.6
Extracorporeal circulation time (mean \pm SD)	118.8 \pm 54.4 (179/342)
Aortic clamping time (mean \pm SD)	89.5 \pm 44.4 (175/342)
Dead patients	37/342 (10.8%)

LVEF = left ventricular ejection fraction (mmHg).

Table 4, we can appreciate the distribution of patients according to risk. As can be observed, only 16.9% of patients were low risk. The remainder were medium (44.4%) or high risk (38.6%). This is in agreement with the greater prevalence of more than 50% of the risk factors

of our studied population in comparison with the EuroSCORE study population. There were 27 deaths, which yielded a crude mortality rate of 10.8%. In Table 4, we may observe the risk-adjusted mortality as follows: low-risk group 1/58 (expected, < 2%; observed, 1.7%), intermediate-risk group 7/152 (expected, < 5%; observed, 4.6%), and the high-risk group 29/132 (expected, > 6%; observed, 22.0%). Table 5 presents the comparison of the variables of the following: the EuroSCORE model; type of surgery; the additive and logistic EuroSCORE score; and time of extracorporeal circulation between the group of patients who died and the group who survived. We observed a higher age in the group of patients who died (53.11 ± 13.87 vs. 49.72 ± 17.5 years, $p = ns$), as well as a notably greater prevalence in 10 variables of the model (extracardiac arteriopathy, neurological dysfunction, prior cardiac surgery, serum creatinine of > 2.2 mg/dL, active endocarditis, critical preoperative status, LVEF 30-50%, LVEF of < 30%, PAH > 60 mmHg, and emergent surgery). Consequently, the additive EuroSCORE model (7.92 ± 3.63 vs. 5.11 ± 3.04 , $p = 0.0005$) and the logistic EuroSCORE model (15.62 ± 15.99 vs. 6.70 ± 8.65 , $p = 0.0005$) were greater in the group of patients who died. The extracorporeal circulation pump time (ECPT) was also higher in the group of patients who died (106.33 ± 73.76 vs. 82.91 ± 48 , $p = 0.0005$).

Validation of the EuroSCORE model in our studied population yielded the following results: for the additive version, the area under the receiver operating curve (ROC) was 0.763 (Figure 1), and the χ^2 of the H-L test was 5.30, with $p = 0.62$ (Figure 2). For the logistic version, the area under the ROC curve was 0.761 (Figure 3) and H-L test was 8.78, with $p = 0.36$ (Figure 4).

Table 3: Prevalence of risk factors. EuroSCORE vs HRAEB.

Variable	EuroSCORE (n = 19,030)	HRAEB (n = 342)
Age (mean \pm SD)	62.5 \pm 10.7	50.02 \pm 16.6
< 60 years (%)	33.2	65.2
60-64	17.8	14.3
65-69	20.7	9.3
70-74	17.9	6.7
\geq 75	9.6	4.3
Female gender	27.8	47.0
Pulmonary disease	3.9	12.86
Extracardiac arteriopathy	11.3	2.63
Neurological dysfunction	1.4	2.04
Previous cardiac surgery	7.3	10.23
Serum creatinine > 200 μ mol/L	1.8	8.47
Active endocarditis	1.1	4.97
Critical preoperative status	4.1	14.61
Unstable angina	8.0	11.69
LVEF 30-50%	25.6	24.26
LVEF < 30%	5.8	2.04
Recent myocardial infarct (< 90 days)	9.7	9.06
Systolic pulmonary pressure > 60 mmHg	2.0	9.64
Emergency operation	4.9	16.9
Non-coronary surgery	36.4	71.05
Thoracic aortic surgery	2.4	3.80
Ventricular septal rupture	0.2	0.29

LVEF = left ventricular ejection fraction (%).

Table 4: Expected and observed mortality by the additive EuroSCORE model.

Risk groups	Patients (n)	Mortality	Observed (IC 95%)	Expected (IC 95%)
Low (0-2)	58	1 (1.7%)	1.91-3.51	1.01-3.51
Medium (3-5)	152	7 (4.6%)	2.51-9.78	2.87-11.17
High (\geq 6)	132	29 (22.0%)	9.44-39.40	9.79-38.04

Table 5: Comparison of the EuroSCORE variables between the group of surviving patients and the group of deceased patients.

Variables	Survivor group n = 305	Deceased group n = 37
Age (mean ± SD)	49.72 ± 17.5	53.11 ± 13.87
% Female	46.22	54.05
Pulmonary disease	12.78	13.51
Extracardiac arteriopathy	1.96	8.10
Neurological dysfunction	0.98	10.81
Previous cardiac surgery	7.54	32.40
Serum creatinine > 200 µmol/L	6.88	21.62
Active endocarditis	4.59	8.10
Critical preoperative state	12.45	32.42
Unstable angina	11.80	10.80
LVEF 30-50%	22.95	35.13
LVEF < 30%	1.31	8.10
Recent myocardial infarct (< 90 days)	9.18	8.10
Systolic pulmonary pressure > 60 mmHg	8.85	16.21
Emergency operation	15.40	29.72
Non-coronary surgery	70.16	75.67
Thoracic aortic surgery	3.60	5.40
Ventricular septal rupture	0.32	0
% Valve surgery (n = 117)	84.61	15.39
Isolated coronary surgery (n = 101)	94.05	5.94
Congenital surgery (n = 50)	96	4
Others surgery (n = 74)	85.13	14.86
Additive EuroSCORE (mean ± SD)	5.11 ± 3.04	7.92 ± 3.63
Logistic EuroSCORE (mean ± SD)	6.70 ± 8.65	15.62 ± 15.99
Extracorporeal circulation time (mean ± SD)	82.91 ± 48.00	106.33 ± 73.76

DISCUSSION

The additive version of the EuroSCORE model was evaluated in a sample of 1,497 patients obtained from the population in which the model was developed, noting discrimination with an area under the ROC curve of 0.76 and a calibration measured with the H-L test of 7.5 ($p < 0.68$), confirming good internal validation.⁴ Later, the equation employed for the calculation of its logistic version was revealed.⁵ The goodness-of-fit of the additive and logistic EuroSCORE model has been evaluated in multiple and different populations, in general cardiac surgery and in its subgroups, with and without the extracorporeal circulation pump. Perhaps this is the probabilistic model that is most evaluated to predict mortality

in heart surgery worldwide, with studies conducted in Europe, Asia, Australia, and in America, with different results.⁶⁻¹⁸ In Europe, in a study that evaluated the performance of the score in populations of the six countries that contributed more than 500 patients to develop the EuroSCORE model, the authors found that, despite the notable epidemiological differences among these populations, the adjustment of the score for predicting mortality was good, with a discrimination evaluated with the area under the ROC curve of 0.82 and calibration evaluated with the H-L goodness-of-fit test of 0.59 ($p \geq 0.05$).⁶ In the U.K., the logistic version of the model was evaluated in nearly 10,000 patients, 67.5% with coronary bypass, 15.33% with isolated valvular surgery, and 15.77% valvular aortic surgery + coronary

bypass, observing good discrimination, with an area under the ROC curve of 0.79 for the whole group, of 0.77 for the coronary-bypass sub-group, and of 0.79 for valvular surgery; however, calibration was not good. The authors concluded that the logistic version model possesses good discrimination, but that calibration varies among the different risk sub-groups, with overestimation of the mortality




Figure 1: Receiver operating characteristic (ROC) curve. Area under the curve = 0.76, compatible with good discrimination of EuroSCORE additive model.




Figure 2: The Hosmer-Lemeshow goodness-of-fit test = 5.30, with $p = 0.62$. The mortality observed was very similar to the mortality estimated in the three risk groups.




Figure 3: Receiver operating characteristic (ROC) curve. Area under the curve = 0.76, compatible with good discrimination for the EuroSCORE logistic model.




Figure 4: The data divided into quintiles show the result of the Hosmer-Lemeshow goodness-of fit test at 8.78, with $p = 0.36$, compatible with good calibration for EuroSCORE logistic model.

observed. These authors recommended recalibration of the model.⁷ In Spain, in a study that included nearly 500 patients submitted to coronary bypass, adjustment of the model was evaluated and the additive and logistic versions were compared. The model's discrimination was very good, with an area under the ROC curve of 0.830 for the logistic version and of 0.839 for the additive version. The value for the statistical C of the χ^2 equation of H-L was 11.51 with $p = 0.32$ for the logistic version. The authors concluded that the logistic model is in greater approximation to the real mortality in the high-risk patient sub-group.⁸ Other studies have also compared the additive and logistic versions, observing that both versions possess satisfactory discrimination, with areas under the ROC curve of 0.80 (logistic) and 0.79 (additive). However, calibration was better with the logistic version ($p = 0.12$) than with the additive version ($p = 0.001$).⁹ In the population with highest risk, better performance of the logistic version of the model was observed.⁵ The results obtained in other studies have also found limitations in the model's additive version. For example, in a study that included six large international samples, the result observed strongly suggest that the additive version overestimates mortality in low-risk groups (EuroSCORE ≤ 6 points) and underestimates mortality in high-risk groups (EuroSCORE ≥ 13 points).¹⁰

In Australia, in a study conducted with 8,331 patients (valvular, coronary bypass, and aortic surgery) to assess the goodness-of-fit of the model in its population, it was found that the discriminatory power of both variants of the model were very good, with an area under the ROC curve for the whole cohort of 0.83, and for the coronary-bypass sub-group, one of 0.82. However, calibration of both variants was poor in terms of predicting mortality in risk sub-groups, in that this was underestimated in nearly all of the risk deciles (χ^2 of H-L with a $p < 0.05$).¹¹ In the U.S., based on the STS data of the EuroSCORE model in its additive version, it revealed good to very good discrimination in general cardiac surgery and in coronary-bypass surgery, with areas under the ROC curve of 0.75 and 0.78, respectively. Calibration was evaluated in five risk sub-groups, noting that the mortality observed was nearly identical to

that foreseen for the model in all of the sub-groups. The authors concluded that, despite the epidemiological and demographic differences between Europe and North America, the EuroSCORE model can be recommended as a simple and uncomplicated instrument for application on both side of the Atlantic.¹² In Latin America, the EuroSCORE model has been evaluated in diverse studies. In Brazil, Moraes de Carvalho et al., in a study of four hospitals of the city of Rio de Janeiro with a randomized sample of 546 patients of a population submitted to coronary bypass, the model demonstrated poor discriminatory power, with an ROC curve of 0.62.¹³ In Argentina, Scaro and collaborators conducted a study with 123 patients to test the discriminatory power of the EuroSCORE model, concluding that the model was not useful for predicting mortality, in particular in sub-groups of intermediate and high-risk.¹⁴ In Colombia, Parga-Gómez et al., in a study of 498 patients, found good discrimination for the additive as well as for the logistic version (area under the ROC curve of 0.85 in both versions), in addition to good calibration (statistic C of χ^2 of H-L 3.39 additive version, and 9.99 for the logistic version with p of 0.8 and 0.2, respectively).¹⁵ In Cuba, Chao-García and collaborators, in a study that included 158 patients submitted to mitral valve surgery, reported an area under the ROC curve of 0.97, that is, excellent discrimination.¹⁶ In Mexico, Careaga et al., in a study of 206 patients submitted to valvular surgery, the authors found an area under the ROC curve of 0.77 for the additive version and of 0.97 for the logistic version, values that are compatible with good and excellent discrimination, respectively. The χ^2 of H-L was 6.7 ($p = 0.034$) and 2.86 ($p = 0.99$), the latter compatible with very good calibration.¹⁷ Rodríguez Chávez et al., in another study in Mexico, evaluated the model in 1,188 patients submitted to valvular surgery and found an Area Under the ROC Curve of 0.707 and of 0.694 for the additive and logistic versions, respectively, while the value of the χ^2 of H-L was 63.15 ($p < 0.001$) and 45.6 ($p < 0.001$) for the corresponding additive and logistic versions. These authors concluded that prediction of mortality in this population of patients, in the additive as well as in the logistic

version, was inadequate and that discrimination was scarcely borderline with the additive and poor for the logistic version.¹⁸

In our study, we observed that, while the population evaluated was 12 years younger than the European population from which the model arose, the prevalence of diabetes mellitus (DM) was greater, as well as of the majority of the prognostic variables of the score already cited in the results section. However, despite the epidemiological differences between the EuroSCORE development group and those of our population, the results obtained in the present study are compatible with good discrimination and good calibration of the model, in its additive as well as in its logistic version, and are similar to the results reported in Europe,^{6,8} the U.S.,¹² Colombia,¹⁵ and Careaga in our environment.¹⁷ It is noteworthy that our evaluated population included vascular surgery, coronary bypass, and congenital surgery in adults, and also other types of surgery that included infrequent cardiac pathologies such as those produced by non-shooting weapons and/or by firearms, that is, it was a population submitted to cardiac surgery for all of the pathologies.

Why are there discrepancies in the results in different studies? The explanations can be several. On the one hand, the continuous progress in cardiovascular surgery (technological, better organization of medical groups, acquisition of experts of surgical groups and of the groups charged with perioperative management, among others) and, on the other hand, improvement in the coverage of the health systems in countries on various continents, as well as a population with a higher degree of education in health, have yielded as results the progressive diminution of perioperative mortality in these patients (despite their being older and having more comorbidities), more notorious in developed countries, which has originated that a determined score that has shown a good prognostic performance during the time in which it was developed, can, a few years later, result in poor discrimination and/or calibration, as suggested in some studies.¹⁹ From this arises the suggestion to update these instruments, as has already been done in some models (in the Parssonnet in its different

versions and in the EuroSCORE model, which has already developed model II, and in the STS score, which performs this periodically and in a dynamic manner).³ Perhaps it would also be convenient to develop models for some surgical groups in particular, such as valvular surgery²⁰ and infectious endocarditis surgery,²¹⁻²³ as has been carried out in some places, with the purpose of possessing more reliable instruments for utilization in these specific pathologies.

CONCLUSIONS

The EuroSCORE model showed to be, at present, a reliable predictive instrument for estimating the probabilities of death in patients submitted to cardiac surgery with extracorporeal pump or without the latter in adult population cared for at the HRAEB.

REFERENCES

1. López-Sendón J, González-Juanatey JR, Pinto F, Cuenca-Castillo J, Badimón L, Dalmau R et al. Indicadores de calidad en cardiología. Principales indicadores para medir la calidad de los resultados (indicadores de resultados) y parámetros de calidad relacionados con mejores resultados en la práctica clínica (indicadores de práctica asistencial). INCARDIO (Indicadores de Calidad en Unidades Asistenciales del Área del Corazón): Declaración de posicionamiento de consenso de SEC/SECTCV. *Rev Esp Cardiol*. 2015; 68 (11): 976-995.e10.
2. Roques F, Nashef SA, Michel P, Gauducheau E, de Vincentiis C, Baudet E et al. Risk factors and outcome in European cardiac surgery: analysis of the EuroSCORE multinational database of 19030 patients. *Eur J Cardiothorac Surg*. 1999; 15 (6): 816-822; discussion 822-823.
3. Cortina-Romero JM, López-Gude MJ, Pérez de la Sota E. Capítulo 2. Estratificación y análisis del riesgo quirúrgico. En: Otero E, Rupilanchas JJ, Belda FJ. Riesgo y complicaciones en cirugía cardíaca. Madrid, España: Editorial Médica Panamericana; 2005. pp. 15-27.
4. Nashef SA, Roques F, Michel P, Gauducheau E, Lemeshow S, Salamon R. European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardiothorac Surg*. 1999; 16 (1): 9-13.
5. Roques F, Michel P, Goldstone AR, Nashef SA. The logistic EuroSCORE. *Eur Heart J*. 2003; 24 (9): 881-882.
6. Roques F, Nashef SA, Michel P, Pinna Pintor P, David M, Baudet E et al. Does EuroSCORE work in individual European countries? *Eur J Cardiothorac Surg*. 2000; 18 (1): 27-30.
7. Bhatti F, Grayson AD, Grotte G, Fabri BM, Au J, Jones M et al. The logistic EuroSCORE in cardiac surgery: how well does it predict operative risk? *Heart*. 2006; 92 (12): 1817-1820.

8. Lafuente S, Trilla A, Bruni L, González R, Bertrán A, Pomar JL et al. Validación del modelo probabilístico EuroSCORE en pacientes intervenidos de injerto coronario. *Rev Esp Cardiol*. 2008; 61 (6): 589-594.
9. Zingone B, Pappalardo A, Dreas L. Logistic versus additive EuroSCORE. A comparative assessment of the two models in an independent population sample. *Eur J Cardiothorac Surg*. 2004; 26 (6): 1134-1140.
10. Gogbashian A, Sedrakyan A, Treasure T. EuroSCORE: a systematic review of international performance. *Eur J Cardiothorac Surg*. 2004; 25 (5): 695-700.
11. Yap CH, Reid C, Yui M, Rowland MA, Mohajeri M, Skillington PD et al. Validation of the EuroSCORE model in Australia. *Eur J Cardiothorac Surg*. 2006; 29 (4): 441-446; discussion 446.
12. Nashif SA, Roques F, Hammill BG, Peterson ED, Michel P, Grover FL et al. Validation of European System for Cardiac Operative Risk Evaluation (EuroSCORE) in North American cardiac surgery. *Eur J Cardiothorac Surg*. 2002; 22 (1): 101-105.
13. Carvalho MR, Souza e Silva NA, Klein CH, Oliveira GM. Application of the EuroSCORE in coronary artery bypass surgery in public hospitals in Rio de Janeiro, Brazil. *Rev Bras Cir Cardiovasc*. 2010; 25 (2): 209-217.
14. Scaro G, Shireishi P, Gerbaudo L, Gallo C, Moisés-Azize G, Guzmán L et al. Valor predictivo del "EUROSCORE" en el hospital público. 4o Congreso Virtual de Cardiología. Available in: <http://www.fac.org.ar/ccvc/llave/tl447/tl447.pdf>.
15. Parga-Gómez R, Buitrago-Gutiérrez G, Roldán-Henao J. Validación del euroSCORE en la valoración del riesgo quirúrgico en un centro de referencia cardiovascular en Colombia. *Rev Mex Cardiol*. 2013; 24 (3): 138-143.
16. Chao GJL, Vázquez RFJ, Mirabal RR, Bermúdez YGJ, Chaljub BE, Medrano PY et al. Validación temporal del modelo EuroSCORE para la evaluación de los resultados de la cirugía de sustitución valvular mitral. *CorSalud*. 2013; 5 (2): 170-175.
17. Careaga-Reyna G, Martínez-Carballo G, María Anza-Costabile L, Ávila-Funés A. Euroscore para predecir morbimortalidad en cirugía cardíaca valvular. *Cir Cir*. 2008; 76 (6): 497-505.
18. Rodríguez-Chávez LL, Figueroa-Solano J, Muñoz-Consuegra CE, Avila-Vanzzini N, Kuri-Alfaro J. EuroSCORE subestima el riesgo de mortalidad en cirugía cardíaca valvular de población mexicana. *Arch Cardiol Méx*. 2017; 87 (1): 18-25.
19. Borracci RA, Rubio M, Cortés y Tristán G, Mémoli R, Giorgi M, Ahuad-Guerrero RA. Validez temporal de los sistemas de estratificación por riesgo para la monitorización continua de los resultados de la cirugía cardíaca. *Rev Argent Cardiol*. 2005; 73: 1-5.
20. Vieira Da Costa GJC, Bodanese LC, Lucas BF, Goldani MA. Propuesta de score de riesgo preoperatorio para pacientes candidatos a cirugía cardíaca valvular. *Arq Bras Cardiol*. 2010; 94 (4): 523-530.
21. Hasbun R, Vikram HR, Barakat LA, Buenconsejo J, Quagliarello VJ. Complicated left-sided native valve endocarditis in adults: risk classification for mortality. *JAMA*. 2003; 289 (15): 1933-1940.
22. Gaca JG, Sheng S, Daneshmand MA, O'Brien S, Rankin JS, Brennan JM et al. Outcomes for endocarditis surgery in North America: a simplified risk scoring system. *J Thorac Cardiovasc Surg*. 2011; 141 (1): 98-106.e1-2.
23. De Feo M, Cotrufo M, Carozza A, De Santo LS, Amendolara F, Giordano S et al. The need for a specific risk prediction system in native valve infective endocarditis surgery. *Scientific World Journal*. 2012; 2012: 307571.

Correspondence to:

Raúl Teniente-Valente

Av. Paseo de Jerez # 405-6,
Colonia Jardines de Jerez,
León, Guanajuato, México.
Tel: (477) 7150043
Móvil: (477) 1379007
E-mail: dr.teniente@yahoo.com