

# Protolith age of the Altar and Carnero complexes and Late Cretaceous–Miocene deformation in the Caborca–Altar region of northwestern Sonora, Mexico

Carl E. Jacobson<sup>1,2,\*</sup>, César Jacques-Ayala<sup>3</sup>, Andrew P. Barth<sup>4</sup>,  
Juan Carlos García y Barragán<sup>3</sup>, Jane N. Pedrick<sup>1</sup>, and Joseph L. Wooden<sup>5</sup>

<sup>1</sup>Department of Geological and Atmospheric Sciences, Iowa State University, 253 Science Hall, 2237 Osborn Drive, Ames, IA 50011, USA.

<sup>2</sup>Department of Earth and Space Sciences, 207 Merion Science Center, 720 S. Church Street, West Chester University of Pennsylvania, West Chester, PA 19355, USA.

<sup>3</sup>Instituto de Geología, Universidad Nacional Autónoma de México, Blvd. Colosio s/n y Madrid, Apartado Postal 1039, Hermosillo, Son. 83000, Mexico.

<sup>4</sup>Department of Earth Sciences, Indiana University-Purdue University Indianapolis, 723 W. Michigan Street, SL118, Indianapolis, IN 46202, USA.

<sup>5</sup>U.S. Geological Survey, 345 Middlefield Road, Menlo Park, CA 94025, USA.

\*cejac@iastate.edu

## ABSTRACT

In the Caborca–Altar area of northwest Sonora, variably deformed and metamorphosed sedimentary and volcanic rocks crop out in a northwest-southeast-trending belt (El Batamote belt) at least 70 km long. We obtained detrital zircon U–Pb ages from two distinctive components of the belt near Altar, here termed the Altar complex and Carnero complex. Zircon ages for metasandstone and metaconglomerate matrix from the Altar complex indicate a Late Cretaceous maximum age of sedimentation, with at least part of the complex no older than  $77.5 \pm 2.5$  (2 $\sigma$ ). Pre-Cretaceous detrital zircons in the complex were derived largely from local sources, including Proterozoic basement, the Neoproterozoic–Cambrian miogeocline and the Jurassic arc. The detrital zircon ages and lithologic character of the Altar complex suggest correlation with the Escalante Formation, the uppermost unit of the Upper Cretaceous El Chanate Group. In contrast, one sample from the Carnero complex yielded a Middle Jurassic maximum depositional age and a detrital zircon age distribution like that of the Jurassic eolianites of the North American Cordillera. The Carnero complex may correlate with the Middle Jurassic Rancho San Martín Formation but could also be a metamorphosed equivalent of the Upper Jurassic Cucurpe Formation, Upper Jurassic to Lower Cretaceous Bisbee Group, or El Chanate Group derived by recycling of Jurassic erg sandstones. The Late Cretaceous age for the Altar complex protolith contradicts models that relate deposition of the entire El Batamote protolith to a basin formed by oblique slip along the Late Jurassic Mojave–Sonora megashear. Instead, the belt is best explained as an assemblage of Middle Jurassic to Upper Cretaceous formations deformed and locally metamorphosed beneath a northeast-directed Laramide thrust complex. Potassium–argon and <sup>40</sup>Ar/<sup>39</sup>Ar ages confirm previous inferences that deformation of El Batamote belt occurred between the Late Cretaceous and late Eocene. A second phase of deformation, involving low-angle normal faults, occurred

during and/or after intrusion of the ~22–21 Ma Rancho Herradura granodiorite.

Key words: Altar complex; Carnero complex; El Batamote belt; El Chanate Group; Mojave–Sonora megashear; Laramide orogeny.

## RESUMEN

En el área de Caborca–Altar, noroeste de Sonora, aflora una franja orientada noroeste-sureste de, cuando menos, 70 km de longitud, a la que se ha denominado franja El Batamote; esta franja está constituida por rocas sedimentarias e ígneas deformadas y metamorizadas en grados muy variables. Obtuvimos edades de U–Pb de circones detríticos de dos componentes distintivos de la franja cercana a Altar que denominaremos complejo Altar y complejo Carnero. Edades obtenidas de circones de metareniscas y de la matriz de metaconglomerados del complejo Altar indican una edad máxima de sedimentación del Cretácico Tardío; al menos una parte del complejo no es mayor que  $77.5 \pm 2.5$  (2 $\sigma$ ). Circones pre-cretácicos del complejo se derivaron de fuentes locales, incluyendo el basamento proterozoico, el miogeoclinal neoproterozoico–cámbrico y el arco jurásico. Las edades de los circones detríticos y el carácter litológico del complejo Altar sugieren una correlación con la Formación Escalante, la unidad superior del Grupo El Chanate del Cretácico Tardío. En contraste, una muestra del complejo Carnero aportó circones que indican una edad máxima de depósito del Jurásico Medio y una distribución de edades semejante a las de las eolianitas jurásicas de la provincia Cordillerana de Norteamérica. El complejo Carnero puede correlacionarse con la Formación Rancho San Martín del Jurásico Tardío, pero podría ser también un equivalente metamórfico del Grupo Bisbee del Jurásico Tardío–Cretácico Temprano, o del Grupo El Chanate, derivados del reciclado de las areniscas de ergs jurásicos. La edad Cretácico Tardío del protolito del complejo Altar contradice los modelos que relacionan

el depósito de todo el protolito de la franja El Batamote en una cuenca formada por deslizamiento oblicuo a lo largo de la megacizalla Mojave-Sonora del Jurásico Tardío. En su lugar, la franja se explica mejor como un conjunto de formaciones del Jurásico Medio al Cretácico Superior deformadas y localmente metamorfoseadas debajo de un complejo cabalgante laramídico dirigido hacia el noreste. Edades de potasio-argón y de  $^{40}\text{Ar}/^{39}\text{Ar}$  confirman inferencias previas de que la deformación de la franja El Batamote ocurrió entre el Cretácico Tardío y el Eoceno tardío. Una segunda fase de deformación, que implicó fallas normales de bajo ángulo, ocurrió durante o después de la intrusión de la granodiorita del Rancho Herradura de ~22–21 Ma.

Palabras clave: complejo Altar; complejo Carnero; franja El Batamote; Grupo El Chanate; megacizalla Mojave-Sonora; orogenia laramídica.

INTRODUCTION

The ranges north and east of Caborca and Altar in northern Sonora expose a narrow, northwest-southeast-trending belt of deformed and

locally metamorphosed rocks at least 70 km long, and possibly over 100 km in length, here termed “El Batamote belt” (Figure 1; Damon et al., 1962; Hayama et al., 1984; Jacques-Ayala et al., 1990, 2009; García y Barragán, 1992; Nourse, 1995, 2001; García y Barragán et al., 1998; Jacques-Ayala, 1999; García y Barragán and Jacques-Ayala, 2010). These rocks have also been referred to as “El Batamote Structural Complex” and “Altar Formation”; however, “El Batamote belt” provides better consistency with other terminology that we introduce below. El Batamote belt was derived from a protolith rich in sandstone and conglomerate with minor shale, limestone and volcanic rocks. Distinctive attributes of the belt include southwest-dipping axial-plane cleavage, stretched pebbles in the conglomerates, northeast-verging folds and local greenschist-facies metamorphism. Rocks assigned to El Batamote belt crop out from the Altar area in the southeast to Cerro Basura and Sierra La Gloria in the northwest. Potentially equivalent rocks have been mapped in ranges even farther northwest. The belt is flanked on the southwest by Proterozoic crystalline rocks and the Neoproterozoic–Cambrian miogeoclinal assemblages of the Caborca block (Jacques-Ayala et al., 1990; Anderson and Silver, 2005; Molina-Garza and Iriondo, 2007). Dominant rocks exposed directly




Figure 1. Generalized geologic map of north-central Sonora and south-central Arizona after Haxel et al. (2008) with modifications based on Jacques-Ayala (1999), Terán-Martínez (1999), Nourse (2001), Castro-Escárcega (2002a, 2002b) and García-Cortez (2002). Labeled areas denote regions of sedimentary rocks deformed, and in many cases also metamorphosed, during the Late Cretaceous–early Cenozoic Laramide orogeny, including the Baboquivari Mountains (BM), Cerro Basura (CB), Cerro Carnero (CA), Cerros La Batellera (LB), Sierra El Batamote (EB), Sierra El Chanate (EC), Sierra La Gloria (SG), Sierra La Vibora (SV). Arrows show approximate distances in kilometers (from the tip of the arrow) to additional locations of Laramide tectonism outside the map area: Arizpe (AZ), Imuris (IM), Naco (NA), Quitovac (Q), Quitobaquito Hills (QH).

northeast of the belt include the Upper Jurassic–Lower Cretaceous Bisbee Group and Upper Cretaceous El Chanate Group (Jacques-Ayala *et al.*, 1990; García y Barragán, 1992; Jacques-Ayala, 1993, 1995, 1999; García y Barragán *et al.* 1998; Nourse, 2001) (we use the geologic time scale of Walker *et al.*, 2018). El Batamote belt lies at the southern end of the Papago domain, a region characterized by almost complete absence of exposures of Precambrian basement (Figure 1; Haxel *et al.*, 1980, 1984). Proterozoic rocks, however, are inferred to be present at depth within the Papago domain and to be part of the Mazatzal basement province (Iriondo *et al.*, 2004; Anderson *et al.*, 2005; Haxel *et al.*, 2008).

Most workers consider that deformation and metamorphism of El Batamote belt occurred during northeast-southwest-directed compression associated with the Late Cretaceous to Paleogene Laramide orogeny (Damon *et al.*, 1962; Hayama *et al.*, 1984; Jacques-Ayala *et al.*, 1990, 2009; Jacques-Ayala and DeJong, 1996; García y Barragán *et al.*, 1998; Nourse, 2001; Nourse *et al.*, 2007). The structural style resembles that observed in areas of Laramide tectonism throughout northern Sonora, southern Arizona and southwestern New Mexico (Haxel *et al.*, 1984, 2008; Nourse, 1995; González-León *et al.*, 2000, 2017; Iriondo *et al.*, 2005; Clinkscales and Lawton, 2015). The timing of the event in El Batamote belt is only loosely constrained. At least part of the deformation postdates the 72 Ma El Charro volcanic complex, which is folded within the center of the range-scale El Chanate syncline (Jacques-Ayala 1993, 1999; Nourse, 2001). The event was probably largely over by the late Paleocene or early Eocene, as indicated by a biotite K-Ar cooling age of  $58 \pm 3$  Ma from the Altar complex east of Altar (Damon *et al.*, 1962) and extrusion of the postkinematic San Jacinto volcanics at  $51 \pm 2$  Ma (Jacques-Ayala, 1999, 2000). Laramide contraction was followed in the early Miocene by intrusion of granitoid igneous bodies and exhumation along low-angle normal faults (Nourse, 1995, 2001; Jacques-Ayala, 1999).

In contrast to the broad consensus regarding Laramide and younger history, the protolith age of El Batamote belt has long been a matter of debate. Jacques-Ayala *et al.* (1990, 2009), García y Barragán *et al.* (1998) and Jacques-Ayala (1999, 2000) concluded that a significant fraction of the protolith was Upper Cretaceous; *i.e.* younger than the Bisbee Group and either correlative with or younger than El Chanate Group. In their view, the Upper Cretaceous component of the protolith represents, at least in part, a synorogenic succession deposited in front of, and eventually overridden by, a Laramide thrust complex. The proposed thrust faults, however, have not been definitively identified; they may be buried beneath younger deposits and/or have been excised by middle Cenozoic extensional faults. Alternatively, Nourse (1995; 2001) and Anderson and Nourse (2005) considered that the protolith of El Batamote belt was entirely Upper Jurassic to Lower Cretaceous and correlative with the Glimpse Conglomerate, the lowest formation of the Bisbee Group. Nourse (1995; 2001) and Anderson and Nourse (2005) further argued that the elongate geometry of El Batamote belt was inherited from deposition in a narrow transtensional (pull-apart) basin adjacent to the inferred Late Jurassic Mojave–Sonora megashear. As such, their proposed Late Jurassic–Early Cretaceous protolith age for the complex represents a primary line of evidence for the megashear in the Caborca–Altar area.

The debate regarding the protolith age of El Batamote belt has added significant uncertainty to both local and regional tectonic syntheses. Recent studies, however, shed new light on this controversy and suggest that the belt was assembled from multiple supracrustal units of various ages (García y Barragán and Jacques-Ayala, 2010). For example, detrital zircon U–Pb ages from El Batamote belt east of Altar, described in preliminary form by Barth *et al.* (2008), confirm that part of the protolith is Upper Cretaceous. On the other hand, Mauel (2008)

obtained a latest Middle Jurassic zircon U–Pb age ( $164 \pm 1$  Ma) from a siliceous tuff (?) within the belt northwest of Altar, between Sierra El Batamote and Sierra El Chanate. In addition, Mauel *et al.* (2011) noted strong lithologic similarity between a phyllite near Altar and the Upper Jurassic Cucurpe Formation located ~100 km southeast of Altar. Here, we investigate the origin of El Batamote belt based on detailed examination of the detrital zircon results of Barth *et al.* (2008). This dataset includes samples from two components of El Batamote belt (Altar complex and Carnero complex, defined below). In addition to detrital zircon ages, we present several new  $^{40}\text{Ar}/^{39}\text{Ar}$  ages for the Altar complex, as well as a U–Pb zircon crystallization age for a Miocene granodiorite that intruded the Carnero complex. The  $^{40}\text{Ar}/^{39}\text{Ar}$  ages and granodiorite age help to constrain the Cenozoic history of the area, including the timing of Miocene low-angle normal faulting.

## GEOLOGIC SETTING AND PREVIOUS WORK

We collected samples from Cerro San Judas northeast of Altar; Cerros La Batellera, southeast of Altar and northeast of Highway 2; and Cerro El Amol and Cerro Carnero, southeast of Altar and southwest of Highway 2 (Figure 2). The rocks underlying these areas have commonly been referred to collectively as the Altar Schist, Altar Massif or Altar Formation (Cooper and Arellano, 1946; Damon *et al.*, 1962; Hayama *et al.*, 1984; García y Barragán *et al.*, 1998). However, lithologic differences between the rocks northeast and southwest of Highway 2 have led others to propose that they are separate units (Jacques-Ayala *et al.*, 1990), a conclusion reinforced by our detrital zircon results. Specifically, the rocks of Cerros La Batellera and Cerro San Judas were derived at least partly from an Upper Cretaceous protolith likely correlative with El Chanate Group. In contrast, the protolith of the rocks at Cerro Carnero and Cerro El Amol is likely, although not unequivocally, Jurassic. Based on these differences, we divided the rocks of these two areas into the (informal) Altar complex and Carnero complex, which we tentatively infer to be separated by a buried thrust fault (Figure 2). Both are subunits of El Batamote belt (previously known as “El Batamote structural complex” but here renamed to avoid using “complex” at two different hierarchical levels).

### Altar complex at Cerros La Batellera

The Altar complex at Cerros La Batellera and Cerro San Judas (to be referred to collectively as “Cerros La Batellera”) is part of a ~2–3-km-thick sequence derived from sandstone, conglomerate and rare limestone (García y Barragán, 1992; Jacques-Ayala *et al.*, 1990; García y Barragán *et al.*, 1998). The rocks exhibit southwest-dipping foliation (generally bedding-parallel) and southwest-plunging lineation. Clasts in the conglomerate are stretched to varying degrees dependent on strength relative to the matrix (Figure 3a). Metamorphic grade decreases from greenschist facies in the west to incipiently recrystallized in the east. Intensity of deformation also decreases from west to east. The boundary between the more- and less-highly-deformed and metamorphosed parts of the complex is gradational, but, for simplicity, is depicted as a discrete contact in Figure 2.

East of the map area, the lesser-deformed/metamorphosed portion of the Altar complex is in contact with rocks interpreted as El Chanate Group, although the stratigraphy does not match in detail with that of the type El Chanate Group of the Sierra El Chanate (García y Barragán *et al.*, 1998). This contact has been interpreted as either depositional or a thrust fault (Jacques-Ayala *et al.*, 1990; García y Barragán *et al.*, 1998; Nourse, 2001). The fault interpretation appears most plausible considering the evidence presented here that the Altar complex is a metamorphosed and deformed equivalent of El Chanate Group (see




Figure 2. Geologic map of the study area (modified after García y Barragán, 1992). Numbered circles, squares and diamonds on the map show locations of K-Ar samples from Damon et al. (1962) and Hayama et al. (1984) and <sup>40</sup>Ar/<sup>39</sup>Ar and detrital zircon samples from this study. The legend shows the ages and errors (2σ) for the K-Ar and <sup>40</sup>Ar/<sup>39</sup>Ar ages and sample numbers for the zircon samples. Note that some previous publications have applied the name “Cerros El Amol” to hills northeast of Highway 2 and east of Cerros La Batellera. In contrast, we follow INEGI (2001) in applying the name “Cerro El Amol” to the small set of hills 3–5 km southeast of Altar.

also Barth *et al.*, 2008; Jacques-Ayala *et al.*, 2009). Even farther east, El Chanate Group is in depositional contact with the Bisbee Group. Rocks of the Bisbee Group are also exposed to the north of Cerros La Batellera in the upper plate of a low-angle normal fault of inferred Miocene age.

### Carnero complex at Cerro Carnero

The Carnero complex of Cerro Carnero and Cerro El Amol (to be referred to collectively as “Cerro Carnero”) was derived from sandstone and minor conglomerate, mudstone and limestone. The rocks exhibit a well-developed east- to northeast-dipping foliation and east-trending lineation (Figures 2 and 3b) (Damon *et al.*, 1962; Hayama *et al.*, 1984; Jacques-Ayala *et al.*, 1990; García y Barragán *et al.*, 1998). A thin band of black phyllite is present at the north end of the Carnero complex. Mauel *et al.* (2011) suggested that the phyllite was derived from marine deposits correlative with the Upper Jurassic Cucurpe Formation. A Jurassic age for the protolith of the Carnero complex is consistent with the detrital zircon results described below.

The Carnero complex is intruded by the Rancho Herradura granodiorite (Figure 3c) and the Carnero leucogranite. The granodiorite is foliated and exhibits a gently west-plunging lineation parallel to lineation in the metasedimentary rocks. Asymmetric (“S-C”) fabrics record top-west extension.

### Thermal history

The cooling and exhumation history of the area is constrained by a limited number of K-Ar ages. Damon *et al.* (1962) reported a biotite K-Ar age of  $58 \pm 3$  Ma from the Altar complex of Cerros La Batellera 2 km southeast of Altar, whereas Hayama *et al.* (1984) obtained a significantly younger biotite K-Ar age of  $16.9 \pm 0.6$  Ma nearby (Figure 2). For Cerro Carnero, Hayama *et al.* (1984) reported K-Ar ages for the metasedimentary rocks of  $55 \pm 3$  Ma for hornblende and  $16.2 \pm 0.5$  Ma and  $16.1 \pm 0.5$  Ma for muscovite. These workers also obtained biotite K-Ar ages of  $14.8 \pm 0.5$  Ma and  $16.6 \pm 0.4$  Ma for the Rancho Herradura granodiorite and a K-Ar age for fine-grained white mica (sericite) of  $16 \pm 0.4$  Ma for the Carnero leucogranite. We consider that the Paleogene K-Ar ages reflect cooling following Laramide underthrusting of the protoliths. Thermal resetting during Laramide magmatism is also possible, although we deem this option less likely, as no Laramide plutons are exposed near the dated localities. The ~17–15 Ma ages presumably reflect a second phase of cooling associated with detachment faulting and/or conductive cooling following intrusion of the Rancho Herradura granodiorite and Carnero leucogranite.

## ANALYTICAL METHODS

### Zircon U-Pb analyses

We analyzed eight samples for detrital zircon, seven from the Altar complex and one from the Carnero complex (Figure 2; see Table A1 of the Electronic Supplement for coordinates of all sample localities). The Altar samples include five of metasandstone (662, 664, 666, 669 and CJ13-15), the (metamorphosed) sand matrix of a metaconglomerate (663) and a quartzite clast from the metaconglomerate (663A). The Carnero sample is a metasandstone (668).

All detrital samples except CJ13-15 were collected and processed during an early phase of the study. We crushed approximately 1 kg of each sample using a jaw crusher and disc mill. For the metaconglomerate clast, only fragments from the interior were used to avoid matrix material adhering to the outer surface. The crushed material was sieved and ~50 ml volumetrically of the fraction finer than 150  $\mu\text{m}$  was processed in heavy liquids (tetrabromoethane followed by methylene iodide) after removing iron filings with a magnet. The final sink was



Figure 3. (a) Metaconglomerate within Altar complex at Cerros La Batellera. Site of sample 663. (b) Carnero complex at Cerro Carnero. Site of sample 668. (c) Granodiorite of Ranch Herradura at Cerro Carnero. Site of sample 667.

further cleaned with a Frantz® magnetic separator. We obtained a few hundred or more zircon grains for all samples except sample 669, which yielded relatively few grains, most of which were quite small. As a result, we were able to obtain only five ages from this sample.

Polished epoxy mounts of the above samples were analyzed at the Arizona Laserchron Center, University of Arizona, using a 193-nm excimer laser and a GVI Isoprobe multicollector-inductively coupled

plasma-mass spectrometer (MC-ICP-MS) following the procedures of Gehrels *et al.* (2008). The ions  $^{206}\text{Pb}$ ,  $^{207}\text{Pb}$ ,  $^{208}\text{Pb}$ ,  $^{232}\text{Th}$  and  $^{238}\text{U}$  were measured with Faraday collectors. A channeltron multiplier was used for  $^{204}\text{Pb}$ . Sri Lanka zircon (SL2;  $563.5 \text{ Ma} \pm 3.2 \text{ Ma } 2\sigma$ ) was used as the primary standard (Gehrels *et al.*, 2008). Common lead was corrected using the  $^{204}\text{Pb}$  method. For unknowns, we targeted grain interiors to minimize the effects of lead loss during metamorphism. We avoided visible inclusions and fractures, but otherwise selected grains randomly, without using cathodoluminescence (CL) imagery. Analytical results are presented in Table A2 of the Electronic Supplement.

We use the  $^{206}\text{Pb}/^{238}\text{U}$  age for grains younger than 800 Ma and the  $^{207}\text{Pb}/^{206}\text{Pb}$  age for grains with  $^{206}\text{Pb}/^{238}\text{U}$  ages  $>800$  Ma. Discordance is relatively minor among the analyses with  $^{206}\text{Pb}/^{238}\text{U}$  ages  $>800$  Ma. For this age group, maximum normal discordance is 25%, with a median of 3%; the maximum reverse discordance is 3%, with a median of 1%. None of these ages were excluded. For ages  $<800$  Ma, particularly Mesozoic ages, evaluation of discordance can be problematic owing to potential large errors associated with  $^{207}\text{Pb}/^{235}\text{U}$  ages obtained by ICP-MS analysis. For these analyses, we evaluated discordance individually for each result in the context of its analytical errors.

Because of the low yield from sample 669, we later collected and processed a second sample (CJ13-15) from the same location (Cerro San Judas, near the shrine to Saint Jude; Figure 2). The sample was crushed in the laboratory and heavy minerals were separated using a Wilfley table and a Frantz<sup>®</sup> magnetic separator. Zircon grains were extracted from the residual non-magnetic fraction, randomly selected under a binocular microscope, mounted in epoxy resin and polished. Intra-grain compositional zoning was identified by CL. Twenty-two zircons were analyzed using a Resonetics Resolution M050 excimer laser workstation coupled to a Thermo ICapQc ICP-MS at the Laboratorio de Estudios Isotópicos (LEI), Centro de Geociencias, Universidad Nacional Autónoma de México, following the methodology reported in Solari *et al.* (2010). Analytical results are presented in Table A3 of the Electronic Supplement.

Zircon from the Rancho Herradura granodiorite from Cerro Carnero was extracted during processing of the initial set of detrital samples. Grains were mounted in epoxy and polished prior to examination using CL. The CL images were used to guide selection of analysis points in zoned zircon grains. Uranium and Pb concentrations and isotopic ratios were measured on the SHRIMP-RG ion microprobe at Stanford University using analytical and data analysis procedures described in Barth and Wooden (2006). Isotopic ratios were standardized against Braintree Complex zircon R33 (419 Ma; Black *et al.*, 2004). Analytical results are presented in Table A4 of the Electronic Supplement.

#### $^{40}\text{Ar}/^{39}\text{Ar}$ analyses

Material generated during crushing of sample CJ13-15 for zircon was used to obtain separates of muscovite, biotite and plagioclase for  $^{40}\text{Ar}/^{39}\text{Ar}$  analysis. The separates were irradiated at the University of McMaster reactor. Analyses were performed at Queen's University, Canada. Step heating was conducted using a Merchantek MIR 10-30  $\text{CO}_2$  laser. A faceted lens was used to diffuse the beam over 3 mm pits. Ions were measured using an MAP-216 mass spectrometer. The J values and errors for samples were determined by polynomial fit to replicate analyses of a flux monitor (biotite GA1550, 98.5 Ma) at measured positions along the length of the irradiation capsule. Analytical results are presented in Tables A5-A7 of the Electronic Supplement.

We report sample ages as total gas ages. In our view, these provide the best estimate for the time of bulk closure of argon diffusion (Jacobson *et al.*, 2002).

## RESULTS

### Detrital zircon ages

#### *Cerros La Batellera*

Probability density plots and cumulative probability plots for the individual samples of the Altar complex from Cerros La Batellera are shown in Figure 4. A plot of the combined age results for the metasandstone and metaconglomerate matrix samples is presented in Figure 5a.

Detrital zircon ages for the metasandstone and metaconglomerate matrix samples from Cerros La Batellera range from Archean (2708 Ma) to latest Cretaceous (72 Ma) (Figures 4a–4f and 5a). Five of the six samples include zircon of latest Early Cretaceous and/or Late Cretaceous age. Two samples include zircon as young as Campanian, five grains with ages of 81–72 Ma in sample 662 and one grain with an age of 77 Ma in sample 663. A weighted mean calculation using Isoplot (Ludwig, 2003) suggests that the six latest Cretaceous grains could all be part of a single population with an average age of  $77.5 \pm 2.5$  Ma (Figure 6). Alternatively, averaging only the three youngest grains (*e.g.*, Dickinson and Gehrels, 2009a) yields a statistically-equivalent weighted mean age of  $76.5 \pm 4.4$  Ma. We take the average of the six grains as the more conservative estimate of the maximum depositional age for the Cerros La Batellera samples.

Grains younger than 285 Ma comprise 72% of the total for the metasandstone and conglomerate matrix samples and are inferred to be derived from Cordilleran magmatic arcs (*cf.* Dickinson and Gehrels, 2009b; González-León *et al.*, 2009, 2017; Mauel *et al.*, 2011; Peryam *et al.*, 2012; Lawton and Molina Garza, 2014):

(1) The latest Cretaceous zircon could have been derived from multiple sources in Sonora, including ranges near Cerros La Batellera (McDowell *et al.*, 2001; Iriondo *et al.*, 2005; Jacques-Ayala *et al.*, 2009; Roldán-Quintana *et al.*, 2009; González-León *et al.*, 2011, 2017; Ortega-Gutiérrez *et al.*, 2014).

(2) Grains of late Early and early Late Cretaceous age ( $\sim 120$ – $90$  Ma) comprise  $\sim 10\%$  of the total zircon population. Igneous rocks of this age are not abundant in Sonora, although the oldest zircons in this group are compatible with a derivation from volcanic rocks within the Aptian–Albian Arroyo Sásabe Formation (Jacques-Ayala *et al.*, 1990; Jacques-Ayala, 1995). Alternatively, late Early and early Late Cretaceous igneous rocks are widespread within the Alisitos and Peninsular Ranges arcs of Baja California, which would have been adjacent to western Sonora prior to opening of the Gulf of California (Wetmore *et al.*, 2003; Busby *et al.*, 2006; Jacques-Ayala *et al.*, 2009; Peryam *et al.*, 2012; Kimbrough *et al.*, 2015). Peryam *et al.* (2012) presented evidence for a fluvial connection between the Late Jurassic–Early Cretaceous Guerrero and Alisitos arcs of Baja California and the Cucurpe area of Sonora during deposition of the Bisbee Group. This drainage system could have remained active into the Late Cretaceous.

(3) Jurassic grains are the most abundant of the arc-derived grains, consistent with the widespread presence of Jurassic igneous rocks in northwest and northcentral Sonora (Figure 1; Nourse, 2001; Anderson *et al.*, 2005; Leggett, 2009).

(4) Four of the six samples include minor Permian–Triassic peaks. Permian–Triassic plutons have been described from northwest Sonora and may be more common than previously recognized (Arvizu *et al.*, 2009; González-León *et al.*, 2009; Arvizu and Iriondo, 2015; Riggs *et al.*, 2016).

Most pre-Permian (“pre-arc”) zircon ages in the above samples fall within three age peaks centered near 1.7, 1.4 and 1.1 Ga, the first being the most prominent (Figures 4a–4f and 5a). Igneous and metamorphic complexes with ages of  $\sim 1.7$  and  $\sim 1.4$  Ga are widespread in the southwestern United States (U.S.) and Sonora (Barth *et al.*,




Figure 4. Relative and cumulative probability plots of detrital zircon ages from the Altar complex at Cerros La Batellera and the Carnero complex at Cerro Carnero. All samples are metasandstone, except sample 663, which is from the sand matrix of a metaconglomerate, and sample 663A, which is a single quartzite clast from that conglomerate. Note age-axis scale break at 300 Ma. For relative probability plots, vertical scales differ to left and right of the age-axis scale break such that equal area represents equal probability throughout the graph. Line color in the cumulative probability plot corresponds to the fill color in relative probability plots. Location of samples shown in Figure 2. n: number of analyses.




Figure 5. Composite detrital zircon age distributions for metasandstone and conglomerate matrix from the Altar complex at Cerros La Batellera (samples 662, 663, 664, 666, 669, CJ13-15) and sandstone from El Chanate Group of Sierra El Chanate (Jacques-Ayala *et al.*, 2009). Scale break and color-coding of relative versus cumulative probability plots as in Figure 4. N: number of samples, n: number of analyses.

2000, 2001a, 2001b; Anderson and Silver, 2005; Farmer *et al.*, 2005; Iriando *et al.*, 2004; Nourse *et al.*, 2005). Those with ages of  $\sim 1.7$  Ga dominate over  $\sim 1.4$  Ga units, consistent with the relative sizes of these two peaks in the Altar samples. In contrast, basement rocks with ages of  $\sim 1.1$  Ga are not common in the southwestern U.S. and Sonora (Iriando *et al.*, 2004; Anderson and Silver, 2005). On the other hand,  $\sim 1.1$  Ga detrital zircon is a significant component of Neoproterozoic and Cambrian cratonal and miogeoclinal quartz arenites from this part of the Cordillera (Gehrels *et al.*, 1995, 2011; Gehrels and Stewart, 1998; Stewart *et al.*, 2001; Gehrels and Pecha, 2014). Hence, we suspect that much, if not most, of the  $\sim 1.1$  Ga zircon in the Altar complex of Cerros La Batellera is recycled from local miogeoclinal sequences (Figure 1). Accordingly, some of the  $\sim 1.7$  and  $\sim 1.4$  Ga zircon in La Batellera samples may also be recycled. However, the ratio of  $\sim 1.7$  Ga to  $\sim 1.1$  Ga zircon in our samples exceeds that typically found in Neoproterozoic–Cambrian sedimentary rocks. This suggests that  $\sim 1.7$  Ga bedrock contributed directly to the protolith of the Altar complex. Potential source areas are present in the Caborca block immediately south of the map area of Figure 1.

Some uncertainty in the interpretation of the Cerros La Batellera section is introduced by the results from sample 669, collected from Cerro San Judas near the shrine to St. Jude. This sample differs from all the others from Cerros La Batellera in that it yielded only Jurassic ages. The dataset, however, is small—only five analyses. As noted in the analytical methods section, zircons obtained from sample 669 were smaller on average than those from the other samples. Only the larger grains could be analyzed, which may have biased the results. In addition, Cretaceous zircon is not abundant in any of the samples of Altar complex from Cerros La Batellera (Figures 4 and 5a) and could have been missed in sample 669 owing to the small number of analyses. Furthermore, Cretaceous zircon is present in sample CJ13-15 from the same locality (Figure 4f). Considering these factors, we conclude that sample 669 likely has the same Late Cretaceous depositional age as the other Cerros La Batellera samples, despite the failure to yield any zircon younger than Jurassic.

In contrast to the dominant arc sources for the sand component of the Altar complex, the 70 ages from the quartzite clast (sample 663A) are all Precambrian—68 Proterozoic (minimum age of  $1060 \pm 37$  Ma) and two Archean (Figure 4g). The Proterozoic ages define the same three age peaks ( $\sim 1.7$ ,  $\sim 1.4$  and  $\sim 1.1$  Ga) as in the sands, although

in more nearly equal proportions than in the latter (Figure 4g *versus* Figure 5a). The age distribution for sample 663A is typical of that for the Neoproterozoic and Cambrian sedimentary sequences from Sonora and adjacent areas (Gehrels and Stewart, 1998; Stewart *et al.*, 2001; Gehrels and Pecha, 2014). Miogeoclinal rocks of this age crop out in ranges west and south of Caborca and comprise a potential source for the quartzite clasts in the Altar complex (Figure 1). The original clasts may have been quartz arenite that was converted to quartzite during Laramide deformation and metamorphism.

Lithologic and map relations led Jacques-Ayala *et al.* (1990), García y Barragán *et al.* (1998) and Jacques-Ayala (1999) to conclude that the protolith of the Altar complex at Cerros La Batellera was either correlative with El Chanate Group or deposited above it. Comparison of composite age distributions for the Altar complex at Cerros La Batellera (this study) and El Chanate Group of the Sierra El Chanate (Jacques-Ayala *et al.*, 2009) favors the former (Figure 5). This conclusion is somewhat speculative, because application of the Kolmogorov-Smirnoff (K-S) test to the two groups yields a P value of 0.005, which implies that the two groups were not derived from the same source




Figure 6. Weighted mean plot of six youngest zircon ages from the Altar complex of Cerros La Batellera. Diagram constructed using Isoplot Version 3.75 (Ludwig, 2003).

(*cf.* Dickinson and Gehrels, 2009b). Nonetheless, we note multiple similarities between the two datasets: (1) Dominance of Jurassic zircon in each, albeit skewed to older ages in El Chanate Group. (2) Similar proportions of  $\sim 1.7$ ,  $\sim 1.4$  and  $\sim 1.1$  Ga populations. (3) Comparable small numbers of Permian–Triassic zircons. (4) Similar distribution of Cretaceous ages, including similar youngest ages ( $\sim 75$  Ma in both groups). These relations lead us to conclude that the protoliths of the Altar complex of Cerros La Batellera and El Chanate Group are correlatives, despite the negative K-S result. We suggest that the low P value derives from the small sample sizes, minor local variation in source area and/or systematic analytical differences between the Altar complex ages, which were obtained by ICP-MS methods, and El Chanate results, which were determined by ion microprobe. Further work is needed to resolve this issue.

Based on the high abundance of quartzite clasts, Jacques-Ayala (1999) and Jacques-Ayala *et al.* (2009) proposed that the Altar complex at Cerros La Batellera correlates with the Pozo Duro Formation, the oldest unit of El Chanate Group. Alternatively, the  $77.5 \pm 2.5$  Ma maximum depositional age of the Cerros La Batellera samples may indicate a correlation with the Escalante Formation, the youngest part of El Chanate Group. The Escalante Formation has a maximum depositional age of  $75 \pm 2$  Ma based on two zircon ages derived from the upper part of the formation (Jacques-Ayala *et al.* (2009). Correlation of the Altar complex of Cerros La Batellera with the Escalante Formation has also been proposed by González-León *et al.* (2017). Nonetheless, the maximum depositional ages of both the Altar and El Chanate

samples are not well constrained, as most of the detrital zircon ages are pre-Late Cretaceous (Figure 5). In other words, even those samples that have so far failed to yield zircon younger than late Early or early Late Cretaceous could, in fact, have Campanian depositional ages. This ambiguity reinforces the need for additional analyses.

#### Cerro Carnero

We determined detrital zircon ages for one sample of metasediment from the Carnero complex (sample 668; Figure 4h). In contrast to the samples from the Altar complex, the Carnero sample includes little arc-derived zircon. Another distinctive aspect of the Carnero sample is the high fraction of Grenville-age zircon (1310–912 age group of Dickinson *et al.*, 2012) relative to the  $\sim 1.7$  and  $\sim 1.4$  Ga peaks. In addition, the Carnero sample includes abundant Neoproterozoic ( $\sim 600$  Ma) and Paleozoic ( $\sim 400$  Ma) zircon, age groups not observed in the samples of Altar complex.

The strong Grenville, Neoproterozoic and Paleozoic components of the Carnero sample suggest affinities with the Jurassic eolian (erg) sandstones of the Colorado Plateau, U.S. (Figure 7b; Dickinson and Gehrels, 2009b). Erg-type sands have been documented in Sonora; *e.g.* in the Middle Jurassic Rancho San Martín Formation near Cucurpe (Leggett, 2009; Mauel *et al.*, 2011). The Rancho San Martín Formation and our sample of Carnero complex exhibit similar detrital zircon age spectra, including similar maximum depositional ages ( $\sim 180$ – $170$  Ma) (Figures 7a and 7c; Leggett, 2009; Mauel *et al.*, 2011). In detail, the Carnero complex and Rancho San Martín Formation display some con-




Figure 7. Comparison of detrital zircon age results from the Carnero complex of Cerro Carnero (sample 668 of Figure 4) with selected Jurassic and Cretaceous formations in Sonora and the southwestern U.S. exhibiting an erg-type detrital zircon signature (*cf.* Dickinson and Gehrels, 2009b). Data sources: (a) this study; (b) Dickinson and Gehrels (2009b); (c) Leggett (2009); (d) Mauel *et al.* (2011); (e) Clinkscales and Lawton (2015). Scale break and color-coding of relative versus cumulative probability plots as in Figure 4. N: number of samples, n: number of zircon ages.

trasts, such as fewer Neoproterozoic grains in the latter. However, variability of this scale is typical within the Jurassic eolianites (Dickinson and Gehrels, 2009b).

The zircon age results for the Carnero sample are consistent with a Middle Jurassic protolith age, but do not rule out a younger age. For example, the basal part of the Upper Jurassic Cucurpe Formation of the Cucurpe area exhibits a strong erg signature, attributed to reworking of eolianite layers in the underlying Rancho San Martín Formation (Figure 7d; Mauel et al., 2011). The basal Cucurpe Formation also includes Late Jurassic zircon, confirming its young age relative to the Rancho San Martín Formation. However, the Late Jurassic zircon is thought to be locally derived (Mauel et al., 2011), and need not be present in all parts of the Cucurpe Formation derived by recycling of eolianites sands.

A strong erg signature has also been observed in the Cintura Formation of the Bisbee Group in southern Arizona, U.S. (Dickinson et al., 2009) and in the Mojado Formation, a correlative of the Cintura Formation in southwestern New Mexico, U.S., (Clinkscapes and Lawton, 2015). The Mojado example is particularly instructive. Clinkscapes and Lawton (2015) obtained 92 zircon ages for one sample of the Mojado Formation. The pre-Permian grains exhibit an age distribution remarkably like that of the Colorado Plateau Jurassic ergs (Figures 7b and 7e). The Mojado sample also includes a modest number of Triassic and Jurassic arc-derived grains. However, none of the ages are younger than early Late Jurassic, despite the well-constrained Albian depositional age.

Erg-style detrital zircon age patterns have also been found locally in Upper Cretaceous strata in Sonora (Jacques-Ayala and García y Barragán, 2018). Nonetheless, most Upper Cretaceous sandstones from Sonora that have been analyzed for detrital zircon include only minor numbers of erg-derived grains (Jacques-Ayala et al., 2009; González-León et al., 2017; this study).

In summary, correlation of the Carnero sample with the Middle Jurassic Rancho San Martín Formation is the most straightforward interpretation, although correlation with the Cucurpe Formation, Bisbee Group or even El Chanate Group is also permissible, with the last option least likely.

#### Rancho Herradura Granodiorite

Zircon grains extracted from the Rancho Herradura granodiorite (Sample 667) are euhedral to subhedral prismatic and up to 200 microns long. About half the grains contain recognizable cores. We analyzed nine grains. Seven yielded Miocene ages, which we interpret as synmagmatic. Two older ages (56 and 946 Ma) are considered premagmatic. The  $^{206}\text{Pb}/^{238}\text{U}$  ages for the synmagmatic zircons range from 23 to 19 Ma and form an array along concordia on a Tera-Wasserburg diagram (Figure 8). We interpret this age range to indicate varying degrees of lead loss, perhaps associated with extensional deformation of the granodiorite. The four oldest grains, which presumably underwent the least lead loss, yield a mean age of  $21 \pm 1$  Ma (95% confidence level). However, even these grains may have lost some Pb, so the true age of the granodiorite could be closer to the upper end of the range of analyses; *i.e.*, ~22 Ma. On the other hand, Nourse et al. (2007) inferred a zircon U-Pb age of  $21.2 \pm 0.2$  (95% confidence level) for this same intrusion, also using the SHRIMP-RG ion microprobe at Stanford University. Considering the two datasets together, we conclude that the age of the granodiorite is ~22–21 Ma.

#### $^{40}\text{Ar}/^{39}\text{Ar}$ Results

The  $^{40}\text{Ar}/^{39}\text{Ar}$  results vary in degree of interpretability (Figure 9). The muscovite spectrum for sample CJ13-15 is the least complicated, with a profile indicative of Ar loss in the low-temperature steps leading to a moderately well-developed plateau for the high-temperature steps. The total gas age is  $45.1 \pm 0.5$  Ma.

The biotite age spectrum is less regular. Nonetheless, even the oldest steps are younger than those on the plateau of the muscovite spectrum, consistent with the lower inferred closure temperature for biotite than muscovite ( $\sim 350 \pm 50$  °C and  $\sim 400 \pm 50$  °C, respectively; McDougall and Harrison, 1999). The total gas age is  $36.0 \pm 0.8$  Ma.

The plagioclase sample exhibits a U-shaped spectrum indicative of excess radiogenic argon (McDougall and Harrison, 1999). The youngest steps yield ages  $>50$  Ma; *i.e.* older than the oldest steps for muscovite and biotite. This confirms that all steps have been impacted by excess radiogenic argon, as the closure temperature for plagioclase (and thus age) is expected to be equal to or less than those for muscovite and biotite (McDougall and Harrison, 1999). We do not believe that a geologically meaningful age can be determined for this sample and do not consider it further.

The muscovite and biotite  $^{40}\text{Ar}/^{39}\text{Ar}$  ages fall in a gap between clusters at 58–55 Ma and 17–15 Ma defined by the K-Ar analyses of Damon et al. (1962) and Hayama et al. (1984) (Figure 2). It is difficult to compare  $^{40}\text{Ar}/^{39}\text{Ar}$  and K-Ar ages directly. Nonetheless, our results are consistent with the evidence from the K-Ar ages for an extended period of cooling from early to mid-Cenozoic time.

## DISCUSSION

### Protolith age of El Batamote belt and tectonic implications

Our detrital zircon analyses confirm the view that much, or all, of the Altar complex at Cerros La Batellera was derived from an Upper Cretaceous protolith likely correlative with El Chanate Group (*e.g.*, Jacques-Ayala et al., 1990, 2009; García y Barragán et al., 1998; Jacques-Ayala, 1999, 2000) and contradict the interpretation that the entire protolith of El Batamote belt is correlative with the Glance Conglomerate (*e.g.*, Nourse, 2001). Some parts of La Batellera section are no older than  $77.5 \pm 2.5$  Ma, which may indicate correlation with the Escalante Formation, the uppermost unit of El Chanate Group (González-León et al., 2017).

Some fraction of El Batamote belt, however, must be derived from units older than Late Cretaceous. The most direct line of evidence is provided by the latest Middle Jurassic (164 Ma) tuff (?) located northwest of Altar (Mauel, 2008; Mauel et al., 2011). An age as old as Middle Jurassic is also suggested by the erg-style detrital zircon




Figure 8. Tera-Wasserburg concordia plot of zircon ages from the Rancho Herradura granodiorite (sample 667).

age spectrum of the sample from the Carnero complex. Alternatively, the Carnero sample could be younger than Middle Jurassic (above), although the most-plausible options nonetheless require that the protolith be pre-Late Cretaceous and thus older than the Altar complex. Similarly, the distinctive lithologic composition of the black phyllite at the north end of the Carnero complex body is consistent with, but not proof of, correlation with the Middle Jurassic Cucurpe Formation (Mauel *et al.*, 2011).

It is apparent from various issues discussed herein that further work is needed to more fully characterize the protolith of El Batamote belt, including additional mapping, petrography, whole-rock geochemical analysis and detrital zircon studies throughout its geographic extent. Nonetheless, based on the available data, we hypothesize that El Batamote belt was derived from a range of Middle Jurassic to Upper Cretaceous supracrustal sequences that were present in the Caborca–Altar area at the time of the Laramide orogeny (see also García y Barragán and Jacques-Ayala, 2010). We envision that the various units were deformed and metamorphosed together beneath a now-hidden Laramide thrust sheet (or sheets). The youngest strata involved in this thrusting (Escalante Formation and parts of the Altar complex) may have been synorogenic deposits shed from the thrust belt as it advanced northeastward toward the Caborca–Altar area (Jacques-Ayala *et al.*, 1990, 2009; García y Barragán *et al.*, 1998; Jacques-Ayala, 1999, 2000). This style of tectonism is analogous to that proposed for much of the Cretaceous–Paleogene Mexican orogen (Fitz-Díaz *et al.*, 2018). Application to the Caborca–Altar area, however, is more speculative, because of the lack of definitive evidence for major thrust faults, either bounding or within El Batamote belt. The inferred thrust fault(s) could be (1) buried beneath basin fill southwest of the exposures of El Batamote belt (Jacques-Ayala and García y Barragán, 2015, 2018); (2) present within the complex, but difficult to recognize because of the lack of good stratigraphic control; and/or (3) overprinted by Miocene detachment faults (Nourse, 2001). Late Cretaceous thrust faults have been recognized within the Sierra La Vibora of the Caborca block southwest of Altar (De Jong *et al.*, 1988; Jacques-Ayala *et al.*, 1990) and could be part of the proposed thrust system.

According to Nourse (1995; 2001) and Anderson and Nourse (2005), the protolith of El Batamote belt was deposited in a northwest-southeast–elongated transtensional basin along the Late Jurassic Mojave–Sonora megashear. Our results, however, indicate that much of the protolith is too young to be related to the megashear. Instead, we propose that the northwest-southeast trend of the belt is the consequence of Late Cretaceous northeast-southwest shortening followed by Miocene extension along approximately the same azimuth. We cannot rule out that some components of the protolith of El Batamote belt were deposited in northwest-southeast–trending basins (*e.g.* The Altar-Cucurpe basin of Mauel *et al.*, 2011). Nonetheless, we consider it unlikely that the present linear outcrop geometry of El Batamote belt is a fundamental signature of Late Jurassic transtension. This conclusion does not by itself disprove the megashear concept, but it greatly diminishes the importance of El Batamote belt as an argument in favor of the megashear.

#### Timing and regional correlations

The contractional event that caused deformation and metamorphism of El Batamote belt and folding of the Bisbee and El Chanate Groups in the Caborca–Altar area must be at least partly younger than the ~78–75 Ma maximum depositional ages of the Altar complex and El Chanate Group, as well as the 72 ± 1 Ma El Charro volcanic complex, which was folded along with El Chanate Group (Jacques-Ayala, 1999). The end of Laramide deformation in this region is constrained by eruption of the post-tectonic San Jacinto volcanics rocks at 51 ± 2 Ma




Figure 9.  $^{40}\text{Ar}/^{39}\text{Ar}$  release spectra for Altar complex sample CJ13-15.

(Jacques-Ayala, 1999, 2000). In addition, deformation was likely over, or at least waning, by the early Eocene based on the biotite K-Ar age of 58 ± 3 Ma (Damon *et al.*, 1962) for Altar complex at Cerros La Batellera. However, our  $^{40}\text{Ar}/^{39}\text{Ar}$  ages of 45.1 ± 0.5 and 36.0 ± 0.8 Ma for muscovite and biotite, respectively, indicate that at least some parts of the section remained at temperatures of 400–350 °C through most of the Eocene.

Deformation similar in style and age to that of the Caborca–Altar area is widespread through northern Sonora and adjoining areas; *e.g.*, near Quitovac, Imuris, Arizpe and Naco in Sonora (Iriondo *et al.*, 2005; González-León *et al.*, 2017) and in the Quitobaquito Hills to Baboquivari Mountains of south-central Arizona (Haxel *et al.*, 1984; Goodwin and Haxel, 1990; Calmus and Sosson, 1995). Common attributes of these areas include: (1) deformation involved the full thickness of Mesozoic sedimentary and volcanic rocks; (2) Upper Cretaceous strata (Cabullona Group, Cocóspera Formation, Fort Crittenden

Formation) were deposited in basins bounded by Laramide thrust and high-angle reverse faults; (3) greenschist facies metamorphism, southwest-dipping cleavage and stretched-pebble conglomerates are locally well developed; (4) magmatism generally outlasted deformation; and (5) both magmatism and deformation appear to have migrated northeastward with time. Among these areas, González-León *et al.* (2017) highlighted similarities between the stratigraphic, structural and age relations for the Cocospera basin near Imuris and Arizpe with those for the Altar complex and El Chanate Group. The Cocospera Formation was thrust beneath the Bisbee Group along the Late Cretaceous San Antonio fault. González-León *et al.* (2017) interpreted the San Antonio fault as a regional Laramide tectonic front that extended westward to the Caborca–Altar area. The San Antonio fault would thus be equivalent to the Altar thrust of Figure 1 (the “Altar sole fault” of Jacques-Ayala and García y Barragán, 2015).

The orogenic system represented by El Batamote belt may have an analog in the McCoy Mountains Formation of southwesternmost Arizona and southeasternmost California, 400 km northwest of the Caborca–Altar area (Harding and Coney, 1985; Barth *et al.*, 2004; Spencer *et al.*, 2011). The McCoy Mountains Formation includes sandstone, siltstone and conglomerate that, like the inferred protoliths of El Batamote belt, range from Middle or Upper Jurassic to Upper Cretaceous. The lower to middle parts of the McCoy Mountains Formation have been linked to the Bisbee Group, including the Upper Jurassic–Lower Cretaceous Glance Conglomerate (Dickinson and Lawton, 2001). The upper part of the McCoy Mountains Formation, in turn, resembles the upper part of the Altar complex, both in terms of its Late Cretaceous depositional age and the abundance of conglomerate. All parts of the McCoy Mountains Formation were affected by Late Cretaceous deformation. These similarities further highlight the extent and continuity of the Laramide orogen in the southwestern U.S. and northwestern Mexico.

### Miocene extension

Laramide deformation probably accounts for the main structures and fabrics in El Batamote belt and the Bisbee and El Chanate Groups of the Caborca–Altar area (see also Nourse, 2001). Nonetheless, top-west extensional fabrics (Jacques-Ayala *et al.*, 1990) in the ~22–21 Ma Rancho Herradura granodiorite at Cerro Carnero provide evidence of an important overprint during Miocene detachment faulting. Biotite and white mica K–Ar ages clustered at 17–15 Ma may indicate that the main phase of extension postdated intrusion of the granodiorite by several million years. In any case, middle Cenozoic extension in the Caborca–Altar is likely part of a much broader metamorphic core-complex event (Nourse, *et al.*, 1994; Wong *et al.*, 2010). The difficulty in recognizing Laramide thrust faults is probably due in large measure to overprinting by middle Cenozoic extensional structures.

### CONCLUSIONS

Detrital zircon from the Altar complex of Cerros La Batellera indicates a Late Cretaceous maximum depositional age with at least part of the protolith no older than  $77.5 \pm 2.5$  (2 $\sigma$ ). The zircon age spectra are consistent with local to somewhat distant Sonoran sources including Proterozoic bedrock, Neoproterozoic to Cambrian miogeoclinal quartz arenites and Permian–Triassic, Jurassic and Cretaceous arcs. The zircon age results and lithology support correlation with the Upper Cretaceous El Chanate Group.

One sample of the Carnero complex from the Cerro Carnero yielded a Middle Jurassic maximum depositional age and an overall age sig-

nature characteristic of the Jurassic eolian (erg) sands of the Colorado Plateau, U.S. This result may indicate a Jurassic depositional age and correlation with the Rancho San Martín Formation. Alternatively, the Cerro Carnero sample could be an equivalent of the Upper Jurassic Cucurpe Formation, Upper Jurassic to Lower Cretaceous Bisbee Group, or even Upper Cretaceous El Chanate Group produced by reworking of Middle Jurassic erg sands.

We infer that the various stratigraphic units incorporated within the Altar and Carnero complexes, specifically, and El Batamote belt, more broadly, were buried beneath a Laramide thrust complex during the Late Cretaceous–Paleogene Laramide orogeny. Initial cooling of the complexes occurred in the late Paleocene to Eocene. A second phase of cooling associated with detachment faulting occurred at ~17–15 Ma following intrusion of the ~22 Ma Rancho Herradura granodiorite.

### ACKNOWLEDGMENTS

CEJ was supported by the U.S. National Science Foundation (NSF) grant EAR 0408580. CJ-A and JCG-B received support from the Consejo Nacional de Ciencia y Tecnología (CONACYT), Research Project 166018. APB was supported by the NSF grant EAR 0408730 and a grant from the International Development Fund at Indiana University–Purdue University Indianapolis. We benefited from discussions with Carlos González-León, Alex Iriondo, Tim Lawton, Dave Mauel, Jon Nourse and Tom Peryam during the work. Special thanks are extended to Tim Lawton, Jon Nourse and an anonymous individual, all of whom provided extremely thorough, fair and helpful reviews of the manuscript.

### SUPPLEMENTARY MATERIAL

The Tables “A1. Sample information and coordinates,” “A2. Results for detrital zircon samples 662, 663, 663A, 664, 666, 668, 669 analyzed by LA-ICP-MS at the Arizona Laserchron Center, University of Arizona,” “A3. Results for detrital zircon sample CJ13-15 analyzed by LA-ICP-MS at the Laboratorio de Estudios Isotópicos (LEI), Centro de Geociencias, UNAM,” “A4. SHRIMP-RG analyses - Rancho Herradura granodiorite,” “A5.  $^{40}\text{Ar}/^{39}\text{Ar}$  results for biotite sample CJ13-15,” “A6.  $^{40}\text{Ar}/^{39}\text{Ar}$  results for muscovite sample CJ13-15,” and “A7.  $^{40}\text{Ar}/^{39}\text{Ar}$  results for plagioclase sample CJ13-15.” of the electronic supplement can be found at the journal web site <<http://rmcg.unam.mx/>>, in the table of contents of this issue.

### REFERENCES

- Anderson, T.H., Nourse, J.A., 2005, Pull-apart basins at releasing bends of the sinistral Late Jurassic Mojave-Sonora fault system, *in* Anderson, T.H., Nourse, J.A., McKee, J.W., Steiner, M.B. (eds.), *The Mojave-Sonora megashear hypothesis: Development, assessment, and alternatives: Geological Society of America Special Paper 393*, 97–122, DOI:10.1130/2005.2393(03).
- Anderson, T.H., Silver, L.T., 2005, The Mojave-Sonora megashear—Field and analytical studies leading to the conception and evolution of the hypothesis, *in* Anderson, T.H., Nourse, J.A., McKee, J.W., Steiner, M.B. (eds.), *The Mojave-Sonora megashear hypothesis: Development, assessment, and alternatives: Geological Society of America Special Paper 393*, 1–50, DOI:10.1130/2005.2393(01).
- Anderson, T.H., Rodríguez-Castañeda, J.L., Silver, L.T., 2005, Jurassic rocks in Sonora, Mexico: Relations to the Mojave-Sonora megashear and its inferred northwestward extension, *in* Anderson, T.H., Nourse, J.A., McKee,

- J.W., Steiner, M.B. (eds.), The Mojave-Sonora megashear hypothesis: Development, assessment, and alternatives: Geological Society of America Special Paper 393, 51-95, DOI:10.1130/2005.2393(02).
- Arvizu, H.E., Iriondo, A., 2015, Control temporal y geología del magmatismo Permo-Triásico en Sierra Los Tanques, NW Sonora, México: Evidencia del inicio del arco magmático cordillerano en el SW de Laurencia: Boletín de la Sociedad Geológica Mexicana, 67(3), 545-586.
- Arvizu, H.E., Iriondo, A., Izaguirre, A., Chávez-Cabello, G., Kamenov, G.D., Solís-Pichardo, G., Foster, D.A., Lozano-Santa Cruz, R., 2009, Rocas graníticas pérmicas en la Sierra Pinta, NW de Sonora, México: Magmatismo de subducción asociado al inicio del margen continental activo del SW de Norteamérica: Revista Mexicana de Ciencias Geológicas, 26(3), 709-728.
- Barth, A.P., Wooden, J.L., 2006, Timing of magmatism following initial convergence at a passive margin, southwestern U.S. Cordillera, and ages of lower crustal magma sources: Journal of Geology, 114, 231-245, DOI:10.1086/499573.
- Barth, A.P., Wooden, J.L., Coleman, D.S., Fanning, C.M., 2000, Geochronology of the Proterozoic basement of southwesternmost North America, and the origin and evolution of the Mojave crustal province: Tectonics, 19, 616-629, DOI:10.1029/1999TC001145.
- Barth, A.P., Jacobson, C.E., Coleman, D.S., Wooden, J.L., 2001a, Construction and tectonic evolution of Cordilleran continental crust: Examples from the San Gabriel and San Bernardino Mountains, in Dunne, G., Cooper, J., compilers, Geologic Excursions in the California Desert and Adjacent Transverse Ranges: Los Angeles, Pacific Section, SEPM (Society of Economic Paleontologists and Mineralogists), Book 88, 17-53.
- Barth, A.P., Wooden, J.L., Coleman, D.S., 2001b, SHRIMP-RG U-Pb zircon geochronology of Mesoproterozoic metamorphism and plutonism in the southwesternmost United States: The Journal of Geology, 109, 319-327, DOI:10.1086/319975.
- Barth, A.P., Wooden, J.L., Jacobson, C.E., Probst, K., 2004, U-Pb geochronology and geochemistry of the McCoy Mountains Formation, southeastern California: A Cretaceous retroarc foreland basin: Geological Society of America Bulletin, 116, 142-153, DOI:10.1130/B25288.1.
- Barth, A., Jacobson, C.E., Jacques-Ayala, C., Roldán, J., Pedrick, J.N., 2008, Age of the Altar complex and Cretaceous deformation in the Caborca-Altar region of northwestern Sonora: Geological Society of America Abstracts with Programs, 40(6), 198.
- Black, L.P., Kamo, S.L., Allen, C.M., Davis, D.W., Aleinikoff, J.N., Valley, J.W., Mundil, R.M., Campbell, I.H., Korsch, R.J., Williams, I.S., Foudoulis, C., 2004, Improved  $^{206}\text{Pb}/^{238}\text{U}$  microprobe geochronology by the monitoring of a trace-element-related matrix effect: SHRIMP, ID-TIMS, ELA-ICP-MS, and oxygen isotope documentation for a series of zircon standards: Chemical Geology, 205, 115-140.
- Busby, C., Fackler Adams, B., Mattinson, J., Deoreo, S., 2006, View of an intact oceanic arc, from surficial to mesozonal levels: Cretaceous Alisitos arc, Baja California: Journal of Volcanology and Geothermal Research, 149, 1-46, DOI:10.1016/j.jvolgeores.2005.06.009.
- Calmus, T., Sosson, M., 1995, Southwestern extension of the Papago Terrane into the Altar Desert region, northwestern Sonora, and its implications, in Jacques-Ayala, C., González-León, C.M., Roldán-Quintana, J. (eds.), Studies on the Mesozoic of Sonora and adjacent areas: Geological Society of America Special Paper, 301, 99-109.
- Castro-Escárcega, J.J., Morales-Morales, H., Salazar-Velázquez, R., 2002a, Carta geológico-minera Nogales H12-2, escala 1:250,000: Pachuca, Hidalgo, México, Consejo de Recursos Minerales, 1 map.
- Castro-Escárcega, J.J., Morales-Morales, H., Salazar-Velázquez, R., 2002b, Carta geológico-minera Puerto Peñasco H12-1, escala 1:250,000: Pachuca, Hidalgo, México, Consejo de Recursos Minerales, 1 map.
- Clinkscales, C.A., Lawton, T.F., 2015, Timing of Late Cretaceous shortening and basin development, Little Hatched Mountains, southwestern New Mexico, USA – implications for regional Laramide tectonics: Basin Research, 27, 453-472, DOI:10.1111/bre.12083.
- Cooper, G.A., Arellano, A.R.V., 1946, Stratigraphy near Caborca, northwest Sonora, Mexico: American Association of Petroleum Geologists Bulletin, 30, 606-610.
- Damon, P.E., Livingston, D.E., Mauger, R.L., Giletti, B.J., Pantoja-Alor, J., 1962, Edad del Precámbrico "Anterior" y de otras rocas del zócalo de la región de Caborca-Altar de la parte noroccidental del Estado de Sonora: Universidad Nacional Autónoma de México, Instituto de Geología Boletín 64, 11-44.
- De Jong, K.A., Escarcega-Escarcega, J.A., Damon, P.E., 1988, Eastward thrusting, southwestward folding, and westward backsliding in the Sierra La Vibora, Sonora, Mexico: Geology, 16, 904-907.
- Dickinson, W.R., Gehrels, G.E., 2009a, Use of U-Pb ages of detrital zircons to infer maximum depositional ages of strata: A test against a Colorado Plateau Mesozoic database: Earth and Planetary Science Letters, 288, 115-125, DOI:10.1016/j.epsl.2009.09.013.
- Dickinson, W.R., Gehrels, G.E., 2009b, U-Pb ages of detrital zircons in Jurassic eolian and associated sandstones of the Colorado Plateau: Evidence for transcontinental dispersal and intraregional recycling of sediment: Geological Society of America Bulletin, 121, 408-433, DOI:10.1130/B26406.1.
- Dickinson, W.R., Lawton, T.F., 2001, Carboniferous to Cretaceous assembly and fragmentation of Mexico: Geological Society of America Bulletin, 113, 1142-1160, DOI:10.1016/S0895-9811(01)00046-3.
- Dickinson, W.R., Lawton, T.F., Gehrels, G.E., 2009, Recycling detrital zircons: A case study from the Cretaceous Bisbee Group of southern Arizona: Geology, 37, 503-506, DOI:10.1130/G25646A.1.
- Dickinson, W.R., Lawton, T.F., Pecha, M., Davis, S.J., Gehrels, G.E., Young, R.A., 2012, Provenance of the Paleogene Colton Formation (Uinta Basin) and Cretaceous–Paleogene provenance evolution in the Utah foreland: Evidence from U-Pb ages of detrital zircons, paleocurrent trends, and sandstone petrofacies: Geosphere, 8, 854-880, DOI:10.1130/GES00763.1.
- Farmer, G.L., Bowring, S.A., Matzel, J., Espinosa Maldonado, G., Fedo, C., Wooden, J., 2005, Paleoproterozoic Mojave Province in northwestern Mexico? Isotopic and U-Pb zircon geochronologic studies of Precambrian and Cambrian crystalline and sedimentary rocks, Caborca, Sonora, in Anderson, T.H., Nourse, J.A., McKee, J.W., Steiner, M.B. (eds.), The Mojave-Sonora megashear hypothesis: Development, assessment, and alternatives: Geological Society of America Special Paper 393, 183-198, DOI:10.1130/2005.2393(05).
- Fitz-Díaz, E., Lawton T.F., Juárez-Arriaga, E., Chávez-Cabello, G., 2018, The Cretaceous-Paleogene Mexican orogen: Structure, basin development, magmatism and tectonics: Earth-Science Reviews, 183, 56-84, DOI:10.1016/j.earscirev.2017.03.002.
- García y Barragán, J.C., 1992, Geology of the western Cerros El Amol, Altar, Sonora, Mexico: Cincinnati, Ohio, University of Cincinnati, M.Sc. Thesis, 80 pp.
- García y Barragán, J.C., Jacques-Ayala, C., 2010, Estratigrafía del Cretácico de Sonora, México, in Calmus, T. (ed.), Contribuciones a la geología de Sonora, México: Universidad Nacional Autónoma de México, Instituto de Geología, Boletín 118, 6, 143-229.
- García y Barragán, J.C., Jacques-Ayala, C., De Jong, K., 1998, Stratigraphy of the Cerros El Amol, Altar, Sonora, Mexico, in Clark, K.F. (ed.), Gold deposits of northern Sonora, Mexico: Society of Economic Geologists Guidebook Series 30, 101-116.
- García-Cortez, J.A., Terán-Martínez, G.E., Zamora-Tovar, E., 2002, Carta geológico-minera Caborca H12-45, escala 1:250,000: Pachuca, Hidalgo, México, Consejo de Recursos Minerales, 1 map.
- Gehrels, G., Pecha, M., 2014, Detrital zircon U-Pb geochronology and Hf isotope geochemistry of Paleozoic and Triassic passive margin strata of western North America: Geosphere, 10, 49-65, DOI:10.1130/GES00889.1.
- Gehrels, G.E., Stewart, J.H., 1998, Detrital zircon U-Pb geochronology of Cambrian to Triassic miogeoclinal and eugeoclinal strata of Sonora, Mexico: Journal of Geophysical Research, 103, 2471-2487.
- Gehrels, G.E., Dickinson, W.R., Ross, G.M., Stewart, J.H., Howell, D.G., 1995, Detrital zircon reference for Cambrian to Triassic miogeoclinal strata of western North America: Geology, 23, 831-834.
- Gehrels, G.E., Valencia, V.A., Ruiz, J., 2008, Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation–multicollector–inductively coupled plasma–mass spectrometry: Geochemistry, Geophysics, Geosystems, 9, Q03017, DOI:10.1029/2007GC001805.
- Gehrels, G.E., Blakey, R., Karlstrom, K.E., Timmons, J.M., Dickinson, W.R., Pecha, M., 2011, Detrital zircon U-Pb geochronology of Paleozoic strata in the Grand Canyon: Lithosphere, 3, 183-200, DOI:10.1130/L121.1.
- González-León, C.M., McIntosh, W.C., Lozano-Santacruz, R., Valencia-Moreno,

- M., Amaya-Martínez, R., Rodríguez-Castañeda, J.L., 2000, Cretaceous and Tertiary sedimentary, magmatic and tectonic evolution of north-central Sonora (Arizpe and Bacanuchi Quadrangles), northwest Mexico: *Geological Society of America Bulletin*, 112, 600-610.
- González-León, C.M., Valencia, V.A., Lawton, T.F., Amato, J.M., Gehrels, G.E., Leggett, E.J., Montijo-Contreras, O., Fernández, M.A., 2009, The lower Mesozoic record of detrital zircon U-Pb geochronology of Sonora, México, and its paleogeographic implications: *Revista Mexicana de Ciencias Geológicas*, 26(2), 301-314.
- González-León, C., Solari, L., Solé, J., Ducea, M.N., Lawton, T.F., Bernal, J.P., González-Becuar, E., Gray, F., López-Martínez, M., Lozano-Santacruz, R., 2011, Stratigraphy, geochronology, and geochemistry of the Laramide magmatic arc in north-central Sonora, Mexico: *Geosphere*, 7, 1392-1418, DOI:10.1130/GES00679.1.
- González-León, C., Solari, L., Madhavaraju, J., 2017, Stratigraphy, geochronology and regional tectonic setting of the Late Cretaceous (ca. 82-70 Ma) Cabullona basin, Sonora, Mexico: *Journal of South American Earth Sciences*, 80, 494-511, DOI:10.1016/j.jsames.2017.09.037.
- Goodwin, L.B., Haxel, G.B., 1990, Structural evolution of the southern Baboquivari Mountains, south-central Arizona and north-central Sonora: *Tectonics*, 9, 1077-1095.
- Harding, L.E., Coney, P.J., 1985, The geology of the McCoy Mountains Formation, southeastern California and southwestern Arizona: *Geological Society of America Bulletin*, 96, 755-769.
- Haxel, G.B., Wright, J.E., May, D.J., Tosdal, R.M., 1980, Reconnaissance geology of the Mesozoic and lower Cenozoic rocks of the southern Papago Indian Reservation, Arizona: A preliminary report, in Jenny, J. P., Stone, C. (eds.), *Studies in western Arizona*: Arizona Geological Society Digest, 12, 17-29.
- Haxel, G.B., Tosdal, R.M., May, D.J., Wright, J.E., 1984, Latest Cretaceous and early Tertiary orogenesis in south-central Arizona: Thrust faulting, regional metamorphism, and granitic plutonism: *Geological Society of America Bulletin*, 95, 631-653.
- Haxel, G.B., Anderson, T.H., Briskey, J.A., Tosdal, R.M., Wright, J.E., May, D.J., 2008, Late Jurassic igneous rocks in south-central Arizona and north-central Sonora: Magmatic accompaniment of crustal extension, in Spencer, J.E., Titley, S.R. (eds.), *Ores and orogenesis: Circum-Pacific tectonics, geologic evolution, and ore deposits*: Arizona Geological Society Digest, 22, 333-355.
- Hayama, Y., Shibata, K., Takeda, H., 1984, K-Ar ages of the low-grade metamorphic rocks in the Altar Massif, northwest Sonora, Mexico: *Journal of the Geological Society of Japan*, 90, 589-596.
- INEGI (Instituto Nacional de Estadística y Geografía), 2001, Carta topográfica 1:50,000 Altar H12A67, México: Aguascalientes, Mexico, 1 map.
- Iriondo, A., Premo, W.R., Martínez-Torres, L.M., Budahn, J.R., Atkinson, W.W. Jr., Siems, D.F., Guaras-Gonzalez, B., 2004, Isotopic, geochemical, and temporal characterization of Proterozoic basement rocks in the Quitovac region, northwestern Sonora, Mexico: Implications for the reconstruction of the southwestern margin of Laurentia: *Geological Society of America Bulletin*, 116, 154-170, DOI:10.1130/B25288.1.
- Iriondo, A., Martínez-Torres, L.M., Kunk, M.J., Atkinson, W.W. Jr., Premo, W.R., McIntosh, W.C., 2005, Northward Laramide thrusting in the Quitovac region, northwestern Sonora, Mexico: Implications for the juxtaposition of Paleoproterozoic basement blocks and the Mojave-Sonora megashear hypothesis, in Anderson, T.H., Nourse, J.A., McKee, J.W., Steiner, M.B. (eds.), *The Mojave-Sonora megashear hypothesis: Development, assessment, and alternatives*: Geological Society of America Special Paper 393, 631-669, DOI:10.1130/2005.2393(24).
- Jacobson, C.E., Grove, M., Stamp, M.M., Vučić, A., Oyarzabal, F.R., Haxel, G.B., Tosdal, R.M., Sherrod, D.R., 2002, Exhumation history of the Orocochia Schist and related rocks in the Gavilan Hills area of southeasternmost California, in Barth, A. (ed.), *Contributions to crustal evolution of the southwestern United States*: Geological Society of America Special Paper 365, 129-154.
- Jacques-Ayala, C., 1993, The Cretaceous in the Caborca-Santa Ana region, northern Sonora, Mexico: Cincinnati, U.S.A., University of Cincinnati, Ph.D. Thesis, 146 pp.
- Jacques-Ayala, C., 1995, Paleogeography and provenance of the Lower Cretaceous Bisbee Group in the Caborca-Santa Ana area, northwestern Sonora, in Jacques-Ayala, C., González-León, C.M., Roldán-Quintana, J. (eds.), *Studies on the Mesozoic of Sonora and adjacent areas*: Geological Society of America Special Paper 301, 79-98.
- Jacques-Ayala, C., 1999, Stratigraphy of El Chanate Group (Late Cretaceous) and its implications for the tectonic evolution of northwestern Sonora, Mexico: *Revista Mexicana de Ciencias Geológicas*, 16(2), 97-120.
- Jacques-Ayala, C., 2000, Deletion of the Mojave-Sonora megashear as a terrane boundary—stratigraphic, sedimentologic, and structural evidence, in Calmuss, T., Pérez-Segura, E. (eds.), *Cuarta Reunión sobre la Geología del Noroeste de México y Áreas Adyacentes*, Libro de Resúmenes: Instituto de Geología, UNAM, Estación Regional del Noroeste, Publicaciones Ocasionales, 2, 46-47.
- Jacques-Ayala, C., DeJong, K.A., 1996, Extension of the Orocochia Schist belt into northwestern Sonora, Mexico: A working hypothesis: *Geological Society of America Abstracts with Programs*, 28(5), 78.
- Jacques-Ayala, C., García y Barragán, J.C., 2015, The Altar sole fault, boundary of the latest Cretaceous-early Eocene foreland/hinterland domains in northwestern Sonora, Mexico: *Geos* 35, 70.
- Jacques-Ayala, C., García y Barragán, J.C., 2018, Late Cretaceous foreland/hinterland basins in the Caborca area: a new approach to understand the tectonic evolution of NW Sonora, Mexico: *Geological Society of America Abstracts with Programs*, 50(5), DOI:10.1130/abs/2018RM-313885.
- Jacques-Ayala, C., García y Barragán, J.C., DeJong, K.A., 1990, Caborca-Altar geology: Cretaceous sedimentation and compression, Tertiary uplift and extension, in Gehrels, G.E., Spencer, J.E. (eds.), *Geologic excursions through the Sonoran Desert region, Arizona and Sonora*: Arizona Geological Survey Special Paper 7, 165-182.
- Jacques-Ayala, C., Barth, A.P., Wooden, J.L., Jacobson, C.E., 2009, Provenance and U-Pb geochronology of El Chanate Group (Late Cretaceous), northwestern Sonora, Mexico, and its tectonic significance: *International Geology Review*, 51, 1051-1077, DOI:10.1080/00206810902945066.
- Kimbrough, D.L., Grove, G., Morton, D.M., 2015, Timing and significance of gabbro emplacement within two distinct plutonic domains of the Peninsular Ranges batholith, southern and Baja California: *Geological Society of America Bulletin*, 127, 19-37, DOI:10.1130/B30914.1.
- Lawton, T.F., Molina Garza, R.S., 2014, U-Pb geochronology of the type Nazas Formation and superjacent strata, northeastern Durango, Mexico: Implications of a Jurassic age for continental-arc magmatism in north-central Mexico: *Geological Society of America Bulletin*, 126, 1181-1199, DOI:10.1130/B30827.1.
- Leggett, W.J., 2009, Stratigraphy, Sedimentology and Geochronology of Lower and Middle Jurassic Rocks Near Rancho San Martin del Rincon, North-Central Sonora, Mexico: Las Cruces, New Mexico, U.S.A., New Mexico State University, M.S. Thesis, 203 pp.
- Ludwig, K.R., 2003, Isoplot/Ex 3.00: A geochronological toolkit for Microsoft Excel: Berkeley Geochronology Center, Special Publication 4, 70 pp.
- Mauel, D.J., 2008, Stratigraphy, sedimentology, and geochronology of Upper Jurassic rocks of the Altar-Cucurpe Basin in the vicinity of Cucurpe, Northcentral Sonora, Mexico: Las Cruces, New Mexico, U.S.A., New Mexico State University, M.S. Thesis, 174 pp.
- Mauel, D.J., Lawton, T.F., González-León, C., Iriondo, A., Amato, J.M., 2011, Stratigraphy and age of Upper Jurassic strata in north-central Sonora, Mexico: Southwestern Laurentian record of crustal extension and tectonic transition: *Geosphere*, 7, 390-414, DOI:10.1130/GES00600.1.
- McDougall, I., Harrison, T.M., 1999, *Geochronology and Thermochronology by the <sup>40</sup>Ar/<sup>39</sup>Ar Method* (2nd edition): New York, Oxford University Press, 269 pp.
- McDowell, F.W., Roldán-Quintana, J., Connelly, J., 2001, Duration of Late Cretaceous-early Tertiary magmatism in east-central Sonora, México: *Geological Society of America Bulletin*, 113, 521-531.
- Molina-Garza, R.S., Iriondo, A., 2007, The Mojave-Sonora megashear: The hypothesis, the controversy, and the current state of knowledge, in Alaniz-Álvarez, S.A., Nieto-Samaniego, Á.F. (eds.), *Geology of México: Celebrating the centenary of the Geological Society of México*: Geological Society of America Special Paper 422, 233-259, DOI:10.1130/2007.2422(07).
- Nourse, J.A., 1995, Jurassic-Cretaceous paleogeography of the Magdalena region, northern Sonora, and its influence on the positioning of Tertiary metamorphic core complexes, in Jacques-Ayala, C., González-León, C.M., Roldán-Quintana, J. (eds.), *Studies on the Mesozoic of Sonora and adjacent areas*: Geological Society of America Special Paper 301, 59-78.

- Nourse, J.A., 2001, Tectonic insights from an Upper Jurassic–Lower Cretaceous stretched-clast conglomerate, Caborca–Altar region, Sonora, Mexico: *Journal of South American Earth Sciences*, 14, 453–474.
- Nourse, J.A., Anderson, T.H., Silver, L.T., 1994, Tertiary metamorphic core complexes in Sonora, northwestern Mexico: *Tectonics*, 13(5), 1161–1182.
- Nourse, J.A., Premo, W.R., Iriondo, A., Stahl, E.R., 2005, Constraining Proterozoic basement complexes near the truncated margin of Laurentia, northwestern Sonora–Arizona international border region, in Anderson, T.H., Nourse, J.A., McKee, J.W., Steiner, M.B. (eds.), *The Mojave–Sonora megashear hypothesis: Development, assessment and alternatives*: Geological Society of America Special Paper 393, 123–182 DOI:10.1130/2005.2393(04).
- Nourse, J.A., Oskin, M.E., Iriondo, A., Premo, W.R., 2007, Laramide fold–thrust belt overprinted by Middle Miocene detachment faults, Caborca region, Sonora, Mexico, in *Ores and Orogenesis; A symposium honoring the career of William R. Dickinson*: Tucson, Arizona, Arizona Geological Society, Program with abstracts, 96–97.
- Ortega-Gutiérrez, F., Elías-Herrera, M., Morán-Zenteno, J.D., Solari, L., Luna-González, L., Schaaf, P., 2014, A review of batholiths and other plutonic intrusions of Mexico, *Gondwana Research*, 26, 834–868, DOI:10.1016/j.gr.2014.05.002.
- Peryam, T.C., Lawton, T.F., Amato, J.M., González-León, C.M., Mauel, D.J., 2012, Lower Cretaceous strata of the Sonora Bisbee Basin: A record of the tectonomagmatic evolution of northwestern Mexico: *Geological Society of America Bulletin*, 124, 532–548, DOI:10.1130/B30456.1.
- Riggs, N.R., Barth, A.P., Walker, J.D., Lindner, P., González-León, C., Cecil, M.R., Marsaglia, K., 2016, Implications of Permian magmatism in southwest Laurentia for tectonic reconstructions: *Geological Society of America Abstracts with Programs*, 48(7), DOI:10.1130/abs/2016AM-279985.
- Roldán-Quintana, J., McDowell, F.W., Delgado-Granados, H., Valencia-Moreno, M., 2009, East–west variations in age, chemical and isotopic composition of the Laramide batholith in southern Sonora, Mexico: *Revista Mexicana de Ciencias Geológicas*, 26(3), 543–563.
- Solari, L.A., Gómez-Tuena, A., Bernal, J.P., Pérez-Arvizu, O., Tanner, M., 2010, U–Pb zircon geochronology by an integrated LA–ICPMS microanalytical workstation: Achievements in precision and accuracy: *Geostandards and Geoanalytical Research*, 34, 5–18, DOI:10.1111/j.1751-908X.2009.00027.x.
- Spencer, J.E., Richard, S.M., Gehrels, G.E., Gleason, J.D., Dickinson, W.R., 2011, Age and tectonic setting of the Mesozoic McCoy Mountains Formation in western Arizona, USA: *Geological Society of America Bulletin*, 123, 1258–1274, DOI:10.1130/B30206.1.
- Stewart, J.H., Gehrels, G.E., Barth, A.P., Link, P.K., Christie-Blick, N., Wrucke, C.T., 2001, Detrital zircon provenance of Mesoproterozoic to Cambrian arenites in the western United States and northwestern Mexico: *Geological Society of America Bulletin* 113, 1343–1356.
- Terán-Martínez, G.E., Cerecero-Luna, M., Mendoza-Romero, M., Castro-Escárcega, J.J., García-Cortez, J.A., 1999, Carta geológico-minera Cananea H12-5, escala 1:250,000: Pachuca, Hidalgo, México, Consejo de Recursos Minerales, 1 map.
- Walker, J.D., Geissman, J.W., Bowring, S.A., Babcock, L.E., compilers, 2018, *Geologic Time Scale v. 5.0*: Geological Society of America, <https://DOI.org/10.1130/2018.CTS005R3C>.
- Wetmore, P.H., Herzig, Ch., Alseben, H., Sutherland, M., Schmidt, K.L., Schultz, P.W., Paterson, S.R., 2003, Mesozoic tectonic evolution of the Peninsular Ranges of southern and Baja California, in Johnson, S.E., Paterson, S.R., Fletcher, J.M., Girty, G.H., Kimbrough, D.L., Martín-Barajas, A. (eds.), *Tectonic evolution of northwestern Mexico and the southwestern USA*: Geological Society of America Special Paper 374, 93–116.
- Wong, M.S., Gans, P.B., Scheier, J., 2010, The  $^{40}\text{Ar}/^{39}\text{Ar}$  thermochronology of core complexes and other basement rocks in Sonora, Mexico: Implications for Cenozoic tectonic evolution of northwestern Mexico: *Journal of Geophysical Research*, 115, B07414, DOI:10.1029/2009JB007032.

Manuscript received: april 3, 2018

Corrected manuscript received: november 2, 2018

Manuscript accepted: november 21, 2018