



Scientific Article

Morphological characterization, phylogeny and pathogenesis of *Setophoma terrestris* causing corky and pink roots of tomato (*Solanum lycopersicum*) in Sinaloa, Mexico

Ana María López-López, Juan Manuel Tovar-Pedraza, Josefina León-Félix, ¹Centro de Investigación en Alimentación y Desarrollo, Coordinación Regional Culiacán, 80110, Sinaloa, México; Raúl Allende-Molar, Universidad Veracruzana, Facultad de Ciencias Biológicas y Agropecuarias, Tuxpan, 92895, Veracruz, México; Nelson Bernardi Lima, Universidad Nacional de Catamarca, Departamento de Ingeniería Agronómica, 4700, Catamarca, Argentina; ¹Isidro Márquez-Zequera, ¹Raymundo Saúl García-Estrada.

***Corresponding Author:**

Raymundo Saúl
García Estrada
rsgarcia@ciad.mx

Section:
Periodical Issue

Received:
27 September, 2023

Accepted:
28 January, 2024

Published:
13 February, 2024

Citation:

López-López AM, Tovar-Pedraza JM, León-Félix J, Allende-Molar R, Bernardi LN, Márquez-Zequera I and García-Estrada RS. 2024. Morphological characterization, phylogeny and pathogenesis of *Setophoma terrestris* causing corky and pink roots of tomato (*Solanum lycopersicum*) in Sinaloa, Mexico. Mexican Journal of Phytopathology 42(2): 12. <https://doi.org/10.18781/R.MEX.FIT.2309-5>



ABSTRACT

Objective / Background. Tomato (*Solanum lycopersicum*) is one of Mexico's main crops. In the years 2017 and 2018, symptoms of corky and pink roots were observed with an incidence of 10 to 20% in Culiacan, Sinaloa, Mexico. In the foliage, plants presented a generalized chlorosis, with stunted growth and senescence in the leaves. In the roots, brown and pink lesions were formed, as well as a corky texture. The objective of this study was to morphologically and molecularly characterize fungal isolates associated to corky and pink root in tomato orchards in Culiacan, Sinaloa, as well as to evaluate their pathogenicity.

Materials and methods. Monoconidial isolates were obtained and they were identified as *Setophoma terrestris*, based on their morphological characteristics. To confirm the identity, the area of the internal transcribed spacers (ITS) of the rDNA was amplified and sequenced, along with a fragment of the gene 28S of the rRNA (LSU).

Results. Using the sequences obtained, a phylogenetic tree was created using the Bayesian Inference and it was found that the sequences were grouped with the ex-type sequences of *Setophoma terrestris*. The pathogenicity of the isolates was verified by inoculating mycelial discs into the root of 10 one-month-old tomato seedlings. The roots of the seedlings inoculated with PDA discs without mycelium served as a control. Thirty days after inoculation, corky and pink root symptoms appeared, whereas the roots of control plants remained healthy.

Conclusion. According to the morphological characterization, the molecular identification and the pathogenicity tests, *Setophoma terrestris* was confirmed to be the causal agent of corky and pink root in agricultural tomato orchards in Culiacan, Sinaloa.

Keywords: *Setophoma*, fungus, tomato, bayesian inference, root disease.

INTRODUCTION

Mexico is the ninth most important producer of tomato (*Solanum lycopersicum*) in the world. In the 2018-2019 agricultural cycle, the country produced a total of 938,009 t, with Sinaloa being the state at the top of the list of nationwide producers with 605,261 t (SIAP, 2020). However, pests and diseases are the main problems that impact this crop (Jones *et al.* 2016).

Out of the diseases in tomato, corky root is scarcely studied, despite being very troublesome; it creates brown lesions and corky roots. Initially, the lesions are limited and later extends progressively and surrounds the roots, developing lesions that range from a few milimeters to several centimeters (Blancard, 2011). In this context, the disease called corky root is one of the most sever problems in recent years in agricultural plots dedicated to the tomato crop in Mexico (Pardo, 1999; Martínez-Ruiz *et al.*, 2016).

Although the corky root disease is often found in greenhouse crops and its distribution was initially reported in Europe (Schneider, 1966), United States (Campbell *et al.*, 1982), Lebanon (Davet, 1973) and Chile (Olavarria, 1991), there are not enough studies related with the causal agent of the disease (USDA, 2017).

Generally, corky root in tomato is associated with *Pseudopyrenochaeta lycopersici* (sin. *Pyrenochaeta lycopersici*) as its causal agent (Golzar, 2009). However, a study carried out in Mexico proved, with morphological bases, phylogenetic analyses and pathogenicity tests, that corky root in Sinaloa fields is caused by *Setophoma terrestris* and additionally, it could cause pink root in the crop (López-López *et al.*, 2020).

According to De Gruyter *et al.* (2010), *S. terrestris* and *P. lycopersici* share very similar morphological characteristics, which were taxonomically differentiated according to phylogenetic analyses with linked sequenesces from the ITS area and fragments of the LSU gene (Valenzuela-López *et al.*, 2018).

Setophoma is a fungal genus that forms pycnidia, which are asexual reproductive structures. The pycnidium can have a globular to semi-globular structure, almost setae-like, with an ostiolar neck around it. Inside the pycnidia are the spores called conidia, which are hyaline, cylindrical, septated and mainly guttulate (De Gruyter *et al.*, 2010).

Setophoma terrestris has been mostly studied in onion (*Allium cepa*) crops, since they are its main host, causing the disease known as pink root, which is important and has a large global economic impact. This fungus attacks the root system first, and progressively affects the entire plant until its death (Castillo and Plata, 2010). It has also been reported to cause damage in other hosts such as canola (*Brassica napus*) in Canada (Yang *et al.*, 2017), zucchini (*Curcubita maxima*) in the USA (Rivedal *et al.*, 2018), as well as garlic (*Allium sativum*) and tea (*Camellia sinensis*) plants in China (Liu *et al.*, 2019; Zhang *et al.*, 2019).

An earlier study (López-López *et al.*, 2020) proved, for the first time, the infectious nature of *Setophoma terrestris* in tomato plants (*S. lycopersicum*) in Mexico, based on one single isolate. The aim of this study was to morphologically and molecularly characterize the fungal isolations associated with the symptoms of corky root and pink root in tomato orchards in Culiacán, Sinaloa, as well as to evaluate their pathogenicity.

MATERIALS AND METHODS

Sample collection. Samples were taken from five sites (Figure 1), which were distributed in five locations in the valley of Culiacán, as well as on the limits of Navolato, Sinaloa, with an intensive tomato production with undetermined growth in the production stage (age three months). These sites had a history of corky and pink root in tomato. In some plants at the harvest stage, different types of symptoms were observed in the foliar area, such as chlorosis and reduction in growth. Therefore, targeted sampling was conducted according to the symptoms mentioned. In addition, the roots of the plants were examined by extraction and observation of the damage along them. Once the symptoms related to the disease of the crop were confirmed, 15 samples of diseased tomato roots were gathered from each site. The collection sites were shade house-type greenhouses, each one approximately 5 ha. Finally, they were placed in plastic bags and transported to the Phytopathology Laboratory of the Coordinación Culiacán of CIAD for analysis.

Isolation, purification and conservation of fungi. The diseased roots were submerged in drinking water to remove excess soil without harming the plant tissue. Subsequently, from necrotic lesions developed in tomato roots, fragments of 5 mm in length were cut, then surface-disinfested by immersion in a 2% sodium hypochlorite solution for 2 minutes. They were rinsed in sterile distilled water and placed in Petri dishes containing acidified potato dextrose agar (APDA) and acidified V8 agar medium (V8A) and incubated at 25 °C in continuous darkness. Afterwards, the fungal isolates were transferred to new Petri dishes with V8A,



Figure 1. Location of the sites of recollection of plants with corky root symptoms in agricultural systems with tomato production in the valley of Culiacán, Sinaloa. Site 1 (24°42'56.58"N, 107°26'42.86"W), Site 2 (24°35'21.67"N, 107°24'56.50"W), Site 3 (24°46'2.77"N, 107°32'51.30"W), Site 4 (24°48'47.44"N, 107°39'26.19"W) and Site 5 (24°32'34.07"N, 107°26'44.93"W).

and after 10 days in incubation at 25 °C, the isolations were purified using the monospore culture technique (Crous *et al.*, 2019), which consisted of placing the pycnidia on microscope slides and the conidia were placed under a compound microscope. From these conidia, monospore cultures were obtained in Petri dishes with V8A medium, and 7 days after planting, a pink mycelial growth was observed. The fungal isolates were preserved by placing discs, 4 mm in diameter, from colonies with 5 days of growth into sterile 2 mL tubes containing sterile distilled water and stored at room temperature in the strain collection of the Phytopathology Laboratory in the Culiacán Unit of CIAD.

Morphological characterization. In order to stimulate the production of asexual structures, approximately three tomato roots were placed in moist chambers at a temperature of 20 °C. After 30 days, they were removed and observed under a stereoscopic microscope, locating the pycnidia and placing them on microscope slides with sterile distilled water for their observation and characterization.

The morphological characterization of the isolates was performed using keys and descriptions reported by De Gruyter *et al.* (2010). In order to do this, the fungus was cultivated in Petri dishes with an V8A medium for 10 days. The isolations that did not form pycnids in the V8A medium were stimulated using the method described by Clerjeau (1973), which consists in placing melon (*Cucumis melo*) seedling roots on the mycelium of the fungus of interest with 5-7 days of growth and later incubate once again, at room temperature (25–30 °C) for 15 to 20 days for the artificial production of pycnidia. Once produced, the characteristics of the pycnidia, conidia and setae were examined under a compound microscope (Zeiss®, Germany) with a built-in digital camera (AxioCam) and 50 measurements were taken of each one of the different asexual fungal structures of each isolate.

DNA extraction, PCR amplification and sequencing. Total DNA extraction as carried out from seven-day-old fungal colonies, cultivated in an V8A medium. The mycelium of each isolation was scraped with a sterile microscope slide, macerated with a sterile mortar using liquid nitrogen and transferred to a 1.5 mL microcentrifuge tube with 500 µL of hexadecyltrimethylammonium bromide solution, CTAB (Tris 100 mM, pH 8; 20 mM of EDTA, pH 8; NaCl, 1.4 M and 3% of CTAB), mixed in a vortex for 10 s and incubated for 60 min at 65 °C. Later, 700 µL of isolamilic chlorophorm-alcohol (24:1v/v) was added, run through a vortex for 10 s and placed in a 5810 R centrifuge (Eppendorf®, EE. UU.) at 13000 g for 10 min. Next, the supernatant was transferred to a 1.5 mL microcentrifuge tube and 700 µL of isopropanol were added. The tubes were mixed by inversion 4 to 5 times and left at -20 °C for 10 min, and were then centrifuges at 13000 g for 10 min to precipitate the DNA. Subsequently, the precipitate was washed twice with 500 µL of ethanol at 70%, centrifuging at 13000 g for 5 min and the supernatant was discarded every time. Afterwards, the DNA pellet was left to dry by placing the inverted tubes on absorbent paper for no more than 15 min to allow the evaporation of the ethanol and finally, the DNA was resuspended to allow the ethanol to evaporate and finally, the DNA was resuspended in 100 µL of sterile water, DNase- and RNase-free. The quality and concentration of the DNA was evaluated in a Q 3000 UV spectrophotometer (Quawell, EE. UU.) and electrophoresis in 1% agarose gel, with DNA being considered adequate when it had purity levels (ratio of A260/280) ranging between 1.8 and 2.0 and a concentration (A260) of approximately 2.0 µg µL⁻¹, as well as intact high-molecular weight bands, respectively. The DNA was stored at -20 °C for later use.

The amplification of the ITS region of the rDNA and the long subunit (LSU) was performed using endpoint polymerase chain reaction (PCR). The ITS region (ITS1–5.8S–ITS2) was amplified using primers ITS1 (TCCGTAGGTGAACCTGCGG) and ITS4 (TCCTCCGCTTATTGA TATGC) with 480 pb, approximately (White

et al., 1990), whereas the amplified band of the LSU gene (including domains D1 and D2) was performed with a nested PCR using primers PM3 (Takamatsu and Kano, 2001) /TW14 (Mori *et al.*, 2000) and NL1/TW14 (Mori *et al.*, 2000) for the first and second reaction, respectively. The reaction mixture was prepared at a final volume of 25 μ L, containing 1X PCR buffer, 2.5 mM of $MgCl_2$, 0.5 mM of dNTPs, 1 μ M of each primer, 1U of polymerase DNA (Promega[®], USA) and 100 ng of DNA. The PCR was carried out under the following thermocycling conditions (C-1000 thermocycler, Bio-Rad[®], USA): initial denaturalization at 95 °C for 5 min, 35 cycles at 95 °C for 30 s, 54 °C for 30 s, 72 °C for 30 s, with a final extension of 72 °C for 10 min. For the amplification of the 28S gene, the same thermocycling conditions were used, except for the temperature of alignment, which was 52 °C. The amplified products were analyzed by electrophoresis in 1% agarose gel dyed with ethidium bromide. Electrophoresis was carried out at 90 V for 40 min and the results were viewed in an M-26X transilluminator in a GelDoc-ItTM 300 image system (UVP[®], USA). The PCR products were purified using the DNA Clean & Concentrator kit (Zymo Research[®], USA), following the recommendations by the manufacturer and they were sequenced in the Macrogen company (www.macrogen.com) in Seoul, South Korea.

Phylogenetic analysis. The quality of the nucleotide sequences and the assembly method were carried out using the Staden package (Staden *et al.*, 2000). The sequences were cleaned and those generated from the ITS region and the large subunit (LSU) of the rRNA were aligned with sequences from different species of the *Setophoma* genus obtained from the GenBank and from published reference data (De Gruyter *et al.*, 2010; Liu *et al.*, 2019) (Table 1). Multiple alignments were generated from sequences using ClustalW as implemented in MEGA 11 (Tamura *et al.*, 2021) and they were improved manually to allow the maximum similarity between the dataset. The Bayesian phylogenetic estimations were inferred with MrBayes 3.2.6 (Ronquist *et al.*, 2012) implemented in the cluster CIPRES (<https://www.phylo.org/portal2/home.action>) and selecting the nucleotide substitution model that best fit the data of this investigation, according to the Akaike Information Criterion (AIC) obtained with MrModeltest 2.3 (Nylander, 2004). The number of generations was 10,000,000 and the subsequent probabilities were calculated after discarding the first 25% of the iterations. The phylogenetic trees were viewed using TreeView (Page, 1996).

Pathogenicity tests. The pathogenicity of seven fungal isolates was determined by inoculating them in the main roots of 30-day-old cv. 8444 tomato seedlings. Ten plants were used for each one of the isolations, along with 10 non-inoculated plants there served as a control. V8A plugs, 4 mm in diameter, with an active growth

Table 1. Information of fungal isolates and GenBank accession numbers of *Setophoma* species used in the phylogeneti analysis.

Species	Collection code	Host	Country	Access number	
				LSU	ITS
<i>Setophoma antiqua</i>	LC6594	<i>Camellia sinensis</i>	China	MK511947	MK511909
	LC6595	<i>C. sinensis</i>	China	MK511948	MK511910
<i>S. chromolaenae</i>	CBS 135105 ^x	<i>Chromolaena odorata</i>	Brazil	KF251747	KF251244
<i>S. cyperi</i>	CBS 141450 ^x	<i>Cyperus sphaerocephala</i>	South Africa	KX228337	KX228286
<i>S. endophytica</i>	CGMCC 3.19528	<i>C. sinensis</i>	China	MK511956	MK511931
	LC3164	<i>C. sinensis</i>	China	MK511957	MK511932
<i>S. longinqua</i>	LC6593 ^x	<i>C. sinensis</i>	China	MK511946	MK511908
<i>S. poaeicola</i>	MFLUCC 16-0880 ^x	-	Thailand	KY550386	KY568988
<i>Setophoma</i> sp.	CGMCC 3.19526	Carbonatite in a cave	China	MK511965	MK511944
<i>S. sacchari</i>	MFLUCC 12-0241	<i>Saccharum officinarum</i>	Thailand	KJ476147	KJ476145
	CBS 333.39 ^x		Brazil	MH867535	MH856038
<i>S. terrestris</i>	LC6449 ^x	<i>Allium sativum</i>	USA	KF251749	KF251246
	CBS 335.87	<i>Allium cepa</i>	Senegal	KF251750	KF251247
<i>S. terrestris</i>	CCLF01	<i>Solanum lycopersicum</i>	Mexico	MN596167	MN596166
<i>S. terrestris</i>	CCLF02	<i>S. lycopersicum</i>	Mexico	OL960402	OL960206
<i>S. terrestris</i>	CCLF03	<i>S. lycopersicum</i>	Mexico	OL960403	OL960207
<i>S. terrestris</i>	CCLF04	<i>S. lycopersicum</i>	Mexico	OL960404	OL960208
<i>S. terrestris</i>	CCLF05	<i>S. lycopersicum</i>	Mexico	OL960398	OL960202
<i>S. terrestris</i>	CCLF06	<i>S. lycopersicum</i>	Mexico	OL960399	OL960203
<i>S. terrestris</i>	CCLF07	<i>S. lycopersicum</i>	Mexico	OL960400	OL960204
<i>S. terrestris</i>	CCLF08	<i>S. lycopersicum</i>	Mexico	OL960401	OL960205
<i>S. vernoniae</i>	CBS 137988 ^x	<i>Vernonanthura</i> sp.	Brazil	KJ869198	KJ869141
<i>S. yingyisheniae</i>	LC12699	<i>C. sinensis</i>	China	MK511951	MK511915
	LC13479 ^x	<i>C. sinensis</i>	China	MT523031	MK511918
<i>S. yunnanensis</i>	LC6532	<i>C. sinensis</i>	China	MK511945	MK511907
	LC6753 ^x	<i>C. sinensis</i>	China	MK511949	MK511913

^xisolates ex-type. The accession numbers of the sequences generated in this study are in bold.

CBS: Collection of isolates in the Centraalbureau Voor Schimmelcultures, Fungal Biodiversity Centre, Utrecht, Netherlands; IMI: Collection of isolates in CABI Europe UK Centre, Egham, United Kingdom; CPC: Pedro W. Crous work collection, stored in CBS. CGMCC: General Center for the Collection of Microbiological Cultures of China, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China; LC: Lei Cai wok collection, stored in the Institute of Microbiology, Chinese Academy of Sciences, Beijing, China; MUCL: Mycothèque de l'Université catholique de Louvain, Louvain-la-Neuve, Belgium.

of the fungus per isolate were used, which were placed on a lesion made earlier with a toothpick. The mycelial disk was attached to the lesion of the root using Parafilm® paper (USA). In the control plants, the lesion was made and a disc with V8A culture medium was placed, without fungal growth. The plants were kept in a greenhouse at 20–35 °C for 30 días. The inoculated roots were then removed and the fungi were reisolates in an V8A medium. Once the mycelium was developed, the morphological and cultural characteristics were considered, such as pigmentation of the V8A medium, as well as the qualitative and quantitative characteristics of

mushrooms, pycnidia and conidia, according to earlier descriptions. The test was repeated twice.

RESULTS AND DISCUSSION

Symptoms of chlorosis and leaf senescence were observed, as well as the reduction in tomato plant growth (Figures 2A–C) in an incidence that varied from 10 to 20% in the five different commercial agricultural fields sampled in the valley of Culiacan, Sinaloa, Mexico. The symptomatic roots developed dark brown lesions, swollen and with a corky texture (Figure 2D–E) and pink (Figure 2F). These symptoms in the foliar area and root system coincide with those reported by López-López *et al.*



Figure 2. Symptoms of corky and pink roots in tomato. **A–C)** Chlorosis, deficient growth and senescence in plants. **D–F)** Dark brown lesions on the root, with swelling, a corky texture and a pink color.

(2020), whereas the same causal agent affected tomato plants reported in Brazil, Canada, USA and Venezuela (Farr and Rossman, 2022). Likewise, these symptoms were reported in other hosts such as scallions (*Allium fistulosum*) in China (Luong *et al.*, 2008). Similarly, the symptoms of the corky and pink root diseases coincide with those described for *S. terrestris* in onion (Schwartz and Mohan, 2008).

Pink root symptomatology may be similar to that caused by *Pseudopyrenochaeta lycopersici*, which was found in a wide number of hosts such as sorghum (*Sorghum*), maize (*Zea mays*), sugarcane (*Saccharum officinarum*) (Sprague, 1950), grass (*Cynodon dactylon*) (Connors, 1967), leek (*Allium porrum*) (Hall *et al.*, 2007), cantaloupe (*Cucumis melo*), zucchini (*Curcubita pepo*), wheat (*Triticum*) and spinach (*Spinacia oleracea*) (USDA, 1960), as well as willow trees (*Salix*) (Pfleger and Vaughn, 1972). On the other hand, in tomato, it has been reported in Australia (Golzar, 2009), Africa (Testen *et al.*, 2019), Germany (Valenzuela-López *et al.*, 2018) and in the United States as a pathogen associated to the brown rot in tomato root (Vrisman *et al.*, 2017).

Isolation and morphological identification. Out of the 15 samples (plants) of tomato roots with symptoms of corky and pink roots by sites, seven fungal isolates were used, which came from the five sites previously indicated on the map with their respective coordinates in the valley of Culiacan. The technique that worked for the isolation of fungi was to place sections of infected tissue in a humid chamber to stimulate the pycnidia in the root tissues at 4 °C (Gilchrist-Saavedra *et al.*, 2005). The isolates obtained correspond to *Setophoma terrestris*, and the colonies were pink on both sides, with a root growth (80 mm) after 7 days of growing (Figures 3A–B). These characteristics of growth of the fungus in V8A medium were similar to those reported by De Gruyter *et al.* (2010).

The morphological examination of the asexual structures displayed black to dark brown globular to obpyriform pycnidia, (Figure 3C) and measuring 142–220 × 104–140 μm. The presence of dark brown septated setae was observed, measuring 52–114 × 2–8 μm and that surrounded the neck ostilar (Figure 3D). The conidia were hyaline, cylindrical, measuring 2–8 × 1–3 μm, and with two very characteristic biguttulate on its tips (Figure 3E–F). All the morphological characteristics coincided with those reported by other authors (Koenning *et al.*, 2007; Wiriyajitsomboon, 2015; Yang *et al.*, 2017; Rivedal *et al.*, 2018; Zhang *et al.*, 2019) (Table 2). Despite some measurements being variable between each other, these data are not forceful to differentiate between species, therefore molecular tools and phylogenetic analyses are required to confirm at the level of species.

Regarding the ranges of the morphometric values, the seven *S. terrestris* of this study were distributed into three groups, according to upstream measures: group




Figure 3. Colonies and asexual reproductive structures of *Setophoma terrestris*. **A)** Colony of *S. terrestris* in a V&A medium with 7 days of growth. **B)** Colony growth on the reverse of the dish. **D)** Pycnidium. **D)** Setae. **E-F)** Biguttulate conidia.

Table 2. Measurements of asexual structures in *Setophoma terrestris* isolates obtained from tomato plants.

Setae (µm)	Pycnidium (µm)	Conidia (µm)	Reference
∅NA	∅NA	4.5–5.5 × 1.8–2.3	Koenning <i>et al.</i> , 2007
≈ 100–120	≈ 200	≈ 4–5	Wiriyajitsomboon, 2015
≈ 100	≈ 300	≈ 5	Yang <i>et al.</i> , 2017
∅NA	∅NA	5.0 × 2.0	Rivedal <i>et al.</i> , 2018
55–150	160–230	4.0–5.0 × 1.5–2.0	Zhang <i>et al.</i> , 2019
68–84 × 2–3	237–345 × 105–250	5–7 × 2–3	López-López <i>et al.</i> , 2020 (CCLF01)
52–114 × 3–6	150–183 × 133–156	6–7 × 2–3	² CCLF02
60–108 × 2–5	142–160 × 124–132	6–8 × 2–3	² CCLF03
46–102 × 3–5	200–233 × 110–130	4–6 × 2–3	² CCLF04
55–106 × 3–4	220–231 × 105–133	2–3 × 1–1.5	² CCLF05
67–90 × 2.5–4	216–255 × 104–140	3–4 × 1–2	² CCLF06
90–107 × 3–8	175–180 × 140–155	5–8 × 2–3	² CCLF07
60–101 × 3–5	160–175 × 137–153	5–6 × 2–3	² CCLF08

∅NA: Not applicable / Not reported.

²The measurements generated in this study are highlighted in bold in references.

1, composed of isolate CCLF05; group 2, by CCLF06 and CCLF04, and finally, group 3, with isolates CCLF07, CCLF08, CCLF02 and CCLF03. It is important to mention that group 3 also includes the isolate (CCLF01) reported previously in another study in Sinaloa, Mexico (López-López *et al.*, 2020).

In turn, Luong *et al.* (2008) identified *S. terrestris* as the causal agent of pink root in onion, using characteristics of the colony and they observed the pink pigment in the culture medium, as well as the setae in the pycnidia and conidia, typical of this root in some pink roots. Additionally, large guttulates were observed in the conidia of their isolates, which coincide with those reported in this study. These same microscopic characteristics in the conidia were described by De Gruyter and Boerema (2002).

Similarly, Ikeda *et al.* (2012) identified *S. terrestris* as the causal agent of pink root in zucchini and they obtained two isolates that formed brown to dark black pycnidia, subglobular to globular in shape and with a neck ostial with setae around it. These were found to be cylindrical in shape and brown to dark brown, and measured $104\text{--}262 \times 3.5\text{--}6.9 \mu\text{m}$. These characteristics coincide with those obtained in this study.

On the other hand, measurements may vary slightly between the isolates and the above may be due to multiple environmental factors that affect the development of the pathogen, as mentioned by Burton *et al.* (1997), making a relationship with the symptoms of the disease in cantaloupe and watermelon and identifying the causal agent based on morphological structures and analysis of ITS sequences. Unlike our isolates, the pycnidia reported by Duarte and Barreto (2015) do not have the typical setae around the ostiole, indicating that this attribute is not a generic characteristic. Likewise, the characteristics shared by the fungi *P. lycopersici* and *S. terrestris* are very similar to the difference that in *P. lycopersici* the setae of its pycnidia are over 200 μm long, whereas *S. terrestris* regularly range between 50–150 μm , approximately (De Gruyter, 2010).

Phylogenetic analysis. The aligned sequences corresponding to the ITS region and the long subunit (LSU) resulted in a total length of 1,345 nucleotides. The phylogenetic analysis generated with MrBayes based on the data set (Figure 4) indicated that all the isolations obtained in this study were grouped with the species *Setophoma terrestris*, with a high posterior probability support (1.0).

On the other hand, the phylogenetic analysis shows two groupings, which mostly addresses all isolates with 0.88 subsequent probability, except for isolate CCLF05, which was grouped separately and displayed the largest phylogenetic difference. Likewise, with the previously mentioned and according to the morphometry, was grouped in only one group. Based on this, it was possible to verify, once again, that *Pseudopyrenochaeta lycopersici* is not the species that causes the disease known as corky and pink root in Sinaloa, Mexico.




Figure 4. Bayesian Tree obtained with combined data from sequences ITS and LSU. The tree shows the phylogenetic relations of the *Setophoma* species. The Bayesian posterior probability values of over 0.70 are shown in the nodes. The species *Pseudopyrenochaeta lycopersici* was used as an external group and the scale bar indicates the number of expected changes per site.

The phylogenetic of this study coincides with the molecular identity reported by Rivedal *et al.* (2018), who used the ITS region, finding that the DNA of all its isolates displayed a 99% similarity with *Setophoma terrestris*. Likewise, those sequences were also compared with those reported by De Gruyter *et al.* (2010) and similar results were obtained. On the other hand, Duarte and Barreto (2015) used the same Bayesian inference analysis criterion, with the difference that they relied only on LSU sequences, whereas in this study, concatenated ITS and LSU

sequences were used. This is of great importance, since in recent years, it has been proven and supported that these genera are very closely related and can only be distinguished through phylogenetic analysis with concatenated sequences (De Gruyter *et al.*, 2010). This information is extremely important, since the *Phoma* genus has grown to encompass over 3000 species that form pycnidia with hyaline and septated conidia (Aveskamp *et al.*, 2008), thereby reevaluating several studies related to this genus upon correct identification (Boerema *et al.*, 2004).

Pathogenicity tests. Thirty days after inoculation, the leaves displayed yellowing (Figures 5A–B). The inoculated tomato roots displayed a corky texture and pink coloring (Figure 5C), which were similar to the symptoms observed on the field.




Figure 5. Pathogenicity tests in tomato plants. **A–B)** Tomato plants of the 8444 variety inoculated with *S. terrestris* and showing symptoms of yellowing. **C)** Corky root symptom 30 days after inoculation with *S. terrestris*. **D)** Colony obtained from the *in vitro* reisolation of *S. terrestris*.

From the reisolations, colonies (Figure 5D) and fungal structures identical to those of the inoculated fungus were obtained, which confirmed the pathogenicity of the seven inoculated isolates, while the roots of the control plants remained asymptomatic.

The results of the pathogenicity tests of this study coincide with those reported by Zhang *et al.* (2019), who observed typical pink root symptoms in garlic. This pathogen has also been found damaging other hosts such as canola (*Brassica napus*) in Canada (Yang *et al.*, 2017), zucchini (*Curcubita maxima*) in the USA (Rivedal *et al.*, 2018), and tea plants (*Camellia sinensis*) in China (Liu *et al.*, 2019).

The symptoms observed in the pathogenicity tests in the study conducted by Zhang *et al.* (2019) showed that *S. terrestris* caused chlorosis, wilting and death in garlic plants within a span of 40 days, similar to descriptions by López-López *et al.* (2020), which also coincides with the results in this study. This proves that *Pseudopyrenochaeta lycopersici* is not the causal agent of corky root in tomato fields in Sinaloa, Mexico, as it was considered for a long time.

Furthermore, *S. terrestris* was proven to be a slow-growing pathogen, since the pathogenicity tests by Rivedal *et al.* (2018) presented pink root symptoms 12 weeks after inoculation in winter squash, whereas in our tests it was 4 weeks, probably due to the pathogen-host relation being different. Similarly, Yang *et al.* (2017) used mycelial *S. terrestris* plugs on canola, and found symptoms of corky and pink root after 21 days, which coincides with the approximate time of this study. The time taken for the pathogen to colonize the plant tissue may be affected by the conditions it is in, i.e., after multiple reactivations in culture medium, its growth rate varies (de Gruyer and Borema, 2002), possibly thereby losing the original degree of virulence of each isolation, consequently delaying the visibility of disease symptoms under study.

CONCLUSIONS

According to the morphological characterization, molecular identification and pathogenicity tests, *Setophoma terrestris* was confirmed to be the causal agent of the disease known as corky and pink root in agricultural tomato fields distributed in Culiacan, Sinaloa. The morphological analysis showed that in *Setophoma terrestris* presented pycnidia measuring $142\text{--}220 \times 104\text{--}140 \mu\text{m}$, dark brown setae measuring $52\text{--}114 \times 2\text{--}8 \mu\text{m}$ and conidia measuring $2\text{--}8 \times 1\text{--}3 \mu\text{m}$. Likewise, three groups of isolates were obtained, in which variations of morphometric and phylogenetic values were observed. The symptoms presented by infected plants were corkiness, pink root and a reduction in root quantity.

LITERATURE CITED

- Allescher A. 1896. Diagnosen einiger neuer, im Jahre 1895 gesammelter Arten bayerischer Pilze, aus der Abteilung der Fungi imperfecti. Berichte der Bayerischen Botanischen Gesellschaft. 4: 31–40. <https://biotanz.landcareresearch.co.nz/references/ea66d275-f7ff-47c8-8c62-0b41c359618f>.
- Aveskamp MM, Gruyter J and Crous PW. 2008. Biology and recent developments in the systematics of *Phoma*, a complex genus of major quarantine significance. Fungal Divers 31: 1–18. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.516.3365&rep=rep1&type=pdf>
- Blancard D, Laterrot H, Marchoux G y Condresse T. 2011. Enfermedades del tomate: identificar, conocer y controlar. Ediciones Mundi-Prensa. Madrid – México. ISBN: 978-84-8476-427-4.
- Bruton BD, Biles CL and Duthie JA. 1997. Pink root of muskmelon and watermelon caused by *Phoma terrestris*. Subtropical Plant Science Journal. 49: 34–41. <http://www.subplantsci.org/wp-content/uploads/2016/02/SPSJ-49-34-41-Bruton-et-al.pdf>
- Boerema GH, Gruyter J de, Noordeloos ME, Hamers MEC. 2004. *Phoma* identification manual. Wallingford, UK: CABI Digital Library. ISBN: 0851997430, 9780851997438
- Castillo JA y Plata G. 2010. Diagnóstico de raíz rosada utilizando técnicas convencionales y moleculares. Revista de Agricultura. 62: 56-61 https://Diagnosis_of_pink_root_by_using_conventional_and_molecular_techniques
- Campbell R, Schweers V and Hall D. 1982. Corky root in California caused by *Pyrenochaeta lycopersici* and control by soil fumigation. Plant Disease 66: 657-661. <https://doi.org/10.1094/PD-66-657>
- Clerjeau M. 1973. Methode rapide de contamination des jeunes plantules de tomate par *Pyrenochaeta lycopersici* Schneider & Gerlach. Annales de Amelioration des Plantes 5 (2): 143-150.
- Connors IL. 1967. An Annotated Index of Plant Diseases in Canada and Fungi Recorded on Plants in Alaska, Canada and Greenland. Canada Department of Agriculture Publication. 1251: 1-381. urn:oclc:record:1039510843
- Davet P. 1973. Distribution et évolution du complexe parasitaire des racines de tomate dans une région du Liban où prédomine le *Pyrenochaeta lycopersici* Schneider et Gerlach. Annales de Phytopathologie 5: 53-63. <https://www.documentation.ird.fr/hor/fdi:06471>
- Gruyter J & Boerema GH. 2002. Contributions towards a monograph of *Phoma* (Coelomycetes) VIII. Section Paraphoma: Taxa with setose pycnidia. Persoonia - Molecular Phylogeny and Evolution of Fungi 17(4) 541–561. <https://repository.naturalis.nl/pub/532020>
- Gruyter J, Woudenberg JHC, Aveskamp MM, Verkley GJM, Groenewald JZ, Crous PW. 2010. Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. Mycotaxon 102: 1066-1081. <https://doi.org/10.3852/09-240>
- Duarte LL and Barreto RW. 2015. *Phoma canadensis* Allesch.: a synonym of *Pyrenochaeta cava*. Mycotaxon 130(3): 731-738. <https://doi.org/10.5248/130.731>
- Farr DF and Rossman AY. 2022. Funga Databases, Syst. Mycol. Microbiol. Lab., ARS, USDA. <https://nt.ars-grin.gov/fungaldatabases/>. Google Scholar.
- French A M. 1989. California plant disease host index. California Department of Food and Agriculture, Sacramento, 394. <http://www.cdfa.ca.gov/plant/>.
- Gilchrist-Saavedra LG, Fuentes-Dávila C, Martínez-Cano RM, López-Atilano E, Duveiller RP, Singh MH y García A. 2005. Guía práctica para la identificación de algunas enfermedades de trigo y cebada. Segunda edición. México, D.F.: CIMMYT. <http://hdl.handle.net/10883/3217>
- Golzar H. 2009. First report of *Pyrenochaeta lycopersici*, causal agent of tomato corky root rot in Australia. Australasian Plant Disease Notes 4: 126-128. <https://doi.org/10.1071/DN09051>
- Hall BH, Hitch CJ, Oxspring EA and Wicks TJ. 2007. Leek diseases in Australia. Australas. Plant Pathology 36: 383-388. <https://doi.org/10.1071/AP07038>
- Ikeda K, Kuwabara K, Urushibara T, Soyai P, Miki S & Shibata S. 2012. Pink root rot of squash caused by *Setophoma terrestris* in Japan. Journal of General Plant Pathology 78(5): 372–375. <https://doi.org/10.1007/s10327-012-0398-2>
- Jones JB, Zitter TA, Momol TM and Miller SA. 2016. Compendium of tomato diseases and pests. Second edition. American Press Society. ISBN: 978-0-89054-434-1. Pp. 15-119. <https://doi.org/10.1094/9780890544341>

- Liu F, Wang J, Li H, Wang W and Cai L. 2019. *Setophoma* spp. on *Camellia sinensis*. Fungal Systematics and Evolution 4: 43–57. <https://doi.org/10.3114/fuse.2019.04.05>
- López-López AM, León-Félix J, Allende-Molar R, Lima NB, Tovar-Pedraza JM and García-Estrada RS. 2020. First report of *Setophoma terrestris* causing corky and pink root of tomato in Sinaloa, Mexico. Plant Disease 104 (5): 1553 <https://doi.org/10.1094/PDIS-10-19-2226-PDN>.
- Luong TM, Huynh LMT, Tran U, Dau VT, Burgess LW, Phan HT, Aveskamp MM and Vo AD. 2008. First report of *Phoma terrestris* causing pink root rot of Chinese onion in Vietnam. Australasian Plant Disease Notes 3: 147-149. <https://doi.org/10.1007/BF03211273>
- Martínez-Ruiz FE, Cervantes-Díaz L, Aíl-Catzím CE, Hernández-Montiel LG, Del Toro Sánchez CL y Rueda-Puente EO. 2016. Hongos fitopatógenos asociados al tomate (*Solanum lycopersicum* L.) en la zona árida del noroeste de México: la importancia de su diagnóstico. European Scientific Journal 12(18): 1857–7881. <https://doi.org/10.19044/esj.2016.v12n18p232>
- Mori Y, Sato Y & Takamatsu S. 2000. Evolutionary analysis of the powdery mildew fungi using nucleotide sequences of the nuclear ribosomal DNA. Mycologia 92(1): 74-93. <https://doi.org/10.1080/00275514.2000.12061132>
- Nylander J. 2004. MrModeltest v2. Program distributed by the author. Bioinformatics 24. 581-583. <https://www.researchgate.net/publication/285805344>
- Olavarria L. 1991. Caracterización e identificación del agente causal de la enfermedad raíz corchosa en tomate (*Lycopersicon esculentum* Mill). Taller de licenciatura Ing. Agr. Quillota, Universidad Católica de Valparaíso, Facultad de Agronomía. 59. <http://repositorio.ucv.cl/handle/10.4151/78293>
- Page RD. 1996. TREEVIEW: An application to display phylogenetic trees on personal computers. Computer Applications in the Biosciences 12 (4): 357–358. <https://doi.org/10.1093/bioinformatics/12.4.357>
- Pfleger FL and Vaughn EK. 1972. *Pyrenochaeta terrestris* as a naturally occurring component of the soil in Oregon. Plant Disease Reporter 56: 180-182. [sim_plant-disease_1972-01_56_1](https://doi.org/10.1094/PD-1972-01_56_1).
- Quaedvlieg W, Verkley GJM, Shin H-D, Barreto RW, Alfenas AC, Swart WJ, Groenewald JZ and Crous PW. 2013. Sizing up Septoria. Studies in Mycology 75: 307–390. <http://dx.doi.org/10.3114/sim0017>.
- Rivedal HM, Stone AG and Johnson KB. 2018. First report of *Setophoma terrestris* causing pink root rot of winter squash (*Cucurbita maxima*) in Oregon. Plant Disease 102 (12) : 2661. <https://doi.org/10.1094/PDIS-06-18-0921-PDN>
- Rivedal HM, Tabima JF, Stone AG and Johnson KB. 2022. Identity and pathogenicity of fungi associated with root, crown, and vascular symptoms related to winter squash yield decline. Plant disease 106(6):1660-1668. <https://doi.org/10.1094/PDIS-09-20-2090-RE>
- Ronquist F, Teslenko M, Van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA and Huelsenbeck JP. 2012. MrBayes v. 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology. 61 (3): 539–542. <https://doi.org/10.1093/sysbio/sys029>.
- Schneider R and Gerlach W. 1966. *Pyrenochaeta lycopersici* nov. spec., der Erreger der Korkwurzelkrankheit der Tomate. Journal of Phytopathology. 56 (2): 117–122. <https://doi.org/10.1111/j.1439-0434.1966.tb02251.x>
- Schwartz HF and Mohan SK. 2008. Compendium of onion and garlic diseases. 2nd ed. American Phytopathological Society Press: St Paul, MN. <https://doi.org/10.1094/9780890545003>
- SIAP. 2020. <https://www.gob.mx/siap/acciones-y-programas/produccion-agricola-33119>. Fecha de consulta: junio del 2020.
- Sprague R. 1950. Diseases of Cereals and Grasses in North America. Agronomy Journal 43(3): 152. <https://doi.org/10.2134/agronj1951.00021962004300030013x>
- Staden R, Beal KF and Bonfiel JK. 2000. The Staden package, 1998. Bioinformatics Methods and Protocols. Methods in Molecular Biology 132: 115-130. <https://doi.org/10.1385/1-59259-192-2:115>
- Takamatsu S and Kano Y. 2001. PCR primers useful for nucleotide sequencing of rDNA of the powdery mildew fungi. Mycoscience 42 (1): 135-139. <http://doi.org/10.1007/BF0246387>
- Tamura K, Stecher G and Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution 38 (7): 3022-3027. <https://doi.org/10.1093/molbev/msab120>
- Testen AL, Chala A, Azerefege F and Miller SA. 2019. First report of corky root rot of tomato caused by *Pyrenochaeta lycopersici* in Africa. Plant Disease 103(5): 1032. <https://doi.org/10.1094/PDIS-10-18-1887-PDN>

USDA. 1960. Index of Plant Diseases in the United States. Agric. Handb. 165: 1-531. (94)

USDA. 2017. Fecha de consulta: mayo del 2017. <https://nt.ars-grin.gov/fungaldatabases/>

Valenzuela-Lopez N, Cano-Lira JF, Guarro J, Sutton DA, Wiederhold N, Crous PW and Stchigel AM. 2018. Coelomycetous Dothideomycetes with emphasis on the families Cucurbitariaceae and Didymellaceae. *Studies in mycology* 90(1): 1–69. <https://doi.org/10.1016/j.simyco.2017.11.003>

Vrisman CM, Testen AL, Elahi F and Miller SA. 2017. First report of tomato brown root rot complex caused by *Colletotrichum coccodes* and *Pyrenochaeta lycopersici* in Ohio. *Plant Disease* 101(1): 247. <https://doi.org/10.1094/PDIS-05-16-0623-PDN>

Yang Y, Zuzak K, Harding M, Neilson E, Feindel D and Feng J. 2017. First report of pink root rot caused by *Setophoma (Pyrenochaeta) terrestris* on canola. *Canadian Journal of Plant Pathology* 39(3): 354-360. <https://doi.org/10.1080/07060661.2017.1355849>

Zhang FB, Zheng HL, Zhang MQ, Yin YS, Cui M and Gao M. 2019. First report of *Setophoma terrestris* causing pink root of garlic in China. *Plant Disease* 103(3): 584 <https://doi.org/10.1094/PDIS-03-18-0530-PDN>