

Retinal capillary hemangioma

Hemangioma capilar retiniano

Jorge A. Sánchez-Ramos^{1*} and Juan A. Ramírez-Estudillo²

¹MD, Retina and Vitreous Department; ²Head of the Retina and Vitreous Department. Fundación Hospital Nuestra Señora de la Luz, Mexico City, Mexico

Fifty-year-old female, with a family history of unspecified tumors. She began her condition with a 2-month history of visual loss and metamorphopsia in the left eye, finding multiple vascular lesions in the retina, so the study protocol for Von-Hippel-Lindau disease was started (Fig. 1).

Funding

The authors received no specific funding for this work.

Conflicts of interest

The authors declare no conflicts of interest.



Figure 1. Photographic composition of the left eye fundus showing multiple vascular lesions with significant exudation in some of them with partial involvement of the fovea.

Correspondence:

*Jorge Arturo Sánchez-Ramos

E-mail: jorgesr84@hotmail.com

2604-1731/© 2019 Sociedad Mexicana de Oftalmología. Published by Permanyer México SA de CV. This is an Open Access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 12-01-2017

Date of acceptance: 20-01-2017

DOI: 10.24875/RMOE.M19000055

Available online: 01-03-2019

Rev Mex Oftalmol (Eng). 2019;93(2):95-95

www.rmo.com.mx