

Revision of moles in the genus *Scapanus*

SERGIO TICUL ÁLVAREZ-CASTAÑEDA¹*, AND PATRICIA CORTES-CALVA¹

¹ Centro de Investigaciones Biológicas del Noroeste, Instituto Politécnico Nacional 195, CP. 23096. La Paz, Baja California Sur, México. Email: sticul@cibnor.mx (STA-C), pcortes04@cibnor.mx (PCC).

*Corresponding author

Scapanus latimanus is a species with many morphological differences among its populations. This variation is associated with multiple taxonomic changes at the species or subspecies level. This study incorporates genetic analyses and comparisons with previous morphological studies to propose a better understanding of the *latimanus* complex. Mitochondrial markers (cytochrome *b*; cytochrome *c* oxidase subunit I; and cytochrome *c* oxidase subunit III) were sequenced to construct a phylogeny for the subfamily Scalopinae in North America. Genetic distances ranged from 2.49 to 10.50 % among geographic areas. Results identified three monophyletic clades with high bootstrap support values. Based on our phylogenetic analysis and previous morphological analyses, we confirm *S. anthonyi* from San Pedro Martir as a valid species and propose that *S. occultus* from southern California and northern Baja California peninsula be considered as a species.

Scapanus latimanus es una especie con muchas diferencias morfológicas entre sus poblaciones. Esta variación está asociada con múltiples cambios taxonómicos a nivel de especie o subespecie. Para proponer una mejor comprensión del complejo *latimanus*, en este estudio se incorpora la información genética a los estudios previos de morfología. Se secuenciaron genes de origen mitocondrial (citocromo *b*; citocromo *c* oxidasa subunidad I y III) para construir la filogenia para la subfamilia Scalopinae en Norteamérica. Se obtuvieron distancias genéticas con un intervalo entre 2.49 a 10.50 % entre áreas geográficas. Los resultados identifican tres clados monofiléticos con altos valores de soporte. Con base en la comparación del análisis filogenético e información morfológica previa conocida, se confirma como una especie válida a *S. anthonyi* de San Pedro Martir y proponemos que *S. occultus* del sur de California y norte de la península de Baja California también sea considerada como una especie.

Keywords: Molecular markers; moles; North America; phylogeny; Scalopinae; Talpidae; taxonomy.

© 2021 Asociación Mexicana de Mastozoología, www.mastozoologiamexicana.org

Introduction

The family Talpidae includes three subfamilies, Scalopinae, Talpinae, and Uropsilinae, with Scalopinae being restricted to America and containing four genera *Condylura*, *Parascalops*, *Scalopus*, and *Scapanus* ([Shinohara et al. 2003](#); [Hutterer 2005](#)). *Scapanus* is the only genus including more than one species; *S. latimanus*, *S. orarius*, and *S. townsendii* ([Hutterer 2005](#)). A fourth species, *S. anthonyi*, has been considered, although it has undergone many taxonomic changes. *S. anthonyi* was described as a full species by [Allen \(1893\)](#), and later was considered to be a subspecies of *S. latimanus* ([Palmer 1937](#)). [Palmer \(1937\)](#) argued that morphometric characteristics of *S. anthonyi*, such as its smaller size and fewer number of upper premolars, also were present in *S. l. occultus*, and consequently, *S. anthonyi* should be considered a subspecies of *S. latimanus* (see: [Palmer 1937](#); [Hutchinson 1987](#)).

In his review of American moles, [Jackson \(1915\)](#) recognized *S. anthonyi* as a species because *S. anthonyi* has a projection in the braincase between the interparietal and the mastoid, which was absent in *S. l. occultus* ([Jackson 1915](#)). However, [Palmer \(1937\)](#) did not acknowledge these characteristics in the specimens that he examined, and therefore did not consider *S. anthonyi* a valid species. [Huey \(1936\)](#) suggested an additional difference between *S. l. occultus* and *S. anthonyi*; specifically, the manus (part of the pentadactyl limb that includes the metacarpals and phalanges)

in *S. anthonyi* is squarer and smaller, with broader and heavier phalanges and with tips of the pterygoids parallel. In an alternative view, [Hutchinson \(1987\)](#) suggested that *S. anthonyi* shared characteristics with *S. orarius*; however, he continued to recognize *S. anthonyi* as a subspecies of *S. latimanus*. Populations of *S. anthonyi* in San Pedro Martir, Baja California and those of *S. l. grinnelli* and *S. l. occultus* in southern California and northern Baja California peninsula are smaller in size and the skull is wider in relation to all other subspecies of *S. latimanus* from central and northern California ([Yates and Salazar-Bravo 2005](#)). Differences in skull morphology also occur between these two groups. *S. anthonyi* has only two or three upper premolars, and the temporal fossae is larger ([Allen 1893](#); [Jackson 1915](#); [Huey 1936](#); [Yates and Salazar-Bravo 2005](#)).

Previous morphological analyses of all subspecies of *S. latimanus* indicated that some subspecies should be junior synonyms ([Yates and Salazar-Bravo 2005](#)) of other subspecies. For example, *S. l. grinnelli* ([Jackson 1914](#)) of *S. l. occultus* ([Grinnell and Storer 1916](#)); *S. dilatus* ([True 1894](#)), *S. alpinus* ([Merriam 1897](#)), and *S. l. caurinus* ([Palmer 1937](#)) of *S. l. latimanus* ([Bachman 1842](#)); *S. l. sericatus* ([Jackson 1914](#)), *S. l. campi* ([Grinnell and Storer 1916](#)), and *S. l. monoensis* ([Grinnell 1918](#)) of *S. l. minusculus* ([Bangs 1899](#)). However, *S. l. insularis* ([Palmer 1937](#)) and *S. l. parvus* ([Palmer 1937](#)) were not subjected to taxonomic changes ([Yates and Salazar-Bravo 2005](#)).

Based on these previous studies, the taxonomic status of species within the *S. latimanus* group has revealed several inconsistencies. The goal of this study is to better define the phylogenetic relationships of populations within *Scapanus* and combine these relationships with known morphological characteristics to evaluate the potential number of species. To achieve this goal, three mitochondrial genes were sequenced: cytochrome *b* (*Cytb*; $n = 23$); cytochrome *c* oxidase subunit I (*Co1*, $n = 29$); and cytochrome *c* oxidase subunit III (*Co3*; $n = 29$).

Materials and Methods

Sample collection. The dataset included specimens of the genus *Scapanus* ($n = 31$) represented by the species *S. orarius*, *S. townsendi*, *S. latimanus*, and outgroup specimens of *Condylura*, *Neurotrichus*, and *Scalopus* ($n = 6$). Tissue samples were obtained from the Collection of Mammalian tissues at Centro de Investigaciones Biológicas del Noroeste (CIB), Field Museum of Natural History (FMNH), Museum of Southwestern Biology at the University of New Mexico (MBS), and Museum of Vertebrate Zoology at the University of California (MVZ). Information for localities and museum catalog numbers are provided in Table 1. All capture and handling methods followed the animal care and use guidelines of the American Society of Mammalogists ([Sikes et al. 2016](#)). For all analyses, we grouped specimens from these localities into three species, *S. orarius*, *S. townsendi*, and *S. latimanus*, with *S. latimanus* further subdivided into three geographic units: 1) central and northern California (Group A, localities 7-14); 2) southern California and northern Baja California peninsula (Group B, localities 15-17); and 3) Sierra de San Pedro Martir (Group C, locality 18; Figure 1; Table 1). This resulted in 31 geographic samples of Scalopinae.

DNA extraction and PCR conditions. Genomic DNA was extracted from muscle tissue preserved in 95 % ethanol (archived at -20°C) or frozen (archived at -80°C) using the DNAeasy Kit (QIAGEN Inc., Valencia, CA) protocols.

The following conditions were used for the initial double-strand amplification: 12.5 μl of (10 ng) template, 4.4 μl ddH₂O, 2.5 μl of each primer pair (10 nM concentration), 0.474 μl (0.4 nM) dNTPs, 0.5 μl (3 mM) MgCl₂, 0.125 μl Taq polymerase (platinum, Invitrogen, Carlsbad, CA), and 1× Taq buffer, to a final volume of 25 μl . The amplification conditions consisted of an initial denaturation at 94°C for 3 min followed by 37 denaturation cycles at 94°C for 45 s each; 60 s annealing at 50°C (*Cytb*), 51°C (*Co1*), 55°C (*Co3*); and extension at 72°C for 60 s; the products of the PCR amplification were verified in agarose gel, purified and sequenced both ways using the sequencing service of Macrogen Inc, Korea. The first part of the cytochrome *b* (*Cytb*, ~800 bp) gene was amplified using the primers MVZ05/MVZ16 (primer sequences given in [Smith and Patton 1993](#); [Smith 1998](#)), the 658-bp fragment of cytochrome *c* oxidase subunit I (*Co1*) was amplified with the primers LCO1490/HCO2198 ([Ivanova et al. 2007](#)), and the 717-bp fragment of cytochrome *c* oxidase subunit III (*Co3*) was amplified with

the primers L8618/H9323 ([Riddle 1995](#)). We aligned nucleotide sequences in Sequencher ver. 3.1 (Gene Codes Corp., Ann Arbor, Michigan), verified alignments visually, and translated them into amino acids for alignment confirmation. The haplotypes generated and used were deposited in GenBank (Table 1).




Figure 1. Distribution map of the species of the subfamily Scalopinae in North America. *Scapanus townsendi* (solid squares 1-3), *S. orarius* (open squares 4-6), and *S. latimanus* (circles). *S. latimanus* is split into three geographic groups: A) Central and northern California (localities 7-14, light gray circle); B) Southern California and northern Baja California peninsula (localities 15-17, half light/half dark circles); and C) Sierra de San Pedro Martir (locality 18, dark gray circle).

Table 1. List of specimens examined, locations according to Figure 1. #Catalog = museum catalog number of the reference collection. GenBank accession number for mitochondrial marker. Group (Gr), Number of map (M), State (ST), * Zhao and Jian 2015. ** Mouchaty *et al.* 2000.

Gr	M	Species	#Catalog	St	Locality	Lat	Long	Co1	Co3	Cytb					
						GenBank accession numbers									
1		<i>Scapanus t. olympicus</i>	MSB 43550	WA	9.2 Mi S, 2.7 Mi W Port Angeles	47.9851	-123.4878	MZ150455	MZ217155	MZ217129					
1		<i>Scapanus t. olympicus</i>	MSB 43552	WA	9.2 Mi S, 2.7 Mi W Port Angeles	47.9851	-123.4878	MZ150456	MZ217156	MZ217130					
2		<i>Scapanus t. townsendii</i>	MVZ 220251	WA	24303 Se 468th Street, Enumclaw	47.1809	-122.0175	MZ150457	MZ217157						
2		<i>Scapanus t. townsendii</i>	MVZ 220252	WA	24303 Se 468th Street, Enumclaw	47.1809	-122.0175	MZ150458	MZ217158						
4		<i>Scapanus t. townsendii</i>	MSB 40780	OR	9 Mi E Alsea	44.3817	-123.4135	MZ150459	MZ217159	MZ217131					
4		<i>Scapanus t. townsendii</i>	MSB 40781	OR	9 Mi E Alsea	44.3817	-123.4135	MZ150460	MZ217160	MZ217132					
3		<i>Scapanus o. schefferi</i>	MSB 54620	WA	2 Mi W Walla Walla	46.0647	-118.3835	MZ150461	MZ217161	MZ217133					
3		<i>Scapanus o. schefferi</i>	MSB 54621	WA	Country Club, Walla Walla	46.0389	-118.3503	MZ150462	MZ217162	MZ217134					
5		<i>Scapanus o. orarius</i>	MSB 43626	CA	3.8 Mi S, 2.7 Mi E Trinidad	41.004	-124.0916		MZ217163	MZ217135					
5		<i>Scapanus o. orarius</i>	MSB 43627	CA	3.8 Mi S, 2.7 Mi E Trinidad; T7n, R1e, Sec 8	41.004	-124.0916	MZ150463	MZ217164	MZ217136					
5		<i>Scapanus o. orarius</i>	MSB 43628	CA	3.8 Mi S, 2.7 Mi E Trinidad; T7n, R1e, Sec 8	41.004	-124.0916	MZ150464							
7		<i>Scapanus o. orarius</i>	MVZ 224399	CA	11 Mi N Westport On Hwy 1.	39.7506	-123.819	MZ150465	MZ217165						
A	6	<i>Scapanus l. dilatus</i>	MVZ 217713	CA	Eagle Lake Road (Lassen Co. A1), Eagle Lake.	40.6235	-120.8399	MZ150466	MZ217166	MZ217137					
	8	<i>Scapanus l. dilatus</i>	MSB 47919	CA	1 Mi S, 4.5 Mi E Somerset, 2850	38.6334	-120.5984	MZ150467	MZ217167	MZ217138					
	9	<i>Scapanus l. caurinus</i>	MVZ 216930	CA	Easy Sweet Farm, Sebastopol	38.472	-122.8544	MZ150468	MZ217168						
	10	<i>Scapanus l. caurinus</i>	MVZ 199506	CA	2930 Redwood Road, Napa	38.3167	-122.3385	MZ150469	MZ217169	MZ217139					
	11	<i>Scapanus l. latimanus</i>	MVZ 218027	CA	103 Aldarado Rd., Berkeley	37.8579	-122.2396	MZ150470							
	12	<i>Scapanus l. latimanus</i>	MVZ 201320	CA	Forest S of Chapel, Yosemite Valley	37.7408	-119.5907		MZ217170						
	13	<i>Scapanus l. latimanus</i>	MSB 48532	CA	Palo Alto, Stanford University Campus	37.429	-122.1695	MZ150471	MZ217171	MZ217140					
	14	<i>Scapanus l. latimanus</i>	MVZ 222251	CA	Hastings Natural History Reservation	36.3785	-121.5568	MZ150472	MZ217172						
	14	<i>Scapanus l. latimanus</i>	MVZ 228295	CA	Haystack Hill, Hastings Natural History Reservation	36.3847	-121.5627	MZ150473	MZ217173	MZ217141					
B	15	<i>Scapanus l. occultus</i>	MSB 47311	CA	10 Mi Se Big Bear City, Heart Bar campground	34.1586	-116.786	MZ150474	MZ217174	MZ217142					
	16	<i>Scapanus l. occultus</i>	MSB 47317	CA	3.6 Mi N, 9.8 Mi E Hemet, Lake Fulmor	33.8052	-116.7785	MZ150475	MZ217175	MZ217143					
	17	<i>Scapanus l. occultus</i>	MSB 43120	BC	Laguna Hanson	32.0489	-115.9056	MZ150476	MZ217176	MZ217144					
	17	<i>Scapanus l. occultus</i>	MSB 40343	BC	Laguna Hanson	32.0489	-115.9056	MZ150477	MZ217177	MZ217145					
	17	<i>Scapanus l. occultus</i>	MSB 40344	BC	Laguna Hanson	32.0489	-115.9056	MZ150478	MZ217178	MZ217146					
	17	<i>Scapanus l. occultus</i>	MSB 40345	BC	Laguna Hanson	32.0489	-115.9056	MZ150479	MZ217179	MZ217147					
	17	<i>Scapanus l. occultus</i>	MSB 47308	BC	Sierra Juárez, Laguna Hanson	32.0489	-115.9056	MZ150480	MZ217180	MZ217148					
C	18	<i>Scapanus l. anthonyi</i>	MSB 47306	BC	Sierra San Pedro Mártir, 3.9 Mi by Road W Vallecitos	31.0167	-115.5333	MZ150481	MZ217181	MZ217149					
	18	<i>Scapanus l. anthonyi</i>	CIB 32000	BC	Sierra San Pedro Mártir	31.0167	-115.5333	MZ150482	MZ217182	MZ217150					
	18	<i>Scapanus l. anthonyi</i>	MSB 47307	BC	Sierra San Pedro Mártir, 20 Mi S, 10.9 Mi E Vallecitos	31.0167	-115.5333	MZ150483	MZ217183	MZ217151					
Out-group		<i>Condylura cristata</i>	KU144678 *						KU144678						
		<i>Condylura cristata</i>	NC029762 *						NC_029762						
		<i>Neurotrichus g. hyacinthinus</i>	MVZ 200061	CA	Headwaters of Big Austin Creek, N of Cazadero	38.6138	-123.1315	MZ150484	MZ217184	MZ217152					
		<i>Scalopus a. macrurus</i>	FMNH 167212	MI	Fennville	42.5939	-86.1017	MZ150485	MZ217185	MZ217153					
		<i>Scalopus a. macrurus</i>	FMNH 167213	MI	Fennville	42.5939	-86.1017	MZ150486	MZ217186	MZ217154					
		<i>Talpa europaea</i>	Y19192 **						Y19192						

Phylogenetics analysis. The methodology for phylogenetic analysis was similar to that used by [Camargo and Álvarez-Castañeda \(2020\)](#). The most appropriate substitution model for the dataset for each of the three gene regions, as well as for the concatenated series, was determined using

the Akaike information criterion (AIC) as implemented in MrAIC ([Nylander 2004](#)). Four separate Bayesian inference and maximum-likelihood analyses were conducted on the three genes independently; the concatenated series had three partitions with one per gene (Cytb, Co1, and Co3).

Bayesian analyses were implemented in (MrBayes ver. 3.0b4; [Huelsenbeck and Ronquist 2001](#)) with four separate runs with Markov-chain Monte Carlo simulations starting from a random tree. Each run was conducted for 20 million generations and sampled at intervals of 1,000 generations. Of the samples trees, the first 50 % were discarded as burn-in and all remaining trees were analyzed to find the posterior probability of resulting nodes. A consensus tree was generated with the 50 % majority-rule algorithm in PAUP 4.0b10 ([Swofford 2002](#)). The percentage of samples recovered in a particular clade was assumed to be the posterior probability of that clade in PAUP 4.0b10 using a heuristic search with 1,000 replicates and swapping with the TBR algorithm.

Maximum-likelihood (ML) analyses were performed in PAUP ver. 4.0b10 ([Swofford 2002](#)) algorithm ([Felsenstein 1981](#)) using a heuristic search with 1,000 replicates and swapping with the TBR algorithm. Reliability was assessed using each of the three codon positions individually while applying equal weights and nodal support using nonparametric bootstrapping. Members of each genus were used because although some phylogenetic analyses were done using allozymes ([Yates and Greenbaum 1982](#); [Moore 1986](#)) the phylogenetic relationships among moles of North America were not previously examined using gene sequencing. Trees were rooted with Scalopini (*Scalopus aquaticus*), Urotrichini (*Neurotrichus gibbsii*), and Condylurini (*Condylura cristata*; [Motokawa 2004](#)).

Results

Phylogenetic analyses. AIC tests revealed that the best evolution model was a GTR model: *Cytb* (GTR + I + G), *Co1* (GTR + I), *Co3* (GTR + G), and the concatenated genes (GTR + I + G). BI and ML trees for *Cytb*, *Co1*, and *Co3*, and the concatenated data with four partitions converged on an essentially identical topology (Figure 2).

Analyses of the three genes within *Scapanus* resolved five haplogroups with strong bootstrap support (>95 %), as follows. Haplogroup 1: only specimens from San Pedro Martir, Group C of *S. latimanus*; Haplogroup 2: specimens from southern California and northern Baja California peninsula, Group B of *S. latimanus*; Haplogroup 3: all *S. latimanus* specimens from Group A of central and northern California; Haplogroup 4: specimens of the two subspecies of *S. townsendi* with a very low percentage of differences between them; and Haplogroup 5: containing two groups, each with specimens of different subspecies of *S. orarius* (Figure 2).

Scapanus latimanus Group C (Haplogroup 1) is separated from the other *S. latimanus* Groups A and B by two different species, *S. townsendi* (Haplogroup 4) and *S. orarius* (Haplogroup 5). The percentage of pairwise genetic differences (p-distance) for the three genes between Group C (San Pedro Martir) and each of Group A (northern California) and Group B (southern California and northern Baja California) ranged from 7.22 to 10.50 %. The genetic

differences between Group A and Group B ranged from 2.49 to 5.75 % (Table 2).

Discussion

Genetic data revealed that *S. townsendi* and *S. orarius* are monophyletic and sibling taxa, as reported by [Shinohara et al. \(2003\)](#), and are substantially different from *S. latimanus*, as previously reported by [Moore \(1986\)](#). However, the geographic groups of *S. latimanus* do not exhibit a north-south phylogenetic relationship. The *S. latimanus* Group C from San Pedro Martir formed an inconsistent relationship with the other two *S. latimanus* haplogroups from California and the northern Baja California peninsula (Haplogroups 2 and 3). Genes *Co3* (boot = 56) and *Cytb* (boot = 68) show *S. latimanus* Group C (San Pedro Martir) as a sister group to specimens from southern California and Baja California. However, analyses of *Co1* (boot = 95) and the concatenated group (boot = 95) show *S. latimanus* Group C basal to all *Scapanus* clades (boot = 95), including *S. townsendi* and *S. orarius* (Figure 2). Each of the topologies show that *S. latimanus* Group C differs from Groups A and B. Although [Hutchinson \(1987\)](#) reported that *S. anthonyi* shared characteristics with *S. orarius*, this was not supported by the sequence data.




Figure 2. Bayesian tree constructed from three mitochondrial DNA genes (cytochrome *b*, cytochrome oxidase subunit I, and cytochrome oxidase subunit III) from members of the subfamily Scalopinae in North America and one of *Talpa* from Europe. Haplogroup 1 contains a single specimen from San Pedro Martir, Group C of *S. latimanus* (*S. anthonyi*). Haplogroup 2 contains specimens from southern California and northern the Baja California peninsula, Group B of *S. latimanus* (*S. occultus*). Haplogroup 3 contains specimens from central and northern California, Group A of *S. latimanus* (*S. latimanus*). Haplogroup 4 is represented by two subspecies of *S. townsendi*. Haplogroup 5 is represented by two subspecies of *S. orarius*.

We did not perform a morphometric analysis of the craniodental measurements because this previously was reported by [Yates and Salazar-Bravo \(2005\)](#). [Yates and Salazar-Bravo \(2005\)](#) reported statistical differentiation in morphological characters between *S. l. occultus* and *S. anthonyi* separated by a distance > 50 km. In addition, they found significant differences between *S. l. occultus* and *S. l. latimanus* *sensus* ([Yates and Salazar-Bravo 2005](#)), with *S. l. occultus* being smaller overall. No specimens of *S. l. occultus* and *S. l. latimanus* have been collected in sympatry and in the two areas where *S. l. occultus* and *S. l. latimanus* occur, namely the southern part of Sierra Nevada and northern portion of Santa Barbara, it appears that *S. l. occultus* occurs in lower altitudes, and *S. l. latimanus* at higher altitudes.

Based on the genetic distance values between Groups A, B, and C, coupled with the morphological differences between them ([Yates and Salazar-Bravo 2005](#)), these

groups can be considered as different species. The main morphological variations in the specimens of these groups are a smaller size in relation to the northern populations of *S. latimanus* and the variation in the number of upper premolars ([Palmer 1937; Yates and Salazar-Bravo 2005](#)).

Further, based on the sequence data, *Scapanus latimanus* from northern and southern California form two haplogroups. Haplogroup 3 includes all the specimens assigned to *S. latimanus* Group A (northern California) and Haplogroup 2 includes Group B (southern California and north Baja California peninsula). The population from San Pedro Martir previously was considered as a distinct species, *S. anthonyi* ([Allen 1893; Jackson 1915; Huey 1936; Yates and Salazar-Bravo 2005](#)) and later subsumed into *S. latimanus* based on morphological characters (although a large series of specimens was never reviewed, which may have biased the interpretation), based primarily on the smaller size and number of upper premolars ([Palmer 1937; Hutterer 2005](#)). The morphological analyses ([Yates and Salazar-Bravo 2005](#)) and genetic analyses performed in this study support the consideration of *S. anthonyi* as a distinct species and indicates that *S. anthonyi* is restricted in distribution to the San Pedro Martir mountain range.

Based on our phylogenetic analysis and its morphological characteristics ([Allen 1893; Jackson 1915; Huey 1936; Yates and Salazar-Bravo 2005](#)), we support that *S. anthonyi* is a different species from *S. latimanus*. Additionally, we propose that specimens known as *S. latimanus occultus* (including *S. l. grinnelli*) from southern California and northern Baja California peninsula should be considered as a distinct species (*S. occultus*) different from *S. latimanus* from central and north California and from *S. anthonyi* inhabiting San Pedro Martir. Therefore, we consider that the genus *Scapanus* contains five species that should be recognized as *S. anthonyi*, *S. latimanus*, *S. occultus*, *S. orarius*, and *S. townsendi*.

	Group A	Group B	Group C
<i>S. latimanus</i> Group A (central and northern California)			
Cytb	0.13-1.96	4.13-5.75	8.00-9.67
Co1	0.00-1.22	2.59-3.65	9.59-10.50
Co3	0.15-0.61	2.49-3.60	7.89-8.58
Concatenated	0.24-1.23	3.27-4.17	8.53-8.95
<i>S. latimanus</i> Group B (southern California and northern Baja California)			
Cytb	4.13-5.75	0.0-1.50	7.88-9.52
Co1	2.59-3.65	0.00-0.90	8.68-9.98
Co3	2.49-3.60	0.0-0.92	7.22-7.55
Concatenated	3.27-4.17	0.00-0.99	7.82-8.01
<i>S. latimanus</i> Group C (San Pedro Martir)			
Cytb	8.00-9.67	7.88-9.52	0.00-1.75
Co1	9.59-10.50	8.68-8.98	0.00-0.15
Co3	7.89-8.58	7.22-7.55	0.00-0.15
Concatenated	8.53-8.95	7.82-8.01	0.05-0.09
<i>S. orarius</i>			
Cytb	6.75-8.66	7.50-9.50	8.00-9.89
Co1	8.98-10.05	9.52-9.89	9.44-10.20
Co3	8.06-10.14	6.88-9.09	8.40-9.61
Concatenated	8.05-9.00	7.82-8.43	8.72-9.00
<i>S. townsendi</i>			
Cytb	8.00-9.39	8.00-9.50	9.75-11.14
Co1	8.98-8.98	7.91-8.98	9.59-10.20
Co3	8.92-9.79	7.72-8.75	9.09-9.61
Concatenated	8.67-9.19	8.40-8.72	9.43-9.62

Scapanus anthonyi Allen 1893

1893. *Scapanus anthonyi* Allen, Bull Amer. Mus. Nat. Hist., 5:200, August. Type locality: "Sierra San Pedro Martir, 7000 ft, Baja California [México]". Adult male, skin and skull, American Museum of Natural History number 6313, collected by A. W. Anthony.

1937. *Scapanus latimanus anthonyi* Palmer, J. Mamm. 18:312, August. Name combination.

Geographic range. Restricted to the highlands of Sierra San Pedro Martir, Baja California, México.

Diagnosis and comparison. *Scapanus anthonyi* can be differentiated from the other species of *Scapanus* in having fewer than seven unicuspids teeth behind the incisors in the mandible and maxilla and total skull length <32.5 mm. Projection present in the braincase between the interparietal and the mastoid ([Jackson 1915](#)). Manus more square and smaller, with broader and heavier phalanges, tips of the pterygoids bones of the upper palate paral-

lel ([Huey 1936](#)). Smaller in all craniodental and somatic measurements relative to all other subspecies of *S. latimanus*, and teeth larger and crowded ([Yates and Salazar-Bravo 2005](#):494 in table 3). Differing from *S. orarius* and *S. townsendii* in a smaller in size; dorsal coloration darker, almost black; no spaces between all unicuspid teeth, usually crowded; and rostrum short and broad.

Comments. *Scapanus anthonyi* has a distribution restricted to the upper portions of the Sierra San Pedro Martir, within the pine and oak-pine forest. Collecting moles in the region is complex for several reasons. First, the gopher *Thomomys nigricans* is very abundant in the same area, so it is common to find gophers galleries that impinge upon and destroy mole galleries. Second, both species share a sympatric distribution throughout the mountain range. Third, although this region is a protected area, large numbers of cattle graze in the area and destroy the mole galleries. Fourth, specimens of *S. anthonyi* are very small in size, so their galleries also are small and the soil relief (molehill) that results from gallery construction is very difficult to determine. Fifth, galleries have a simple structure, and raise just 3 cm above the ground, and any leaf litter makes these molehills invisible (Cortés-Calva pers. obs.).

In the area, both *Scapanus* and *Thomomys* are named “*topos*” (moles) with no distinction between them, and only old ranchers give different names to them. *Scapanus* are called “*topos de manoplas*” (baseball-gloved moles) in reference to its forefoot size. *Thomomys* are known only as “*topos*”.

Scapanus occultus Grinnell and Swarth 1912

1912. *Scapanus latimanus occultus* Grinnell and Swarth, Univ. California, Publ. Zool., 10:131, April. Type locality: “Santa Ana canyon, 400 ft (12 mi NE Santa Ana), Orange County California”. Subadult female, skin and skull, Museum of Vertebrate Zoology, University of California, Berkeley, number 2369, collected by H. S. Swarth.

1914. *Scapanus latimanus grinnelli* Jackson, Proc. Biol. Soc. Washington, 27:56. Considered as junior synonym.

Geographic range. From Laguna Hanson (Sierra de Juárez) Baja California, México northwestward to Santa Barbara and northward to Yosemite Valley in Mariposa County, California.

Diagnosis and comparison. *Scapanus occultus* can be differentiated from *S. latimanus* in its smaller size and longer and wider skull ([Yates and Salazar-Bravo 2005](#):494 in table 3). Some specimens have fewer than seven unicuspid teeth, but only on a single side of the mandible or maxilla. It differs from *S. orarius* and *S. townsendii* by the same characteristics mentioned in *S. anthonyi*.

Scapanus latimanus (Bachman 1842)

1842. *Scapanus latimanus* Bachman, Boston Jour. Nat. Hist., 4:34. Type locality “probably from Santa Clara, Santa Clara, California” Mounted specimen with imperfect skull, Berlin Museum, collected during October 1834.

1912. *Scapanus latimanus latimanus* Grinnell and Swarth, Univ. California, Publ. Zool., 10:131, April. First use of current name combination.

Geographic range. From Santa Barbara and Yosemite Valley, California, northward to southcentral Oregon.

Diagnosis and comparison. *Scapanus latimanus* can be differentiated from *S. orarius* and *S. townsendii* by the same characteristics mentioned in *S. anthonyi*.

Keys for the species of *Scapanus*

1. Dorsal coloration usually brown to gray. All unicuspid teeth with variable spacing between them and usually crowded; rostrum short and broad 2
- 1a. Dorsal coloration almost black. All unicuspid teeth with regular spacing between them, and not crowded; rostrum long and narrow 4
2. Fewer than seven unicuspid teeth behind the incisors in the mandible and maxilla; total skull length less than 32.5 mm *Scapanus anthonyi*
- 2a. Seven unicuspid teeth behind the incisors in the mandible and maxilla; total skull length >32.5 mm 3
3. Total length >161.0 mm. Skull length >34.0 mm in males and 33.4 mm in females. Ratio of mastoidal breadth to greatest skull length <49%, including the population of Alameda Island, California *Scapanus latimanus*
- 3a. Total length <161.0 mm. Skull length <34.0 mm in males and 33.4 mm in females. Ratio of mastoidal breadth to greatest skull length >49%, not including the population of Alameda Island, California *Scapanus occultus*
4. Total length >200.0 mm on average. Sublacrimo-maxillary ridge well developed; skull > 40.0 mm *Scapanus townsendii*
- 4a. Total length <200.0 mm on average. Sublacrimo-maxillary ridge little developed; skull <40.0 mm *Scapanus orarius*

Acknowledgments

We dedicate this article to “Dave” Schmidly as a small tribute to his great academic and scientific career. As a recognition of how much he has done for the understanding of Mammalogy, especially in México and to the great support that he has always had from his colleagues in the profession. Although, the authors of this article did not have the opportunity to work directly with him, we have been under what he helped create. In fact, Therya is a secondary product of what the impulse to create a few decades ago and today it is an important window for the dissemination of science that is done mainly in México and Latin America. We wish to express our appreciation to Joseph A. Cook, Jonathan L. Dunnum, and Mariel L. Campbell Museum of Southwestern Biology (MSB), The University of New Mexico. Bruce Patterson, Field Museum of Natural History (FMNH); James L. Patton and Christopher J. Conroy, Museum of Vertebrate

Zoology at the University of California (MVZ); Vicky Mingna Zhuang, The University of Texas at El Paso (UTEP). We thank G. De León Girón (Parque Nacional Sierra San Pedro Mártir), for granting permission and logistic support, Fr. Arce and M. de La Paz Cuevas for their valuable assistance in the field; and G. Gallegos-Simental for her assistance in laboratory Node CIBNOR Barcode of Mexican Network of the Barcode of Life (MEXBOL), and C. I. Gutiérrez Rojas in laboratory work. This study received financial support (Grant 280896, 295569-CONACYT) Red temática CONACYT Código de Barras de la Vida, María Elena Sánchez-Salazar translated the manuscript into English.

Literature Cited

ALLEN, J. A. 1893. On a collection of mammals from San Pedro Martir region of Lower California. *Bulletin American Museum Natural History* 5:181-202.

BANGS, O. 1899. Descriptions of some new mammals from western North America. *Proceedings of the New England Zoological Club* 1:65-72.

BACHMAN, J. 1842. Observations on the genus *Scalops*, (shrew moles,) with descriptions of the species found in North America. *Boston Journal of Natural History* 4:26-35.

CAMARGO, I., AND S. T. ÁLVAREZ-CASTAÑEDA. 2020. A new species and three subspecies of the desert shrew (*Notiosorex*) from the Baja California peninsula and California. *Journal of Mammalogy* 101:872-886.

FELSENSTEIN, J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution* 17:368-376.

GRINNELL, J. 1918. Six new mammals from the Mohave Desert and Inyo regions of California. *University of California Publications in Zoology* 17:423-430.

GRINNELL, J., AND T. I. STORER. 1916. Diagnoses of seven new mammals from east-central California. *University of California Publications in Zoology* 17:1-8.

GRINNELL, J., AND H. S. SWARTH. 1912. The mole of southern California. *University of California Publications in Zoology* 10:131-136.

HUELSENBECK, J. P., AND F. RONQUIST. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754-755.

HUEY, L. M. 1936. Notes on the moles of Lower California, México. *Journal of Mammalogy* 17:166-167.

HUTCHINSON, J. H. 1987. Moles of the *Scapanus latimanus* Groups (Talpidae, Insectivora) from the Pliocene of California. *Contributions in Science, Natural History Museum of Los Angeles County* 386:1-15.

HUTTERER, R. 2005. Order Soricomorpha Pp. 220-311, in *Mammal species of the world. A taxonomic and geographic reference* (Wilson, D. E., and D. A. M. Reeder, eds.). Third edition. The Johns Hopkins University Press. Baltimore, U.S.A.

IVANOVA, N. V., S. T. ZEMLAK, R. H. HANNER, AND P. HEBERT. 2007. Universal primer cocktail for fish DNA barcoding. *Molecular Ecology Notes* 7:544-548

JACKSON, H. H. T. 1914. New moles of the genus *Scapanus*. *Proceedings of the Biological Society of Washington* 27:55-56.

JACKSON, H. H. T. 1915. A review of the American moles. *North American Fauna* 38:1-100.

MERRIAM, C. H. 1897. The mammals of mount Mazama, Oregon. *Mazama* 1:204-230.

MOORE, D. W. 1986. Systematic and Biogeographic Relationships among the Talpinae (Insectivora: Talpidae), Ph.D. dissertation, University of New Mexico, Albuquerque.

MOTOKAWA, M. 2004. Phylogenetic relationships with Talpidae (Mammalia: Insectivora). *Journal of Zoology (London)* 263:147-157.

MOUCHATY, S., A. GULLBERG, A. JANKE, AND U. ARNASON. 2000. The phylogenetic position of the Talpidae within Eutheria based on analysis of complete mitochondrial sequences. *Molecular Biology and evolution* 17:60-67.

NYLANDER, J. A. A. 2004. MrModeltest v2.2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University. Uppsala, Sweden.

PALMER, F. G. 1937. Geographic variation in the mole *Scapanus latimanus*. *Journal of Mammalogy* 18:280-314.

RIDDLE, B. R. 1995. Molecular biogeography in the pocket mice (*Perognathus* and *Chaetodipus*) and grasshopper mice (*Onychomys*): the Late Cenozoic development of a North American arid lands rodent guild. *Journal of Mammalogy* 76:283-301.

SHINOHARA, A., K. L., CAMPBELL, AND H. SUZUKI. 2003. Molecular phylogenetic relationships of moles, shrew moles, and desmans from the new and old worlds. *Molecular Phylogenetics and Evolution* 27: 247-258.

SIKES, R. S., AND THE ANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF MAMMALOGISTS. 2016. Guidelines of the American society of mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* 97:663-688.

SMITH, M. F. 1998. Phylogenetic relationships and geographic structure in pocket gophers in the genus *Thomomys*. *Molecular Phylogenetics and Evolution* 9:1-14.

SMITH, M. F., AND J. L. PATTON. 1993. The diversification of South American murid rodents: evidence from mitochondrial DNA sequence data of the akodontine tribe. *Biological Journal of the Linnean Society* 50:149-177.

SWOFFORD, D. L. 2002. PAUP: phylogenetic analysis using parsimony, version 4.0b10. Sinauer Associates. Sunderland, Massachusetts.

TRUE, F. W. 1894. Diagnosis of new North American mammals. *Proceeding U. S. Natural Museum* 19:1-11.

YATES, T. L., AND I. F. GREENBAUM. 1982. Biochemical systematics North American moles (Insectivora:Talpidae). *Journal of Mammalogy* 63:368-374.

YATES, T. L., AND J. SALAZAR-BRAVO. 2005. Revision of the *Scapanus latimanus*, with the revalidation of a species of Mexican mole. Pp. 489-506, in *Contribuciones mastozoológicas en Honor a Bernardo Villa* (V. Sánchez-Cordero, and R. Medellín, eds.). Instituto de Biología, UNAM, Instituto de Ecología UNAM. Ciudad de México, México.

ZHAO, J., AND G. JIAN. 2015. Complete mitochondrial genome of *Condylura cristata* (Unpublished). BGI Education Center, University of Chinese Academy of Sciences, Shenzhen, China.

Associated editor: Lisa and Robert Bradley

Submitted: March 27, 2021; Reviewed: April 27, 2021

Accepted: May 8, 2021; Published on line: May 28, 2021.