

Spatial and temporal distribution of bat mortality on a highway in southeast Brazil

MARCIONE BRITO DE OLIVEIRA¹, AND CECÍLIA BUENO^{2*}

¹ Postgraduate Program in Zoology, National Museum, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista s/n, São Cristóvão, C. P. 20940-040. Rio de Janeiro, RJ, Brazil. Email: oliveira01marcione@gmail.com (MBO).

² Universidade Veiga de Almeida, Núcleo de Estudos de Vertebrados Silvestres, Rua Ibituruna, 108, Maracanã, C. P. 20271-020. Rio de Janeiro, RJ, Brazil. Email: cecilia.bueno2020@gmail.com (CB).

*Corresponding author

Roads and highways can work as barriers to the movement of many species, thereby preventing the individuals from accessing feeding and reproduction sites and the immediate risk of colliding wild species with vehicles. Identifying the impacts of roads on wildlife can contribute to the establishment of actions that promote conservation. In Brazil, data on collisions between bats and vehicles are scarce and underestimated in the literature. We described bat roadkill from 2008 to 2019 on a stretch of the BR-040 highway, which crosses an area of Atlantic Forest. Roadkill species were identified and the sites with high collision frequencies were characterized. A total of 923 individuals of 57 species and five families of chiropterans were identified. Frugivore bats showed the largest number of affected individuals, with *Artibeus lituratus*, a common species in the study region, with the highest number of roadkills. The diet and foraging behaviour were the most likely factors explaining most of the bats killed on the highway. The highest roadkill rate was documented in the fall, and the critical points located nearby the APA Petrópolis and REBIO of Tinguá, environmental protection areas. We reinforce the need to mitigate these roadkills, ensuring that road systems, which constitute municipal, state and federal highways, are built to prevent major disturbance of habitat and displacement routes of these species. We believe in the need for mitigations, and considering the various species involved, we suggest speed bumps construction reducing the speed limit, installing bridges, and signaling the presence of wildlife, before the stretches identified as hotspots.

Los caminos y carreteras pueden funcionar como barreras para el movimiento de muchas especies, impidiendo así el acceso de los individuos a los sitios de alimentación y reproducción y el riesgo inmediato de colisión de especies silvestres con vehículos. Identificar los impactos de los caminos sobre la vida silvestre puede contribuir al establecimiento de acciones que promuevan la conservación. En Brasil, los datos sobre colisiones entre murciélagos y vehículos son escasos y subestimados en la literatura. Describimos murciélagos atropellados entre 2008 y 2019 en un tramo de la carretera BR-040, que atraviesa un área de Mata Atlántica. Se identificaron las especies de animales atropellados y se caracterizaron los sitios con alta frecuencia de colisión. Se identificaron un total de 923 individuos de 57 especies y cinco familias de quirópteros. Los murciélagos frugívoros mostraron el mayor número de individuos afectados, con *Artibeus lituratus*, una especie común en la región de estudio, con el mayor número de atropellamientos. La dieta y el comportamiento de búsqueda de alimento fueron los factores más probables que explicaron la mayoría de los murciélagos muertos en la carretera. La mayor tasa de atropellamientos se registró en otoño, y los puntos críticos se ubicaron en las cercanías de APA Petrópolis y REBIO de Tinguá, áreas de protección ambiental. Reforzamos la necesidad de mitigar estos atropellamientos, asegurando que los sistemas viales, que constituyen las carreteras municipales, estatales y federales, se construyan para evitar que se alteren el hábitat y las rutas de desplazamiento de estas especies. Creemos en la necesidad de mitigaciones y considerando las diversas especies involucradas. Sugerimos la construcción de topes para reducir el límite de velocidad, instalar puentes y señalizando la presencia de fauna, antes de los tramos identificados como hotspots.

Keywords: Chiroptera; collisions; mitigation; road ecology; southeast Brazil.

© 2022 Asociación Mexicana de Mastozoología, www.mastozoologiamexicana.org

Introduction

The human population in recent decades has caused several changes in the environment, including road and highway construction that is currently a significant concern for biodiversity conservation ([do Prado et al. 2006](#); [Gumier-Costa and Sperber 2009](#); [Bueno et al. 2015](#)). Road impacts on wildlife include deaths from vehicle collisions, habitat destruction and fragmentation, barrier effect, edge effects and disturbances caused by light, noise and chemical pollution ([Forman and Alexander 1998](#); [Gibbs and Shriver 2002](#); [Row et al. 2007](#); [Kerth and Melber 2009](#); [Stone et al. 2009](#); [Zurcher et al. 2010](#); [Berthiussen and Altringham 2012](#); [Bueno et al. 2013](#); [Esperandio et al. 2019](#)). For several vertebrate species,

some of the ecological consequences include a reduction in the gene flow between populations and a decrease in both migration rates and genetic diversity ([Corlatti et al. 2009](#); [Esperandio et al. 2019](#)).

The order Chiroptera is the second largest among mammals, with more than 1,400 living species ([Simmons and Cirranello 2020](#)). In Brazil, this order is represented by 68 genera and 181 species subdivided into nine families ([Garbino et al. 2020](#)). The chiropterans play a fundamental role in maintaining the ecological processes of different vegetation domains, being excellent objects of study for characterizing habitats and their quality since they are also sensitive to changes ([Fenton et al. 1992](#); [Medellín et al. 2000](#);

[Gorresen et al. 2005](#)). However, most data on bat deaths on the roads comes from Europe, where several studies have been developed ([Bafaluy 2000](#); [Lesiński 2007](#); [Gaisler et al. 2009](#); [Kerth and Melber 2009](#); [Berthiussen and Altringham 2012](#); [Medinas et al. 2013](#); [Ramalho and Aguiar 2020](#)), while in Brazil, data on bat collisions are scarce ([Pracucci et al. 2012](#); [Pinheiro and Turci 2013](#); [Almeida and Cardoso Junior 2014](#); [Ceron et al. 2017](#); [Valadão et al. 2018](#); [Damásio et al. 2021](#); [de Figueiredo Ramalho et al. 2021](#)). The bat's low body mass, flight speed and nocturnal behavior make it difficult to find bat carcasses, resulting in the subsampling of the group in studies of road ecology ([Bafaluy 2000](#); [Lesiński et al. 2011](#); [Berthiussen and Altringham 2012](#)).

The bats forage using echolocation ([Kalko et al. 1996](#); [Siemers and Schnitzler 2000](#); [Schaub et al. 2008](#)) and hearing the sound of prey such as amphibians and insects ([Marimuthu and Neuweiler 1987](#); [Faure and Barclay 1992](#); [Altringham and Kerth 2016](#)). According to [Schaub et al. \(2008\)](#), the 'passive listening' strategy is adopted by bat species specialized in hunting arthropods in vegetation or on the ground, where other sounds of the environment mask the noises emitted by prey; thus, the sounds produced by cars can compromise the echolocation of bats nearby, reducing their feeding effectiveness. Therefore, roads act as barriers, reducing the bats' ability to access suitable places to forage or shelter, thus being a concerning factor, especially regarding migratory species ([Zurcher et al. 2010](#); [Lesiński et al. 2011](#); [Ceron et al. 2017](#); [Claireau et al. 2019](#)).

Although some studies show that bats recognize vehicles as a threat ([Baxter et al. 2006](#); [Zurcher et al. 2010](#); [Altringham and Kerth 2016](#)), many others show that bats are killed on the roads by vehicle collisions ([Bafaluy 2000](#); [Gaisler et al. 2009](#); [Ceron et al. 2017](#)). Moreover, some studies recent provide only a list of affected species ([Ceron et al. 2017](#); [Novaes et al. 2018](#)). Important variables that affect the collision rates between bats and vehicles, such as the diversity and abundance of specific plants in the landscape, the presence of flight corridors, habitat fragmentation and foraging routes ([Gaisler et al. 2009](#); [Medinas et al. 2013](#)), are poorly documented in Brazil. Despite these factors, bat species moving on the road may vary from region to region, indicating other differences in the structure of the local landscape ([Lesiński 2007](#); [Kerth and Melber 2009](#)).

Due to the lack of studies on collisions between vehicles and bats on the roads, the need to obtain data on mortality from being run over in different regions and landscapes is evident since the impact on natural populations can be severe and may cause a decline due to the cumulative effect of roadkills ([Lesiński et al. 2011](#); [Berthiussen and Altringham 2012](#); [de Figueiredo Ramalho et al. 2021](#)). In that regard, monitoring of the occurrence of collisions between vehicles and bats was carried out on a long stretch on the BR-040 highway in the period from 2008 to 2019, identifying the affected species and the points with the highest collision frequencies.

Materials and Methods

Study area. Data was collected on the BR-040 highway, which covers nine municipalities, from Rio de Janeiro to Juiz de Fora (Figure 1), accounting for 180.4 km. In the State of Rio de Janeiro, it goes from 0 km to 125.2 km, and in the State of Minas Gerais from 773.5 km to 828.7 km. Six municipalities are located in the state of Rio de Janeiro: Rio de Janeiro, Duque de Caxias, Petrópolis, Areal, Três Rios, Comendador Levy Gasparian, and three in the state of Minas Gerais: Simão Pereira, Matias Barbosa and Juiz de Fora. Along the studied stretch, the BR-040 highway crosses three Conservation Units of Atlantic Forest: Petrópolis Environmental Protection Area (APA Petrópolis), the Tinguá Biological Reserve (REBIO for Tinguá), and the Serra da Estrela Wildlife Refuge. The climate along the monitored highway is humid subtropical with a hot temperate summer and a dry winter, according to the Köppen classification, with precipitation concentrated in the summer, in an annual total that varies in the region between 1,300 and 2,200 mm ([Alvares et al. 2013](#)). We plot a final map with the critical areas of fatalities identified from analysis in Siriema software in the ArcGIS Desktop ([Ormsby et al. 2010](#)).

Monitoring and data collection. The data were compiled in the period from 2008 to 2019, and the stretches were travelled weekly at 40 km/h by two observers, in




Figure 1. BR-040, stretch from the municipality of Rio de Janeiro to Juiz de Fora.

addition to highway traffic inspectors monitoring 24 hours a day, seven days a week. The bats found dead were collected; when possible, they were identified at the lowest taxonomic level, stored in individual plastic bags, and frozen at the four User Service Centers. Subsequently, the specimens were sent to the Wild Animal Studies Center located at the Veiga de Almeida University, where they were prepared for deposition in the mammalogy section of the National Museum of Rio de Janeiro, a reference collection. Roadkill locations were georeferenced, and bats were photographed, sexed, measured, and identified more accurately in the laboratory. In this study, we included all individuals that could be identified within the order Chiroptera. Bats were identified with the help of identification keys provided by [Gardner \(2007\)](#), and [Reis et al. \(2017\)](#), and whenever necessary, identification was checked in the laboratory with available literature ([Gregorin and Taddei 2002](#); [Dias and Peracchi 2008](#); [Velazco et al. 2010](#); [Moratelli et al. 2013](#); [Cirranello et al. 2016](#)).

The collections of roadkill specimens were made under SISBIO License, Number: 30727-12. The use of these animals in this study complies with and is under the Operating License, Number: 1187/2013 and Authorization for Capture, Collection, and Transport of Biological Material - Abio (Renewal) No. 514/2014.

Data analysis. Siriema 2.0 software was used ([Coelho et al. 2014](#)) to identify the main stretches of significant groupings within a highway, the hotspots, and the analyses were 2D HotSpots and Ripley 2D K statistics. Ripley's K statistic describes the dispersion of data across a range of spatial scales ([Ripley 1981](#); [Levine 2004](#)). To define the different scales evaluated, we used an initial radius of 100 m and an increment of 400 m for each step. To assess the meaning of possible aggregations, we subtracted the observed K values from the average obtained in 1,000 simulations of random roadkill distributions for each stop. The values above the 95% confidence limits obtained in the simulations indicate scales with significant aggregations ([Levine 2004](#); [Teixeira et al. 2013](#); [Coelho et al. 2014](#)). To find the critical hit-and-run points, we performed a two-dimensional analysis of the identification of the hotspots. Using an average of 1,000 simulations with a 95% confidence interval, we divided the highway into segments of 200 meters each ([Teixeira et al. 2013](#); [Coelho et al. 2014](#)).

Data on the number of individuals run over in each of the four years and comparing the seasons were tested for normality of residues, using the Shapiro-Wilk test and homoscedasticity of residues, using the Bartlett test. Having the data sets in which the assumptions of normality and homoscedasticity of the residuals were accepted, ANOVA (one way) was performed, and a BoxPlot was plotted to verify differences between the results obtained in the present study. The significance level used in the tests was 0.05%, and the R software was used to perform all the statistical tests ([R Core Team 2021](#)).

Results

A total of 923 roadkilled individuals of 57 species of bats were recorded. Five families (Table 1) were observed; amongst them, the Phyllostomidae ($n = 652$; 71%) and Vespertilionidae ($n = 204$; 22 %) were the most common roadkilled. The species with the highest collision frequency were *Artibeus lituratus* ($n = 216$; 23.4 %), *Glossophaga soricina* ($n = 61$; 6.61 %), *Carollia perspicillata* ($n = 55$; 5.96 %), *Platyrrhinus recifinus* ($n = 46$; 4.98 %), *Sturnira lilium* ($n = 41$; 4.44 %), *Platyrrhinus lineatus* ($n = 40$; 4.33 %), *Phyllostomus hastatus* ($n = 38$; 4.12 %), *Nyctinomops laticaudatus* ($n = 33$; 3.58 %), *Artibeus fimbriatus* and *Anoura caudifer* ($n = 31$; 3.36 %); other species had less than 20 records each, and 151 individuals (16.36 %) could only be identified to the order level.

Frugivores represented the most roadkills, represented by 19 species and 52.87 % of the total number of individuals; however, the greatest recorded richness was that of insectivores, represented by 24 species and 14 % of individuals roadkills. The nectarivores were represented by five species and 10.83 % of the total roadkill. The other trophic guilds were hematophagous, with two species, and carnivore, omnivore and piscivore, represented by one species each. These represent a total of 5.9 % of roadkill recorded in the study. In most of the specimens, it was not possible to identify sex and age; however, in those where it was possible, the male/female rate was similar, and most of them were adults (Table 1). Differences were observed in the frequency of bat collisions between the seasons. The highest mortality was documented in the fall with 330 (35.75 %) individuals killed in collisions, followed by 251 (27.19 %) in the summer, 217 (23.51 %) in the winter and 125 (13.54 %) in the spring. Considering the number of individuals run over in relation to the seasons and each monitoring year (2014, 2015, 2016, 2017, and 2018). According to the Shapiro-Wilk test, the data are normally distributed ($W = 0.98471$, $\alpha = 2$, $P = 0.8891$) and are homoscedastic according to the Bartlett test (2.1022 , $\alpha = 3$, $P = 0.5515$). Therefore, the data were submitted to the parametric ANOVA (One Way) test, where significant differences were observed between the data set of individuals hit by cars in relation to the seasons of the year in each monitored year ($F = 1.337$, $\alpha = 3$, $P = 0.297$). The BoxPlot showed a difference, in terms of the number of individuals run over, between the spring season and the other seasons of the year (Figure 2).

In the analyses to determine the scales in which road deaths were significantly aggregated in space, we found different aggregation points for bats. The stretch in the city of Rio de Janeiro, that cover less than 1 km, was the only one that had no record of bat's roadkills. The spatial distribution of roadkill along the BR-040 highway was not random, indicating that the records concentrated on some specific points along the highway. Many stretches had significant aggregations, with the most probable and most significant extension being located between the Duque de Caxias and Petrópolis kilometers. This most critical stretch begins at the ascent of the sierra of Petrópolis. Other peaks of signifi-

Table 1. Species of bats roadkill during the study and the number of collisions according to their age and sexual classification. Undefined sex = SexInd; Undefined age category = AgeInd. Conservation status for species at global level follows IUCN (2020). Data Deficiente (DD); Least Concern (LC); Near Threatened (NT); No identification (S/ID).

Family / Species	Female	Male	SexInd	Adult	Young	AgeInd	IUCN
Emballonuridae							
<i>Saccopteryx leptura</i>	0	0	1	1	0	0	LC
Phyllostomidae							
<i>Anoura caudifer</i>	6	6	19	16	0	15	LC
<i>Anoura geoffroyi</i>	1	1	1	1	0	2	LC
<i>Anoura</i> sp.	0	0	1	0	0	1	
<i>Artibeus cinereus</i>	1	0	2	1	0	2	LC
<i>Artibeus fimbriatus</i>	5	9	17	16	0	15	LC
<i>Artibeus lituratus</i>	43	50	123	105	4	107	LC
<i>Artibeus obscurus</i>	0	0	1	0	0	1	LC
<i>Artibeus planirostris</i>	0	1	0	1	0	0	LC
<i>Artibeus</i> sp.	0	0	17	8	0	9	
<i>Carollia perspicillata</i>	17	11	27	24	2	29	LC
<i>Chiroderma doriae</i>	3	1	2	3	0	3	LC
<i>Chiroderma</i> sp.	0	0	1	1	0	0	
<i>Chiroderma villosum</i>	1	2	1	2	0	2	LC
<i>Choeroniscus minor</i>	1	0	0	1	0	0	LC
<i>Chrotopterus auritus</i>	0	0	4	0	0	4	LC
<i>Desmodus rotundus</i>	2	2	5	6	0	3	LC
<i>Diphylla ecaudata</i>	0	1	1	1	0	1	LC
<i>Glossophaga soricina</i>	18	13	30	35	1	25	LC
<i>Lonchophylla peracchii</i>	0	1	2	1	0	2	LC
<i>Macrophyllum macrophyllum</i>	0	0	1	0	0	1	LC
<i>Micronycteris hirsuta</i>	0	0	1	1	0	0	LC
<i>Micronycteris megalotis</i>	1	1	0	2	0	0	LC
<i>Micronycteris microtis</i>	0	0	1	0	0	1	LC
<i>Micronycteris minuta</i>	1	0	0	1	0	0	LC
<i>Mimon bennettii</i>	2	2	1	4	0	1	LC
<i>Phyllostomus hastatus</i>	5	7	26	24	1	13	LC
<i>Platyrrhinus</i> sp.	0	0	1	0	0	1	
<i>Platyrrhinus lineatus</i>	12	4	24	17	0	23	LC
<i>Platyrrhinus recifinus</i>	10	8	28	20	1	25	LC
<i>Pygoderma bilabiatum</i>	1	2	10	3	0	10	LC
<i>Sturnira lilium</i>	9	8	24	16	3	22	LC
<i>Sturnira tildae</i>	0	0	3	0	0	3	LC
<i>Uroderma</i> sp.	0	0	1	0	0	1	
<i>Vampyressa pusilla</i>	5	1	1	5	0	2	DD
<i>Vampyrodes caraccioli</i>	1	0	0	1	0	0	LC
Noctilionidae							
<i>Noctilio leporinus</i>	1	0	0	1	0	0	LC
Molossidae							
<i>Eumops</i> sp.	0	0	1	0	0	1	
<i>Molossops neglectus</i>	1	0	0	0	0	1	DD
<i>Molossus molossus</i>	1	3	11	5	0	10	LC
<i>Molossus rufus</i>	1	0	5	1	0	5	LC
<i>Nyctinomops aurispinosus</i>	0	0	1	0	0	1	LC
<i>Nyctinomops laticaudatus</i>	1	4	28	18	0	15	LC
<i>Nyctinomops</i> sp.	0	0	5	2	0	3	
<i>Tadarida brasiliensis</i>	0	0	1	0	0	1	LC
Molossidae (unidentified)	0	0	2	2	0	0	

Table 1. Continuation...

Family / Species	Female	Male	SexInd	Adult	Young	AgeInd	IUCN
Vespertilionidae							
<i>Eptesicus brasiliensis</i>	1	1	3	3	0	2	LC
<i>Eptesicus diminutus</i>	1	0	0	1	0	0	LC
<i>Eptesicus</i> sp.	1	0	2	2	0	1	
<i>Histiotus velatus</i>	1	0	0	1	0	0	DD
<i>Dasypterus ega</i>	1	1	4	2	0	4	LC
<i>Myotis albescens</i>	0	0	1	1	0	0	LC
<i>Myotis izecksohni</i>	1	1	1	2	0	1	DD
<i>Myotis nigricans</i>	1	3	5	3	0	6	LC
<i>Myotis riparius</i>	2	0	2	1	0	3	LC
<i>Myotis ruber</i>	0	1	0	1	0	0	NT
<i>Myotis</i> sp.	0	0	16	4	0	12	
Vespertilionidae (unidentified)	0	0	3	0	0	3	
S/ID	0	1	150	27	1	123	
Total	159	146	618	394	13	516	

cant aggregations were observed between the kilometers of Três Rios and Comendador Levy Gasparian (Figure 3). The most critical points on the road, according to the analyses, were in the Atlantic Forest Biodiversity Corridor, which includes the Tinguá Biological Reserve, the Serra da Estrela State Wildlife Refuge and the APA Petrópolis (Figure 4).

Discussion

Bats are valuable indicators of biodiversity and ecosystem health, as a result of his sensitivity to fragmentation, that can decrease the richness and abundance of the order in altered areas (Reis *et al.* 2003; Alurralde and Díaz 2021). Thus, habitat fragmentation could be an aggravating factor in the seed dispersal process carried out by Chiroptera species. Despite the relative richness and abundance observed during the monitoring of the current study, these values can be related to the time and

size of the sample. However, still can be underestimated for the region due to the limitation of the study that only recorded species found dead by collisions. Species richness during this monitoring was similar to other studies in southeastern Brazil (Dias and Peracchi 2008; Esbérard and Bergallo 2008; Tavares *et al.* 2010; Perini *et al.* 2014), yet, we observed a high variation between the richness and diversity of these studies, which is probably due to differences in capture techniques, such as the use of mist nets or active search for shelters in areas with landscape mosaic varying from conserved to degraded environments (Dias and Peracchi 2008; Esbérard and Bergallo 2008; Tavares *et al.* 2010; Perini *et al.* 2014; Alurralde and Díaz 2021), and differences in collision probabilities by bat species in the present study.

Frugivore bats are usually the most frequent group in collisions with vehicles on the roads, according to our data and other studies on South America (Pinheiro and Turci 2013; Ceron *et al.* 2017; Valadão *et al.* 2018; de Figueiredo Ramalho *et al.* 2021), where frugivores are more representative (Fenton *et al.* 1992; Martins *et al.* 2014). Species as *Artibeus lituratus*, *Carollia perspicillata*, and *Sturnira lilium* are not inhibited by fragmentation and are more abundant in disturbed areas than other bat species (Medellín *et al.* 2000; Reis *et al.* 2003; Alurralde and Díaz 2021). However, these species have been most affected by collisions with vehicles on the roads (Pinheiro and Turci 2013; Ceron *et al.* 2017; Valadão *et al.* 2018; de Figueiredo Ramalho *et al.* 2021). These species have been considered abundant in the study region (Dias and Peracchi 2008; Tavares *et al.* 2010; Luz *et al.* 2011), and have characteristics that make them good indicators of damaged areas, providing advantages for plants whose fruits are consumed by them, is a key factor for dispersion and genetic flow of plants in regions affected by fragmentation (Muller and dos Reis 1992; Reis *et al.* 2003; Bianconi *et al.* 2006).




Figure 2. Boxplot (median with upper and lower quartiles) of the number of individuals of bats killed in collisions by season of the year on highway BR-040.




Figure 3. Bat roadkill aggregation intensity (blue line) and 95% confidence limits (black line) along 180.4 km of BR-040.

This higher rate of collisions with frugivores bats is probably due to some associated factors, such as flight height, displacement patterns in foraging, the temporal and spatial distribution of the fruits and the plants, as well as the landscape structure (Bernard and Fenton 2003; Clevenger *et al.* 2003; Bianconi *et al.* 2006; Gumier-Costa and Sperber 2009; Pracucci *et al.* 2012; Gomes *et al.* 2020). An essential factor is that many of these frugivorous species are consumers of pioneer plants, which are usually found in large densities in degraded areas, such as the vicinity and edges of roads (Muller and dos Reis 1992; Reis *et al.* 2003). The species most impacted by the collisions on BR-040, *Artibeus lituratus*, was also the most abundant in studies in the southeastern region (Esbérard and Bergallo 2008; Tavares *et al.* 2010; Luz *et al.* 2011; Perini *et al.* 2014). Although roadkills did not show marked seasonal trends, fewer bat collisions were documented during the spring. Moreover, there is evidence of the difference in the number of collisions by bats in the seasons (Lesiński *et al.* 2011), but little is known about the factors that can influence the number of roadkills by these species, needing further investigations.

Different species of bats can be impacted differently by the fragmentation of habitats resulting from a road network. Some factors may be contributing to this difference in the amount of roadkill by bat species, such as foraging strategies, diet, or resilience of species in degraded areas (Kalko *et al.* 1996; Stone *et al.* 2009; Bhardwaj *et al.* 2017; Claireau *et al.* 2019), and need be investigated. Rare species can potentially be affected even by small mortality rates (Fensome and Mathews 2016; Damásio *et al.* 2021). In the present study, many insectivores considered rare were found roadkilled. The study by Bhardwaj *et al.* (2021) pointed out that the activity of seven of ten insectivorous bats species decreased significantly within the proximity of a highway due to changes created in the surrounding environment by traffic, such as pollution, light, noise, and vibrations. The authors called the surrounding environment a "road effect zone," and in this area, the habitat is degraded or entirely unsuitable for wildlife, which leads to an indirect loss of habitat (Bhardwaj *et al.* 2021).

One of the stretches identified in this study as a highly significant hotspot is in an area of Atlantic Forest in the municipality of Petrópolis, a region inserted in the APA Petrópolis, the Serra da Estrela State Wildlife Refuge, and close to the Biological Reserve (REBIO) of Tinguá. Twenty-eight species of bats have been registered in REBIO so far, the most abundant being *C. perspicillata*, *S. lilium*, and *A. lituratus* (Dias and Peracchi 2008), also the most frequently recorded in the present study. We highlight the species *Myotis ruber*, which had a run-over record close to the REBIO area, and which is listed as almost threatened with extinction in the IUCN Red List (IUCN 2020). The most critical points were on the stretch that crosses Tinguá Biological Reserve, highlighting the need for more urgent conservation measures in this region. However, the simple counting of carcasses found on the roads is an underestimation of the actual rate of bats killed by being run over. Even in regular monitoring, some factors may influence this count. Bat carcasses can be completely destroyed by continuous vehicle traffic, some can be thrown off the road and lost in the vegetation, and some individuals could be removed by other animals for consumption (Pracucci *et al.* 2012; Cunha *et al.* 2015; Ramalho and Aguiar 2020).

The roads that cross parks and forest reserves can have an extremely negative impact, a barrier to the natural movement of the species, and establishing a population decline, as well as decreasing the genetic flow for the wildlife of these regions (Samson *et al.* 2016). Critically, bat collisions with vehicles have been observed mainly in these areas where endangered and endemic bat species occur (Lesiński 2007). Moreover, these regions also harbor other wild species, thus making roadkills more critical for the conservation of species (IUCN 2020). Efforts to mitigate these accidents in the region are carried out with wildlife warning signs, yet these had not shown been proved effective. Roads are already proven to be harmful to many wild animals (Trombulak and Frissell 2000; Fahrig and Rytwinski 2009; Gumier-Costa and Sperber 2009), and there are several approaches that can be used to reduce the effects of




Figure 4. The sections marked in red indicate the critical areas of fatalities (HotSpots) identified from monitoring of run over bat fauna, analysis in Siriema software.

roads and mortality in wild animal populations ([Jackson 2000](#); [Glista et al. 2009](#)). The use of underpasses and ecological bridges for wild fauna probably varies from species to species, depending on their behavior in terms of movement ([Berthinussen and Altringham 2012](#); [Bhardwaj et al. 2017](#)). However, a combination of monitoring and ecological bridges has shown great effectiveness in decreasing roadkill rates ([Bhardwaj et al. 2017](#)).

Our results suggest that bats are particularly vulnerable to the impact of roads. Consequently, they may take a long time to recover from disturbances due to their low fertility, longevity, and foraging activity in large areas of the landscape ([Findley 1993](#)). Roads can affect the amount of area available to wildlife habitat, particularly through the fragmentation, and also affect the quality of this area by traffic of vehicles that cause collisions and deaths of wild animals ([Bafaluy 2000](#); [Bueno et al. 2015](#); [de Figueiredo Ramalho et al. 2021](#)). Other factors that influence this mortality are the lack of proper signs and awareness of drivers who travel on these highways. Hybrid mitigation is applied to reduce the adverse effects of roads on wildlife and comply with environmental legislation. These mitigations should account for several species, considering the broad diversification of habits showed by bats ([Bernard and Fenton 2003](#); [Bianconi et al. 2006](#); [Gomes et al. 2020](#)), and their sensitivity to increased lighting, which can change their activity pattern ([Stone et al. 2009](#); [Appel et al. 2017](#); [Gomes et al. 2020](#)). We suggest speed bumps construction to reduce the speed limit, install bridges, and signaling the presence of wildlife on the stretches identified as hotspots in our analyses, that is, with a higher risk of collision for these species of bats.

Acknowledgments

Marcione B. Oliveira received CNPq fellowship (Proc. 162273/2018-5) and Cecília Bueno thanks the Concer, BR-040 Concessionaire, for support to this research. We thank Igor Pfeifer Coelho, Camila L. Nacif, and Maria Clara S. Ribeiro for their suggestions on improving the manuscript.

Literature Cited

ALMEIDA, V. M., AND J. C. S. CARDOSO JUNIOR. 2014. Silvestres na Rodovia Vicinal Antônio Joaquim de Moura Andrade entre os Municípios de Mogi Guaçu-SP e Itapira-SP. Foco 5:99-116.

ALTRINGHAM, J., AND G. KERTH. 2016. Bats and roads. Pp. 35-62, in *Bats in the Anthropocene: conservation of bats in a changing world* (Voigt, C. C., and T. Kingston, eds.). Springer International Publishing. London, U.K.

ALURRALDE, S. G., AND M. M. DÍAZ. 2021. Assemblage-level responses of Neotropical bats to forest loss and fragmentation. *Basic and Applied Ecology* 50:57-66.

ALVARES, C. A., ET AL. 2013. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift* 22:711-728.

APPEL, G., A. LÓPEZ-BAUCELLS, W. ERNEST-MAGNUSSON, AND P. E. D. BO-BROWIEC. 2017. Aerial insectivorous bat activity in relation to moonlight intensity. *Mammalian Biology* 85:37-46.

BAFALUY, J. 2000. Mortandad de murciélagos por atropello en carreteras del sur de la provincia de Huesca. *Galemys* 12:15-23.

BAXTER, D. J. M., J. M. PSYLLAKIS, M. P. GILLINGHAM, AND E. L. O'BRIEN. 2006. Behavioural response of bats to perceived predation risk while foraging. *Ethology* 112:977-983.

BERNARD, E., AND M. B. FENTON. 2003. Bat mobility and roosts in a fragmented landscape in central Amazonia, Brazil. *Biotropica* 35:262-277.

BERTHINUSSEN, A., AND J. ALTRINGHAM. 2012. Do bat gantries and underpasses help bats cross roads safely? *Plos One* 7:e38775.

BHARDWAJ, M., ET AL. 2021. Insectivorous bats are less active near freeways. *Plos One* 16:e0247400.

BHARDWAJ, M., ET AL. 2017. Differential use of highway underpasses by bats. *Biological Conservation* 212:22-28.

BIANCONI, G. V., S. B. MIKICH, AND W. A. PEDRO. 2006. Movements of bats (Mammalia, Chiroptera) in Atlantic Forest remnants in southern Brazil. *Revista Brasileira de Zoologia* 23:1199-1206.

BUENO, C., C. O. M. SOUSA, AND S. R. FREITAS. 2015. Habitat or matrix: which is more relevant to predict roadkill of vertebrates? *Brazilian Journal of Biology* 75:228-238.

BUENO, C., M. T. FAUSTINO, AND S. R. FREITAS. 2013. Influence of landscape characteristics on capybara roadkill on highway BR-040, southeastern Brazil. *Oecologia Australis* 17:130-137.

CERON, K., ET AL. 2017. Roadkilled bats (Mammalia: Chiroptera) in two highways of Santa Catarina state, Southern Brazil. *Oecologia Australis* 21:207-212.

CIRRANELLO, A., N. B. SIMMONS, S. SOLARI, AND R. J. BAKER. 2016. Morphological diagnoses of higher-level phyllostomid taxa (Chiroptera: Phyllostomidae). *Acta Chiropterologica* 18:39-71.

CLAIREAU, F., ET AL. 2019. Major roads have important negative effects on insectivorous bat activity. *Biological Conservation* 235:53-62.

CLEVINGER, A. P., B. CHRUSZCZ, AND K. E. GUNSON. 2003. Spatial patterns and factors influencing small vertebrate fauna roadkill aggregations. *Biological Conservation* 109:15-26.

COELHO, A. V. P., I. P. COELHO, F. T. TEIXEIRA, AND A. KINDEL. 2014. Siriema: road mortality software. Manual do Usuário V. 2.0. NERF, UFRGS, Porto Alegre, Brasil. <https://www.ufrgs.br/sirimea>. Accessed on 07 July 2020

CORLATTI, L., K. HACKLÄNDER, AND F. FREY-Roos. 2009. Ability of wildlife overpasses to provide connectivity and prevent genetic isolation. *Conservation Biology* 23:548-556.

CUNHA, G. G., M. T. HARTMANN, AND P. A. HARTMANN. 2015. Roadkills of vertebrate species in the Pampa Region, South Brazil. *Ambiência* 11:307-320.

DAMÁSIO, L. *ET AL.* 2021. Diversity and Abundance of Roadkilled Bats in the Brazilian Atlantic Forest. *Diversity* 13:335.

DE FIGUEIREDO RAMALHO, D. *ET AL.* 2021. Factors influencing bat road casualties in a Neotropical savanna. *Perspectives in Ecology and Conservation* 19:189-194.

DIAS, D., AND A. L. PERACCHI. 2008. Quirópteros da Reserva Biológica do Tinguá, estado do Rio de Janeiro, sudeste do Brasil (Mammalia: Chiroptera). *Revista Brasileira de Zoologia* 25:333-369.

DO PRADO, T. R., A. A. FERREIRA, AND Z. F. S. GUIMARÃES. 2006. Efeito da implantação de rodovias no cerrado brasileiro sobre a fauna de vertebrados. *Acta Scientiarum. Biological Sciences* 28:237-241.

ESBÉRARD, C. E. AND H. G. BERGALLO. 2008. Influência do esforço amostral na riqueza de espécies de morcegos no sudeste do Brasil. *Revista Brasileira de Zoologia* 25:67-73.

ESPERANDIO, I. B. *ET AL.* 2019. Do roads act as a barrier to gene flow of subterranean small mammals? A case study with *Ctenomys minutus*. *Conservation Genetics* 20:385-393.

FAHRIG, L., AND T. RYTWINSKI. 2009. Effects of roads on animal abundance: An empirical review and synthesis. *Ecology and Society* 14:21.

FAURE, P. A., AND R. M. BARCLAY. 1992. The sensory basis of prey detection by the long-eared bat, *Myotis evotis*, and the consequences for prey selection. *Animal Behaviour* 44:31-39.

FEN SOME, A. G., AND F. MATHEWS. 2016. Roads and bats: a meta-analysis and review of the evidence on vehicle collisions and barrier effects. *Mammal Review* 46:311-323.

FENTON, M. B. *ET AL.* 1992. Phyllostomid bats (Chiroptera: Phyllostomidae) as indicators of habitat disruption in the neotropics. *Biotropica* 24:440-446.

FINDLEY, J. S. (Eds.). 1993. Bats: a community perspective. Cambridge University Press. Cambridge, U.K.

FORMAN, R. T. T., AND L. E. ALEXANDER. 1998. Roads and their major ecological effects. *Annual Review of Ecology and Systematics* 29:207-231.

GAISLER, J., Z. ŘEHÁK, AND T. BARTONIČKA. 2009. Bat casualties by road traffic (Brno-Vienna). *Acta Theriologica* 54:147-155.

GARBINO, G. S. T. *ET AL.* 2020. Updated checklist of Brazilian bats: versão 2020. Comitê da Lista de Morcegos do Brasil—CLMB. Sociedade Brasileira para o Estudo de Quirópteros (Sbeq). <https://www.sbeq.net/lista-de-especies>. Accessed on 30 September 2020

GARDNER, A. L. (Eds.). 2007. Mammals of South America, Volume I. Marsupials, xenarthrans, shrews, and bats. The University of Chicago Press. Chicago, U.S.A.

GIBBS, J. P., AND W. G. SHRIVER. 2002. Estimating the effects of road mortality on turtle populations. *Conservation Biology* 16:1647-1652.

GLISTA, D. J., T. L. DEVault, AND J. A. DEWOODY. 2009. A review of mitigation measures for reducing wildlife mortality on roadways. *Landscape and Urban Planning* 91:1-7.

GOMES, D. G., G. APPEL, AND J. R. BARBER. 2020. Time of night and moonlight structure vertical space use by insectivorous bats in a Neotropical rainforest: an acoustic monitoring study. *PeerJ* 8:e10591.

GORRESEN, P. M., M. R. WILLIG, AND R. E. STRAUSS. 2005. Multivariate analysis of scale-dependent associations between bats and landscape structure. *Ecological Applications* 15:2126-2136.

GREGORIN, R., AND V. A. TADDEI. 2002. Chave artificial para identificação de molossídeos brasileiros (Mammalia, Chiroptera). *Mastozoologia Neotropical* 9:13-32.

GUMIER-COSTA, F., AND C. F. SPERBER. 2009. Atropelamentos de vertebrados na Floresta Nacional de Carajás, Pará, Brasil. *Acta Amazonica* 39:459-466.

IUCN. 2020. The IUCN Red List of Threatened Species. Version 2020-1. <https://www.iucnredlist.org>. Accessed on 30 June 2020

JACKSON, S. D. 2000. Overview of transportation impacts on wildlife movement and populations. *Wildlife and highways: seeking solutions to an ecological and socio-economic dilemma*. The Wildlife Society 7-20.

KALKO, E. K. V., C. O. HANDLEY, AND D. HANDLEY. 1996. Structure, diversity, and long-term dynamics of a Neotropical bat community. Pp. 503-553, in *Long-Term studies of vertebrate communities* (Cody, M. L., and J. A. Smallwood, eds.). Academic Press. New York, U.S.A.

KERTH, G., AND M. MELBER. 2009. Species-specific barrier effects of a motorway on the habitat use of two threatened forest-living bat species. *Biological Conservation* 142:270-279.

LEŚIŃSKI, G. 2007. Bat road casualties and factors determining their number. *Mammalia* 71:138-142.

LEŚIŃSKI, G., A. SIKORA, AND A. OLSZEWSKI. 2011. Bat casualties on a road crossing a mosaic landscape. *European Journal of Wildlife Research* 57:217-223.

LEVINE, N. 2004. CrimeStat III: A Spatial Statistics Program for the Analysis of Crime Incident Locations. Ned Levine and Associates, Houston, TX, and the National Institute of Justice. Washington, DC, U.S.A.

Luz, J. L., L. D. M. COSTA, E. C. LOURENÇO, AND C. E. L. ESBÉRARD. 2011. Morcegos (Mammalia, Chiroptera) da Reserva Rio das Pedras, Rio de Janeiro, Sudeste do Brasil. *Biota Neotropica* 11:95-101.

MARIMUTHU, G., AND G. NEUWEILER. 1987. The use of acoustical cues for prey detection by the Indian False Vampire Bat, *Megaderma lyra*. *Journal of Comparative Physiology A* 160:509-515.

MARTINS, M. P. V., J. M. TORRES, AND E. A. C. DOS ANJOS. 2014. Dieta de morcegos filostomídeos (Mammalia, Chiroptera, Phyllostomidae) em fragmento urbano do Instituto São Vicente, Campo Grande, Mato Grosso do Sul. *Papeis Avulsos de Zoologia* 54:665-670.

MEDELLÍN, R. A., M. EQUIHUA, AND M. A. AMIN. 2000. Bat diversity and abundance as indicators of disturbance in neotropical rainforest. *Conservation Biology* 14:1666-1675.

MEDINAS, D., J. T. MARQUES, AND A. MIRA. 2013. Assessing road effects on bats: The role of landscape, road features, and bat activity on roadkills. *Ecological Research* 28:227-237.

MORATELLI, R., A. L. GARDNER, J. A. OLIVEIRA, AND D. E. WILSON. 2013. Review of *Myotis* (Chiroptera, Vespertilionidae) from north-

ern South America, including description of a new species. American Museum of Natural History. American Museum Novitates 2013:1-36.

MULLER, M. F., AND N. R. DOS REIS. 1992. Partição de recursos alimentares entre quatro espécies de morcegos frugívoros (Chiroptera, Phyllostomidae). Revista Brasileira de Zoologia 9:345-355.

NOVAES, R. L. M. *ET AL.* 2018. On a collision course: the vulnerability of bats to roadkills in Brazil. Mastozoología Neotropical 25:115-128.

ORMSBY, T. *ET AL.* 2010. Getting to Know ArcGIS Desktop. Redlands, ESRI Press. California, U.S.A.

PERINI, F. A., V. C. TAVARES, AND C. NASCIMENTO. 2014. Bats from the city of Belo Horizonte, Minas Gerais, southeastern Brazil. Chiroptera Neotropical 9:169-173.

PINHEIRO, B. F., AND L. C. B. TURCI. 2013. Vertebrados atropelados na estrada da Variante (BR-307), Cruzeiro do Sul, Acre, Brasil. Natureza on line 11:68-78.

PRACUCCI, A., C. ALVES DA ROSA, AND A. BAGER. 2012. Variação sazonal da fauna selvagem atropelada na rodovia MG 354, Sul de Minas Gerais – Brasil. Biotemas 25:73-79.

RAMALHO, D. F., AND L. AGUIAR. 2020. Bats on the Road—A Review of the Impacts of Roads and Highways on Bats. Acta Chiropterologica 22:417-433.

R CORE TEAM. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

REIS, N. R. *ET AL.* 2017. História Natural dos Morcegos Brasileiros Chave de identificação de Espécies. Technical Books Editora. Rio de Janeiro, Brasil.

REIS, N. R., M. L. S. BARBIERI, I. P. DE LIMA, AND A. L. PERACCHI. 2003. O que é melhor para manter a riqueza de espécies de morcegos (Mammalia, Chiroptera): um fragmento florestal grande ou vários fragmentos de pequeno tamanho? Revista Brasileira de Zoologia 20:225-230.

RIPLEY, B. D. 1981. Spatial Statistics. John Wiley and Sons. New York, U.S.A.

ROW, J. R., G. BLOUIN-DEMERS, AND P. J. WEATHERHEAD. 2007. Demographic effects of road mortality in black ratsnakes (*Elaphe obsoleta*). Biological Conservation 137:117-124.

SAMSON, A. *ET AL.* 2016. Effect of vehicular traffic on wild animals in Sigur Plateau, Tamil Nadu, India. Journal of Threatened Taxa 8:9182-9189.

SCHAUB, A., J. OSTWALD, AND B. M. SIEMERS. 2008. Foraging bats avoid noise. Journal of Experimental Biology 211:3174-3180.

SIEMERS, B., AND H. SCHNITZLER. 2000. Natterer's bat (*Myotis nattereri* Kuhl, 1818) hawks for prey close to vegetation using echolocation signals of very broad bandwidth. Behavioral Ecology and Sociobiology 47:400-412.

SIMMONS, N. B., AND A. L. CIRRANELLO. 2020. Bat species of the world: a taxonomic and geographic database [online]. <https://batnames.org/>. Accessed on 10 December 2020

STONE, E. L., G. JONES, AND S. HARRIS. 2009. Street Lighting Disturbs Commuting Bats. Current Biology 19:1123-1127.

TAVARES, V. C. *ET AL.* 2010. Bats of the state of Minas Gerais, southeastern Brasil. Chiroptera Neotropical 16:675-705.

TEIXEIRA, F. Z. *ET AL.* 2013. Are road-kill hotspots coincident among different vertebrate groups? Oecologia Australis 17:36-47.

TROMBULAK, S. C., AND C. A. FRISSELL. 2000. Review of ecological effects of roads on terrestrial and aquatic communities. Conservation Biology 14:18-30.

VALADÃO, R. M., L. F. BASTOS, AND C. P. DE CASTRO. 2018. Atropelamentos De Vertebrados Silvestres Em Quatro Rodovias No Cerrado, Mato Grosso, Brasil. Multi-Science Journal 1:62-74.

VELAZCO, P. M., A. L. GARDNER, AND B. D. PATTERSON. 2010. Systematics of the *Platyrrhinus helleri* species complex with descriptions of two new species. Zoological Journal of the Linnean Society 159:785-812.

ZURCHER, A. A., D. W. SPARKS, AND V. J. BENNETT. 2010. Why the bat did not cross the road? Acta Chiropterologica 12:337-340.

Associated editor: Sergio Solari

Submitted: January 16, 2022; Reviewed: February 18, 2022

Accepted: March 5, 2022; Published on line: March 31, 2022