

Calcium-carbonate and phytoelements to improve egg production in backyard laying hens under heat stress

Lilia Gutiérrez¹

 0000-0002-4823-0388

Corazón de Jesús Mendoza²

 0000-0002-9796-0301

Zacnité Sánchez¹

 0000-0001-6554-7574

Lizbeth Carrillo¹

 0000-0002-1236-1976

Héctor Sumano^{1*}

 0000-0002-8802-5274

¹ Universidad Nacional Autónoma de México.
Facultad de Medicina Veterinaria y Zootecnia..
Departamento de Fisiología y Farmacología.
Ciudad de México, México.

² Universidad Nacional Autónoma de México.
Facultad de Química.
Departamento de Química Orgánica.
Ciudad de México, México.

***Corresponding author**

Email address:
sumano@unam.mx

Abstract

The bioavailability (F) of calcium and other nutrients from the gastrointestinal tract in the second and third laying cycle hens and in hens under heat-stress is insufficient to sustain the necessary egg production. The administration of additional amounts of any source of calcium has been shown to be insufficient to meet the demand for calcium. However, improved calcium F has already been achieved in these hens if calcium carbonate is pharmaceutically prepared as patented pellets (FOLAs). Hence, clinical challenges were set to test the FOLA pellets as a form to sustain egg production and eggshell thickness in 180-second and 180-third cycle Bovans-White laying hens, randomly divided into six groups, as follows: cycle-2 control group (GCC2); FOLA-group-capsaicin (GFcC2); FOLA-group-capsaicin-turmeric powder-cinnamon powder (GfexC2); control-group for third cycle hens (GCC3); FOLA-group-capsaicin (GFcC3); FOLA-group-capsaicin-turmeric powder-cinnamon powder (GfexC3). This trial was carried out under moderate heat-stress conditions, and a follow-up was carried out utilizing eye thermographic measurements. The results show that all FOLA-fed hens of the second or third laying cycle presented better laying average and eggshell thickness than the untreated control groups, i.e., GFcC2 128 % higher than GCC2; GfexC2 135 % higher than GCC2; GFcC3 168 % higher than GCC3; and GfexC3 was 173 % higher than GCC3 ($P < 0.01$ in all comparisons). FOLA pellet formulation, similar to GfexC2, is postulated as a viable natural solution to counter the decreased egg-laying and shell thickness in second and third-cycle hens under moderate heat stress.

Keywords: Phyto elements; Nutraceuticals; Egg production; Shell-thickness; 2nd and 3rd laying cycle; Laying-hens.

Cite this as:

Gutiérrez L, Mendoza C de J, Sánchez Z, Carrillo L, Sumano H. Calcium-carbonate and phyto elements to improve egg production in backyard laying hens under heat stress. *Veterinaria Mexico OA*. 2024;11. doi: 10.22201/fmvz.24486760e.2024.1253.

Submitted: 2023-08-10
Accepted: 2024-02-19
Published: 2024-05-03

Additional information and declarations
can be found on page 11

© Copyright 2024
Lilia Gutiérrez et al.



Distributed under Creative Commons CC-BY 4.0

Study contribution

Given the threat of global warming to egg production and the carbon footprint associated with the replacement of single-cycle laying hens by flocks, a calcium carbonate-phytoelements pharmaceutical design was studied. Its supplementation could preserve egg production in second and third-cycle egg-producing backyard hens under moderated heat stress.

Introduction

Eggs are of paramount importance for human nutrition. Approximately 80 million tons of chicken eggs are produced worldwide, and growth in this industry is expected to exceed 25 % by 2030.⁽¹⁻³⁾ Hence, it is necessary to improve egg production, and such a challenge should not entail using antibacterial production promoters and must not compromise the health of the laying hens. Even small family farms are emerging to meet this egg demand.⁽⁴⁾ The obvious approach should be to bring producing hens to an optimum state of health without affecting their welfare and in an antibacterial-free manner. This situation has called on researchers to seek alternatives to optimize egg production, i.e., herbal constituents, as few and inconsequential side-effects have been reported with this approach.^(5, 6) Herbal extracts have been shown to be innocuous and contribute to maintaining or improving hen intestinal health while allowing an increase in production without significant environmental impact.^(7, 8)

More than 80 % of the egg-producing farms in Mexico are regarded as intensive producers, which entails one single egg-production cycle. Then, the hens are culled. This decision is primarily based on the fact that egg posture in the second and third cycles markedly decreases, i.e., 40 % or more. It has been pointed out that deterioration in eggshell quality as the flock ages is a sizable physiological puzzle to investigate in laying hens, particularly in second and third-cycle hens. Egg quality also decreases with age, although egg weight increases, a trait that may be of little use as eggshell quality, particularly shell thickness and breaking strength, decreases, rendering many eggs useless.⁽⁹⁾ Consequently replacing such hens is considered more viable than dealing with reduced egg production.⁽¹⁰⁾ In contrast, to non-technified and backyard farms, second and even third laying cycles are a common practice despite the detriment in egg production and quality. The main surrogate parameter for evaluating egg quality is a firm and full-thickness eggshell, resistant to cracking and capable of sustaining its typical shape. A significant inverse relationship between shell thickness and the percentage of cracked eggs has been identified.⁽¹¹⁾ Low-quality eggshells in the second laying cycle have been linked to body changes such as poor intestinal absorption of nutrients, particularly calcium. We have reported that a modified release preparation of various pharmaceuticals and nutraceuticals can be designed to improve the absorption capabilities of the gastrointestinal tract (GIT) of hens and poultry,⁽¹²⁻¹⁴⁾ and that the inclusion of capsaicin and calcium in the FOLA pellets improves the bioavailability of calcium in second cycle hens. A patent has been granted for the pelleted pharmaceutical form (MX/a/2012/013222) under the name of FOLA (F = bioavailability; O = optimal;

LA = Long action) as it can improve the bioavailability of calcium from calcium carbonate, one of the most inexpensive sources of this element.⁽¹²⁻¹⁴⁾

An additional problem to consider is global climate warming. Laying hens are among the most sensitive species to heat stress due to their deficient ability to dissipate body heat.^(15, 16) Egg production and health rapidly deteriorate under heat stress.⁽¹⁷⁻¹⁹⁾ The optimum environmental temperature for laying hens has been set at approximately 20 °C to 25 °C, with minor variations due to hen-line.^(16, 17) If the environmental temperature approaches or exceeds 30 °C, various signs of heat stress appear, such as a decreased feed intake, lower egg weight, and a distinctive drop in egg production observed together with increases in flock mortality.⁽¹⁶⁻¹⁹⁾ Apart from the obvious recommended housing improvement and hen husbandry, a wide variety of studies have shown that the administration of essential oils and plant extracts limits the detrimental effects of heat stress in these birds, mainly when heat stress is not extreme.^(20, 21)

In this study, two FOLA-type pellets have been manufactured in an attempt to sustain both eggshell thickness and egg production in second and even third-cycle hens and also in laying hens exposed to temperatures above their comfort zone (≥ 30 °C) in the central part of Mexico during summer. The FOLA-type pellets contain a gastro-retentive vehicle and capsicum or these constituents, plus cinnamon and turmeric powder. The FDA has already approved cinnamon as phytoadditive.⁽²²⁾ It has potent anti-inflammatory, antimicrobial, and antioxidant properties with free radical scavenging actions and strong inhibitory effects on nitric oxide (NO) production by inhibiting NF κ B activity.⁽⁷⁾ In addition, it has been shown to reduce the effects of heat stress in poultry,⁽²²⁾ reducing oxidative activities in poultry when supplemented in their diets.^(23, 24) Also, antiviral, antibacterial, and antifungal activities^(7, 25) have been demonstrated in broiler chickens.^(26, 27) Capsicum oleoresin improves immunity and overall gastrointestinal health, enhancing growth and meat quality even in broiler chickens under heat stress.^(7, 28-31) Turmeric (*Curcuma* spp.) powder and its main active principle, curcumin (hydrophobic polyphenolic phytochemical), possess several therapeutic benefits as feed-additive in poultry, including disease-resistance featured. Apart from other potentially active principles, turmeric powder contains curcumin, demethoxycurcumin, and bisdemethoxycurcumin.⁽³²⁾ These substances act as antioxidant and anti-inflammatory agents⁽³³⁻³⁵⁾ and have been chosen as an alternative to antibiotics in animal feed to improve production.^(36, 37)

The FOLA formulation manages to maintain the nutrients in the GIT of the birds for a longer time. Hence, as modified-release pharmaceutical system for antibiotics^(12, 38) it was thought possible that it could improve the bioavailability of nutrients and active principles in poultry. Therefore, the working hypothesis for this trial is that egg production and shell quality in second and third-cycle backyard hens subjected to moderated heat-stress, will be better sustained with phytochemicals and calcium carbonate manufactured as for the FOLA design, as compared to unsupplemented hens.

Material and methods

Ethical statement

This study was conducted on a family farm in Texcoco, State Mexico. The Subcommittee Institutional for the Care and Use of Experimental Animals of the Faculty of Veterinary Medicine and Zootechnics at the National Autonomous University of Mexico (CUEA-FVMZ/NAUM) study number: SICUAE-DC2018/2-1.

Development of the preparations

FOLA-pellets were manufactured as described in Patent MX/a/2012/013222. Briefly, FOLA-pellets are obtained by extruding corn and wheat flour (1:1) containing modified release vehicles (mentioned in the referred patent), plus 4.1 % calcium-carbonate.^(12, 14)

Animals

A total of 180 Bovans-White weighing $1\,570 \pm 20$ g, laying hens in a second cycle (> 180 weeks old) and 180 Bovans-White hens in the third cycle (> 270 weeks of age), were included in this trial carried out from August to Mid December 2022. To test FOLA pellet subjected to environmental heat stress, further 90-second and third-cycle Bovans-White hens were studied for egg production and egg-shell thickness during May 2023, the hottest period of the year in the central part of Mexico, reaching temperatures at or above 30 °C.^(20, 21) Hens were allocated in groups of three per California-type cage, 40 cm wide, 45 cm deep, and 45 cm high (600 cm³/hen). Water was provided *ad libitum* through cup drinkers (one per cage), and a 16 light/8 dark h lighting program was established.

Groups

For the posture cycle test, hens were divided into six groups of 30 hens each with the following designation: GCC2 = control group of the second cycle; GFcC2 = FOLA group + 2 ppm oleoresin of 500 000 Scoville Heat Units (SHU) (VEPinsa SA de CV, Sinaloa, Mexico); GfexC2 = FOLA group + 2 ppm oleoresin of 500 000 Scoville Heat Units + 1 % cinnamon powder (McCormick® food grade) and 1 % turmeric powder (McCormick® food grade) were added; GCC3 = control group of the third cycle; GFcC3 = FOLA group + capsicum oleoresin 2 ppm; GfexC3 = FOLA group + capsicum oleoresin 2 ppm + turmeric powder 1 % + 1 % cinnamon powder.

The control groups (GCC2 and GCC3) were fed a balanced diet according to the utilized lineage guidelines in this trial (Table 1). The experimental groups (GFcC2, GfexC2, GFcC3, GfexC3) were deprived of all the fine and coarse calcium carbonate generally added to the chicken diets, and only the calcium contained as orthophosphate, already present in their feed, was maintained. Calcium requirements for the experimental hens were met with calcium carbonate FOLA pellets. The rest of the feed constituents remained the same. Their feed was supplied in a trough feeder (13.3 cm/hen), and the experimental groups received a dose of 8 g of FOLA pellets/hen containing 4.1 % calcium carbonate.

Table 1. Calculated nutrient of the diet in the untreated control groups

Ingredients (kg)	
Corn	595
Soybean meal 48 (%)	267
Soy oil	13.00
Salt	4.00
Calcium carbonate	101.00
Ortophosphate 1 820	11.00
Vitamine ¹ and mineral ² premix	3.00
DL-methionine 99	1.98
L-lysine 2.75	2.75
Vitamin B ₂	0.200
Vitamin B ₁₂	0.200
Ronozyme Phytase (GT)	0.100
Total	1 000
Calculated analysis (g/kg)	
Meatbolizable energy (kcal/kg)	2859
Crude protein (%)	18.792
Total calcium	4.100
Phosphorus (disp)3.80	0.450
Methionine + Cysteine 4.50	0.800
Lysine (%)	0.980
Tryptophan (%)	0.220
Sodium (%)	0.180
Choline (mg/kg)	1 022.63
Chlorine (%)	0.275
Linoleic acid (%)	1.867

¹ Quantity/kg: Retinol 0.9 g, cholecalciferol 0.019 g, d-alpha tocopherol 10.004 g, phylloquinone 1 g, riboflavin 4 g, cyanocobalamin 0.06 g, pyridoxine 3 g, calcium pantothenate 13 g, niacin 25 g, biotin 0.063 g, choline chloride 250 g.

² Quantity/kg: 0.2 g selenium, 0.1 g cobalt, 0.3 g iodine, 10 g copper, 50 g zinc, 100 g iron, 100 g manganese.

In [Table 1](#), the untreated control groups were of second-cycle (GCC2), and third cycle GCC3 laying hens. Calcium for the experimental hens (GFcC2; GFex2; GFcC3; GFex3) was incorporated into the FOLA pellets, also as calcium carbonate. The rest of the feed constituents remained unaltered. To evaluate the possible effect of calcium primed FOLA-pellets on laying hens under moderate heat-stress, the body temperature of the hens was individually taken using a thermographic camera (FLIR®, [Figure 1](#)) aiming the eye. Chicken house environmental temperatures were measured at 13:00, 14:00, and 15:00 h which are considered the hottest months in the area, with May being the hottest month.^(39, 40)



Figure 1. Representative image of the temperature measurement using FLIR camera.

Statistical evaluation

The thickness of the eggshell was evaluated daily, in the middle zone (between the poles) with a precision digital Vernier that admits an error of 0.01 mm. These data were statistically compared using Student's t-tests and no parametric tests (F-test to compare variances). A statistically significant value was considered when $P \leq 0.05$. All statistical analyses were performed with Prism (GraphPad Software, for macOS, LLC).

Results

Figure 2 shows the mean $\pm$ 1 SD of the eggshell thickness of the FOLA-treated and control groups one week before treatment and three weeks after. Considering the values from the fourth day of FOLA administration in the experimental groups, the improvements in thickness were 128 % for GFcC2 and 135 % for GFexC2 as compared to GCC2, and an improvement of 168 % for GFcC3 and 173 % for GFexC3 as compared to the control group (GCC3). However, groups in which no statistical significance was found were GFexC2 vs GFexC3 and GFcC2 vs GFexC3 using ANOVA and Bonferroni t-tests ($P < 0.0500$).

Global laying percentages for all groups are depicted in **Figure 3**, which shows the daily global mean values $\pm$ 1 SD obtained. Also, using ANOVA and Bonferroni t-tests, results reveal higher values of 185 % superior for GFcC2 and 189 % for GFexC2, compared to GCC2. Hens from GFcC3 had 193 % higher laying, and GFexC3 197 %, compared to GCC3. Other comparisons of global laying percentages showed no statistical differences, i.e., GFcC2 vs. GFexC2 and GFcC2 vs. GFcC3 ($P < 0.0500$).




Figure 2. Mean $\pm$ 1 SD of eggshell thickness in control groups, and pellet-FOLA calcium-supplemented groups. Basal data (-1 week) and after treatment (weeks 1-3). GCC2 and GCC3 = control groups for second and third laying cycles; GFcC2 = FOLA group + capsicum oleoresin two ppm; GFexC2 = FOLA group + capsicum oleoresin + turmeric powder + cinnamon powder; GCC3 = control group of cycle 3; GFcC3 = FOLA group + capsicum oleoresin two ppm; GFexC3 = FOLA group + capsaicin + turmeric powder + cinnamon powder.




Figure 3. Mean $\pm$ 1 SD of laid eggs in control groups, and in pellet-FOLA calcium-supplemented groups. Basal data (-1 week) and after treatment (weeks 1-3). GCC2 and GCC3 = control groups for second and third laying cycles; GFcC2 = FOLA group + capsicum oleoresin two ppm; GFexC2 = FOLA group + capsicum oleoresin + turmeric powder + cinnamon powder; GCC3 = control group of cycle 3; GFcC3 = FOLA group + capsicum oleoresin two ppm; GFexC3 = FOLA group + capsaicin + turmeric powder + cinnamon powder.

Figure 4 shows the body temperatures ± 1 SD of the untreated control hens and those of the second and third-cycle hens treated with FOLA. Using ANOVA and Bonferroni and t-tests, It becomes apparent that from day 4 of treatment onwards, mean temperatures become statistically distinguishable from untreated control groups ($P < 0.05$ in all cases). The GfexC2 group had the lowest mean body temperature (37.8 ± 0.2 °C), statistically significant as compared to the GCC3 group, which had the highest mean body temperature of 39.1 ± 0.5 °C. The groups in which no statistical differences were found in body temperatures were GCC2 vs GCC3; GFcC2 vs. GFcC3; GFcC3 vs GFexC3 and GfexC2 vs GFexC3.

Only on some days were the parameters of laying percentage and eggshell thickness achieved in the GfexC2 groups statistically different from other FOLA-supplemented groups. As already demonstrated in the literature, egg weight increases in second and third-cycle laying hens^(10, 11). Hence, measuring this parameter was omitted in this study. The mean highest in-chicken house temperature during May 2023 was 31 ± 2 °C.




Figure 4. Mean ± 1 SD of body temperatures measured by FLIR thermographic camera in the eye of egg-producing hens subjected to moderate heat stress (approximately 30 °C) with the following groups: untreated second and third cycle hens (GCC2 and GCC3) and in pellet-FOLA calcium-supplemented groups: GFcC2 = FOLA group + capsicum oleoresin two ppm; GfexC2 = FOLA group + capsicum oleoresin + turmeric powder + cinnamon powder; GFcC3 = FOLA group + capsicum oleoresin two ppm; GFexC3 = FOLA group + capsaicin + turmeric powder + cinnamon powder.

Discussion

The commercially viable lifespan of laying hens in intensive farming enterprises is generally limited to a single cycle of approximately one year⁽⁴¹⁾ This is due to both commercial commitments with hatcheries for replacements and the sharp drop in egg production in hens during the second and third cycles, having drops of approximately 30 %.^(41, 42) In a previous study, it was shown that adding calcium carbonate, as in FOLA pellets, improved egg quality in second-cycle laying hens as for the parameters measured in this trial. Hence, production was maintained as efficiently as that observed in first cycle hens.^(12, 14) Nevertheless, it may be possible that the supplementation with turmeric powder and cinnamon powder could have assisted the formulation of FOLA-pellets described here, as it has been shown that these phytoelements improve various performance parameters in poultry.^(43, 44) To ponder these results, and despite the benefits outlined, it will be of particular interest to carry out economic studies in hens that produce eggs for human consumption, as well as in hens producing fertile eggs for broiler production. In contrast, in the backyard or small-production egg-producing units, layer hens can be benefited and have their productive lives extended. Particularly when their replacement is not possible or unprofitable. Hence, this study can have immediate benefit to this important sector of the egg-industry. It is essential to add that the production cost of FOLA pellets is not burdensome, and it is easily offset by the production obtained from commercially viable eggs.

The production of quality layer eggs depends on many factors, including environmental stress such as noise, diet adequacy, and whether or not there is heat stress.^(11, 19) However, the key features that stand out for their successful marketing and handling are their integrity and eggshell thickness. To optimize the former features, it is critical to enhance blood calcium bioavailability.^(12, 14, 15) In this context, it is noteworthy that one of the primary sources of calcium in the hens' diet is calcium carbonate, as it is the most economical source of calcium. Additionally, the added phytoelements, turmeric and cinnamon powder, are inexpensive enough to be included in the FOLA treatments without affecting the cost-benefit ratio of the cycle.

Calcium carbonate is the most used ingredient of diets in poultry production. However, it is the least water-soluble source of calcium, and therefore, absorption of calcium can be low and unpredictable, particularly in second and third-cycle laying hens.^(12, 14) Additionally, it is recognized that not only calcium is deficiently absorbed, but other nutrients also seem to be affected during the second and third production cycles, generating a deficient egg-lying performance, as has been shown in previous works.^(10, 14, 45) In this study, it became clear that medication of second and third-cycle hens with calcium carbonate incorporated in FOLA pellets and fortified with turmeric and cinnamon powder, achieves a remarkable recovery of egg production under moderate heat stress conditions. Phytogetic additives have been proven in multiple trials to be of great importance in improving or contributing to the health of the gastrointestinal tract of poultry and, through this effect, can significantly improve both their production and welfare.^(7, 46) Furthermore, it has been shown that essential oils and extracts can mitigate the environmental impact of egg production by optimizing the relationship between egg production and a hen's food intake.^(47, 48) Thus, the pellet / FOLA design of calcium carbonate with added phytochemicals meets the demand to design the referred calcium source that remains available in the gastrointestinal tract of egg-producing hens when it is

required, e.g., during the shell formation process at night. This is, in turn, achieved by the programmed release of calcium carbonate facilitated by vehicles that also enhance bioavailability.⁽¹²⁾ It is also possible to postulate that including phytoadditives in the FOLA pellets contributes to maintaining better intestinal health, enabling the modern genetic line of hens to meet their production nutritional requirements. The theoretical bases of the FOLAs described here have already been published in part.⁽¹²⁾ Nevertheless, assuming that the ideal formula or formulas still require additional research is feasible. Also, it would be important to carry out studies on an industrial scale to confirm or discard these results.

Enabling economically viable second and third cycle laying hens with the FOLA-calcium and phytoelements pellets complies well with breeding companies that are seeking the development of “long life” laying hens, whose goal is ending up with laying hens producing 500 eggs in a single continuous cycle of 100 weeks.⁽⁴¹⁾ This goal has been conceived using selection programs based on phenotype, cross breed progeny testing, and genotype information derived from DNA markers associated with phenotypic traits.⁽⁴⁹⁾ Nevertheless, proper nutrient supplementation must be anticipated to meet new physiological demands, and the formulation of FOLA-calcium and phytoelements here studied may result adequate.⁽⁴¹⁾

Data availability

The original datasets used in this research and if applicable, supporting information files, are deposited and available for download at the SciELO Dataverse repository (doi:_____).

Funding statement

This project was partly financed by a grant from Universidad Nacional Autónoma de México (www.unam.mx): PAPIIT IT200322.

Conflicts of interest

The authors confirm that they do not have any conflicts of interest.

Author contributions

Conceptualization: H Sumano.

Investigation: L Gutiérrez, Z Sánchez, L Carrillo, CJ Mendoza.

Writing-original draft: L Gutiérrez, H Sumano.

Writing-review and editing: L Gutiérrez, H Sumano.

References

1. SIAP. Escenario mensual de productos agroalimentarios. Ciudad de México, México: Dirección de Análisis Estratégico. https://www.gob.mx/cms/uploads/attachment/file/759467/Huevo_Agosto_2022.pdf
2. FAOSTAT. Statistics division: poultry production and products. Food and Agriculture Organization of the United Nations; 2022. <https://www.fao.org/poultry-production-products/production/es/>
3. UNA. México, promedio de consumo anual: 345 huevos por habitante – Unión Nacional de Avicultores; 2023. <https://una.org.mx/mexico-promedio-de-consumo-anual-345-huevos-por-habitante/>
4. Indarani S, Siddhartha N, Pradip KD. Chapter 14 - Biosecurity strategies for backyard poultry: a Controlled Way for Safe Food Production. In: Holban AM, Grumezescu AM, editors. Food Control and Biosecurity. Academic Press, USA; 2018. pp.481-517. doi: 10.1016/B978-0-12-811445-2.00014-3.
5. Mehdi Y, Létourneau-Montminy MP, Gaucher ML, Chorfi Y, Suresh G, Rouissi T, et al. Use of antibiotics in broiler production: global impacts and alternatives. Animal Nutrition. 2018(Jun);4(2):170–178. doi: 10.1016/j.aninu.2018.03.002.
6. Sureshbabu A, Smirnova E, Karthikeyan A, Moniruzzaman M, Kalaiselvi S, Nam K, et al. The impact of curcumin on livestock and poultry animal's performance and management of insect pests. Frontiers in Veterinary Science. 2023;10:1048067. doi: 10.3389/fvets.2023.1048067.
7. Melkamu BZ. Phytochemicals as feed additives in poultry production: a review. International Journal of Extensive Research. 2015;3:49–60.
8. Küçükylmaz K, Bozkurt M, Herken EN, Çınar M, Çatlı A, Bintaş Y, et al. Effects of rearing systems on performance, egg characteristics, and immune response in two layer hen genotype. Asian-Australasian Journal of Animal Science. 2012;25(4):559–568. doi: 10.5713/ajas.2011.11382.
9. Liu Y, Uyanga VA, Jiao H, Wang X, Zhao J, Zhou Y, Lin H. Effect of feeding strategies on eggshell quality of laying hens during late laying period. Poultry Science. 2022;102(2):102406. doi: 10.1016/j.psj.2022.102406.

10. Bell DD, Adams CJ. First and second cycle egg production characteristics in commercial table egg flocks. *Poultry Science*. 1992;71:448–459. doi: 10.3382/ps.0710448.
11. Roberts J. Factors affecting egg internal quality and egg shell quality in laying hens. *The Journal of Poultry Science*. 2004;41(3):161–177. doi: 10.2141/jpsa.41.161.
12. Carrillo L, Bernad M-J, Monroy-Barreto M, Coello CL, Sumano H, Gutiérrez L. Higher bioavailability of calcium in chickens with a novel in-feed pharmaceutical formulation. *Frontiers in Veterinary Science*. 2020;7:343. doi: 10.3389/fvets.2020.00343.
13. Gutierrez OL, Alcala Y, Bernard MJ, Sumano LH. Increased bioavailability of tylosin phosphate as in-feed medication formulation for long-action pellets in broilers chickens. *Journal of Applied Poultry Research*. 2017;27(1):16–22. doi: 10.3382/japr/pfx035.
14. Sumano H, Carrillo L, Monroy M, Tapia G, Gutierrez L. Bioavailability of four calcium sources in the second-cycle egg-producing hens. *Journal of Applied Poultry Research*. 2021;30(1):100–106. doi: 10.1016/j.japr.2020.10.005.
15. Abdulhakim S, Salam YA, Reza T. Egg quality and safety with an overview of edible coating application for egg preservation. *Food Chemistry*. 2019(Oct)30;296:29–39. doi: 10.1016/j.foodchem.2019.05.182.
16. Oluwagbenga EM, Fraley GS. Heat stress and poultry production: a comprehensive review. *Poultry Science*. 2023;103:141. doi: 10.1016/j.psj.2023.103141.
17. Wasti S, Sah N, Mishra B. Impact of heat stress on poultry health and performances, and potential mitigation strategies. *Animals*. 2020;10:1–19. doi: 10.3390/ani10081266.
18. Roberts J. Factors affecting egg internal quality and egg shell quality in laying hens. *The Journal of Poultry Science*. 2004;41(3):161–177. doi: 10.2141/jpsa.41.161.
19. Ahmadi F, Rahimi F. Factors affecting quality and quantity of egg production in laying hens: a review. *World Applied Sciences Journal*. 2011;12(3):72–384.
20. Kim D-H, Lee K-W. An update on heat stress in laying hens. *Worlds Poultry Science Journal*. 2023;1–24. doi: 10.1080/00439339.2023.2239769.
21. Kilic I, Simsek E. The effects of heat stress on egg production and quality of laying hens. *Journal of Animal and Veterinary Advances*. 2013;12(1). doi: 10.3923/javaa.2013.42.47.
22. Ali A, Ponnampalam EN, Pushpakumara G, Cottrell JJ, Suleria HAR, Dunshea FR. Cinnamon: A natural feed additive for poultry health and production: a review. *Animals*. 2021;11(7):2026. doi: 10.3390/ani11072026.
23. Yang Y, Zhao L, Shao Y, Liao X, Zhang L, Lu YL. Effects of dietary graded levels of cinnamon essential oil and its combination with bamboo leaf flavonoid on immune function, antioxidative ability and intestinal microbiota of broilers. *Journal of Integrative Agriculture*. 2019;18(9):2123–2132. doi: 10.1016/S20953119(19)62566-9.
24. Yaqoob A, Razzaq A, Iqbal S, Hassan O, Ishtiaq H, Hussain S, et al. Cinnamon bioactives and their impact on poultry nutrition and meat quality - impact on human health. *Acta Scientific Nutritional Health*. 2022;6(2):29–38. doi: 10.31080/asnh.2022.06.0990.

25. Fariba E, Masoumeh Z, Yousefpoor Y, Alipanah H, Safari E, Osanloo M. Anti-inflammatory and anti-nociceptive effects of Cinnamon and Clove essential oils nanogels: an in vivo study. BMC Complementary Medicine and Therapies. 2022;22(1):1–10.
26. Rao P, Gan S. Cinnamon: a multifaceted medicinal plant. Evidenced-Based Complementary and Alternative Medicine. 2014;2014. doi: 10.1155/2014/642942.
27. Muhammad D, Dewettinck K. Cinnamon and its derivatives as potential ingredient in functional food-a review. International Journal of Food Properties. 2017;20(Suppl2):2237–2263. doi: 10.1080/10942912.2017.1369102.
28. Puvača N, Ljubojević D, Spasevski N, Duragić O, Nikolova N, Prodanović R, Bošković J. Effects of turmeric powder (*Curcuma longa*) in laying hens nutrition: table eggs production, quality and lipid profile. Concepts of Dairy & Veterinary Sciences. 2018;2(1):162–163. doi: 10.32474/cdvs.2018.01.000129.
29. Khan R, Naz M, Nikousefat M, Selvaggi M, Tufarelli V, Laudadio V. The use of Turmeric (*Curcuma longa*) in poultry feed. World's. Poultry Science Journal. 2012;68(1):97–103. doi: 10.1017/S0043933912000104.
30. Manjunath M, Yashodhar P. Natural antibiotic effect of turmeric in poultry management. International Journal of Poultry and Fisheries Sciences. 2018;2(2):1–2. doi: 10.15226/2578-1898/2/2/00109.
31. Laganá C, Saldanha ESPB, Sartori JR, Turco PHN, Gonzales E, Luciano RL, et al. Turmeric on poultry production: a review. Agricultural Science. 2019;10(12):1592–1601. doi: 10.4236/as.2019.1012117.
32. Esatbeyoglu T, Huebbe P, Ernst IMA, Chin D, Wagner AE, Rimbach G. Curcumin-from molecule to biological function. Angewandte Chemie International Edition. 2012;51(22):5308–5332. doi: 10.1002/anie.201107724.
33. Hewlings S, Kalman D. Curcumin: a review of its effects on human health. Foods. 2017;6(10):92. doi: 10.3390/foods6100092.
34. Gera M, Sharma N, Ghosh M, Huynh DL, Lee SJ, Min T, et al. Nanoformulations of curcumin: an emerging paradigm for improved remedial application. Oncotarget. 2017;8(39):66680–66698. doi: 10.18632/oncotarget.19164.
35. Karthikeyan A, Young KN, Moniruzzaman M, Beyene AM, Do K, Kalaiselvi S, et al. Curcumin and its modified formulations on inflammatory bowel disease (IBD): the story so far and future outlook. Pharmaceutics. 2021;13(4):484. doi: 10.3390/pharmaceutics13040484.
36. Placha I, Gai F, Simonová MP. Natural feed additives in animal nutrition-their potential as functional feed. Frontiers in Veterinary Science. 2022;9:1062724 doi: 10.3389/978-2-8325-0843-5.
37. Shehata AA, Yalçın S, Latorre JD, Basiouni S, Attia YA, Abd El-Wahab A, et al. Probiotics, prebiotics, and phytogenic substances for optimizing gut health in poultry. Microorganisms. 2022;10(2):395. doi: 10.3390/microorganisms10020395.
38. Gutierrez L, Zermeño J, Alcalá YY, Sumano H. Higher bioavailability of doxycycline in broiler chickens with a novel in-feed pharmaceutical formulation. Poultry Science. 2017;96(8):2662–2669. doi: 10.3382/ps/pex036.
39. Murray-Tortarolo GN. Seven decades of climate change across Mexico. Atmósfera. 2021;34(2):217–226. doi: 10.20937/atm.52803.
40. Weather Spark. El clima en Texcoco, el tiempo por mes, temperatura promedio México. https://es.weatherspark.com/y/6599/Clima-promedio-en-Te-xco-co-M%C3%A9xico-durante-todo-el-a%C3%B1o#google_vignette

41. Bain MM, Dunn IC. Increasing persistency in lay and stabilising egg quality in longer laying cycles. What are the challenges? *British Poultry Science*. 2016;57(3):330–338. doi: 10.1080/00071668.2016.1161727.
42. Arulnathan V, Turner I, Bamber N, Grassauer F, Doyon M, Pelletier N. A systematic review of potential productivity, egg quality, and animal welfare implications of extended lay cycles in commercial laying hens in Canada. *Poultry Science*. 2024;103(4):103475. doi: 10.1016/j.psj.2024.103475.
43. Foluka AA, Olufemi MA. Turmeric (*Curcuma longa*): an alternative to antibiotics in poultry nutrition. *Translational Animal Science*. 2023;1:txad133. doi: 10.1093/tas/txad133.
44. Ali A, Ponnampalam EN, Pushpakumara G, Cottrell JJ, Suleria HAR, Dunshea FR. Cinnamon: a natural feed additive for poultry health and production-a review. 2021;11(7):2026. doi: 10.3390/ani11072026.
45. Pelicia K, Garcia A, Faitarone G, Silva P, Berto A, Molino B, et al. Calcium and available phosphorus levels for laying hens in the second production cycle. *Brazilian Journal of Poultry Science*. 2009;11:39–49. doi: 10.1590/S1516635X2009000100007.
46. Xiao G, Zheng L, Yan X, Gong L, Yang Y, Qui Q, et al. Effects of dietary essential oils supplementation on egg quality, Biochemical Parameters, and Gut Microbiota of Late-Laying Hens. *Animals*. 2022;12(19):2561. doi: 10.3390/ani12192561.
47. Zhang L, Gao F, Ge J, Li H, Xia F, Bai H, et al. Potential of aromatic plant-derived essential oils for the control of foodborne bacteria and antibiotic resistance in animal production: a review. *Antibiotics*. 2022;11(11):1673. doi: 10.3390/antibiotics11111673.
48. Gopi M, Karthik K, Manjunathachar V, Tamilmahan P, Kesavan M, Dashprakash M, et al. Essential oils as a feed additive in poultry nutrition. *Advances in Animal and Veterinary Sciences*. 2014;2(1):1–7.
49. O'Sullivan NP. What are the egg traits that further sustainable production? *International Poultry Production*. 2009;17(6):15–16.